

Mass angular momentum inequality for axisymmetric vacuum data with small trace

Xin Zhou

Abstract: In this paper, we proved the mass angular momentum inequality [16][12][23] for axisymmetric, asymptotically flat, vacuum constraint data sets with small trace. Given an initial data set with small trace, we construct a boost evolution spacetime of the Einstein vacuum equations as [10]. Then a perturbation method is used to solve the maximal surface equation in the spacetime under certain growing condition at infinity. When the initial data set is axisymmetric, we get an axisymmetric maximal graph with the same ADM mass and angular momentum as the given data. The inequality follows from the known results [16][12][23] about the maximal graph.

1 Introduction

Based on the gravitational collapse pictures [14], it is conjectured that the angular momentum should be bounded by the mass for physically reasonable solutions of the Einstein equations. It is true for Kerr black hole solutions which are stationary. For dynamical, axisymmetric solutions some progresses have been made over the past few years. Dain [16] first proved such an inequality for Brill data (See Definition 2.1 of [16]), which is a class of specialized axisymmetric, maximal, asymptotically flat vacuum data. Later, Chruściel, Li and Weinstein [11][12] generalized it to a class of axisymmetric, maximal data admitting an Ernst potential with positive mass density and certain asymptotically flatness conditions. Recently R. Schoen and the author [23] gave a simplified proof for more general asymptotic conditions and an L^6 norm bound.

All the existing results require the solutions to be maximal, which restricts the data to be a special time-slice in a spacetime. However it should be unnecessary according to the gravitational collapse pictures¹. It is natural and interesting to study the non-maximal case. In this paper we will prove the mass angular momentum inequality for non-maximal vacuum data with small trace by exploring the Einstein equations and a perturbation method. Using notations in Section 1.2, our main theorem is

Theorem 1.1. (Main Theorem 1) *Suppose (Σ, e) is a simply connected 3-manifold, which is Euclidean at infinity with two ends and axisymmetric in the sense of Definition 1.3. Given*

¹The axisymmetric condition is indeed necessary, since otherwise vacuum counterexamples were constructed by Huang, Schoen and Wang [18]

an asymptotically flat, axisymmetric vacuum data $(g, k) \in \mathcal{VC}_{s+2, \delta+\frac{1}{2}}^a(\Sigma)$ (see Definition 1.5) with $s \in \mathbb{N}$, $s \geq 7$, $\delta \in \mathbb{R}$, $-\frac{3}{2} < \delta < -1$, if $\|tr_g k\|_{H_{s-2, \delta+\frac{3}{2}}(\Sigma)} \leq \epsilon$ with ϵ given in Theorem 1.6, we have

$$m \geq \sqrt{|J|}, \quad (1.1)$$

where m and J are the ADM mass (1.6) and angular momentum (1.8) of (Σ, g, k) respectively.

Our method comes from a question suggested by R. Schoen:

(Q) : *Is there a canonical way to deform a non-maximal, axisymmetric, vacuum data to a unique maximal, vacuum data with the same physical quantities, i.e. the mass and angular momentum, which also preserves the axially symmetry?*

A definite answer of the above question will imply the mass angular momentum inequality in the non-maximal case. In fact, there are already some works about the deformation of vacuum constraint equations [4][13]. But it is hard to maintain the symmetries and physical quantities at the same time. So the main difficulty is to maintain the symmetries and the physical quantities simultaneously when deforming the vacuum constraint equations. We overcome this difficulty by using certain conversation laws of the Einstein equations.

1.1 General Relativity backgrounds

In Einstein's theory for General Relativity², we use $(\mathcal{V}^{3,1}, \gamma)$ to denote a spacetime, where $\mathcal{V}^{3,1}$ is a 4-dimensional oriented smooth manifold, and γ is a Lorentzian metric of signature $(3, 1)$. The Einstein equation, which predicts the evolution of the spacetime, is given by

$$Ric_\gamma - \frac{1}{2}R_\gamma\gamma = 8\pi T, \quad (1.2)$$

where Ric_γ is the Ricci curvature of γ , and R_γ the scalar curvature of γ . T is the stress-energy tensor. In the vacuum case, $T \equiv 0$, so the *Einstein vacuum equation*, abbreviated as **(EVE)** in the following, reduces to

$$Ric_\gamma = 0. \quad (1.3)$$

A *vacuum constraint initial data set* or abbreviated as *vacuum data* for the Einstein vacuum equations is a triple (Σ, g, k) , where Σ is a connected complete 3-dimensional manifold, g a Riemmanian metric, and k a symmetric two tensor on Σ , satisfying the *vacuum constrain equations*, abbreviated as **(VCE)**,

$$\begin{cases} R_g - |k|_g^2 + (tr_g k)^2 = 0, \\ div_g(k - (tr_g k)g) = 0. \end{cases} \quad (1.4)$$

By the famous initial value formulation for the Einstein equations by Y. Choquet-Bruhat in 1952(see [9][24]), we can always think the vacuum data (Σ, g, k) as been embedded in some

²We refer to [24] for all the concepts.

spacetime (\mathcal{V}, γ) satisfying (EVE), where g is the restriction of γ to Σ , and k is the second fundamental form of the embedding.

(Σ, e) is called *Euclidean at infinity*, where e is a Riemannian metric on Σ , if there is a compact subset $\Sigma_{int} \subset \Sigma$, such that the complement $\Sigma_{ext} = \Sigma \setminus \Sigma_{int}$ is a disjoint union of finitely many open sets $\Sigma_{ext} = \cup_i E_i$, and each E_i is diffeomorphic to \mathbb{R}^3 cutting off a ball B_R , and on each E_i , e is the pull back of the standard Euclidean metric on \mathbb{R}^3 . Here Σ_{int} is called the *interior region*, Σ_{ext} the *exterior region*, and each E_i an *end*. Each end E has a coordinate system $\{x_i : i = 1, 2, 3\}$ inherited from \mathbb{R}^3 . Let $r = \sqrt{\sum_i (x_i)^2}$. (Σ, g, k) is said to be *asymptotically flat*, abbreviated as **(AF)**, if (Σ, e) is Euclidean at infinity for some e , and there exists an $\alpha > \frac{1}{2}$, such that under coordinates $\{x_i : i = 1, 2, 3\}$,

$$g_{ij} = \delta_{ij} + o_2(r^{-\alpha}), \quad k_{ij} = o_1(r^{-1-\alpha}). \quad (1.5)$$

Under these conditions, the ADM mass is defined as,

$$m = \lim_{r \rightarrow \infty} \frac{1}{16\pi} \int_{S_r} (g_{ij,i} - g_{ii,j}) \nu^j d\sigma(r), \quad (1.6)$$

where $g_{ij,k} = \frac{\partial g_{ij}}{\partial x_k}$ and ν^j is the Euclidean unit outer normal of S_r with $d\sigma(r)$ the surface element of S_r . The famous positive mass theorem by Schoen and Yau [21][22] and Witten [25] says that $m \geq 0$ under the dominant energy condition.

If the initial data set (Σ, g, k) is axisymmetric([16][11]) under a Killing vector field ξ , i.e.

$$\mathcal{L}_\xi g = 0, \quad \mathcal{L}_\xi k = 0, \quad (1.7)$$

where \mathcal{L} denotes the Lie derivative, we also have a well-defined angular momentum J ([16][24]) of a close 2-surface $S \subset \Sigma$

$$J(S) = \frac{1}{8\pi} \int_S \pi_{ij} \xi^i \nu^j d\sigma_g, \quad (1.8)$$

where $\pi_{ij} = k_{ij} - \text{tr}_g(k)g_{ij}$ is divergence free by (1.4), and ν , $d\sigma_g$ are, respectively the unit outer normal of S and surface element w.r.t. g .

1.2 Ideas and main results

In this paper, we will prove the mass angular momentum inequality for certain axisymmetric, AF vacuum data (Σ, g, k) with small $\text{tr}_g k$, especially we partially solved the question asked by Schoen. We will use the full Einstein equations and a perturbation method. Given an AF vacuum data (Σ, g, k) , we will solve the boost problem of (EVE) for (Σ, g, k) as [10][3] to get a spacetime (\mathcal{V}, γ) , where \mathcal{V} is a subset of $\Sigma \times \mathbb{R}$ which grows linearly at infinity. Given a function u defined on Σ , the graph $\text{Graph}_u = \{(x, u(x)) \in \Sigma \times \mathbb{R}, x \in \Sigma\}$ of u lies inside \mathcal{V} , when $|u|$ has roughly sub-linear growth. We want to find a solution to $H_u = 0$, where H_u is the mean curvature of Graph_u w.r.t. (\mathcal{V}, γ) . Now fix a 3-manifold (Σ, e) Euclidean at infinity, we can construct a mapping \mathcal{H} which takes the triple (g, k, u) to the mean curvature H_u , i.e.

$\mathcal{H} : (g, k, u) \rightarrow H_u$. Viewing (g, k) as parameters and u as unknown function, our equation changes to

$$\mathcal{H}(g, k, u) = 0. \quad (1.9)$$

When (g, k) is maximal, $u \equiv 0$ is a solution to (1.9). So we can use the inverse function theorem to solve $\mathcal{H}(g, k, u) = 0$ when $tr_g k$ is small enough. From now on, we always assume $s \in \mathbb{N}$ and $\delta \in \mathbb{R}$. Using notations from Section 2, we have

Definition 1.2. Fix a 3-dimensional manifold (Σ, e) which is Euclidean at infinity.

- (1) The *vacuum constraint data sets* $\mathcal{VC}_{s+1, \delta + \frac{1}{2}}(\Sigma)$ is defined to be the set of solutions (g, k) to (1.4), such that $(g - e, k) \in H_{s+1, \delta + \frac{1}{2}}(\Sigma) \times H_{s, \delta + \frac{3}{2}}(\Sigma)$.
- (2) The *maximal vacuum constraint data sets* $\mathcal{MVC}_{s+1, \delta + \frac{1}{2}}(\Sigma)$ is defined to be the subset of $\mathcal{VC}_{s+1, \delta}(\Sigma)$ satisfying $tr_g k = 0$.

Inside $\mathcal{VC}_{s+1, \delta + \frac{1}{2}}(\Sigma)$ and $\mathcal{MVC}_{s+1, \delta + \frac{1}{2}}(\Sigma)$, we use the topology induced by the Sobolev norms of $H_{s+1, \delta + \frac{1}{2}}(\Sigma) \times H_{s, \delta + \frac{3}{2}}(\Sigma)$ as in Definition 2.3.

Definition 1.3. A simply connected 3-manifold (Σ, e) which is Euclidean at infinity is called *axisymmetric*, if

- (1) Σ is diffeomorphic to \mathbb{R}^3 minus some points $\{a_k\}_{k=1}^{l-1}$ on the z -axis $\Gamma = \{(\rho, \varphi, z) \in \mathbb{R}^3 : \rho = 0\}$, with one end modeled by a neighborhood of ∞ , and other ends by a neighborhood of a_k with coordinates given by a Kelvin transformation: $\{x' = \frac{x - a_k}{|x - a_k|^2}\}$;
- (2) $\mathcal{L}_{\partial_\varphi} e = 0$, where φ is the azimuth of the cylindrical coordinate $\{\rho, \varphi, z\}$.

Remark 1.4. Near ∞ , e is given by the Euclidean metric ds_0^2 , and near each puncture a_k , e is the pull back of the Euclidean metric by the Kelvin transformation, i.e. $e = \frac{1}{|x|^4} ds_0^2$. In fact, by Chruściel's reduction in [11], any simply connected, axisymmetric, AF vacuum data (Σ, g) has the underlying topology Σ given by \mathbb{R}^3 minus finitely many points on the z axis, with the Killing vector field $\frac{\partial}{\partial \varphi}$.

Definition 1.5. Given (Σ, e) as in Definition 1.3.

- (1) An initial data set (g, k) is called *axisymmetric*, if symmetry conditions (1.7) hold for the Killing vector field $\xi = \frac{\partial}{\partial \varphi}$.
- (2) $\mathcal{VC}_{s+1, \delta + \frac{1}{2}}^a(\Sigma)$ and $\mathcal{MVC}_{s+1, \delta + \frac{1}{2}}^a(\Sigma)$ are the axisymmetric subset of $\mathcal{VC}_{s+1, \delta + \frac{1}{2}}(\Sigma)$ and $\mathcal{MVC}_{s+1, \delta + \frac{1}{2}}(\Sigma)$ respectively.

The following Theorem is one of our main results, which is a summarization of Theorem 4.12, Lemma 4.13, Lemma 4.14 and Theorem 4.16.

Theorem 1.6. (Main Theorem 2) *Given $s \geq 4$, $-2 < \delta < -1$.*

- (i) *Let (Σ, e) be a 3-dimensional manifold which is Euclidean at infinity. For any $(g, k) \in \mathcal{VC}_{s+2, \delta + \frac{1}{2}}(\Sigma)$, where $\lambda e \leq g \leq \lambda^{-1}e$ for some $\lambda > 0$, there exists a small number ϵ depending only on λ and $\|g - e\|_{H_{s+2, \delta + \frac{1}{2}}(\Sigma)} + \|k\|_{H_{s+1, \delta + \frac{3}{2}}(\Sigma)}$, such that if $\|tr_g k\|_{H_{s-2, \delta + \frac{3}{2}}(\Sigma)} \leq \epsilon$, then there exists a spacetime (\mathcal{V}, γ) solving the (EVE), and a function $u \in H_{s+2, \delta - \frac{1}{2}}(\Sigma)$ solving the maximal surface equation (1.9) inside (\mathcal{V}, γ) . The induced metric g_u and second fundamental form k_u of Graph_u satisfy $(g_u, k_u) \in \mathcal{MVC}_{s+1, \delta + \frac{1}{2}}(\Sigma)$.*

- (ii) If $-\frac{3}{2} < \delta < -1$, the ADM mass of (Σ, g_u, k_u) is the same as that of (Σ, g, k) .
 (iii) If (Σ, e, g, k) is simply connected, axisymmetric, then u can be chosen to be axisymmetric, hence (Σ, g_u, k_u) is axisymmetric, and has the same angular momentum as (Σ, g, k) .

Remark 1.7. The weight δ corresponds the decay $g \sim e + o(r^{-(\delta+2)})$ and $k \sim o(r^{-(\delta+3)})$ by the Sobolev embedding lemma 2.5. (g_u, k_u) is always assumed to be pulled back to Σ by the graphical map $F_u : x \rightarrow (x, u(x))$.

Remark 1.8. The order of regularity of our final solution (g_u, k_u) decreases by 1 than our starting data (g, k) . This is due to the fact that the restriction of H_s Sobolev functions on a spacetime to a hypersurface decreases the regularity by 1 (see Lemma 2.8).

Our main Theorem 1.1 is then a corollary of the above theorem.

Proof of Theorem 1.1: Let u be the solution given in part (iii) of Theorem 1.6. Then the induced maximal data $(g_u, k_u) \in \mathcal{MVC}_{s+1, \delta+\frac{1}{2}}^a(\Sigma)$, and the ADM mass m and angular momentum J of (g, k) and (g_u, k_u) are the same. Now by Sobolev embedding lemma 2.5, $(g_u - e, k_u) \in C_{\beta}^{s-1}(\Sigma) \times C_{\beta+1}^{s-2}(\Sigma)$ for some $\frac{1}{2} < \beta < \delta+2 < 1$. So (Σ, g_u, k_u) is an axisymmetric, maximal vacuum data, with asymptotic conditions $g_u = \delta + O_{s-1}(\frac{1}{r^{\beta}})$ and $k_u = O_{s-2}(\frac{1}{r^{\beta+1}})$, so the mass angular momentum inequality in [23] holds on (Σ, g_u, k_u) . Hence $m \geq \sqrt{|J|}$. \square

The paper is organized as follows: In Section 2 we will review the weighted Sobolev space theories covered by [10][8][7][2] and the geometry of hypersurfaces in 3+1 dimension Lorentzian spaces. In Section 3 we will extend the boost theory in [10][8] to the case of multi-ends. In Section 4 we will set up a perturbation problem for the mean curvature of graphs. We will take initial data sets as parameters and use linear theory in [2][19] and [7] and the Quantitative Inverse Function Theorem 4.10. Finally we will prove the main results in Section 4.4.

Acknowledgement: The author would like to express his gratitude to his advisor Professor Richard Schoen for all of his helpful guidance and constant encouragement. He would like to thank Professor Rafe Mazzeo and Professor Leon Simon for lots of useful talks. He would also like to thank his friend Pin Yu for talking a lot about the hyperbolic equations.

2 Preliminaries

In this section, we give some preliminary results on the weighted Sobolev space theories and the geometry of hypersurfaces in Lorentz spaces.

2.1 Weighted Sobolev space theories

Here we give our definition of the weighted sobolev space. Most of the results here can be found in [7][10] and [8]. We will mainly talk about two types of domains.

Type 1 domain: sub-domain of \mathbb{R}^3 .

Let U be an open set in \mathbb{R}^n , $\sigma(x) = (1 + |x|^2)^{1/2}$ for $x \in \mathbb{R}^n$, and V a finite dimensional vector space. Given $s \in \mathbb{N}$, $\delta \in \mathbb{R}$.

Definition 2.1. $C_\delta^s(U)$ is the Banach space of C^s functions $u : U \rightarrow V$, with finite norm

$$\|u\|_{C_\delta^s(U)} = \sup_U \left\{ \sum_{|\alpha| \leq s} \sigma^{\delta+|\alpha|} |D^\alpha u| \right\}.$$

$H_{s,\delta}(U)$ is the class of functions $u : U \rightarrow V$, with weak derivatives up to order s , such that $\sigma^{\delta+|\alpha|} D^\alpha u \in L^2(U)$ for all $|\alpha| \leq s$. $H_{s,\delta}(U)$ is a Hilbert space with inner product:

$$\langle u_1, u_2 \rangle_{H_{s,\delta}(U)} = \sum_{|\alpha| \leq s} \langle \sigma^{\delta+|\alpha|} D^\alpha u_1, \sigma^{\delta+|\alpha|} D^\alpha u_2 \rangle_{L^2(U)}.$$

Then the norm is: $\|u\|_{H_{s,\delta}(U)} = \langle u, u \rangle_{H_{s,\delta}(U)}^{1/2}$.

Now we will list some properties of $H_{s,\delta}(U)$, which can be found in [10][7] and [8]. Given $0 < \epsilon \leq 1$, and $\phi_\epsilon : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by $\phi_\epsilon(x) = \frac{x}{(\sigma(x))^{1-\epsilon}}$. An open subset $U \subset \mathbb{R}^3$ is said to have the *extended cone property* if $\phi_\epsilon(U)$ has the cone property³ for each $0 < \epsilon \leq 1$.

Lemma 2.2. *Given U satisfying the extended cone property,*

- (i) (embedding). *If $s' < s - \frac{n}{2}$ and $\delta' < \delta + \frac{n}{2}$, the inclusion $H_{s,\delta}(U) \subset C_{\delta'}^{s'}(U)$ is continuous;*
- (ii) (multiplication). *If $s \leq s_1, s_2$, $s < s_1 + s_2 - \frac{n}{2}$ and $\delta < \delta_1 + \delta_2 + \frac{n}{2}$, the multiplication $(f_1, f_2) \rightarrow f_1 f_2$ is continuous: $H_{s_1,\delta_1}(U) \times H_{s_2,\delta_2}(U) \rightarrow H_{s,\delta}(U)$.*

Hence $H_{s,\delta}(U)$ is a Banach algebra if $s > \frac{n}{2}$ and $\delta > -\frac{n}{2}$.

Type 2 domain: manifold which is Euclidean at infinity.

Let (Σ, e) be an n dimensional manifold which is Euclidean at infinity. Let $x = \{x^i\}$ be the local coordinates, where $\{x^i\}$ is the pull back of the standard coordinates from $\mathbb{R}^n \setminus B_R$ when restricted to E_i , and $e = ds_0^2 = \sum_{i=1}^n (dx^i)^2$ on E_i . Fix a point $O \in \Sigma_{int}$, and define a function on Σ by

$$\sigma_e(x) = (1 + d_e^2(x, O))^{1/2}.$$

Clearly $\sigma_e(x)$ is equivalent to $\sigma(x) = (1 + |x|^2)^{1/2}$ on each end E_i .

When we use Σ to model an initial data set, the spacetime should have topology as a sub-domain of $\Sigma \times \mathbb{R}$. Using coordinates (x^i, t) on $\Sigma \times \mathbb{R}$, it has a natural reference metric

$$\tilde{e} = dt^2 + e. \tag{2.1}$$

For $\theta \in (0, 1]$, the *boost region* Ω_θ is defined as,

$$\Omega_\theta = \{(x, t) \in \Sigma \times \mathbb{R} : |t| \leq \theta \sigma_e(x)\}. \tag{2.2}$$

On Ω_θ , the distance function $d_{\tilde{e}}(\cdot, O)$ is equivalent to $d_e(\cdot, O)$, so we can use σ_e to define the weighted Sobolev space on Ω_θ . Given a smooth tensor bundle $E \rightarrow \Sigma$ or $E \rightarrow \Omega_\theta$ and $s \in \mathbb{N}$, $\delta \in \mathbb{R}$.

³See the remark under Definition 2.3 of [10]

Definition 2.3. $C_\delta^s(\Sigma)$ or $C_\delta^s(\Omega_\theta)$ is the Banach space of C^s sections $u : \Sigma \rightarrow E$, or $u : \Omega_\theta \rightarrow E$, with finite norm

$$\|u\|_{C_\delta^s(\Sigma(\text{or } \Omega_\theta))} = \sup_{\Sigma(\text{or } \Omega_\theta)} \left\{ \sum_{|\alpha| \leq s} \sigma_e^{\delta+|\alpha|} |D^\alpha u|_{e(\text{or } \bar{e})} \right\}.$$

$H_{s,\delta}(\Sigma)$ or $H_{s,\delta}(\Omega_\theta)$ is the class of sections $u : \Sigma \rightarrow E$, or $u : \Omega_\theta \rightarrow E$ with weak derivatives up to order s , such that $\sigma_e^{\delta+|\alpha|} D^\alpha u \in L^2(\Sigma, e)(\text{or } L^2(\Omega_\theta, \bar{e}))$ for all $\alpha \leq s$. $H_{s,\delta}(\Sigma)$ or $H_{s,\delta}(\Omega_\theta)$ is a Hilbert space with inner product:

$$\langle u_1, u_2 \rangle_{H_{s,\delta}(\Sigma)(\text{or } H_{s,\delta}(\Omega_\theta))} = \sum_{|\alpha| \leq s} \langle \sigma_e^{\delta+|\alpha|} D^\alpha u_1, \sigma_e^{\delta+|\alpha|} D^\alpha u_2 \rangle_{L^2(\Sigma, e)(\text{or } L^2(\Omega_\theta, \bar{e}))}.$$

Then the norm is: $\|u\|_{H_{s,\delta}(\Sigma)(\text{or } H_{s,\delta}(\Omega_\theta))} = \langle u, u \rangle_{H_{s,\delta}(\Sigma)(\text{or } H_{s,\delta}(\Omega_\theta))}^{1/2}$.

Remark 2.4. In fact, the definitions are independent of the choice of e on Σ_{int} .

Lemma 2.5. (Lemma 2.4, 2.5 in [7], Appendix 1 in [6])

- (i) (embedding). If $s' < s - \frac{n}{2}$, $\delta' < \delta + \frac{n}{2}$, the inclusion $H_{s,\delta}(\Sigma) \subset C_{\delta'}^{s'}(\Sigma)$ is continuous;
- (ii) (multiplication). If $s \leq s_1, s_2$, $s < s_1 + s_2 - \frac{n}{2}$, $\delta < \delta_1 + \delta_2 + \frac{n}{2}$, the multiplication $(f_1, f_2) \rightarrow f_1 f_2$ is a continuous map: $H_{s_1, \delta_1}(\Sigma) \times H_{s_2, \delta_2}(\Sigma) \rightarrow H_{s, \delta}(\Sigma)$, hence $H_{s, \delta}(\Sigma)$ is a Banach algebra if $s > \frac{n}{2}$, $\delta > -\frac{n}{2}$. Furthermore,

$$\|f_1 f_2\|_{H_{s, \delta}(\Sigma)} \leq C \|f_1\|_{H_{s_1, \delta_1}(\Sigma)} \|f_2\|_{H_{s_2, \delta_2}(\Sigma)}, \quad (2.3)$$

where C is a constant depending only on $\{n, s_1, s_2, \delta_1, \delta_2\}$.

Divide Ω_θ as $\Omega_\theta = (\Omega_\theta)_{int} \cup_{i=1}^l (\Omega_\theta)_i$, where $(\Omega_\theta)_i = \{(x, t) \in \Omega_\theta : x \in E_i\}$, and $(\Omega_\theta)_{int}$ the compliment. Now $(\Omega_\theta)_{int}$ is a compact manifold, and $(\Omega_\theta)_i \subset \mathbb{R}^{n+1}$ satisfies the extended cone property in the above section, and hence Lemma 2.2. By working separately on $(\Omega_\theta)_i$ and $(\Omega_\theta)_{int}$ as in [7] using Lemma 2.2, we have similar results,

- Lemma 2.6.** (i) (embedding). If $s' < s - \frac{n+1}{2}$, $\delta' < \delta + \frac{n+1}{2}$, the inclusion is $H_{s+1, \delta}(\Omega_\theta) \subset C_{\delta'}^{s'}(\Omega_\theta)$ is continuous;
- (ii) (multiplication). If $s \leq s_1, s_2$, $s < s_1 + s_2 - \frac{n+1}{2}$, $\delta < \delta_1 + \delta_2 + \frac{n+1}{2}$, then the multiplication $(f_1, f_2) \rightarrow f_1 f_2$ is a continuous map: $H_{s_1, \delta_1}(\Omega_\theta) \times H_{s_2, \delta_2}(\Omega_\theta) \rightarrow H_{s, \delta}(\Omega_\theta)$, hence $H_{s, \delta}(\Omega_\theta)$ is a Banach algebra if $s > \frac{n+1}{2}$, $\delta > -\frac{n+1}{2}$.

Using ideas similar to the proof of Theorem 2.3 in [10] and Lemma 2.6, we have

Lemma 2.7. (composition). Given $\Omega_\theta, \Omega_{\theta'}$ as above and $f : \Omega_\theta \rightarrow \Omega_{\theta'}$ a differentiable map, such that $|Df|_{\bar{e}} \geq c > 0$ and $f - id \in H_{s+1, \delta-1}(\Omega_\theta)$ with $s > \frac{n+1}{2}$ and $\delta > -\frac{n+1}{2}$, then for any $s' \leq s+1$, $\delta' \in \mathbb{R}$, the composition $u \rightarrow u \circ f$ is an isomorphism as a map:

$$H_{s', \delta'}(f(\Omega_\theta)) \rightarrow H_{s', \delta'}(\Omega_{\theta'}).$$

Define the function $\tau(x, t) = \frac{t}{\sigma_e(x)}$. Denote the level surface of τ by $\Sigma_\tau = \{(x, t) \in \Sigma \times \mathbb{R} : \tau(x, t) = \tau\}$. Then Ω_θ has a foliation $\Omega_\theta = \cup_{\tau \in (-\theta, \theta)} \Sigma_\tau$. The restriction norm is defined as:

$$\|u\|_{H_{s,\delta}(\Sigma_\tau, \Omega_\theta)} = \left(\sum_{k=0}^s \|D_t^k u|_{\Sigma_\tau}\|_{H_{s-k,\delta+k}(\Sigma)}^2 \right)^{1/2}. \quad (2.4)$$

Using ideas similar to the proof of Lemma 3.1 in [8], we can get,

Lemma 2.8. (restriction). $\forall \tau \in (-\theta, \theta)$, we have the following continuous inclusion:

$$H_{s+1,\delta}(\Omega_\theta) \subset H_{s,\delta+\frac{1}{2}}(\Sigma_\tau, \Omega_\theta),$$

for every $s \in \mathbb{N}$ and $\delta \in \mathbb{R}$.

2.2 Geometry of hypersurface in Lorentzian space

In this section, we will review the geometry of hypersurfaces in a Lorentzian space. We will mainly focus on the mean curvature of the hypersurface. Notation and part of the results here trace back to [1], and all concepts of Lorentzian space can be found in [24]. Let (\mathcal{V}, γ) be a (3+1) dimensional Lorentzian space, with $\langle \cdot, \cdot \rangle$ the metric pairing and ∇ the connection. A smooth function $t \in C^\infty(\mathcal{V})$ is called a *time function* if ∇t is nonzero, and everywhere *timelike*, i.e. $\langle \nabla t, \nabla t \rangle < 0$. We call a hypersurface Σ *spacelike* if the restriction of γ to Σ is Riemannian. In a local coordinate system $\{x^i, t\}$, where t is a time function, the metric can be written as (See equation (2.12) of [1]):

$$\gamma = -(\alpha^2 - \beta^2)dt^2 + 2\beta_i dx^i dt + g_{ij} dx^i dx^j, \quad (2.5)$$

where α is the lapse function, i.e. $\alpha^2 = -\langle \nabla t, \nabla t \rangle$, g_{ij} a Riemannian metric, and $\beta = g^{ij} \beta_i \partial_j$ the shift vector⁴. Here we use $\partial_t = \frac{\partial}{\partial t}$ and $\partial_i = \frac{\partial}{\partial x^i}$ as coordinate vectors. The inverse metric γ^{-1} is given by:

$$\gamma^{\mu\nu} = \begin{bmatrix} -\frac{1}{\alpha^2} & \frac{\beta^i}{\alpha^2} \\ \frac{\beta^j}{\alpha^2} & g^{ij} - \frac{1}{\alpha^2} \beta^i \beta^j \end{bmatrix}, \quad (2.6)$$

under coordinate system $\{t, x^1, x^2, x^3\}$.

We will denote the level surface of the time function t by $\Sigma_t = \{p \in \mathcal{V} : t(p) = t\}$. Let D be the gradient operator on Σ_t , and div^0 the divergence operator on Σ_t . The future-directed timelike unite normal T of Σ_t is given by⁵:

$$T = -\alpha \nabla t = \alpha^{-1}(\partial_t - \beta), \quad (2.7)$$

and the second fundamental form k_{ij}^0 and the mean curvature H^0 of the slice Σ_t are given by,

$$k_{ij}^0 = \langle \partial_i, \nabla_{\partial_j} T \rangle = \frac{1}{2} \alpha^{-1} \partial_t g_{ij} - \frac{1}{2} \alpha^{-1} \mathcal{L}_\beta g_{ij}, \quad (2.8)$$

⁴See Chap 10 of [24] for details.

⁵See Appendix 5 for details.

$$H^0 = g^{ij} A_{ij}^0 = \frac{1}{2} \alpha^{-1} g^{ij} \partial_t g_{ij} - \alpha^{-1} \text{div}^0(\beta). \quad (2.9)$$

Given a spacelike hypersurface Σ , we can always choose local coordinates $\{x^i, t\}$, such that Σ is locally the $t = 0$ level surface Σ_0 . Given a smooth function $u \in C^\infty(\Sigma)$, we can study the graph of u , i.e. $\text{Graph}_u = \{(x^i, u(x))\}$ in local coordinates. So we call this u the *height function*. By extending u parallel to \mathcal{V} requiring that

$$\partial_t u = 0, \quad (2.10)$$

Graph_u can be viewed as level surface of $(u - t) = 0$. The unit normal of Graph_u is⁶:

$$N = \nu(U + T), \quad (2.11)$$

where

$$U = \frac{\alpha Du}{1 + \langle \beta, Du \rangle}, \text{ and } \nu = \frac{1}{(1 - |U|_g^2)^{1/2}}. \quad (2.12)$$

So Graph_u is spacelike iff $1 - |U|_g^2 > 0$, i.e. ν well-defined. Define the canonical graphical diffeomorphism $F : \Sigma \rightarrow \text{Graph}_u$ by $F(x) = (x, u(x))$. Then Graph_u has a local coordinate system $\{x^i : i = 1, 2, 3\}$. The coordinate vector frame $\{\partial_i\}$ on Σ is passed by F to a local frame

$$\alpha_i = \partial_i + u_i \partial_t : i = 1, 2, 3, \quad (2.13)$$

on Graph_u . Now denote $M = \text{Graph}_u$. Using this local coordinates, the restriction $\gamma|_M$ of γ to Graph_u , denoting by $g_M = (g_M)_{ij} dx^i dx^j$, is given by

$$(g_M)_{ij} = g_{ij} + \beta_i u_j + u_i \beta_j - (\alpha^2 - \beta^2) u_i u_j. \quad (2.14)$$

Then the inverse metric matrix is calculated in the Appendix 5 by equation (5.12) as:

$$\begin{aligned} (g_M)^{ij} &= g^{ij} - \frac{1}{\alpha^2} \beta^i \beta^j + \frac{\nu^2}{\alpha^2} (\beta - \alpha U)^i (\beta - \alpha U)^j \\ &= \gamma^{ij} + \frac{\nu^2}{\alpha^2} (\beta - \alpha U)^i (\beta - \alpha U)^j. \end{aligned} \quad (2.15)$$

So the mean curvature H_u of the graph M is given by

$$H_u = (g_M)^{ij} \langle \nabla_{\alpha_i} N, \alpha_j \rangle_\gamma. \quad (2.16)$$

3 Boost evolution

Fix a 3-manifold (Σ, e) , which is Euclidean at infinity. Let $\tilde{e} = dt^2 + e$ be the reference metric(2.1) on $\Sigma \times \mathbb{R}$. Given integer $s \geq 4$, and real number $\delta > -2$, we consider vacuum constraint initial data sets (Σ, g, k) , such that $(g, k) \in \mathcal{V}_{s, \delta + \frac{1}{2}}(\Sigma)$. Here boost evolution means

⁶See Appendix 5 for details.

that in the spacetime (\mathcal{V}, γ) which is evolved by (EVE) taking (Σ, g, k) as initial data set, where $\mathcal{V} \subset \Sigma \times \mathbb{R}$, both the future and past temporal distance $\chi_{\pm}(x)$ ⁷ to the boundary of \mathcal{V} is proportional to the space distance $\sigma(x)$ for $x \in \Sigma$, i.e. $\chi_{\pm}(x) \geq c\sigma(x)$ for $c > 0$. We will extend the boost evolution on \mathbb{R}^3 in [10] to the case of Σ .

3.1 Reduced Einstein equation and results on compact domain

Let us review the reduction using harmonic gauge initially introduced by Y. Choquet-bruhat (see [6]). Using $\{x^i : i = 1, 2, 3\}$ as local coordinates on Σ , and $x^\mu = (x^0, x^i)$, with $x^0 = t$ as coordinates on $\mathcal{V} \subset \Sigma \times \mathbb{R}$, the Ricci curvature can be expressed as⁸:

$$Ric^{\mu\nu} = R_h^{\mu\nu} + \frac{1}{2}(\gamma^{\mu\alpha} D_\alpha \Gamma^\nu + \gamma^{\nu\alpha} D_\alpha \Gamma^\mu),$$

where $\Gamma_{\alpha\beta}^\mu$ is the Christoffel symbol of γ , $\Gamma^\mu = \gamma^{\alpha\beta} \Gamma_{\alpha\beta}^\mu$, and

$$R_h^{\mu\nu} = \frac{1}{2}\{\gamma^{\alpha\beta} D_\alpha D_\beta \gamma^{\mu\nu} - B^{\mu\nu}(\gamma, D\gamma)\},$$

with $B^{\mu\nu} = P_{\alpha\beta, \kappa\lambda}^{\mu\nu, \rho\sigma} D_\rho \gamma^{\alpha\beta} D_\sigma \gamma^{\kappa\lambda}$, and P is a rational function of $\gamma^{\alpha\beta}$. In fact, The Einstein vacuum equation $Ric_\gamma = 0$ is a degenerated differential equation system due to its invariance under diffeomorphic transformations. *Harmonic gauge* is used to fix this gauge freedom by Y. Choquet-bruhat, which means that we can choose $id : (\mathcal{V}, \gamma) \rightarrow (\mathcal{V}, \tilde{e})$ to be a wave map, i.e. $\square_{(\gamma, e)} id = 0$ ⁹. Denote

$$f^\mu = \Gamma^\mu - \gamma^{\alpha\beta} \tilde{\Gamma}_{\alpha\beta}^\mu, \quad (3.1)$$

to be the *harmonic gauge vector*, where $\tilde{\Gamma}_{\alpha\beta}^\mu$ the Christoffel symbol of \tilde{e} . f^μ is the difference of two connections, hence a tensor, then harmonic gauge condition reduces to $f^\mu = 0$, or:

$$\square_\gamma x^\mu = -\gamma^{\alpha\beta} \tilde{\Gamma}_{\alpha\beta}^\mu, \quad (3.2)$$

where \square_γ is the Laplacian operator of the Lorentzian metric γ , and $\square_\gamma x^\mu = -\Gamma^\mu$. Now under harmonic gauge (3.2), the (EVE)(1.3) reduced to¹⁰

$$\gamma^{\alpha\beta} D_\alpha D_\beta \gamma^{\mu\nu} = B^{\mu\nu}(\gamma, D\gamma) + \frac{1}{2} \gamma^{\alpha\beta} \{\gamma^{\mu\rho} \tilde{R}_{\beta\alpha\rho}^\nu + \gamma^{\nu\rho} \tilde{R}_{\beta\alpha\rho}^\mu\}, \quad (3.3)$$

where \tilde{R} is the curvature of \tilde{e} . The Cauchy data for these equations consist of:

$$\gamma|_\Sigma = \phi, \quad D_t \gamma|_\Sigma = \psi. \quad (3.4)$$

⁷See [10] for reference.

⁸See Sec. 4 of [10] and Chap. 10.2 of [24].

⁹See Section 7.4 of Chap 6 in [6]

¹⁰See page 163 in [6].

For given initial data set (g, k) , we need to construct Cauchy data (ϕ, ψ) by requiring $f^\mu|_\Sigma = (\Gamma^\mu - \gamma^{\alpha\beta} \tilde{\Gamma}_{\alpha\beta}^\mu)|_\Sigma = 0$. To fix the freedom in choosing a harmonic gauge, we require the coordinate system of \mathcal{V} is Gaussian on Σ , which means:

$$\phi^{00} = -1, \quad \phi^{0i} = 0, \quad \phi^{ij} = g^{ij}. \quad (3.5)$$

The condition $(\Gamma^\mu - \gamma^{\alpha\beta} \tilde{\Gamma}_{\alpha\beta}^\mu)|_\Sigma = 0$ requires¹¹:

$$\psi^{00} = -4tr_g k, \quad \psi^{0i} = -(\Gamma_g^i - g^{kj} \tilde{\Gamma}_{kj}^i), \quad \psi^{ij} = 2g^{ik} g^{jl} k_{kl}. \quad (3.6)$$

Define a reference Lorentzian metric by

$$\tilde{\eta} = -dt^2 + e. \quad (3.7)$$

When the initial data $(g - e, k) \in H_{s, \delta + \frac{1}{2}}(\Sigma) \times H_{s-1, \delta + \frac{3}{2}}(\Sigma)$, the Cauchy data (3.5)(3.6) satisfy $(\phi - \tilde{\eta}, \psi) \in H_{s, \delta + \frac{1}{2}}(\Sigma) \times H_{s-1, \delta + \frac{3}{2}}(\Sigma)$. In fact, by multiplication lemma 2.5, $(g - e, k) \rightarrow (\phi - \tilde{\eta}, \psi)$ is a continuous map $H_{s, \delta + \frac{1}{2}}(\Sigma) \times H_{s-1, \delta + \frac{3}{2}}(\Sigma) \rightarrow H_{s, \delta + \frac{1}{2}}(\Sigma) \times H_{s-1, \delta + \frac{3}{2}}(\Sigma)$.

To solve (EVE)(1.3), we can first solve the reduced equation (3.3) by quasilinear theory (see Appendix 3 in [6], and Section 5 in [10]), and then show that the harmonic gauge is preserved. In fact, Bianchi identity and the reduced equation (3.3) imply that the harmonic gauge vector f^μ satisfies a linear equation¹²:

$$\square_\gamma f^\mu + A(\gamma, D\gamma) Df = 0. \quad (3.8)$$

So we can use uniqueness of linear equations to show that $f^\mu \equiv 0$ since we chose $f^\mu|_\Sigma = 0$, and the constraint equations (1.4) implies that $\partial_t f^\mu|_\Sigma = 0$ ¹³.

Now we summarize a local version of the existence and causal uniqueness theorem based on the interior region Σ_{int} of (Σ, e) , which has dimension $n = 3$. We can extend the interior region Σ_{int} to contain the annuli $B_{2R} \setminus B_R$ of each end E_i of (Σ, e) . Now define a causal set $(V_{int})_{\theta, \lambda}$ based on Σ_{int} as follows:

$$(V_{int})_{\theta, \lambda} = \{(x, t) \in \Sigma_{int} \times [-\theta, \theta] : |x| \leq 2R - \lambda|t|, \text{ if } x \in E_i\}, \quad (3.9)$$

where $\theta \in (0, 1]$ and $\lambda \geq 2$ is a positive number. Now $(V_{int})_{\theta, \lambda}$ has a lateral boundary $L_{\theta, \lambda} = \{(x, t) \in (V_{int})_{\theta, \lambda} : |x| = 2R - \lambda|t|\}$. When λ is large enough depending only on e , $L_{\theta, \lambda}^+ = L_{\theta, \lambda} \cap \{t \geq 0\}$ (or $L_{\theta, \lambda}^- = L_{\theta, \lambda} \cap \{t \leq 0\}$) is spacelike and ingoing (or outgoing) w.r.t. $\tilde{\eta}$, hence $((V_{int})_{\theta, \lambda}, \tilde{\eta})$ is causal¹⁴.

Combining Theorem 7.4, Theorem 8.3 of Chap 6, and Corollary 4.8, Theorem 4.11, Theorem 4.13 of Appendix 3 in [6], and using a cutoff argument as in Theorem 3.7, we have the following well-known local existence and uniqueness theorem,

¹¹See page 164 in [6].

¹²See page 167 in [6] and Section 4 in [10]

¹³See page 167 in [6].

¹⁴See Definition 2.11 of Appendix 3 in [6].

Theorem 3.1. *Given an integer $s \geq 4$. For a vacuum constraint data set (Σ_{int}, g, k) , with $(g - e, k) \in H_s(\Sigma_{int}) \times H_{s-1}(\Sigma_{int})$, and $g \geq \lambda_0 e$ for some $\lambda_0 > 0$, there exists $\theta > 0$, $\lambda \geq 2$ and $C_0 > 0$ depending only on λ_0 and $\|g - e\|_{H_s(\Sigma_{int})} + \|k\|_{H_{s-1}(\Sigma_{int})}$, and a unique regularly sliced¹⁵ Lorentzian metric γ solving the reduced EVE(3.3) on $(V_{int})_{\theta, \lambda}$, taking (3.5)(3.6) as initial value which is given by (g, k) , such that $(\gamma - \tilde{\eta}) \in H_s((V_{int})_{\theta, \lambda})$, with $\|\gamma - \tilde{\eta}\|_{H_s((V_{int})_{\theta, \lambda})} \leq C_0$, and $L_{\theta, \lambda}^+$ (or $L_{\theta, \lambda}^-$) is spacelike and ingoing (or outgoing) w.r.t γ . Furthermore, γ is a solution of (EVE)(1.3) under harmonic gauge.*

3.2 Boost evolution on manifold Euclidean at infinity

We first modify the linear boost theory in [10] to the case based on an Euclidean end $E \cong \mathbb{R}^{n-1} \setminus B_R$. Let us fix a special type of boost regions. Denote $\bar{x} = (x^1, \dots, x^{n-1}) \in \mathbb{R}^{n-1}$, such that $x = (\bar{x}, t) \in \mathbb{R}^n$. Later on, we will denote the index for t -coordinates as 0, while index for \bar{x} as i with $i = 1, \dots, n-1$. Let $\bar{\sigma}(\bar{x}) = (1 + |\bar{x}|^2)^{1/2}$. For $\theta \in (0, 1/2]$, $\lambda \geq 2$ and a given end $E \cong \mathbb{R}^{n-1} \setminus B_R$, the boost region $V_{\theta, \lambda}$ based on E is defined as:

$$V_{\theta, \lambda} = \{(\bar{x}, t) \in \mathbb{R}^n, \frac{|t|}{\bar{\sigma}(\bar{x})} < \theta, |\bar{x}| \geq R + \lambda|t|\}, \quad (3.10)$$

Define function τ as $\tau(x) = \frac{t}{\bar{\sigma}(\bar{x})}$. Then the level surface of τ is $E_\tau = \{x \in V_{\theta, \lambda} : \tau(x) \equiv \tau\}$. $V_{\theta, \lambda}$ has a foliation:

$$V_{\theta, \lambda} = \cup_{\tau \in (-\theta, \theta)} E_\tau.$$

The lateral boundary of $V_{\theta, \lambda}$ is defined as,

$$L_{\theta, \lambda} = \{(\bar{x}, t) \in V_{\theta, \lambda} : |\bar{x}| = R + \lambda|t|\}. \quad (3.11)$$

Denote the upper part of $V_{\theta, \lambda}$ by $V_{\theta, \lambda}^+ = \{(\bar{x}, t) \in V_{\theta, \lambda} : t \geq 0\}$, then the boundary $\partial V_{\theta, \lambda}^+$ is constituted by E , E_θ and the upper lateral boundary $L_{\theta, \lambda}^+ = L_{\theta, \lambda} \cap V_{\theta, \lambda}^+$. Similarly, we have $V_{\theta, \lambda}^- = \{(\bar{x}, t) \in V_{\theta, \lambda} : t \leq 0\}$ and lower lateral boundary $L_{\theta, \lambda}^- = L_{\theta, \lambda} \cap V_{\theta, \lambda}^-$. Clearly $V_{\theta, \lambda}^\pm$ and the slices E_τ satisfy the extended cone property in \mathbb{R}^n and \mathbb{R}^{n-1} respectively as in Section 2.1, and hence satisfy Lemma 2.2.

We introduce a class of hyperbolic metrics on $V_{\theta, \lambda}$ using the foliation $\{E_\tau\}_{\tau \in (-\theta, \theta)}$. The function τ is in fact a time function on $(V_{\theta, \lambda}, \eta)$, where $\eta = -dt^2 + \sum_{i=1}^{n-1} (dx^i)^2$ is the Minkowski metric. Let \tilde{n}_μ be the unit future co-normal of the foliation $\{E_\tau : \tau \in (-\theta, \theta)\}$, given by

$$\tilde{n} = \tilde{N} D\tau = \frac{1}{\sqrt{1 - \tau^2 |\bar{x}|^2 \sigma^{-2}(\bar{x})}} (dt - \frac{\tau}{\sigma(\bar{x})} x^i dx^i), \quad (3.12)$$

where \tilde{N} is the lapse function for the foliation $\{E_\tau\}$, defined by: $\tilde{N}^{-2} = -\langle D\tau, D\tau \rangle_\eta = \frac{1 - \tau^2 |\bar{x}|^2 \sigma^{-2}(\bar{x})}{\sigma^2(\bar{x})}$. \tilde{n} can be viewed as a standard calibration for the foliation $V_{\theta, \lambda} = \cup E_\tau$, which is used to define the “regularity” of hyperbolicity. Denoting $|\cdot|$ as the standard Euclidean norm for tensors on V_θ , we have¹⁶:

¹⁵See page 397 and page 585 in [6]

¹⁶See also Definition 4.1 in [10].

Definition 3.2. A C^0 covariant symmetric 2-tensor $\gamma^{\mu\nu}$ on $V_{\theta,\lambda}$ is called *regularly hyperbolic*, if there exist positive numbers a , b and C such that in $V_{\theta,\lambda}$:

- (1) $-\gamma^{\mu\nu}\tilde{n}_\mu\tilde{n}_\nu \geq a$;
- (2) for all tangent covectors ζ_μ of E_τ , i.e. $\gamma^{\mu\nu}\zeta_\mu\tilde{n}_\nu = 0$, we have $\gamma^{\mu\nu}\zeta_\mu\zeta_\nu \geq b|\zeta|^2$;
- (3) $|\gamma| \leq C$;
- (4) The upper(or lower) lateral boundary $L_{\theta,\lambda}^+$ (or $L_{\theta,\lambda}^-$) is spacelike and ingoing(or out going) w.r.t. γ , i.e. every timelike curve entering $V_{\theta,\lambda}^+$ (or every timeline curve exiting $V_{\theta,\lambda}^-$) is past directed.

The *coefficient of regular hyperbolicity* of γ is defined as,

$$h = \max\left\{\frac{1}{a}, \frac{1}{b}, C\right\}. \quad (3.13)$$

Remark 3.3. Condition (4) implies that this type of $V_{\theta,\lambda}$ is a *causal subset* based on E w.r.t γ ¹⁷. Here we briefly talk about the criterion for Condition (4) to be true. We mainly discuss the case $L_{\theta,\lambda}^+$, and $L_{\theta,\lambda}^-$ is similar. The defining function of L^+ is given by $l(\bar{x}, t) = \lambda t + R - |\bar{x}|$, so the normal co-vector of L^+ is given by $dl = \lambda dt - d\bar{r}$, where $\bar{r} = |\bar{x}|$. Now $dl = \lambda(dt - \frac{\tau\bar{r}}{\sigma(\bar{x})}d\bar{r}) + (\lambda\frac{\tau\bar{r}}{\sigma(\bar{x})} - 1)d\bar{r} = \lambda\sqrt{1 - \tau^2|\bar{x}|^2\sigma^{-2}(\bar{x})}\tilde{n} + (\lambda\frac{\tau\bar{r}}{\sigma(\bar{x})} - 1)d\bar{r}$. So using the regularly hyperbolicity, we have $\gamma(dl, dl) \leq \lambda^2(1 - \theta^2)\gamma(\tilde{n}, \tilde{n}) + \lambda(\theta\lambda - 1)C \leq -a\lambda^2(1 - \theta^2) + C\lambda(\theta\lambda - 1) < 0$, when λ is chosen large enough depending only on a and C , hence on h .

Remark 3.4. The set of regularly hyperbolic metrics on $V_{\theta,\lambda}$ is open in the space $C^0(V_{\theta,\lambda})$ of bounded continuous covariant symmetric 2-tensors. In fact, η is regular hyperbolic with $a = 1$, $b = 1 - \theta^2$ and $C = \sqrt{n}$, and $L_{\theta,\lambda}$ is space-like and ingoing w.r.t η when $\lambda \geq 2$. Since the space-like and ingoing condition for $L_{\theta,\lambda}$ is an open condition, there exist a small $\epsilon > 0$, depending only on θ , λ and n , such that any C^0 covariant symmetric 2-tensor γ , with $|\gamma - \eta| \leq \epsilon$, is regularly hyperbolic in $V_{\theta,\lambda}$.

Now consider a family of linear differential operators of second order in $V_{\theta,\lambda}$:

$$Lu = \sum_{k=0}^2 a_k \cdot D^k u, \quad (3.14)$$

where u and Lu are \mathbb{R}^N valued functions on $V_{\theta,\lambda}$, and a_k are matrix valued functions. The following hypotheses are required to the existence theory:

- *Hypothesis (1)*(weak coupling and hyperbolicity). $a_2 = \gamma Id$, i.e. $(a_2)^{\mu\nu I}_J = \gamma^{\mu\nu} \delta^I_J$, $\mu, \nu = 0, \dots, n-1$, $I, J = 1, \dots, N$, where γ is a regularly hyperbolic metric on $V_{\theta,\lambda}$.
- *Hypothesis (2)*(regularity). There exist integers s_k and real numbers δ_k , such that: $s_k > \frac{n}{2} + k - 1$, $\delta_k > 2 - k - \frac{n}{2}$: $0 \leq k \leq 2$, and (1): $a_k \in H_{s_k, \delta_k}(V_{\theta,\lambda})$ for $k = 0, 1$; (2) $\gamma - \eta \in H_{s_2, \delta_2}(V_{\theta,\lambda})$.

¹⁷ See Definition 2.11 of Appendix 3 in [6].

Remark 3.5. Now denote

$$s' = \min_{0 \leq k \leq 2} \{s_k\} + 1, \quad (3.15)$$

$$m = \|\gamma - \eta\|_{H_{s_2, \delta_2}(V_{\theta, \lambda})} + \sum_{k=0}^1 \|a_k\|_{H_{s_k, \delta_k}(V_{\theta, \lambda})}, \quad (3.16)$$

$$\mu = \|\gamma - \eta\|_{H_{s_2-1, \delta_2+1/2}(E, V_{\theta, \lambda})} + \sum_{k=0}^1 \|a_k\|_{H_{s_k-1, \delta_k+1/2}(E, V_{\theta, \lambda})}. \quad (3.17)$$

By the restriction Lemma 2.8, $\mu \leq cm$. Using the multiplication Lemma 2.2, the regularity hypothesis (2) implies that

$$L : H_{s+1, \delta}(V_{\theta, \lambda}) \rightarrow H_{s-1, \delta+2}(V_{\theta, \lambda}),$$

is a continuous map for $1 \leq s \leq s'$ and $\delta \in \mathbb{R}$.

Then we have the existence and uniqueness theorem for linear systems.

Theorem 3.6. *Let L be a differential operator defined by (3.14) in $V_{\theta, \lambda}$, satisfying Hypotheses (1) and (2). Let $\beta \in H_{s-1, \delta+2}(V_{\theta, \lambda})$, $\phi \in H_{s, \delta+\frac{1}{2}}(E)$ and $\psi \in H_{s-1, \delta+\frac{3}{2}}(E)$, with $2 \leq s \leq s'$, $\delta \in \mathbb{R}$. Then the Cauchy problem:*

$$Lu = \beta, \quad u|_{\Sigma} = \phi, \quad D_t u|_{\Sigma} = \psi, \quad (3.18)$$

has a unique solution $u \in H_{s, \delta}(V_{\theta, \lambda})$, and satisfies the estimates:

$$\|u\|_{H_{s, \delta}(V_{\theta, \lambda})} \leq c\theta^{\frac{1}{2}} \{ \|\phi\|_{H_{s, \delta+\frac{1}{2}}(E)} + \|\psi\|_{H_{s-1, \delta+\frac{3}{2}}(E)} + \|\beta\|_{H_{s-1, \delta+2}(V_{\theta, \lambda})} \}, \quad (3.19)$$

where c is a continuous increasing function of (θ, h, m) , and h, m are defined by equations (3.13)(3.16) respectively.

Proof. It follows from the energy estimates Theorem 5.8 in Appendix 5.2, and similar approximation argument as in the proof of Theorem 5.1 in [8] and Theorem 4.1 in [10]. \square

Now we extend the existence theory for the boost problem in [10] to Σ . Let Ω_{θ} be the boost region based on Σ as defined in (2.2). We will construct a solution to the reduced EVE(3.3) in Ω_{θ} under the harmonic gauge. We deal with the boost evolution separately on the interior region Σ_{int} and on each end E_i . On compact set Σ_{int} , we can use Theorem 3.1. On each end E , we can complete the initial data $(g, k)|_E$ to \mathbb{R}^3 and apply the boost theory in [10] to get existence. Then we can cut off the solution in the causal set based on the end E by our linear Theorem 3.6. Causal uniqueness (see Corollary 4.8 of Appendix 3 in [6]) tells us that the solutions we got based on Σ_{int} and E_i 's match together to a global solution.

Theorem 3.7. *For $s \geq 4$, $\delta > -2$. Given vacuum data $(g, k) \in \mathcal{VC}_{s, \delta+\frac{1}{2}}(\Sigma)$, with $g \geq \lambda_0 e$ for some $\lambda_0 > 0$, there exists a $\theta \in (0, 1)$ and a $C_0 > 0$ depending only on λ_0 , $\|g - e\|_{H_{s, \delta+\frac{1}{2}}(\Sigma)} + \|k\|_{H_{s-1, \delta+\frac{3}{2}}(\Sigma)}$, and a unique Lorentzian metric γ solving the reduced EVE (3.3) on Ω_{θ} , which has Cauchy data (ϕ, ψ) on Σ given by (g, k) in (3.5)(3.6), such that $(\gamma - \tilde{\eta}) \in H_{s, \delta}(\Omega_{\theta})$, and $\|\gamma - \tilde{\eta}\|_{H_{s, \delta}(\Omega_{\theta})} \leq C_0$. Furthermore γ is the solution to EVE (1.3) under harmonic gauge.*

Proof. We first focus on a fixed end E . In fact, we can extend $(g, k)|_E$ to (\bar{g}, \bar{k}) on \mathbb{R}^3 by a cut and paste method, such that $(\bar{g}, \bar{k}) = (g, k)$ on E with $\bar{g} \geq \bar{\lambda}\delta$, where $\bar{\lambda} \geq c^{-1}\lambda_0$ and $\|\bar{g} - \delta\|_{H_{s, \delta + \frac{1}{2}}(\mathbb{R}^3)} + \|\bar{k}\|_{H_{s-1, \delta + \frac{3}{2}}(\mathbb{R}^3)} \leq c(\|g - e\|_{H_{s, \delta + \frac{1}{2}}(E)} + \|k\|_{H_{s-1, \delta + \frac{3}{2}}(E)})$ for some fixed $c > 1$. By Lemma 5.1 and Theorem 6.1 in [10], there exist $C_1 > 0$ and $\theta_1 \in (0, 1)$ depending only on $\bar{\lambda}$ and $\|\bar{g} - \delta\|_{H_{s, \delta + \frac{1}{2}}(\mathbb{R}^3)} + \|\bar{k}\|_{H_{s-1, \delta + \frac{3}{2}}(\mathbb{R}^3)}$, and a unique solution $\bar{\gamma}$ to the reduce EVE (3.3) on Ω_{θ_1} , taking on \mathbb{R}^3 the Cauchy data $(\bar{\phi}, \bar{\psi})$ given by (\bar{g}, \bar{k}) as in (3.5)(3.6) where the Christoffel symbol for \mathbb{R}^3 is $\tilde{\Gamma}|_{\mathbb{R}^3} = 0$, and $\|\bar{\gamma} - \eta\|_{H_{s, \delta}(\Omega_{\theta_1})} < C_1$. Here Ω_{θ_1} is the boost region (2.2) when $\Sigma = R^3$. Furthermore, $\bar{\gamma}$ is regularly hyperbolic¹⁸, with the coefficient of regularly hyperbolicity h_1 depending only on $\bar{\lambda}$ and $\|\bar{g} - \delta\|_{H_{s, \delta + \frac{1}{2}}(\mathbb{R}^3)} + \|\bar{k}\|_{H_{s-1, \delta + \frac{3}{2}}(\mathbb{R}^3)}$.

We claim that there exists a $\lambda_1 > 2$ depending only on h_1 , such that $\bar{\gamma}$ is regularly hyperbolic on V_{θ_1, λ_1} . The first three conditions in Definition 3.2 are naturally satisfied since $\bar{\gamma}$ is regularly hyperbolic in Ω_{θ_1} (See Definition 4.1 in [10]). Condition (4) is true if we take λ_1 large enough depending only on the regularly hyperbolicity h_1 of $\bar{\gamma}$ as discussed in Remark 3.3.

Then we claim that $\bar{\gamma}$ is a solution of (EVE)(1.3) in harmonic gauge inside the causal set V_{θ_1, λ_1} . In fact, since (g, k) is a solution of (VCE)(1.4) on E , the harmonic gauge condition $f^\mu = \Gamma_{\bar{\gamma}}^\mu = 0$ and $\partial_t f^\mu = 0$ on E are satisfied by the choice of initial conditions (3.5)(3.6). Notice that f satisfies a linear equation (3.8), which satisfies the requirement of Theorem 3.6 by argument on page 293 in [10]. Hence the harmonic gauge vector $f = 0$ on V_{θ_1, λ_1} by the estimate (3.19) in Theorem 3.6, hence $\bar{\gamma}$ is a solution of EVE (1.3) on V_{θ_1, λ_1} .

Now denote the restriction $\bar{\gamma}$ to V_{θ_1, λ_1} by γ . We claim that $(V_{\theta_1, \lambda_1}, \gamma)$ is uniquely determined by $(g, k)|_E$ when γ is regularly hyperbolic on V_{θ_1, λ_1} . Suppose γ_1 and γ_2 are two such solutions of reduced EVE (3.3) as above with initial value given by (3.5)(3.6) from vacuum data (g_1, k_1) and (g_2, k_2) respectively. Then $\|\gamma_i - \eta\|_{H_{s, \delta}(V)}$ are uniformly bounded by the corresponding norm of $(g_i - \eta, k_i)$. Now subtract the reduced EVE (3.3) satisfied by γ_1 and γ_2 :

$$\gamma_1^{\alpha\beta} D_\alpha D_\beta (\gamma_1^{\mu\nu} - \gamma_2^{\mu\nu}) - (D^2 \gamma_2)(\gamma_2 - \gamma_1) - (B(\gamma_1, D\gamma_1) - B(\gamma_2, D\gamma_2)) = 0, \quad (3.20)$$

where (see equations (4.4)(4.5) in [10])

$$\begin{aligned} B(\gamma_1, D\gamma_1) - B(\gamma_2, D\gamma_2) &= P(\gamma_1)(D\gamma_1)^2 - P(\gamma_2)(D\gamma_2)^2 \\ &= (P(\gamma_1) - P(\gamma_2))(D\gamma_1)^2 + P(\gamma_2)(D\gamma_1 + D\gamma_2)(D\gamma_1 - D\gamma_2). \end{aligned}$$

Here P is a rational function of γ . Using the multiplication lemma 2.2, $(D\gamma_1)^2$, $P(\gamma_2)(D\gamma_1 + D\gamma_2) \in H_{s-1, \delta+1}(V)$. Using the mean value inequality, and the Soblev embedding lemma 2.2, we have the pointwise estimates:

$$|P(\gamma_1) - P(\gamma_2)| \leq C|\gamma_1 - \gamma_2|,$$

¹⁸Here regularly hyperbolicity is given in Definition 4.1 in [10], which only requires the first three conditions in Definition 3.2.

where C depends only on $\|\gamma_i - \eta\|_{H_{s,\delta}(V)}$, $i = 1, 2$. Now viewing equation (3.20) as a differential equation for $(\gamma_1 - \gamma_2)$, and using the first energy estimate Lemma 5.4 in Appendix 5.2, we have

$$\|\gamma_1 - \gamma_2\|_{H_{1,\delta+\frac{1}{2}}(E_\tau,V)} \leq C\|\gamma_1 - \gamma_2\|_{H_{1,\delta+\frac{1}{2}}(E,V)} \leq C(\|g_1 - g_2\|_{H_{1,\delta+\frac{1}{2}}(E)} + \|k_1 - k_2\|_{H_{0,\delta+\frac{3}{2}}(E)}).$$

Hence the uniqueness is true.

Combing all the above, we get a unique regularly hyperbolic solution γ to the (EVE) under harmonic gauge on V_{θ_1,λ_1} , where θ_1 , λ_1 and $\|\gamma - \eta\|_{H_{s,\delta}(V)}$ ¹⁹ depend only on λ_0 and $\|g - e\|_{H_{s,\delta+\frac{1}{2}}(E)} + \|k\|_{H_{s-1,\delta+\frac{3}{2}}(E)}$.

Now extend Σ_{int} to include annuli $B_r \setminus B_R \subset E_i$, and take the solution γ inside the causal set $(V_{int})_{\theta_0,\lambda_0}$ based on Σ_{int} by Theorem 3.1. We can combine it with all the solutions $(V_{\theta_i,\lambda_i}, \gamma)$ on each end E_i . Now causal uniqueness(See Corollary 4.8 of Appendix 3 in [6]) implies that they coincide in the intersection of $(V_{int})_{\theta_0,\lambda_0}$ and V_{θ_i,λ_i} , since $(V_{int})_{\theta_0,\lambda_0} \cap V_{\theta_i,\lambda_i}$ is a causal set based on $\Sigma_{int} \cap E$ w.r.t. γ by our construction. So by choosing the smallest θ , such that $\Omega_\theta \subset (V_{int})_{\theta_0,\lambda_0} \cup_{i=1}^l V_{\theta_i,\lambda_i}$, we get the conclusion. \square

4 Perturbation method

Here we will apply the Inverse Function Theorem(See [5][20]) to get maximal graphs in the spacetime evolution of given AF vacuum data sets with small trace. Fix a 3-manifold (Σ, e) which is Euclidean at infinity. We always assume $s \in \mathbb{N}$, $s \geq 4$, and $\delta > -2$. Consider the vacuum data sets (Σ, g, k) , with $(g, k) \in \mathcal{VC}_{s+1,\delta+\frac{1}{2}}(\Sigma)$. Let (\mathcal{V}, γ) be the boost evolution of (g, k) given by Theorem 3.7, then we will study the graph of given function u in the spacetime (\mathcal{V}, γ) . We will take (g, k) as parameters, and study the perturbation problem for the mean curvature H_u of this graph. We will show that for appropriately chosen weighted Sobolev spaces, the linearization of H_u with respect to u is invertible in certain sense.

4.1 Differentiability of mean curvature operator

Given a vacuum data set $(g, k) \in \mathcal{VC}_{s+1,\delta+\frac{1}{2}}(\Sigma)$, with $g \geq \lambda e$ for some $\lambda > 0$. By Theorem 3.7, there exists a uniform $\theta \in (0, 1)$ and a uniform $C > 0$, depending only on λ and $\|g - e\|_{H_{s+1,\delta+\frac{1}{2}}(\Sigma)} + \|k\|_{H_{s,\delta+\frac{3}{2}}(\Sigma)}$, and a unique Lorentzian solution γ of the reduced EVE (3.3) on Ω_θ , taking (g, k) as initial data, and $\|\gamma - \tilde{\eta}\|_{H_{s+1,\delta}(\Omega_\theta)} \leq C$. Moreover, from the proof of Theorem 3.7, the regularly hyperbolic coefficient h of γ in each boost end V_{θ_i,λ_i} , and the regularly sliced coefficient²⁰ of γ in $(V_{int})_{\theta_0,\lambda_0}$ are all uniformly bounded by a constant depending only on λ and the norm of (g, k) . Hence the determinant of $\gamma^{\mu\nu}$ is bounded away from 0 by a constant depending only on λ and the norm of (g, k) .

Now let us summarize some properties of metric components of γ .

¹⁹The bound for $(\gamma - \eta)$ also comes directly by Theorem 3.6.

²⁰See the constant N , A and B in Definition 11.8 in page 397 in [6]

Lemma 4.1. *For $s \geq 3$, $\delta > -2$. Given a $(3+1)$ Lorentz metric $\gamma^{\mu\nu}$ of form (2.6) in Ω_θ with $(\gamma - \tilde{\eta})^{\mu\nu} \in H_{s,\delta}(\Omega_\theta)$, if the determinant $\det(\gamma^{\mu\nu}) \leq -\tilde{\lambda}$ for some $\tilde{\lambda} > 0$, then $(\gamma - \tilde{\eta})_{\mu\nu}$ lies in $H_{s,\delta}(\Omega_\theta)$, and in the metric form (2.5)(2.6) of γ , the components $(\alpha^{-2} - 1), (\alpha - 1), \beta^i, \beta_i, g^{ij} - e^{ij}, g_{ij} - e_{ij}$ all lie in $H_{s,\delta}(\Omega_\theta)$. Furthermore, their norms are all bounded by a constant depending only on $\tilde{\lambda}$ and $\|\gamma - \tilde{\eta}\|_{H_{s,\delta}(\Omega_\theta)}$.*

Proof. The inverse matrix $\gamma_{\mu\nu} = \det(\gamma^{\mu\nu}) \text{adj}(\gamma^{\mu\nu})$, where $\text{adj}(\gamma^{\mu\nu})$ is the adjoint matrix of $\gamma^{\mu\nu}$. Since $\det(\gamma^{\mu\nu})$ is bounded away from 0 by $\tilde{\lambda}$, the Banach algebra property (Lemma 2.6) of $H_{s,\delta}(\Omega_\theta)$ implies that $\gamma_{\mu\nu} - \tilde{\eta}_{\mu\nu}$ also lies in $H_{s,\delta}(\Omega_\theta)$, with $\|\gamma_{\mu\nu} - \tilde{\eta}_{\mu\nu}\|_{H_{s,\delta}(\Omega_\theta)}$ bounded by a constant depending only on $\tilde{\lambda}$ and $\|(\gamma - \tilde{\eta})^{\mu\nu}\|_{H_{s,\delta}(\Omega_\theta)}$. From the expression (2.5)(2.6) of γ and the fact that $(\gamma - \tilde{\eta})^{\mu\nu}, (\gamma - \tilde{\eta})_{\mu\nu} \in H_{s,\delta}(\Omega_\theta)$, we know that $(\alpha^2 - 1), (\frac{1}{\alpha^2} - 1), \beta^i, \frac{\beta^i}{\alpha^2}, (g_{ij} - e_{ij}), (g^{ij} - \frac{\beta^i \beta^j}{\alpha^2} - e^{ij}) \in H_{s,\delta}(\Omega_\theta)$ with their norms bounded by $\|(\gamma - \tilde{\eta})\|_{H_{s,\delta}(\Omega_\theta)}$. So α^2 is bounded both from below and above by certain constant. By Taylor's expansion $|\alpha - 1| = |\sqrt{1 + (\alpha^2 - 1)} - 1| \leq C|\alpha^2 - 1|$, hence is L_δ^2 integrable. For higher order derivatives of $(\alpha - 1)$, we can use the multiplication Lemma 2.6 and the bound of $(\alpha^2 - 1)$ to show that $D^\mu(\alpha - 1)$ lies in $L_{s-|\mu|, \delta+|\mu|}(\Omega_\theta)$. So $(\alpha - 1)$ lies in $H_{s,\delta}(\Omega_\theta)$ and has the norm bounded by a constant depending only on $\tilde{\lambda}$ and $\|(\gamma - \tilde{\eta})^{\mu\nu}\|_{H_{s,\delta}(\Omega_\theta)}$. \square

So the metric coefficients of our boost solution γ satisfy that $\{(\alpha - 1), \beta^i, \beta_i, g^{ij} - e^{ij}, g_{ij} - e_{ij}\} \in H_{s+1,\delta}(\Omega_\theta)$ with norms bounded by a constant depending only on the elliptic constant λ of g and $\|g - e\|_{H_{s+1,\delta+\frac{1}{2}}(\Sigma)} + \|k\|_{H_{s,\delta+\frac{3}{2}}(\Sigma)}$. By the Sobolev embedding $H_{s+1,\delta}(\Omega_\theta) \subset C_\kappa^2(\Omega_\theta)$ for some $0 < \kappa < \delta + 2$, all the terms above are uniformly bounded.

Given $s_1 \geq 3$ and $\delta_1 > -2$. Let \mathcal{B}_ρ be a ball of radius ρ containing scalar functions in $H_{s_1+1,\delta_1-\frac{1}{2}}(\Sigma)$ with $\|u\|_{H_{s_1+1,\delta_1-\frac{1}{2}}(\Sigma)} \leq \rho$. We can choose ρ small enough, such that after embedding $\|u\|_{C_\kappa^2(\Sigma)} \leq C\rho \leq \theta/2$ for some $-1 < \kappa < \delta_1 + 1$, and,

$$\text{Condition (A): } \|u(x)\|_{L^\infty} \leq (\theta/2)(\sigma(x))^{-\kappa} < (\theta/2)\sigma(x). \quad (4.1)$$

So $\text{Graph}_u = \{(x, u(x)) : x \in \Sigma\}$ is a submanifold in Ω_θ . Furthermore, $|Du|_e \leq C\rho(\sigma(x))^{-(\kappa+1)}$. As $(\alpha - 1), \beta, (g - e)$ are all uniformly bounded, we can then choose ρ small enough satisfying:

$$\text{Condition (B): } |Du|_e \leq \frac{1}{100}, |\langle \beta, Du \rangle_g| \leq \frac{1}{2}, |U| = \left| \frac{\alpha |Du|_g}{1 + \langle \beta, Du \rangle_g} \right| \leq \frac{1}{2}, \quad (4.2)$$

where U is defined in (2.12). Then Graph_u is spacelike and $\nu = \sqrt{1 - |U|^2}$ is well-defined. So we can study the operator

$$\mathcal{H} : u \rightarrow H_u, \quad (4.3)$$

where H_u is the mean curvature of Graph_u given by (2.16).

Now we will show that composition is continuous as follows,

Lemma 4.2. *Given $s_1 \geq 3$, $\delta_1 > -2$ and $\theta \in (0, 1)$. Consider $\mathcal{B}_\rho \subset H_{s_1+1,\delta_1-\frac{1}{2}}(\Sigma)$ with ρ small enough satisfying Conditions (A) as above for the θ . Then the composition map:*

$$(f, u) \rightarrow \tilde{f} = f(x, u(x) + t),$$

is a continuous map $H_{s',\delta'}(\Omega_\theta) \times \mathcal{B}_\rho \rightarrow H_{s',\delta'}(\Omega_{\theta/2})$, for $s' \leq s_1 + 1$ and $\delta' \in \mathbb{R}$. Furthermore, when restricted to Graph_u ,

$$(f, u) \rightarrow f(x, u(x))$$

is a continuous map $H_{s',\delta'}(\Omega_\theta) \times \mathcal{B}_\rho \rightarrow H_{s'-1,\delta'+\frac{1}{2}}(\Sigma, \Omega_{\theta/2})$.

Proof. Condition(A) (4.1) implies that $|u(x)| \leq (\theta_0/2)\sigma(x)^{-\kappa}$ for some $-1 < \kappa < \delta_1 + 1$, so we can consider a well-defined map $F : \Omega_\theta \rightarrow \Omega_{\frac{3}{2}\theta}$, where $F : (x, t) \rightarrow (x, u(x) + t)$. Then $|DF| = 1$, so F is a diffeomorphism $\Omega_\theta \rightarrow F(\Omega_\theta)$. Furthermore, $(F - id)(x, t) = (0, u(x)) \in H_{s_1+1,\delta_1}(\Omega_\theta)$. Now we can apply lemma 2.7 to the mapping F , so $f \rightarrow \tilde{f} = f \circ F$ is an isomorphism $H_{s',\delta'}(\Omega_\theta) \rightarrow H_{s',\delta'}(F(\Omega_\theta))$. In fact, by the bound of u , we know that $F(\Omega_\theta)$ contains $\Omega_{\theta/2}$, so clearly \tilde{f} lies in $H_{s',\delta'}(\Omega_{\theta/2})$, and we have the continuity for the first factor f . For the second factor u , we only need to show that $u \rightarrow f(x, u(x) + t)$ is continuous $H_{s_1+1,\delta_1-\frac{1}{2}}(\Sigma) \rightarrow L_{\delta'}^2(\Omega_{\theta/2})$ for fixed $f \in L_{\delta'}^2(\Omega_\theta)$. Using multiplication lemma 2.6 recursively to higher derivatives as in the proof of Theorem 2.3 in [10] gives the continuity in $H_{s',\delta'}$. Suppose $u_n \rightarrow u$ in $H_{s_1+1,\delta_1-\frac{1}{2}}$, hence $u_n \rightarrow u$ in C_κ^0 for some $-1 < \kappa < \delta_1 + 1$. To show the $L_{\delta'}^2$ continuity, we can approximate f by compactly supported smooth function g in $L_{\delta'}^2$, then $|f(x, u_n(x) + t) - f(x, u(x) + t)| \leq |f(x, u_n(x) + t) - g(x, u_n(x) + t)| + |g(x, u_n(x) + t) - g(x, u(x) + t)| + |g(x, u(x) + t) - f(x, u(x) + t)|$. The first and second terms can be chosen very small in $L_{\delta'}^2$, and the third one converge to 0 in $L_{\delta'}^2$. So we get the continuity. For the restriction, we can directly apply the restriction lemma 2.8 to \tilde{f} . \square

Moreover, we also have the differentiability w.r.t. u .

Lemma 4.3. *Given $s_1 \geq 3$, $\delta_1 > -2$, $\theta \in (0, 1)$, $\delta' \in \mathbb{R}$ and $f \in H_{s_1+1,\delta'}(\Omega_\theta)$. Consider $\mathcal{B}_\rho \subset H_{s_1+1,\delta_1-\frac{1}{2}}(\Sigma)$ with ρ chosen to satisfy Condition(A) in (4.1) for the θ . Then*

$$\mathcal{F} : u \rightarrow f(x, u(x)), \quad (4.4)$$

is continuous Fréchet differentiable as a map $\mathcal{B}_\rho \rightarrow H_{s_1-1,\delta'+\frac{1}{2}}(\Sigma)$. Furthermore, the Fréchet derivative is given by formal derivatives,

$$D_u \mathcal{F}(v) = \partial_t f(x, u(x)) \cdot v, \quad (4.5)$$

where $v \in H_{s_1+1,\delta_1-\frac{1}{2}}(\Sigma)$.

Proof. Using lemma 4.2, we know that $f(x, u(x) + t)$ lies in $H_{s_1+1,\delta'}(\Omega_{\theta/2})$, and $f(x, u(x)) \in H_{s_1,\delta'+\frac{1}{2}}(\Sigma, \Omega_{\theta/2})$. Hence $\partial_t f(x, t) \in H_{s_1,\delta'+1}(\Omega_{\theta/2})$ and $\partial_t f(x, u(x)) \in H_{s_1-1,\delta'+\frac{3}{2}}(\Sigma)$. To show that \mathcal{F} is Fréchet differentiable (See Definition 1.1.1 in [5]), we can first show Gateaux differentiable (See Definition 1.1.2 in [5]), i.e.

$$\lim_{\tau \rightarrow 0} \frac{\|f(x, u(x) + \tau v(x)) - f(x, u(x)) - \partial_t f(x, u(x))(\tau v(x))\|_{H_{s_1-1,\delta'+\frac{1}{2}}(\Sigma)}}{\tau \|v(x)\|_{H_{s_1+1,\delta_1-\frac{1}{2}}(\Sigma)}} = 0, \quad (4.6)$$

for any $v \in H_{s_1+1, \delta_1-\frac{1}{2}}(\Sigma)$. Using Newton-Leibniz formula,

$$f(x, u(x) + \tau v(x)) - f(x, u(x)) = \left(\int_{s=0}^1 \partial_t f(x, u(x) + s\tau v(x)) ds \right) (\tau v(x)), \quad (4.7)$$

Using the multiplication lemma (2.3) in the case $H_{s_1-1, \delta'+\frac{3}{2}}(\Sigma) \times H_{s_1+1, \delta_1-\frac{1}{2}}(\Sigma) \rightarrow H_{s_1-1, \delta'+\frac{1}{2}}(\Sigma)$, we only need to show,

$$\lim_{\tau \rightarrow 0} \|\partial_t f(x, u(x) + \tau v(x)) - \partial_t f(x, u(x))\|_{H_{s_1-1, \delta'+\frac{3}{2}}(\Sigma)} = 0.$$

This convergence follows from the continuity of $(\partial_t f, u) \rightarrow \partial_t f(x, u(x))$ as a map $H_{s_1, \delta'+1}(\Omega_\theta) \times H_{s_1+1, \delta_1-\frac{1}{2}}(\Sigma) \rightarrow H_{s_1-1, \delta'+\frac{3}{2}}(\Sigma)$ in lemma 4.2. Now the multiplication operator $L_u : v \rightarrow \partial_t f(x, u(x)) \cdot v$ is a bounded linear operator $L(H_{s_1+1, \delta_1-\frac{1}{2}}(\Sigma), H_{s_1-1, \delta'+\frac{1}{2}}(\Sigma))$ with

$$\|L_u\|_{L(H_{s_1+1, \delta_1-\frac{1}{2}}(\Sigma), H_{s_1-1, \delta'+\frac{1}{2}}(\Sigma))} \leq C \|\partial_t f(x, u(x))\|_{H_{s_1-1, \delta'+\frac{3}{2}}(\Sigma)}$$

by inequality (2.3). The operator L_u is also continuous w.r.t u by lemma 4.2, so we know that \mathcal{F} is Fréchet differentiable by Theorem 1.1.3 in [5], and $D_u \mathcal{F}(v) = \partial_t f(x, u(x)) \cdot v$. \square

Now we can prove the differentiability of H_u w.r.t. u .

Proposition 4.4. *For $s \geq 4$, $\delta > -2$. Given a vacuum data $(g, k) \in \mathcal{VC}_{s+1, \delta+\frac{1}{2}}(\Sigma)$ and θ the boost ratio as in the beginning of this section. If $\mathcal{B}_\rho \subset H_{s, \delta-\frac{1}{2}}(\Sigma)$ with ρ satisfying Conditions (A)(B) as in (4.1)(4.2) for the θ , then the mean curvature operator (4.3) $\mathcal{H} : \mathcal{B}_\rho \rightarrow H_{s-2, \delta+\frac{3}{2}}(\Sigma)$ is continuous differentiable w.r.t. u , i.e. $(D_u \mathcal{H}) \in C(\mathcal{B}_\rho, L(H_{s, \delta-\frac{1}{2}}(\Sigma), H_{s-2, \delta+\frac{3}{2}}(\Sigma)))$. Furthermore, $D_u \mathcal{H}$ is given by the formal variational formula.*

Proof. By the choice of ρ , \mathcal{H} is well-defined. Write out the expression for H_u in (2.16) in local coordinates $\{(t, x^i) : i = 1, 2, 3\}$ of Ω_θ as follows:

$$\begin{aligned} H_u &= (g_M)^{ij} \langle \nabla_{\alpha_i} N, \alpha_j \rangle_\gamma = \nu \cdot (g_M)^{ij} \langle \nabla_{\partial_i + u_i \partial_t} (U + T), \partial_j + u_j \partial_t \rangle_\gamma \\ &= \nu \cdot (g_M)^{ij} \{ (\partial_i + u_i \partial_t)(U + T)^\mu \langle \partial_\mu, \partial_j + u_j \partial_t \rangle_\gamma + (U + T)^\mu \langle \nabla_{\partial_i + u_i \partial_t} \partial_\mu, \partial_j + u_j \partial_t \rangle_\gamma \} \\ &= \nu \cdot (\gamma^{ij} + \frac{\nu^2}{\alpha^2} (\beta^i - \alpha U^i)(\beta^j - \alpha U^j)) \{ (\partial_i + u_i \partial_t)(U^\mu + T^\mu) \cdot (\gamma_{\mu j} + u_j \gamma_{\mu t}) \\ &\quad + (U^\mu + T^\mu)(\Gamma_{i\mu, j} + u_i \Gamma_{t\mu, j} + u_j \Gamma_{i\mu, t} + u_i u_j \Gamma_{t\mu, t}) \}, \end{aligned} \quad (4.8)$$

where $\Gamma_{\mu\nu, \sigma}$ is the Christoffel symbol for γ , and all coefficients of γ are evaluated at $(x, u(x))$. Except for the term ν , H_u is an algebraic expression containing two type of terms in (4.8). One type of terms are the composition of the coefficients of $(\gamma - \tilde{\eta})$ and $\partial\gamma$ with $(x, u(x))$, and the other terms contains ∂u and $\partial^2 u$. The only term appears in the denominator is $1 + \langle \beta, Du \rangle_g$, and $|\langle \beta, Du \rangle_g| \leq \frac{1}{2}$ by the choice of ρ as in Condition(B).

Since $(\gamma - \tilde{\eta}) \in H_{s+1, \delta}(\Omega_\theta)$, the composition of the metric coefficients of $(\gamma - \tilde{\eta})$ with $(x, u(x))$, i.e. $\{(\gamma^{\mu\nu} - \tilde{\eta}^{\mu\nu}), (\gamma_{\mu\nu} - \tilde{\eta}_{\mu\nu}), (\alpha - 1), \beta^i, \beta_i, (g^{ij} - e^{ij}), (g_{ij} - e_{ij})\}(x, u(x))$ are

continuous differentiable w.r.t. u as maps $H_{s,\delta-\frac{1}{2}}(\Sigma) \rightarrow H_{s-2,\delta+\frac{1}{2}}(\Sigma)$ by lemma 4.3. Similarly the composition of the coefficients of $\partial\gamma$ with $(x, u(x))$, i.e. $(\partial\gamma)(x, u(x))$ are also continuous differentiable w.r.t. u as maps $H_{s,\delta-\frac{1}{2}}(\Sigma) \rightarrow H_{s-2,\delta+\frac{3}{2}}(\Sigma)$. The terms ∂u and $\partial^2 u$ are trivially continuous differentiable w.r.t. u as maps $H_{s,\delta-\frac{1}{2}}(\Sigma) \rightarrow H_{s-1,\delta+\frac{1}{2}}(\Sigma)$ and $H_{s,\delta-\frac{1}{2}}(\Sigma) \rightarrow H_{s-2,\delta+\frac{3}{2}}(\Sigma)$ respectively. So $U = \frac{\alpha Du}{1+\langle\beta, Du\rangle} \in H_{s-1,\delta+\frac{1}{2}}(\Sigma)$ and is continuous differentiable w.r.t. u , hence is $\nu^2 - 1 = \frac{|U|^2}{1-|U|^2} \in H_{s-1,\delta+\frac{3}{2}}(\Sigma)$, since $|U| \leq \frac{1}{2}$. So by similar argument as that for α in lemma 4.1, $(\nu - 1)$ is also continuous differentiable w.r.t. u as $H_{s,\delta-\frac{1}{2}}(\Sigma) \rightarrow H_{s-2,\delta+\frac{3}{2}}(\Sigma)$. Combing all them together, H_u is continuously differentiable w.r.t. u by the multiplication lemma 2.6. \square

4.2 Linear theory

Given a 3-dimensional manifold (Σ, e) which is Euclidean at infinity. Let us give some results about linear elliptic operators which are asymptotic to the Laplacian Δ_e on (Σ, e) . Such type of elliptic operators have been widely studied in [2][7][10][19].

Let L be an operator on (Σ, e) of form:

$$Lu = \sum_{k=0}^2 a_k \partial^k u,$$

with u and Lu functions on Σ , satisfying:

$$\begin{aligned} \lambda e &\leq a_2 \leq \lambda^{-1} e \text{ as metrics, with } \lambda \text{ the elliptic coefficient;} \\ (a_2 - e) &\in H_{s_0+1,\delta_0}(\Sigma), \quad a_1 \in H_{s_0,\delta_0+1}(\Sigma), \quad a_0 \in H_{s_0-1,\delta_0+2}(\Sigma), \end{aligned} \quad (4.9)$$

where $s_0 \geq 4$, $\delta_0 > -\frac{3}{2}$. We will show that in certain weighted spaces, such L have uniformly bounded inverse on the orthogonal compliment of $\ker(L)$ depending only on the norms of the coefficients. First we have,

Lemma 4.5. *Given $s \leq s_0$, $-\frac{3}{2} < \delta < -\frac{1}{2}$. There exists a constant C and a large $r > R$, depending only on s_0, δ_0 , the elliptic coefficient λ and the norms $\|a_2 - e\|_{H_{s_0,\delta_0}(\Sigma)}$, $\|a_1\|_{H_{s_0-1,\delta_0+1}(\Sigma)}$ and $\|a_0\|_{H_{s_0-2,\delta_0+2}(\Sigma)}$, such that for any $u \in H_{s,\delta-1}(\Sigma)$,*

$$\|u\|_{H_{s,\delta-1}(\Sigma)} \leq C(\|Lu\|_{H_{s-2,\delta+1}(\Sigma)} + \|u\|_{H_{s-2}(\Sigma_{int,2r})}), \quad (4.10)$$

where $\Sigma_{int,2r}$ is the union of Σ_{int} with all the annuli $B_{2r} \setminus B_R$ inside each end Σ_i , and H_{s-2} is the standard L^2 Sobolev space on $\Sigma_{int,2r}$.

Proof. Let $\Sigma = \Sigma_{int} \cup_{i=1}^l E_i$. Given a function $\chi \in C_c^\infty(\mathbb{R}^3 \setminus B_1)$, such that $0 \leq \chi \leq 1$ and $\chi = 1$ on $\mathbb{R}^3 \setminus B_2$. We can find a partition of unity $\{\chi_{i,r}\}_{i=0}^l$ of Σ for $r > R$, with $\chi_{i,r}(x) = \chi(|x|/r)$ for $x \in E_i \cong \mathbb{R}^3 \setminus B_R$, and $\chi_{i,r}(x) = 0$ for $x \in \Sigma \setminus E_i$, and $\chi_{0,r}(x) = 1 - \sum_{i=1}^l \chi_{i,r}(x)$. Then $u = \sum_{i=1}^l u_{i,r}$, with $u_{i,r} = \chi_{i,r} u$. Let us fix an end E_i and $u_{i,r}$ and forget the sub-index i now.

Since $-\frac{3}{2} < \delta < -\frac{1}{2}$ corresponds to non-exceptional value in [2], we can apply Theorem 1.7 in [2] with $p = 2$ here,

$$\begin{aligned} \|u_r\|_{H_{s,\delta-1}(\mathbb{R}^3)} &\leq C_1 \|\Delta u_r\|_{H_{s-2,\delta+1}(\mathbb{R}^3)} \\ &\leq C_1 \{ \|Lu_r\|_{H_{s-2,\delta+1}(E)} + \|(L - \Delta)u_r\|_{H_{s-2,\delta+1}(E)} \}, \end{aligned} \quad (4.11)$$

where Δ is the laplacian operator w.r.t. δ_{ij} and C_1 a uniform constant.

$$\begin{aligned} \|Lu_r\|_{H_{s-2,\delta+1}(E)} &\leq \|\chi_r Lu\|_{H_{s-2,\delta+1}(E)} + \|2a_2^{ij} \partial_i u \partial_j \chi_r + (a_2^{ij} \partial^2 \chi_r + a_1^i \partial \chi_r) u\|_{H_{s-2,\delta+1}(E)} \\ &\leq C_2(r) (\|Lu\|_{H_{s-2,\delta+1}(E)} + \|u\|_{H_{s-1}(A_r)}), \end{aligned} \quad (4.12)$$

with $A_r = B_{2r} \setminus B_r$, and $C_2(r)$ is a constant depending only on r and $\|a_2 - e\|_{H_{s_0,\delta_0}(A_r)}$, $\|a_1\|_{H_{s_0-1,\delta_0+1}(A_r)}$. Since $\delta_0 > -\frac{3}{2}$, we can assume $\delta_0 - \epsilon = \delta_1 > -\frac{3}{2}$ for some $\epsilon > 0$. Using multiplication lemma 2.5,

$$\begin{aligned} \|(L - \Delta)u_r\|_{H_{s-2,\delta+1}(E_r)} &= \|(a_2^{ij} - \delta^{ij}) \partial^2 u_r + a_1^i \partial u_r + a_0 u_r\|_{H_{s-2,\delta+1}(E_r)} \\ &\leq C_3 (\|a_2 - e\|_{H_{s,\delta_1}(E_r)} + \|a_1\|_{H_{s-1,\delta_1+1}(E_r)} + \|a_0\|_{H_{s-2,\delta_1+2}(E_r)}) \|u_r\|_{H_{s,\delta-1}(E_r)}, \end{aligned} \quad (4.13)$$

where $E_r = \mathbb{R}^3 \setminus B_r$ and C_3 a uniform constant. Now $\|a_2 - e\|_{H_{s,\delta_1}(E_r)} + \|a_1\|_{H_{s-1,\delta_1+1}(E_r)} + \|a_0\|_{H_{s-2,\delta_1+2}(E_r)} \leq (\|a_2 - e\|_{H_{s,\delta_0}(E_r)} + \|a_1\|_{H_{s-1,\delta_0+1}(E_r)} + \|a_0\|_{H_{s-2,\delta_0+2}(E_r)}) r^{-\epsilon}$ for r large enough. So we can always choose a $r > R$, depending only on δ_0 and $(\|a_2 - e\|_{H_{s,\delta_0}(E_r)} + \|a_1\|_{H_{s-1,\delta_0+1}(E_r)} + \|a_0\|_{H_{s-2,\delta_0+2}(E_r)})$, such that $\|(L - \Delta)u_r\|_{H_{s-2,\delta+1}(E_r)} \leq \frac{1}{2C_1} \|u_r\|_{H_{s,\delta-1}(E_r)}$. Putting them back to inequality (4.11),

$$\|u_r\|_{H_{s,\delta-1}} \leq C_4 \{ \|Lu\|_{H_{s-2,\delta+1}(E)} + \|u\|_{H_{s-1}(A_r)} \}, \quad (4.14)$$

where C_4 depends only on $C_2(r)$. Using an interpolation inequality (see Lemma 2.2 in [7]) to $\|u\|_{H_{s-1}(A_r)}$, we can get the estimate of (4.10) on each end. Applying the standard L^2 estimates to $u_{0,r}$ on $\Sigma_{int,2r}$ (See Corollary 2.2 on page 547 in [6]),

$$\|u_{0,r}\|_{H_s(\Sigma_{int,2r})} \leq C_5 \{ \|Lu_{0,r}\|_{H_{s-2}(\Sigma_{int,2r})} + \|u_{0,r}\|_{H_{s-2}(\Sigma_{int,2r})} \}, \quad (4.15)$$

where C_5 depends only on s_0 , the elliptic coefficient λ and the norms $\|a_2 - e\|_{H_s(V_{int,2r})}$, $\|a_1\|_{H_{s-1}(V_{int,2r})}$ and $\|a_0\|_{H_{s-2}(V_{int,2r})}$. Combing results on all ends $E_{i,r}$ and $V_{int,2r}$ together, we can get (4.10) with r and constant C satisfying the requirement. \square

Now we can prove a lemma similar to Theorem 1.10 in [2] and Theorem 5.6 in [19].

Lemma 4.6. *Given $s \leq s_0$, $-\frac{3}{2} < \delta < -\frac{1}{2}$, the operator L is a Fredholm operator:*

$$H_{s,\delta-1}(\Sigma) \rightarrow H_{s-2,\delta+1}(\Sigma),$$

i.e. L has finite-dimensional kernel $\ker(L, \delta - 1) = \{v \in H_{s,\delta-1}(\Sigma) : Lv = 0\}$, and finite-dimensional co-kernel $\text{coker}(L, \delta - 1)$.

Proof. From the multiplication lemma 2.5, we know that L is a bounded linear map $H_{s,\delta-1}(\Sigma) \rightarrow H_{s-2,\delta+1}(\Sigma)$. Standard argument using inequality (4.10) as in Theorem 1.10 in [2] shows that $N(L)$ is finite-dimensional and L has close range. So L is semi-Fredholm.

To show that L has finite-dimensional co-kernel, we will borrow the techniques in Theorem 5.6 of [19]. First, inequality (4.13) shows that the operator norm of $(L - \Delta) : H_{s,\delta-1}(E_r) \rightarrow H_{s-2,\delta+1}(E_r)$ is $o(1)$ as $r \rightarrow \infty$. So for large enough r , the fact that Δ is Fredholm by Theorem 1.7 in [2] and that the Fredholm property is open w.r.t operator norms show that $L_i = \Delta + \chi_{i,r}(L - \Delta)$ is Fredholm on \mathbb{R}^3 , where clearly $L_i = L$ on E_{2r} . So there exists a bounded linear operator $S_i : H_{s-2,\delta+1}(\mathbb{R}^3) \rightarrow H_{s,\delta-1}(\mathbb{R}^3)$, such that $L_i S_i = id + K_i$ with K_i a compact operator. Now $L : H_{s,\delta-1}(\Sigma_{int,8r}) \rightarrow H_{s-2,\delta+1}(\Sigma_{int,8r})$ is Fredholm since $\Sigma_{int,8r}$ is compact, so there exists a Fredholm inverse $S_0 : H_{s-2,\delta+1}(\Sigma_{int,8r}) \rightarrow H_{s,\delta-1}(\Sigma_{int,8r})$, such that $LS_0 = id + K_0$, for K_0 compact operator. Define

$$Su = \chi_{0,4r} S_0 u_{0,8r} + \sum_{i=1}^l \chi_{i,2r} S_i u_{i,r}, \quad (4.16)$$

which is a bounded linear operator $H_{s-2,\delta+1}(\Sigma) \rightarrow H_{s,\delta-1}(\Sigma)$. Then a calculation as in (5.6.5) of [19] shows that $LS = id + K$ for K compact operator. So L has finite-dimensional co-kernel. \square

The Fredholm index of L is defined to be:

$$i(L, \delta - 1) = \dim \ker(L, \delta - 1) - \dim \operatorname{coker}(L, \delta - 1).$$

By comparing the index of L to the laplacian Δ_e of e , we can show that L is surjective when $a_0 \leq 0$.

Lemma 4.7. *Given $s \leq s_0$, $-\frac{3}{2} < \delta < -\frac{1}{2}$. Suppose $a_0 \leq 0$, then L is surjective. Furthermore, $\dim \ker(L, \delta - 1) = d_{\delta-1} = \dim \ker(\Delta_e, \delta - 1)$. If we denote $\ker(L, \delta - 1)^\perp$ to be the orthogonal compliment of $\ker(L, \delta - 1)$ w.r.t the $L_{\delta-1}^2$ inner product $\langle \cdot, \cdot \rangle_{L_{\delta-1}^2(\Sigma)}$ as in definition 2.3, then:*

$$L : \ker(L, \delta - 1)^\perp \rightarrow H_{s-2,\delta+1}(\Sigma),$$

is an isomorphism.

Proof. Since L can be joint continuously to Δ_e by $L_t = tL + (1-t)\Delta_e$, we know $i(L, \delta - 1) = i(\Delta_e, \delta - 1)$. Theorem 6.2 in [19] says that Δ_e is surjective when $\delta - 1 < -\frac{1}{2}$. In order to show L is surjective, or equivalently $\dim \operatorname{coker}(L, \delta - 1) = 0$, we only need to show $\dim \ker(L, \delta - 1) \leq \dim \ker(\Delta_e, \delta - 1)$. This comes from the asymptotical expansion given in [2]. For $u \in \ker(L, \delta - 1)$, by Theorem 1.17 in [2], $Lu = 0$ implies that on each end E_i , there exists a harmonic homogenous function h_k of order $k \leq k(\delta)$, where $k(\delta) = \max\{k \in \mathbb{Z} : k \leq -(\delta + \frac{3}{2})\}^{21}$, such that $u = h_k + o(r^{k-\beta})$ for $0 < \beta < \delta + \frac{3}{2}$. In our case, $k(\delta) = 0$. In fact, if $u \neq 0$, there must exist at one end, on which $k \geq 0$. Or the decay implies $u = o(1)$ at infinity

²¹See the definition for $k(\delta)$ in [2]. Their δ is the same as $-(\delta + \frac{3}{2})$ here.

on Σ , so $u = 0$ by maximum principle since $a_0 \leq 0$. So $\dim \ker(L, \delta - 1)$ is less or equal to the number of linearly independent harmonic polynomials of order $\leq k(\delta)$ multiplied with the number of ends. It is easy to see that the basis of $\ker(\Delta_e, \delta - 1)$ is consisted just by functions which have main part the harmonic polynomial on one end, and $O(1/r)$ parts in other ends. So the leading terms shows that $\dim \ker(L, \delta - 1) \leq \dim \ker(\Delta_e, \delta - 1)$. The isomorphism on orthogonal complement is direct when L is surjective. \square

In fact, we can show a uniform norm bound for the inverse of L on $\ker(L, \delta - 1)^\perp$.

Lemma 4.8. *Given $s \leq s_0$, $-\frac{3}{2} < \delta < -\frac{1}{2}$. Suppose $a_0 \leq 0$. Denote the inverse of $L : \ker(L, \delta - 1)^\perp \rightarrow H_{s-2, \delta+1}(\Sigma)$ by L^{-1} , then there exists a constant C depending only on s_0 , δ_0 , the elliptic coefficient λ and the norms $\|a_2 - e\|_{H_{s_0+1, \delta_0}(\Sigma)}$, $\|a_1\|_{H_{s_0, \delta_0+1}(\Sigma)}$ and $\|a_0\|_{H_{s_0-1, \delta_0+2}(\Sigma)}$, such that for any $v \in H_{s-2, \delta+1}(\Sigma)$,*

$$\|L^{-1}v\|_{H_{s, \delta-1}(\Sigma)} \leq C\|v\|_{H_{s-2, \delta+1}(\Sigma)}. \quad (4.17)$$

Proof. We only need to show that for any $u \in \ker(L, \delta - 1)^\perp$,

$$\|u\|_{H_{s, \delta-1}(\Sigma)} \leq C_1\|Lu\|_{H_{s-2, \delta+1}(\Sigma)}$$

for a uniform constant C_1 depending only on s_0 , δ_0 , the elliptic coefficient λ and the norms $\|a_2 - e\|_{H_{s_0+1, \delta_0}(\Sigma)}$, $\|a_1\|_{H_{s_0, \delta_0+1}(\Sigma)}$ and $\|a_0\|_{H_{s_0-1, \delta_0+2}(\Sigma)}$. By contradiction argument, suppose that the statement is wrong, which means that there exists a sequence of operators L_i with $a_{i,0} \leq 0$, uniformly bounded elliptic coefficient $\lambda_i \geq \lambda_0 > 0$ and uniformly bounded coefficients $\|a_{i,2} - e\|_{H_{s_0+1, \delta_0}(\Sigma)}$, $\|a_{i,1}\|_{H_{s_0, \delta_0+1}(\Sigma)}$, $\|a_{i,0}\|_{H_{s_0-1, \delta_0+2}(\Sigma)} \leq C_0$, and a sequence of functions $u_i \in \ker(L_i, \delta - 1)^\perp$, such that $\|u_i\|_{H_{s, \delta-1}(\Sigma)} \geq i\|L_i u_i\|_{H_{s-2, \delta+1}(\Sigma)}$. By re-normalizing, we get a sequence of functions u_i , with $\|u_i\|_{H_{s, \delta-1}(\Sigma)} = 1$, while $\|L_i u_i\|_{H_{s-2, \delta+1}(\Sigma)} \rightarrow 0$. By weak compactness, there exists a subsequence, which we still denote by L_i , such that the coefficients of L_i converges weakly to that of a linear operator L_∞ with $\lambda_0 e \leq a_{\infty,2} \leq \lambda_0^{-1} e$, $a_{\infty,0} \leq 0$ and $\|a_{\infty,2} - e\|_{H_{s_0+1, \delta_0}(\Sigma)}$, $\|a_{\infty,1}\|_{H_{s_0, \delta_0+1}(\Sigma)}$, $\|a_{\infty,0}\|_{H_{s_0-1, \delta_0+2}(\Sigma)} \leq C_0$. Using inequality (4.10), there is a uniform constant C_2 ,

$$\begin{aligned} \|u_i - u_j\|_{H_{s, \delta-1}(\Sigma)} &\leq C_2(\|L_i(u_i - u_j)\|_{H_{s-2, \delta+1}(\Sigma)} + \|u_i - u_j\|_{H_{s-2}(\Sigma_{int, 2r})}) \\ &\leq C_2(\|L_i u_i\|_{H_{s-2, \delta+1}(\Sigma)} + \|(L_i - L_j)u_j\|_{H_{s-2, \delta+1}(\Sigma)} \\ &\quad + \|L_j u_j\|_{H_{s-2, \delta+1}(\Sigma)} + \|u_i - u_j\|_{H_{s-2}(\Sigma_{int, 2r})}). \end{aligned} \quad (4.18)$$

Now $\|(L_i - L_j)u_j\|_{H_{s-2, \delta+1}(\Sigma)} \leq C(\|a_{2,i} - a_{2,j}\|_{H_{s_0, \delta'-1}(\Sigma)} + \|a_{1,i} - a_{1,j}\|_{H_{s_0-1, \delta'}(\Sigma)} + \|a_{0,i} - a_{0,j}\|_{H_{s_0-2, \delta'+1}(\Sigma)})$ for some $\delta_0 > \delta' > -\frac{3}{2}$ by multiplication lemma 2.6. The compact embedding (Lemma 2.1 in [7]) of $H_{s_0+1-i, \delta_0-1+i}(\Sigma) \subset H_{s_0-i, \delta'-1+i}(\Sigma)$ for $i = 0, 1, 2$ imply that $\|(L_i - L_j)u_j\|_{H_{s-2, \delta+1}(\Sigma)} \rightarrow 0$ for a subsequence of $\{L_i\}$. Together with the compactness of $H_{s, \delta-1}(\Sigma) \subset H_{s-2}(\Sigma_{int, 2r})$, there exists a subsequence, which we still denote by u_i , such that u_i converge strongly in $H_{s, \delta-1}(\Sigma)$ to a function u_∞ , with $\|u_\infty\|_{H_{s, \delta-1}(\Sigma)} = 1$. Furthermore we

have that $L_\infty u_\infty = 0$ weakly by the weak convergence, and hence strongly in $H_{s-2,\delta+1}(\Sigma)$ by elliptic regularity.

By Lemma 4.7, we know that $\dim \ker(L_i, \delta - 1) \equiv d_{\delta-1}$. We claim that $\ker(L_i, \delta - 1)$ converge to a $d_{\delta-1}$ dimensional linear subspace of $\ker(L_\infty, \delta - 1)$. Let $\{v_{i,a}\}_{a=1}^{d_{\delta-1}}$ be an $L_{\delta-1}^2$ orthogonal basis for $\ker(L_i, \delta - 1)$, with $\|v_{i,a}\|_{H_{s,\delta-1}(\Sigma)} = 1$. By equation (4.18),

$$\|v_{i,a} - v_{j,a}\|_{H_{s,\delta-1}(\Sigma)} \leq C(\|(L_i - L_j)v_{j,a}\|_{H_{s-2,\delta+1}(\Sigma)} + \|v_{i,a} - v_{j,a}\|_{H_{s-2}(\Sigma_{int,2r})}).$$

Similar argument as above implies that a subsequence of $v_{i,a}$ converge strongly in $H_{s,\delta-1}(\Sigma)$ to some $v_{\infty,a}$. Hence $v_{\infty,a} \in \ker(L_\infty, \delta - 1)$, and $\{v_{\infty,a}\}_{a=1}^{d_{\delta-1}}$ are also orthogonal in $L_{\delta-1}^2$ with $\|v_{\infty,a}\|_{H_{s,\delta-1}(\Sigma)} = 1$. Since L_∞ satisfies all the requirement of Lemma 4.7, $\dim \ker(L_\infty, \delta - 1) = d_{\delta-1}$. Hence the limit of $\ker(L_i, \delta - 1)$ is exactly the entire $\ker(L_\infty, \delta - 1)$. As u_i is perpendicular to $\ker(L_i, \delta - 1)$ in $L_{\delta-1}^2$, passing to the limit, we know that u_∞ is perpendicular to $\ker(L_\infty, \delta - 1)$ in $L_{\delta-1}^2$ too, which is a contradiction to that $\|u_\infty\|_{H_{s,\delta-1}(\Sigma)} = 1$ and $L_\infty u_\infty = 0$. So we finish the proof. \square

4.3 Existence of maximal data

Now let us calculate the linearization of \mathcal{H} with respect to u at $(g, k, 0)$. Fix a vacuum data $(g, k) \in \mathcal{VC}_{s+2,\delta+\frac{1}{2}}(\Sigma)$ with the unique boost solution (\mathcal{V}, γ) given by Theorem 3.7. Recall the form (2.5) of γ in local coordinates (x^i, t) of Ω_θ . According to the initial data equations (3.5)(3.6) for γ , the coefficients restricted to $t = 0$ slice are given by:

$$\alpha|_\Sigma \equiv 1; \quad \beta|_\Sigma \equiv 0. \quad (4.19)$$

In fact, our choice of $\alpha|_\Sigma$ and $\beta|_\Sigma$ implies that $\partial_t|_\Sigma$ is the unit normal of Σ . Now recall the second variational formula for the mean curvature in section 2 of [1]. Let X be a vector field in a neighborhood of Σ with associated flow $\phi_s : \mathcal{V} \rightarrow \mathcal{V}$. Denote $H(s)$ by the mean curvature of $\phi_s(\Sigma)$, then

$$\partial_s(H(s))|_{s=0} = -\Delta_g \langle X, N \rangle + \langle X, N \rangle (|k|_g^2 + Ric_\gamma(N, N)) + \langle X, \nabla_g H \rangle, \quad (4.20)$$

where N is the unit normal of Σ insider \mathcal{V} , and Ric_γ the Ricci curvature of γ . In our case, $Ric_\gamma \equiv 0$ by (1.3) since our (\mathcal{V}, γ) is vacuum, and the unit normal $N = \partial_t$ on Σ . We can choose the vector field to be $X = v\partial_t$, where v is a compactly supported smooth scalar function, so $\langle X, \nabla_g H \rangle = 0$. Then $\partial_s H(s)|_{s=0}$ is the linearization of H w.r.t u , and $\langle X, N \rangle = -v$. Now combining all and using Proposition 4.4, we have,

Lemma 4.9. *Using notations in Proposition 4.4, the Fréchet derivative of $\mathcal{H}(g, k, u)$ with respect to factor u at a vacuum data $(g, k, 0)$ is a linear operator $L_0 : H_{s,\delta-\frac{1}{2}}(\Sigma) \rightarrow H_{s-2,\delta+\frac{3}{2}}(\Sigma)$ given by:*

$$(D_v \mathcal{H})_{(g,k,0)} = L_0 v = (\Delta_g - |k|_g^2)v. \quad (4.21)$$

Now let us focus on the operator L_0 . L_0 is in fact Fredholm and surjective by Lemma 4.6 and Lemma 4.7. By making use the fact that L_0 has finite-dimensional kernel and is surjective, we can get the existence of solutions of $\mathcal{H}(g, k, u) = 0$ for (g, k) with small trace $tr_g k$ by a perturbation method, but no uniqueness due to the existence of non-trivial kernel $\ker(L_0, \delta - \frac{1}{2})$. We will give an existence and uniqueness theorem in the orthogonal complement of the kernel in order to find symmetry preserving solutions in the following section. Let us first give a Quantitative Inverse Function Theorem motivated by [20].

Theorem 4.10. *Let X, Y be Banach spaces, and $U \subset X$ an open set. Suppose $F : U \rightarrow Y$ is a continuous map, and has Fréchet derivative w.r.t x , such that $\frac{\partial F}{\partial x}$ is continuous. For a point $x_0 \in U$, with $F(x_0) = y_0$. Suppose $\frac{\partial F}{\partial x}(x_0) : X \rightarrow Y$ is invertible, and $\|[\frac{\partial F}{\partial x}(x_0)]^{-1}\| \leq C$. Assume that we can find $r_0 > 0$, such that for any $x \in B_{r_0}(x_0) \subset U$,*

$$\|\frac{\partial F}{\partial x}(x) - \frac{\partial F}{\partial x}(x_0)\| \leq \frac{1}{2C}. \quad (4.22)$$

Then for any $y \in Y$ with

$$|y - y_0|_Y < \frac{r_0}{2C},$$

there exist a unique $x \in B_{r_0}(x_0)$, such that $F(x) = y$.

Proof. Fix a $y \in B_{r_0/2C}(y_0) \subset Y$. Let us consider the map $T : B_{r_0}(0) \subset X \rightarrow Y$, defined by

$$T(x) = x - [\frac{\partial F}{\partial x}(x_0)]^{-1}(F(x_0 + x) - y).$$

x is a fixed point if and only if $F(x_0 + x) = y$. So let us use the fixed point theorem to find a fixed point for T on $B_{r_0}(0)$. first, for any $x_1, x_2 \in B_{r_0}(0)$,

$$\begin{aligned} |T(x_1) - T(x_2)|_X &= |(x_1 - x_2) - [\frac{\partial F}{\partial x}(x_0)]^{-1}(F(x_0 + x_1) - F(x_0 + x_2))|_X \\ &\leq \|[\frac{\partial F}{\partial x}(x_0)]^{-1}\| \cdot |\frac{\partial F}{\partial x}(x_0)(x_1 - x_2) - \frac{\partial F}{\partial x}(x_0 + \bar{x})(x_1 - x_2)|_Y \\ &\leq C \|\frac{\partial F}{\partial x}(x_0) - \frac{\partial F}{\partial x}(x_0 + \bar{x})\| \cdot |x_1 - x_2|_X \\ &\leq C \frac{1}{2C} |x_1 - x_2|_X \leq \frac{1}{2} |x_1 - x_2|_X, \end{aligned} \quad (4.23)$$

where we used the mean value theorem in the first “ \leq ”, and condition (4.22) in the third “ \leq ”. So T is a contraction map on $B_{r_0}(0)$. Next, for any $x \in B_{r_0}(0)$, and $|y - y_0|_Y < \frac{r_0}{2C}$,

$$\begin{aligned} |T(x)|_X &\leq \|[\frac{\partial F}{\partial x}(x_0)]^{-1}\| \cdot |\frac{\partial F}{\partial x}(x_0)x - (F(x_0 + x) - F(x_0)) - (y - F(x_0))|_Y \\ &\leq C(|(\frac{\partial F}{\partial x}(x_0) - \frac{\partial F}{\partial x}(x_0 + \bar{x}))x|_X + |y - F(x_0)|_Y) \\ &\leq C(\|\frac{\partial F}{\partial x}(x_0) - \frac{\partial F}{\partial x}(x_0 + \bar{x})\| \cdot |x|_X + |y - F(x_0)|_Y) \\ &< C(\frac{1}{2C}r_0 + \frac{r_0}{2C}) < r_0, \end{aligned} \quad (4.24)$$

where we use condition (4.22) in the last “ $<$ ”. So T maps $B_{r_0}(0)$ to $B_{r_0}(0)$. By applying the Contraction Mapping Theorem to $T : B_{r_0}(0) \rightarrow B_{r_0}(0)$, we finish the proof. \square

Remark 4.11. This can be viewed as a carefully reworking of the proof of Theorem 1.2.1 in [5]. Theorem 3.1 and Theorem 3.2 in [20] also gave a proof about the quantitative inverse function theorem.

Theorem 4.12. *For $s \geq 4$, $-2 < \delta < -1$. Fix a 3-manifold (Σ, e) which is Euclidean at infinity and a $\lambda > 0$. Given a vacuum data $(g, k) \in \mathcal{VC}_{s+2, \delta + \frac{1}{2}}(\Sigma)$, with $g \geq \lambda e$, there is an $\epsilon > 0$ and a $\rho' > 0$ small enough, depending only on the norms $\|g - e\|_{H_{s+2, \delta + \frac{1}{2}}(\Sigma)} + \|k\|_{H_{s+1, \delta + \frac{3}{2}}(\Sigma)}$ and the elliptic constant λ , such that if $\|tr_g k\|_{H_{s-2, \delta + \frac{3}{2}}(\Sigma)} \leq \epsilon$, there exists a unique function $u \in \ker(L_0, \delta - \frac{1}{2})^\perp$ with $\|u\|_{H_{s, \delta - \frac{1}{2}}(\Sigma)} \leq \rho'$, such that u is a solution of the maximal surface equation (1.9).*

Proof. For the given $(g, k) \in \mathcal{VC}_{s+2, \delta + \frac{1}{2}}(\Sigma)$ with θ the boost ratio, we can choose a ρ ball $\mathcal{B}_\rho \subset H_{s, \delta - \frac{1}{2}}(\Sigma)$, with ρ small enough depending only on θ , $\|g - e\|_{H_{s+2, \delta + \frac{1}{2}}(\Sigma)} + \|k\|_{H_{s+1, \delta + \frac{3}{2}}(\Sigma)}$ and λ as in Proposition 4.4. Then the map \mathcal{H} is continuously differentiable w.r.t. u as a map $\mathcal{B}_\rho \cap \ker(L_0, \delta - \frac{1}{2})^\perp \rightarrow H_{s-2, \delta + \frac{3}{2}}(\Sigma)$, and the Fréchet derivative is $(D_u \mathcal{H})_{(g, k, 0)} = L_0 v = (\Delta_g - |k|_g^2)v$ by Lemma 4.9. The coefficient of L_0 satisfies the hypothesis (4.9), where $s_0 = s + 1$ and $\delta_0 = \delta + \frac{1}{2}$ by multiplication lemma 2.5, the elliptic constant equals to λ and $\|a_{0,2} - e\|_{H_{s+2, \delta + \frac{1}{2}}(\Sigma)}$, $\|a_{0,1}\|_{H_{s+1, \delta + \frac{3}{2}}(\Sigma)}$, $\|a_{0,0}\|_{H_{s, \delta + \frac{5}{2}}(\Sigma)}$ are bounded from above by a constant depending only on $\|g - e\|_{H_{s+2, \delta + \frac{1}{2}}(\Sigma)}$ and $\|k\|_{H_{s+1, \delta + \frac{3}{2}}(\Sigma)}$. So $(D_u \mathcal{H})_{(g, k, 0)}$ is an isomorphism $\ker(L_0, \delta - \frac{1}{2})^\perp \rightarrow H_{s-2, \delta + \frac{3}{2}}(\Sigma)$ by Lemma 4.7, since $a_{0,0} = -|k|_g^2 \leq 0$. Now we will show that the conditions in the quantitative Inverse Function Theorem 4.10 are satisfied. By Lemma 4.8, there exists a constant C_0 depending only on λ , $\|a_{0,2} - e\|_{H_{s+2, \delta + \frac{1}{2}}(\Sigma)}$, $\|a_{0,1}\|_{H_{s+1, \delta + \frac{3}{2}}(\Sigma)}$, $\|a_{0,0}\|_{H_{s, \delta + \frac{5}{2}}(\Sigma)}$, such that,

$$\|L_0^{-1}\|_{L(H_{s-2, \delta + \frac{3}{2}}(\Sigma), \ker(L_0, \delta - \frac{1}{2})^\perp)} \leq C_0.$$

Abbreviate the operator norm $\|\cdot\|_{L(H_{s, \delta - \frac{1}{2}}(\Sigma), H_{s-2, \delta + \frac{3}{2}}(\Sigma))} = \|\cdot\|$. Let us study $\|D_u \mathcal{H}(g, k, u) - D_u \mathcal{H}(g, k, 0)\|$. Fix the boost evolution (Ω_θ, γ) of (g, k) , with $\|\gamma - \tilde{\eta}\|_{H_{s+2, \delta}(\Omega_\theta)}$ uniformly bounded by a constant depending only on λ and $\|g - e\|_{H_{s+2, \delta + \frac{1}{2}}(\Sigma)} + \|k\|_{H_{s+1, \delta + \frac{3}{2}}(\Sigma)}$. Then $D_u \mathcal{H}(g, k, u)$ is the first variation $D_u(H_u)$ of H_u w.r.t. u inside (Ω_θ, γ) . From the formula of H_u in (4.8), we know that $D_u(H_u)$ is a second order differential operator. The coefficients of $D_u(H_u)$ are constituted by algebraic expressions of ∂u , $\partial^2 u$ and components of γ , $\partial \gamma$, $\partial^2 \gamma$ evaluated at $(x, u(x))$. Let a be any component of $\partial^2 \gamma$ (similar for γ and $\partial \gamma$), using the Newton-Leibniz formula,

$$a(x, u(x)) - a(x, 0) = \left(\int_{\tau=0}^1 \partial_t a(x, \tau u(x)) d\tau \right) u(x),$$

where $\partial_t a(x, u(x))$ has uniform $H_{s-2, \delta+\frac{7}{2}}(\Sigma)$ norm depending only on $\|\partial^3 \gamma\|_{H_{s-1, \delta+3}(\Omega_\theta)}$ and ρ by Lemma 4.2. So $\|a(x, u(x)) - a(x, 0)\|_{H_{s-2, \delta+\frac{5}{2}}(\Sigma)} \leq C_3 \|u\|_{H_{s, \delta-\frac{1}{2}}(\Sigma)}$ by multiplication lemma 2.6, where C_3 depends only on $\|\gamma - \tilde{\eta}\|_{H_{s+2, \delta}(\Omega_\theta)}$ and ρ . Hence by comparing the coefficients of $D_u \mathcal{H}(g, k, u)$ and $D_u \mathcal{H}(g, k, 0)$, we can choose $\|u\|_{H_{s, \delta-\frac{1}{2}}(\Sigma)} \leq \rho'$ with ρ' small enough, depending only on $\|\gamma - \tilde{\eta}\|_{H_{s+2, \delta}(\Omega_\theta)}$ and C_0 , such that,

$$\|D_u \mathcal{H}(g, k, u) - D_u \mathcal{H}(g, k, 0)\| \leq \frac{1}{2C_0}. \quad (4.25)$$

For the ρ' chosen above, if we take $\epsilon < \frac{\rho'}{2C_0}$, then

$$\|0 - \mathcal{H}(g, k, 0)\|_{H_{s-2, \delta+\frac{3}{2}}(\Sigma)} = \|tr_g k\|_{H_{s-2, \delta+\frac{3}{2}}(\Sigma)} < \frac{\rho'}{2C_0}.$$

Now by the Quantitative Inverse Function Theorem 4.10, if we choose the ϵ and ρ' as above, where ϵ and ρ' depend only on λ , $\|g - e\|_{H_{s+2, \delta+\frac{1}{2}}(\Sigma)}$ and $\|k\|_{H_{s+1, \delta+\frac{3}{2}}(\Sigma)}$, there exists a unique $u \in \mathcal{B}_{\rho'} \cap \ker(L_0, \delta - \frac{1}{2})^\perp$, such that u solves $\mathcal{H}(u) = 0$. \square

4.4 Proof of the main Theorems

Here we will study the properties of the maximal graph gotten above. We will improve the regularity of the solution using a bootstrap argument, and show that the ADM mass of the maximal graph is the same as the given data. Moreover the maximal graph can be chosen to be axisymmetric if (g, k) is axisymmetric, and the angular momentum of the maximal graph is the same as (g, k) .

In Theorem 4.12, the solution u has only s weak derivatives due to the contraction mapping principal. In fact, by exploring the structure of the mean curvature operator (4.8), we can gain more regularity for u .

Lemma 4.13. (Regularity analysis). *In Theorem 4.12, the solution $u \in H_{s+2, \delta-\frac{1}{2}}(\Sigma)$. Denote $M = \text{Graph}_u$, and let g_M be the metric and k_M the second fundamental form induced by $M \subset (\mathcal{V}, \gamma)$, then $(g_M, k_M) \in \mathcal{VC}_{s+1, \delta+\frac{1}{2}}(\Sigma)$.*

Proof. In the local coordinates formula (4.8), we can collect together all the terms containing $\partial_{ij}^2 u$, then the maximal surface equation $\mathcal{H}(u) = 0$ can be rewritten as a linear second order elliptic equation for u with ∂u and u terms as coefficients:

$$(g^M)^{ij}(x, u(x))u_{ij} = f(x),$$

where $f(x)$ is a polynomial of $g^M(x, u(x))$, ∂u , $\gamma(x, u(x))$ and $(\partial \gamma)(x, u(x))$. First the spacelike property of $M = \text{Graph}_u$ implies that $(g^M)^{ij}$ is elliptic. Furthermore, $(g^M)^{ij}(x, u(x)) - e^{ij}(x) = \gamma^{ij} - e^{ij} + \frac{\nu^2}{\alpha^2}(\beta^i - \alpha U^i)(\beta^j - \alpha U^j) \in H_{s-1, \delta+\frac{1}{2}}(\Sigma)$, $f(x) \in H_{s-1, \delta+\frac{3}{2}}(\Sigma)$ by the argument in the proof of Proposition 4.4, Lemma 4.2 and the Banach algebra property in

Lemma 2.6. Since $(g^M)^{ij}$ lie in C^0 and H_{s-1} locally, $u \in (H_{s+1})_{loc}(\Sigma)$ by standard elliptic regularity theory. Furthermore, the linear operator $Lu = (g^M)^{ij} \partial_{ij}^2 u$ satisfies the hypothesis of the weighted elliptic regularity Theorem 6.1 in [7] since $s \geq 4$, hence $u \in H_{s+1, \delta - \frac{1}{2}}(\Sigma)$ by Theorem 6.1 in [7]. Now we can bootstrap this process. In fact, by the composition Lemma 4.2, the right hand side $f(x)$ lies in at most $H_{s, \delta + \frac{3}{2}}(\Sigma)$ since there are $\partial\gamma(x, u(x))$ terms. So bootstrap ends when $u \in H_{s+2, \delta - \frac{1}{2}}(\Sigma)$.

On the graph M , $(g_M)_{ij} = (g_{ij} + \beta_i u_j + \beta_j u_i - (\alpha^2 - \beta^2) u_i u_j)(x, u(x))$ by (2.14),

$$(k_M)_{ij} = \nu \cdot \{ (\partial_i + u_i \partial_t)(U^\mu + T^\mu) \cdot (\gamma_{\mu j} + u_j \gamma_{\mu t}) \\ + (U^\mu + T^\mu)(\Gamma_{i\mu, j} + u_i \Gamma_{t\mu, j} + u_j \Gamma_{i\mu, t} + u_i u_j \Gamma_{t\mu, t}) \}, \quad (4.26)$$

by formula (4.8). So by the proof of Proposition 4.4, $((g_M)_{ij} - e_{ij}) \in H_{s+1, \delta + \frac{1}{2}}(\Sigma)$ and $(k_M)_{ij} \in H_{s, \delta + \frac{3}{2}}(\Sigma)$. \square

In order to define the ADM mass and linear momentum, we need to assume $-\frac{3}{2} < \delta < -1$, then by the embedding lemma 2.5, $(g_M - e) \in C_\beta^{s-1}(\Sigma)$ and $k_M \in C_{\beta+1}^{s-2}(\Sigma)$ for some $\frac{1}{2} < \beta < 1$, which satisfy the conditions (1.5). Similar conditions are also satisfied by $(g - e, k)$. We can defined the ADM mass m , m_M for (g, k) and (g_M, k_M) respectively.

Lemma 4.14. *For $-\frac{3}{2} < \delta < -1$, in Theorem 4.12, $m = m_M$.*

Proof. We will use the multiplication lemma 2.5 frequently when we multiply two Soblev functions. $(g_M)_{ij}(x) - g_{ij}(x, u(x)) = (\beta_i u_j + \beta_j u_i - (\alpha^2 - \beta^2) u_i u_j)(x, u(x))$ by (2.14). Now $\beta(x, u(x)) \in H_{s+1, \delta + \frac{1}{2}}(\Sigma)$ and $\partial u \in H_{s+1, \delta + \frac{1}{2}}(\Sigma)$ imply $(g_M)_{ij}(x) - g_{ij}(x, u(x)) \in H_{s+1, \delta+1}(\Sigma)$.

$$g_{ij}(x, u(x)) - g_{ij}(x) = \left\{ \int_{s=0}^1 \partial_t g_{ij}(x, su(x)) ds \right\} \cdot u(x),$$

which shows $\{g_{ij}(x, u(x)) - g_{ij}(x)\} \in H_{s+1, \delta+1}(\Sigma)$, since $\partial_t g_{ij}(x, su(x)) \in H_{s+1, \delta + \frac{3}{2}}(\Sigma)$ and $u \in H_{s+2, \delta - \frac{1}{2}}(\Sigma)$. Hence $\{(g_M)_{ij}(x) - g_{ij}(x)\} \in H_{s+1, \delta+1}(\Sigma) \subset C_\beta^{s-1}(\Sigma)$, for some $1 < \beta < \delta + \frac{5}{2}$ by the embedding lemma 2.5. By checking the definition (1.6), we know that a error term of decay rate $o(r^{-1})$ will not change the mass, so $m = m_M$. \square

Now we will study the preservation of symmetry by this constructions. We need a lemma about symmetry preserving by the reduced EVE (3.3).

Lemma 4.15. *Given a vacuum data $(g, k) \in \mathcal{VC}_{s+2, \delta + \frac{1}{2}}(\Sigma)$, and (Ω_θ, γ) the boost evolution of (g, k) given by Theorem 3.7. Suppose that both (g, k) and e are symmetric under a Killing vector field ξ on Σ , i.e. (g, k) satisfy (1.7), and $\mathcal{L}_\xi e = 0$, where e is the canonical metric on Σ . Then the parallel translation $\tilde{\xi}$ of ξ into Ω_θ is a Killing vector field of γ .*

Proof. Now let $\phi_s : \Sigma \rightarrow \Sigma$ be the one parameter group of diffeomorphisms corresponding to ξ . Then $(\phi_s)^* g = g$, $(\phi_s)^* k = k$ and $(\phi_s)^* e = e$. Now let us extend ϕ_s to a diffeomorphism $\tilde{\phi}_s : \Omega_\theta \rightarrow \Omega_\theta$ by

$$\tilde{\phi}_s(x, t) = (\phi_s(x), t). \quad (4.27)$$

Then $(\tilde{\phi}_s)^*\tilde{e} = \tilde{e}$ where \tilde{e} is defined by (2.1). By the initial conditions (3.5)(3.6) for γ , we know that $\gamma_s = (\tilde{\phi}_s)^*\gamma$ has the same initial conditions as γ on Σ . If we can show that γ_s also solves the reduced (EVE) (3.3), the uniqueness in Theorem 3.7 implies that $\gamma_s = \gamma$. Since γ_s is Ricci flat, we only need to show that $(\Omega_\theta, \gamma_s)$ is also in a harmonic gauge, or equivalently, $id : (\Omega_\theta, \gamma_s) \rightarrow (\Omega_\theta, \tilde{e})$ is a wave map. By pulling back the wave map equation $\square_{(\gamma, \tilde{e})}id = 0$ by $\tilde{\phi}$, we get $\square_{((\tilde{\phi}_s)^*\gamma, (\tilde{\phi}_s)^*\tilde{e})}id = 0$, which reduces to $\square_{(\gamma_s, \tilde{e})}id = 0$. This means that γ_s is also in a harmonic gauge, hence $\gamma_s = \gamma$. Now the vector field corresponding to $\tilde{\phi}_s$ is clearly the parallel translation of ξ into Ω_θ . \square

Now we can prove the preservation of symmetry for the maximal surface.

Theorem 4.16. *Given $s \geq 4$, $-2 < \delta < -1$. Suppose (Σ, e) is a 3-manifold, which is Euclidean at infinity and axisymmetric in the sense of Definition 1.3. If $(g, k) \in \mathcal{VC}_{s+2, \delta+\frac{1}{2}}^a(\Sigma)$ is axisymmetric, and $\|tr_g k\|_{H_{s-2, \delta+\frac{3}{2}}(\Sigma)} \leq \epsilon$ with ϵ given by Theorem 4.12, then the solution u of the maximal surface equation (1.9) given in Theorem 4.12 can be chosen to be axisymmetric, i.e. $\partial_\varphi u = 0$. Hence (Σ, g_u, k_u) is axisymmetric, and the angular momentum of (g_u, k_u) equals that of (g, k) .*

Proof. By Theorem 4.12, $\mathcal{H}(g, k, u) = 0$ has a unique solution $u \in \mathcal{B}_{\rho'} \cap \ker(L_0, \delta - \frac{1}{2})^\perp$. Let ϕ_s be the diffeomorphism corresponding to the Killing vector field $\xi = \frac{\partial}{\partial \varphi}$ in Definition 1.3, and $\tilde{\phi}_s$ the extension given in (4.27). When (g, k) is also axisymmetric, the boost solution (Ω_θ, γ) is invariant under $\tilde{\phi}_s$ by Lemma 4.15. Now pulling back $\mathcal{H}(g, k, u) = 0$ by $\tilde{\phi}_s$, we can see that ϕ_s^*u is a solution of $\mathcal{H}(\phi_s^*g, \phi_s^*k, \phi_s^*u) = 0$, hence $\mathcal{H}(g, k, \phi_s^*u) = 0$. Since (Σ, e) and (g, k) are all invariant under ϕ_s , $\ker(L_0, \delta - \frac{1}{2})$ and hence $\ker(L_0, \delta - \frac{1}{2})^\perp$ are also invariant under ϕ_s , which means that $(\phi_s)^*u \in \mathcal{B}_{\rho'} \cap \ker(L_0, \delta - \frac{1}{2})^\perp$, then uniqueness in Theorem 4.12 implies that $(\phi_s)^*u = u$. So u is axisymmetric, hence is (g_u, k_u) since γ is also axisymmetric.

For the angular momentum, we have another formular, which is called Komar integral (see section 11.2 in [24] for definition and equivalence with (1.8)),

$$J(S) = \frac{1}{16\pi} \int_S *d\xi, \quad (4.28)$$

where $*$ is the Hodge star operator w.r.t γ , and ξ the killing vector field. Since $*d\xi$ is a close form, we know that $J(S)$ is invariant for any two spacelike close surface S and S' which are homologous equivalent. So (Σ, g, k) and $(Graph_u, g_u, k_u)$ have the same angular momentum. \square

5 Appendix

5.1 Geometry of hypersurface

Here we show the detailed calculation of the mean curvature of a level surface. Part of the results here already appeared in [1]. First let us calculate the future-directed timelike unit

normal vector of Σ_t defined by $T = -\frac{\nabla t}{|\nabla t|}$, which is given by:

$$T = -\alpha \nabla t = -\alpha(\gamma^{tt} \partial_t + \gamma^{ti} \partial_i) = \alpha^{-1}(\partial_t - \beta). \quad (5.1)$$

$Graph_u$ can be viewed as level surface of $(u - t) = 0$, so the unit normal of $Graph_u$ is $N = \frac{\nabla(u-t)}{|\nabla(u-t)|}$. Now

$$\begin{aligned} \nabla u &= \gamma^{tj} u_j \partial_t + \gamma^{ij} u_j \partial_i = \frac{1}{\alpha^2} \langle \beta, Du \rangle \partial_t + Du - \frac{1}{\alpha^2} \langle \beta, Du \rangle \beta^i \partial_i \\ &= \frac{1}{\alpha} \langle \beta, Du \rangle T + Du. \end{aligned} \quad (5.2)$$

So N is calculated as

$$\begin{aligned} \nabla(u - t) &= \frac{1}{\alpha^2} \langle \beta, Du \rangle \partial_t + Du - \frac{1}{\alpha^2} \langle \beta, Du \rangle \beta^i \partial_i + \frac{1}{\alpha^2} \partial_t - \frac{1}{\alpha^2} \beta^i \partial_i \\ &= Du + \frac{1 + \langle \beta, Du \rangle}{\alpha^2} (\partial_t - \beta) \\ &= \alpha^{-1} (1 + \langle \beta, Du \rangle) \left(\frac{\alpha Du}{1 + \langle \beta, Du \rangle} + T \right). \end{aligned} \quad (5.3)$$

Writing $U = \frac{\alpha Du}{1 + \langle \beta, Du \rangle}$, then $N = \frac{U+T}{|U+T|}$, where $|U + T| = (1 - |U|^2)^{1/2}$, so we get equation (2.11).

Denoting $M = Graph_u$, let us calculate the mean curvature. For completeness we give the inverse metric matrix $(g_M)^{-1}$ of g_M in (2.14). First we need to calculate the co-frame of (2.13). Denoting them by $\alpha^i = a_k^i dx^k + a_t^i dt : i, k = 1, 2, 3..$ Then they should satisfy:

$$\alpha^i(\alpha_k) = \delta_k^i, \quad \alpha^i(N) = 0. \quad (5.4)$$

The last equation gives

$$\begin{aligned} (a_k^i dx^k + a_t^i dt)[\alpha(U + T)] &= (a_k^i dx^k + a_t^i dt) \left(\frac{\alpha^2 Du}{1 + \langle \beta, Du \rangle} + (\partial_t - \beta) \right) \\ &= a_k^i \left(\frac{\alpha^2 u^k}{1 + \langle \beta, Du \rangle} - \beta^k \right) + a_t^i = 0. \end{aligned} \quad (5.5)$$

So

$$a_t^i = a_k^i \left(\beta^k - \frac{\alpha^2 Du}{1 + \langle \beta, Du \rangle} \right) = a_k^i (\beta^k - \alpha U^k). \quad (5.6)$$

Putting into the first on in (5.4), we have

$$(a_k^i dx^k + a_l^i (\beta^l - \alpha U^l) dt) (\partial_k + u_k \partial_t) = a_k^i + a_l^i (\beta^l - \alpha U^l) u_k = \delta_k^i. \quad (5.7)$$

Denoting matrix $A = (a_k^i)$, then the above equations change to the matrix equation

$$A \cdot [id + (\beta - \alpha U)(Du)^t] = id. \quad (5.8)$$

Solving the last matrix equation²², we get

$$\begin{aligned} a_k^i &= Id - \frac{(\beta^i - \alpha U^i)u_k}{1 + \langle \beta - \alpha U, Du \rangle} = Id - \frac{(\beta^i - \alpha U^i)u_k}{1 + \langle \beta, Du \rangle - (1 + \langle \beta, Du \rangle)|U|^2} \\ &= Id - \nu^2 \frac{(\beta^i - \alpha U^i)u_k}{1 + \langle \beta, Du \rangle} = Id - \nu^2 (\beta/\alpha - U)^i U_k, \end{aligned} \quad (5.9)$$

where we have used $U = \frac{\alpha Du}{1 + \langle \beta, Du \rangle}$, and $\nu^{-2} = 1 - |U|^2$. Then

$$\begin{aligned} a_t^i &= a_k^i (\beta - \alpha U)^k = (\delta_k^i - \nu^2 \frac{(\beta^i - \alpha U^i)u_k}{1 + \langle \beta, Du \rangle}) (\beta^k - \alpha U^k) \\ &= \beta^i - \nu^2 \frac{(\beta^i - \alpha U^i) \langle \beta, Du \rangle}{1 + \langle \beta, Du \rangle} - \alpha U^i + \nu^2 (\beta^i - \alpha U^i) |U|^2 \\ &= (1 + \nu^2 |U|^2) (\beta^i - \alpha U^i) - \nu^2 \frac{\langle \beta, Du \rangle}{1 + \langle \beta, Du \rangle} (\beta^i - \alpha U^i) \\ &= \nu^2 (\beta^i - \alpha U^i) - \nu^2 \frac{\langle \beta, Du \rangle}{1 + \langle \beta, Du \rangle} (\beta^i - \alpha U^i) \\ &= \frac{\nu^2}{1 + \langle \beta, Du \rangle} (\beta - \alpha U)^i. \end{aligned} \quad (5.10)$$

So the co-frame is given by

$$\alpha^i = (\delta_k^i - \nu^2 (\beta/\alpha - U)^i U_k) dx^k + \frac{\nu^2}{1 + \langle \beta, Du \rangle} (\beta - \alpha U)^i dt. \quad (5.11)$$

²²The inverse of $Id + \mathbf{u}\mathbf{v}^t$ is given by $Id - \frac{\mathbf{u}\mathbf{v}^t}{1 + \mathbf{u} \cdot \mathbf{v}}$

Taking inner product of the co-frame with respect to γ^{-1} , we can calculate g_M^{-1} .

$$\begin{aligned}
& (g_M)^{ij} \\
&= \left\langle (\delta_k^i - \nu^2(\beta/\alpha - U)^i U_k) dx^k + \frac{\nu^2(\beta - \alpha U)^i}{1 + \langle \beta, Du \rangle} dt, (\delta_l^j - \nu^2(\beta/\alpha - U)^j U_l) dx^l + \frac{\nu^2(\beta - \alpha U)^j}{1 + \langle \beta, Du \rangle} dt \right\rangle_\gamma \\
&= (\delta_k^i - \nu^2(\beta/\alpha - U)^i U_k)(\delta_l^j - \nu^2(\beta/\alpha - U)^j U_l)(g^{kl} - \frac{1}{\alpha^2} \beta^k \beta^l) \\
&+ (\delta_k^i - \nu^2(\beta/\alpha - U)^i U_k) \frac{\nu^2(\beta - \alpha U)^j}{1 + \langle \beta, Du \rangle} \frac{\beta^k}{\alpha^2} + (\delta_k^j - \nu^2(\beta/\alpha - U)^j U_k) \frac{\nu^2(\beta - \alpha U)^i}{1 + \langle \beta, Du \rangle} \frac{\beta^k}{\alpha^2} \\
&- \frac{1}{\alpha^2} \frac{\nu^4(\beta - \alpha U)^i(\beta - \alpha U)^j}{(1 + \langle \beta, Du \rangle)^2} \\
&= g^{ij} - \frac{1}{\alpha^2} \beta^i \beta^j - \nu^2(\beta/\alpha - U)^j U^i - \nu^2(\beta/\alpha - U)^i U^j + \frac{\nu^2(\beta - \alpha U)^j \beta^i \langle \beta, Du \rangle}{\alpha^2(1 + \langle \beta, Du \rangle)} \\
&+ \frac{\nu^2(\beta - \alpha U)^i \beta^j \langle \beta, Du \rangle}{\alpha^2(1 + \langle \beta, Du \rangle)} + \nu^4(\beta/\alpha - U)^i(\beta/\alpha - U)^j |U|^2 \\
&- \frac{\nu^4(\beta - \alpha U)^i(\beta - \alpha U)^j}{\alpha^2(1 + \langle \beta, Du \rangle)^2} \langle \beta, Du \rangle^2 + \frac{\nu^2}{\alpha^2} \frac{\beta^i(\beta - \alpha U)^j + \beta^j(\beta - \alpha U)^i}{1 + \langle \beta, Du \rangle} \\
&- 2 \frac{\nu^4(\beta - \alpha U)^i(\beta - \alpha U)^j}{\alpha^2(1 + \langle \beta, Du \rangle)^2} \langle \beta, Du \rangle - \frac{\nu^4(\beta - \alpha U)^i(\beta - \alpha U)^j}{\alpha^2(1 + \langle \beta, Du \rangle)^2} \\
&= g^{ij} - \frac{1}{\alpha^2} \beta^i \beta^j - \nu^2(\beta/\alpha - U)^j U^i - \nu^2(\beta/\alpha - U)^i U^j + \nu^4(\beta/\alpha - U)^i(\beta/\alpha - U)^j |U|^2 \\
&+ \frac{\nu^2}{\alpha^2} [\beta^i(\beta - \alpha U)^j + \beta^j(\beta - \alpha U)^i] - \frac{\nu^4}{\alpha^2} (\beta - \alpha U)^i(\beta - \alpha U)^j \\
&= g^{ij} - \frac{1}{\alpha^2} \beta^i \beta^j + \frac{\nu^2}{\alpha^2} (\beta - \alpha U)^i(\beta - \alpha U)^j.
\end{aligned} \tag{5.12}$$

5.2 Linear boost estimates on an end

Here we will give a detailed version of linear boost estimates on an Euclidean end using method in [8] and [10]. It was also mentioned in [3]. We will mainly give the energy estimates needed to prove Theorem 3.6. For convenience, we sometime abbreviate $V_{\theta, \lambda} = V$ in this section. Given a regularly hyperbolic metric $\gamma^{\mu\nu}$ and a \mathbb{R}^N -valued function u in $V_{\theta, \lambda}$, we can associate it with the *energy-momentum tensor* $T^{\mu\nu 23}$:

$$T^{\mu\nu} = G^{\mu\nu\rho\sigma} D_\rho u \cdot D_\sigma u^{24}, \tag{5.13}$$

where

$$G^{\mu\nu\rho\sigma} = \gamma^{\mu\rho} \gamma^{\nu\sigma} + \gamma^{\mu\sigma} \gamma^{\nu\rho} - \gamma^{\mu\nu} \gamma^{\rho\sigma}.$$

²³See also equation (4.6) in [8].

²⁴Here the inner product of $D_\rho u \cdot D_\sigma u = \sum_{k=1}^N D_\rho u^k D_\sigma u^k$.

Given the unit normal \tilde{n} of $\{E_\tau\}$ defined in (3.12), the *momentum vector field* relative to \tilde{n} is

$$P^\mu = T^{\mu\nu} \tilde{n}_\nu. \quad (5.14)$$

Furthermore, the divergence of P^μ is

$$D_\mu P^\mu = 2(\gamma^{\rho\sigma} \tilde{n}_\rho D_\sigma u) \cdot \gamma^{\mu\nu} D_{\mu\nu}^2 u + Q, \quad (5.15)$$

where

$$Q = \Lambda^{\mu\nu} D_\mu u \cdot D_\nu u, \text{ with } \Lambda^{\mu\nu} = D_\rho(G^{\mu\nu\rho\sigma} \tilde{n}_\sigma).$$

Let $N^{-2} = -\langle D\tau, D\tau \rangle_\gamma$ be the lapse function for τ w.r.t. γ and $n = ND\tau$ the unit co-normal of $\{E_\tau\}$ w.r.t. γ . We introduce an orthonormal frame $\{e_0, e_1, \dots, e_{n-1}\}$ w.r.t. γ , such that e_0 is along the direction of $\tilde{n}^\mu = \gamma^{\mu\nu} \tilde{n}_\nu$, i.e. $e_0 = \frac{N}{|\tilde{n}|_\gamma} \tilde{n}^\mu$, where $(\frac{N}{|\tilde{n}|_\gamma})^{-2} = |\tilde{n}|_\gamma^2$, and e_i perpendicular to \tilde{n}^μ . According to Section 2 in [8], we know that $|\tilde{n}|_\gamma^2 = \gamma^{\mu\nu} \tilde{n}_\mu \tilde{n}_\nu = (\frac{N}{|\tilde{n}|_\gamma})^{-2}$ is bounded from both above and below by some constants depending only on θ and h .

Lemma 5.1. *When γ is regularly hyperbolic, P^μ is past time-like w.r.t γ .*

Proof. $T^{\mu\nu} = 2D^\mu u \cdot D^\nu u - |Du|_\gamma^2 \gamma^{\mu\nu}$, so $P^\mu = T^{\mu\nu} \tilde{n}_\nu = 2D^\mu u \cdot D^\nu u \tilde{n}_\nu - |Du|_\gamma^2 \tilde{n}^\mu$, and

$$\begin{aligned} \gamma_{\mu\nu} P^\mu P^\nu &= 4\gamma_{\mu\nu} (D^\mu u \cdot D^\rho u \tilde{n}_\rho) (D^\nu u \cdot D^\sigma u \tilde{n}_\sigma) - 4|Du|_\gamma^2 (D^\mu u \tilde{n}_\mu \cdot D^\nu u \tilde{n}_\nu) + |Du|_\gamma^4 |\tilde{n}|_\gamma^2 \\ &\leq |Du|_\gamma^4 |\tilde{n}|_\gamma^2 \leq 0. \end{aligned}$$

The first “ \leq ” comes from Cauchy-Schwartz inequality, and the second comes from the fact that \tilde{n} is time-like w.r.t. γ .

Take l^μ as a future like vector field, then in the orthonormal frame $\{e_0, e_1, \dots, e_{n-1}\}$ as above, $l^0 > \sqrt{\sum_{i=1}^{n-1} (l^i)^2}$, and

$$\begin{aligned} \gamma_{\mu\nu} P^\mu l^\nu &= 2[(D_0 u) l^0 + (D_i u) l^i] (D_0 u) (\tilde{N}/N) - [(D_0 u)^2 + \sum (D_i u)^2] (-l_0) (\tilde{N}/N) \\ &= [(D_0 u)^2 + \sum (D_i u)^2] l^0 (\tilde{N}/N) + 2D_i u D_0 u l^i (\tilde{N}/N) \\ &\geq (\tilde{N}/N) [(D_0 u)^2 + \sum (D_i u)^2] (l^0 - \sqrt{\sum (l^i)^2}) \geq 0. \end{aligned}$$

The first “ \geq ” comes from the Cauchy-Schwartz inequality. So it shows that P is past time-like w.r.t. γ . \square

Now we introduce the restriction norm and restriction lemma similar to (2.4) and Lemma 2.8. Given $u \in H_{s,\delta}(V_{\theta,\lambda})$, the restriction norm to hypersurface E_τ is defined as:

$$\|u\|_{H_{s,\delta}(E_\tau, V_{\theta,\lambda})} = (\sum_{k=0}^s \|D_t^k u|_{E_\tau}\|_{H_{s-k,\delta+k}(E)}^2)^{1/2}. \quad (5.16)$$

The following restriction lemma follows similar from Lemma 3.1 in [8]:

Lemma 5.2. (restriction). $\forall \tau \in (-\theta, \theta)$, we have the following continuous inclusion:

$$H_{s+1,\delta}(V_{\theta,\lambda}) \subset H_{s,\delta+\frac{1}{2}}(E_\tau, V_{\theta,\lambda}),$$

for every $s \in \mathbb{N}$ and $\delta \in \mathbb{R}$.

Now we have the first energy estimates.

Lemma 5.3. (First Energy Estimates). Assume that $\gamma^{\mu\nu}$ is regularly hyperbolic, and $(\gamma - \eta) \in C^\infty \cap C^{1,0}(V)$, $a_1 \in C^\infty \cap C^{0,1}(V)$ and $a_0 \in C^\infty \cap C^{0,2}(V)$. For L defined in (3.14), with $a_2 = \gamma Id$, every $u \in C_0^\infty(V)$ satisfies the fundamental energy estimates:

$$\|u\|_{H_{1,\delta+\frac{1}{2}}(E_\tau,V)} \leq c(\|u\|_{H_{1,\delta+\frac{1}{2}}(E,V)} + \|\beta\|_{H_{0,\delta+2}(V)}), \quad (5.17)$$

where $0 \leq \tau \leq \theta$, $\beta = Lu$, and c is a constant depending only on θ , the coefficients h of regular hyperbolicity (3.13) of γ , and $\|D\gamma\|_{C^{0,1}} + \|a_1\|_{C^{0,1}} + \|a_2\|_{C^{0,2}}$.

Proof. Let $\tilde{P}^\mu = \sigma^{2(\delta+\frac{3}{2})} P^\mu$. Multiply (5.15) by $\sigma^{2(\delta+\frac{3}{2})}$, we get:

$$D_\mu \tilde{P}^\mu = 2\sigma^{2(\delta+\frac{3}{2})} (\gamma^{\rho\sigma} \tilde{n}_\rho D_\sigma u) \cdot \gamma^{\mu\nu} D_{\mu\nu}^2 u + \tilde{Q}, \quad (5.18)$$

where

$$\tilde{Q} = \sigma^{2(\delta+\frac{3}{2})} Q',$$

with

$$Q' = Q + 2(\delta + 3/2)x^i / \sigma^2 P^i \simeq (D\gamma * \gamma + \sigma^{-1} \gamma * \gamma) Du * Du.$$

Plug in $Lu = \beta$,

$$D_\mu \tilde{P}^\mu = \sigma^{2(\delta+\frac{3}{2})} [2(\gamma^{\rho\sigma} \tilde{n}_\rho D_\sigma u) \cdot (\beta - a_1 Du - a_0 u) + Q'].$$

Now we integrate on the upper part $V_{\tau,\lambda}^+ = \{x \in V_{\tau,\lambda} : t \geq 0\}$ for $\tau \leq \theta$. Since P is compactly supported, the divergence theorem of $(V_{\tau,\lambda}^+, \eta)$ gives,

$$\begin{aligned} \int_{E_\tau} \tilde{P}^\mu \tilde{n}_\mu d\Sigma_\tau - \int_E \tilde{P}^\mu \tilde{n}_\mu d\Sigma + \int_{L_{\tau,\lambda}^+} \tilde{P}^\mu \tilde{\nu}_\mu d\sigma &= \int_{V_{\tau,\lambda}^+} \tilde{P}^\mu \tilde{n}_\mu dx \\ &= \int_{V_{\tau,\lambda}^+} \sigma^{2(\delta+\frac{3}{2})} [2(\gamma^{\rho\sigma} \tilde{n}_\rho D_\sigma u) \cdot (\beta - a_1 Du - a_0 u) + Q'] dx, \end{aligned} \quad (5.19)$$

where $\tilde{\nu}_\mu$ is the unit outer co-normal of the upper lateral boundary $L_{\tau,\lambda}^+ = L_{\tau,\lambda} \cap V_{\tau,\lambda}^+$ under η , which is future timelike w.r.t. γ by property (4) of the regular hyperbolicity (3.2). Using the fact that P is past time-like (Lemma 5.1), we know that

$$\tilde{P}^\mu \tilde{\nu}_\mu = \sigma^{2(\delta+\frac{3}{2})} P^\mu \tilde{\nu}_\mu \geq 0, \text{ on } L_{\tau,\lambda}^+.$$

Now define:

$$x_1(\tau) = \int_{E_\tau} |\sigma^{\delta+3/2} Du|^2 d\Sigma = \|Du\|_{H_{0,\delta+\frac{3}{2}}(E_\tau,V)}^2. \quad (5.20)$$

Since $\tilde{n}_\mu = \tilde{N}D_\mu\tau = \frac{\tilde{N}}{N}n_\mu$,

$$\begin{aligned} P^\mu \tilde{n}_\mu &= T^{\mu\nu} \tilde{n}_\mu \tilde{n}_\nu = 2(\gamma^{\mu\sigma} D_\sigma u \tilde{n}_\mu)^2 - |Du|_\gamma^2 |\tilde{n}|_\gamma^2 \\ &= \left(\frac{\tilde{N}}{N}\right)^2 (2n^\mu n^\nu + \gamma^{\mu\nu}) D_\mu u D_\nu u. \end{aligned}$$

Using Proposition 2.3 in [8], $\Gamma^{\mu\nu} = 2n^\mu n^\nu + \gamma^{\mu\nu}$ is uniformly elliptic, with the elliptic coefficient depending only on the coefficient of regular hyperbolicity h . Using equation (2.8)(2.13) of [8], $d\Sigma_\tau \simeq cd\Sigma$, with c depending only on θ , so we have:

$$\begin{aligned} \int_{E_\tau} \tilde{P}^\mu \tilde{n}_\mu d\Sigma_\tau &\geq c_1^{-1} x_1(\tau), \\ \int_E \tilde{P}^\mu \tilde{n}_\mu d\Sigma_0 &\leq c_1 x_1(0), \end{aligned}$$

where c_1 is a constant depending only on θ and the regular hyperbolicity coefficient h . Now using Cauchy-Schwartz inequality and the fact $dx = \sigma d\tau d\Sigma$ to the right hand side of (5.19),

$$\begin{aligned} \left| \int_{V_{\tau,\lambda}^+} 2\sigma^{2(\delta+\frac{3}{2})} (\gamma^{\rho\sigma} \tilde{n}_\rho D_\sigma u) \cdot \beta dx \right| &\leq c_1 \int_0^\tau \|Du\|_{H_{0,\delta+\frac{3}{2}}(E_{\tau'},V)} \|\beta\|_{H_{0,\delta+\frac{5}{2}}(E_{\tau'},V)} d\tau'; \\ \left| \int_{V_{\tau,\lambda}^+} 2\sigma^{2(\delta+\frac{3}{2})} (\gamma^{\rho\sigma} \tilde{n}_\rho D_\sigma u) \cdot a_1 Du dx \right| &\leq c_1 \|a_1\|_{C^{0,1}} \int_0^\tau \|Du\|_{H_{0,\delta+\frac{3}{2}}(E_{\tau'},V)}^2 d\tau'; \\ \left| \int_{V_{\tau,\lambda}^+} 2\sigma^{2(\delta+\frac{3}{2})} (\gamma^{\rho\sigma} \tilde{n}_\rho D_\sigma u) \cdot a_0 u dx \right| &\leq c_1 \|a_0\|_{C^{0,2}} \int_0^\tau \|Du\|_{H_{0,\delta+\frac{3}{2}}(E_{\tau'},V)} \|u\|_{H_{0,\delta+\frac{1}{2}}(E_{\tau'},V)} d\tau'; \\ \left| \int_{V_{\tau,\lambda}^+} 2\sigma^{2(\delta+\frac{3}{2})} Q' dx \right| &\leq c_1 (1 + \|D\gamma\|_{C^{0,1}}) \int_0^\tau \|Du\|_{H_{0,\delta+\frac{3}{2}}(E_{\tau'},V)}^2 d\tau', \end{aligned}$$

where c_1 denotes a constant depending only on the regular hyperbolicity coefficient h . Now define:

$$x_0(\tau) = \int_{E_\tau} |\sigma^{\delta+1/2} u|^2 d\Sigma = \|u\|_{H_{0,\delta+\frac{1}{2}}(E_\tau,V)}^2, \quad (5.21)$$

then (5.19) can be changed to

$$x_1(\tau) \leq c_2 \left\{ x_1(0) + \int_0^\tau \|\beta\|_{H_{0,\delta+\frac{5}{2}}(E_{\tau'},V)} x_1(\tau')^{1/2} d\tau' + m_1 \int_0^\tau y_1(\tau') d\tau' \right\}, \quad (5.22)$$

where c_2 is a constant depending only on θ and the regular hyperbolicity coefficient h , and

$$m_1 = \|D\gamma\|_{C^{0,1}} + \|a_1\|_{C^{0,1}} + \|a_0\|_{C^{0,2}}, \quad (5.23)$$

$$y_1(\tau) = x_1(\tau) + x_0(\tau) = \|u\|_{H_{1,\delta+\frac{1}{2}}(E_\tau,V)}^2. \quad (5.24)$$

Using Cauchy-Schwartz inequality,

$$(u(\tau) - u(0))^2 = \left(\int_0^\tau \frac{\partial u}{\partial \tau'} d\tau' \right)^2 \leq \tau \int_0^\tau \left(\frac{\partial u}{\partial \tau'} \right)^2 d\tau'.$$

Consider the projection map $\pi : V_{\theta,\lambda} \rightarrow E$ defined by $\pi(\bar{x}, t) = \bar{x}$, then $E'_\tau = \pi(E_\tau) \subset E'_{\tau'}$ if $\tau' < \tau$, then

$$\int_{E'_\tau} |\sigma^{\delta+1/2}(u(\tau) - u(0))|^2 d\Sigma \leq \tau \int_0^\tau \left\{ \int_{E'_{\tau'}} |\sigma^{\delta+3/2} \frac{\partial u}{\partial t}|^2 d\Sigma \right\} d\tau' \leq \tau \int_0^\tau x_1(\tau') d\tau'.$$

So,

$$x_0(\tau) \leq 2x_0(0) + 2\tau \int_0^\tau x_1(\tau') d\tau'. \quad (5.25)$$

Adding (5.22) and (5.25), we can get the integral inequality,

$$y_1(\tau) \leq c_2 \{y_1(0) + \int_0^\tau \|\beta\|_{H_{0,\delta+\frac{5}{2}}(E_{\tau'},V)} y_1^{1/2}(\tau') d\tau' + m_1 \int_0^\tau y_1(\tau') d\tau'\} \quad (5.26)$$

Using the Gronwall lemma,

$$y_1^{1/2}(\tau) \leq \exp\left(\frac{1}{2}c_2 m_1 \tau\right) \{y_1^{1/2}(0) + \frac{1}{2} \int_0^\tau e^{\frac{1}{2}c_2 m_1 \tau'} c_2 \|\beta\|_{H_{0,\delta+\frac{5}{2}}(E_{\tau'},V)} d\tau'\}. \quad (5.27)$$

Hence we finished the proof by using $y_1^{1/2}(\tau) = \|u\|_{H_{1,\delta+\frac{1}{2}}(E_\tau,V)}$. \square

This result can be weaken to the case of rough coefficients by approximation methods.

Lemma 5.4. *If γ is regularly hyperbolic on V , $(\gamma - \eta) \in C^{1,0}(V)$, $a_1 \in C^{0,1}(V)$ and $a_0 \in C^{0,2}(V)$, then every $u \in H_{2,\delta}(V)$ satisfies the fundamental energy estimates (5.17), with $\beta = Lu$.*

Proof. This comes from an approximation argument exactly the same as Lemma 4.2 in [8]. \square

Using more differentiability of the coefficients, we can improve the energy estimates containing high order derivatives.

Lemma 5.5. (High Order Estimates). *Given $s \leq s'$ with s' defined in (3.15). If γ is regularly hyperbolic, $(\gamma - \eta) \in C^\infty(V)$, $a_1 \in C^\infty(V)$ and $a_0 \in C^\infty(V)$, then every $u \in C_0^\infty(V)$ satisfies the main energy estimates:*

$$\|u\|_{H_{s,\delta+\frac{1}{2}}(E_\tau,V)} \leq c(\|u\|_{H_{s,\delta+\frac{1}{2}}(E,V)} + \|\beta\|_{H_{s-1,\delta+2}(V)}), \quad (5.28)$$

where $0 \leq \tau \leq \theta$, $\beta = Lu$, and c is a constant depending only on θ , the coefficient of regular hyperbolicity h and m (defined in (3.16)).

Proof. Apply D^{i-1} for $2 \leq i \leq s$ to $Lu = \beta$, we can get

$$\gamma^{\mu\nu} D_{\mu\nu}^2 u^{[i-1]} = \beta^{[i-1]}, \quad (5.29)$$

where $u^{[i-1]} = D^{i-1}u$, and

$$\beta^{[i-1]} = D^{i-1}\beta - \sum_{p=1}^{i-1} \binom{i-1}{p} D^p \gamma D^{i+1-p} u - \sum_{p=0}^{i-1} \binom{i-1}{p} (D^p a_1 D^{i-p} u + D^p a_0 D^{i-1-p} u).$$

Now define

$$x_i(\tau) = \int_{E_\tau} |\sigma^{\delta+i+\frac{1}{2}} D^i u|^2 d\Sigma = \|D^i u\|_{H_{0,\delta+i+1/2}(E_\tau,V)}^2, \quad (5.30)$$

and apply (5.22) in Lemma 5.3 to (5.29) with δ replaced by $\delta + i - 1$, then

$$x_i(\tau) \leq c_1 \{x_i(0) + \int_0^\tau \|\beta^{[i-1]}\|_{H_{0,\delta+i+3/2}(E_{\tau'},V)} x_i^{1/2}(\tau') d\tau' + m_1 \int_0^\tau x_i(\tau') d\tau'\},$$

with c_1 depending only on the coefficient of regular hyperbolicity h and m_1 defined in (5.23). Compared to (5.22), we have only $x_i(\tau')$ in the third term since there is no first order term a_0 in (5.29). Now using the multiplication lemma 2.2 and restriction lemma 5.2 in the case

$$H_{s_2-p-1,\delta_2+p+1/2}(E_\tau,V) \times H_{p-1,\delta+i+3/2-p}(E_\tau,V) \rightarrow H_{0,\delta+i+3/2}(E_\tau,V),$$

we get

$$\|D^p \gamma D^{i+1-p} u\|_{H_{0,\delta+i+3/2}(E_\tau,V)} \leq c_3 \|D\gamma\|_{H_{s_2-1,\delta_2+1}(V)} \|Du\|_{H_{i-1,\delta+3/2}(E_\tau,V)},$$

with c_3 a constant depending only on i and δ . Similarly,

$$\|D^p a_1 D^{i-p} u + D^p a_0 D^{i-1-p} u\|_{H_\tau(E_\tau,V)} \leq c_3 (\|a_1\|_{H_{s_1,\delta_1}(V)} + \|a_0\|_{H_{s_0,\delta_0}(V)}) \|u\|_{H_{i,\delta+1/2}(E_\tau,V)}.$$

So

$$\|\beta^{[i-1]}\|_{H_{0,\delta+i+\frac{3}{2}}(E_{\tau'},V)} \leq \|D^{i-1}\beta\|_{H_{0,\delta+i+\frac{3}{2}}(E_{\tau'},V)} + c_4 m \|u\|_{H_{i,\delta+1/2}(E_\tau,V)},$$

where c_4 is a constant depending on s , δ , and m is given by (3.16). Now define:

$$y_i(\tau) = y_1(\tau) + \sum_{j=2}^i x_j(\tau) = \|u\|_{H_{i,\delta+1/2}(E_\tau,V)}^2. \quad (5.31)$$

We have

$$x_i(\tau) \leq c_1 \{x_i(0) + \int_0^\tau \|D^{i-1}\beta\|_{H_{0,\delta+i+\frac{3}{2}}(E_{\tau'},V)} x_i^{1/2}(\tau') d\tau' + c_4(m+m_1) \int_0^\tau y_i(\tau') d\tau'\}.$$

Summing all i from 1, we can get

$$y_i(\tau) \leq c_1 \{y_i(0) + \int_0^\tau \|\beta\|_{H_{i-1,\delta+5/2}(E_{\tau'},V)} y_i^{1/2}(\tau') d\tau' + c_4(m+m_1) \int_0^\tau y_i(\tau') d\tau'\}. \quad (5.32)$$

Using the Gronwall lemma,

$$y_i^{1/2}(\tau) \leq \exp(c_5(m+m_1)\tau) \{y_i^{1/2}(0) + c_1 \int_0^\tau e^{c_5(m+m_1)\tau'} \|\beta\|_{H_{i-1,\delta+5/2}(E_{\tau'},V)} d\tau'\}, \quad (5.33)$$

where $c_5 = \frac{1}{2}c_1c_4$. Hence we finish the proof realizing $m_1 \leq c_6m$ by the imbedding lemma 2.2. \square

Using the equation $Lu = \beta$ and an argument similar to Lemma 4.4 in [8], we can estimate $\|u\|_{H_{s,\delta+1/2}(E,V)}$ by the spatial norms $\|\phi\|_{H_{s,\delta+1/2}(E)}$, $\|\psi\|_{H_{s-1,\delta+3/2}(E)}$ and $\|\beta\|_{H_{s-2,\delta+5/2}(E,V)}$, where $\phi = u|_E$ and $\psi = D_t u|_E$. We need the following technical lemma which says that we can take the division in the Banach algebra $H_{s,\delta}(U)$, when $s > \frac{n}{2}$ and $\delta > -\frac{n}{2}$.

Lemma 5.6. *Given U satisfied the extended cone property, $s > \frac{n}{2}$, $\delta > -\frac{n}{2}$ and a function f , if $(f-1) \in H_{s,\delta}(U)$, and $|f| \geq c > 0$, then $(f^{-1}-1) \in H_{s,\delta}(U)$, furthermore, $\|f^{-1}-1\|_{H_{s,\delta}(U)}$ is bounded by a constant depending only on n , s , δ and $\|f-1\|_{H_{s,\delta}(U)}$.*

Proof. Since $|f| \geq c > 0$, f^{-1} is well defined. Since $f^{-1}-1 = -\frac{f-1}{f}$ and $|f|^{-1} \leq c^{-1}$ uniformly bounded, $(f^{-1}-1) \in H_{0,\delta}(U)$. Now $D^\alpha(f^{-1}-1) = \sum_{\alpha_1+\dots+\alpha_l=\alpha} \frac{D^{\alpha_1}f \dots D^{\alpha_l}f}{f^{|\alpha|+1}}$, where α is multi-indexes, with $1 \leq |\alpha| \leq s$. Since $(f^{|\alpha|})^{-1}$ is uniformly bounded, and using the multiplication Lemma 2.2, $D^{\alpha_1}f \dots D^{\alpha_l}f \in H_{0,\delta+|\alpha|}(U)$, hence $D^\alpha(f^{-1}-1) \in H_{0,\delta+|\alpha|}(U)$. So $(f^{-1}-1) \in H_{s,\delta}(U)$. The norm bounds follows from the norm bounds of each $D^\alpha(f^{-1}-1)$. \square

Lemma 5.7. *Given an operator L defined in (3.14) satisfying Hypothesis (1) and (2), then every $u \in H_{s+1,\delta}(V)$ with $2 \leq s \leq s'$, which solves $Lu = \beta$ satisfies:*

$$\|u\|_{H_{s,\delta+1/2}(E,V)} \leq c(\|\phi\|_{H_{s,\delta+1/2}(E)} + \|\psi\|_{H_{s-1,\delta+3/2}(E)} + \|\beta\|_{H_{s-2,\delta+5/2}(E,V)}), \quad (5.34)$$

where $\phi = u|_E$, $\psi = D_t u|_E$ and c is a constant depending only on s , δ and μ (defined in (3.17)).

Proof. By the restriction Lemma 5.2, $u \in H_{s+1,\delta}(V)$ implies that $\phi \in H_{s,\delta+1/2}(E)$ and $\psi \in H_{s-1,\delta+3/2}(E)$. Now define the following functions on E :

$$\psi^{[p]} = D_t^p u, \quad 0 \leq p \leq s.$$

Since

$$\|u\|_{H_{s,\delta+1/2}(E,V)}^2 = \sum_{p=0}^s \|\psi^{[p]}\|_{H_{s-p,\delta+p+1/2}(E)}^2,$$

we only need to prove that:

$$\|\psi^{[p]}\|_{H_{s-p,\delta+p+1/2}(E)} \leq c_p(\|\phi\|_{H_{s,\delta+1/2}(E)} + \|\psi\|_{H_{s-1,\delta+3/2}(E)} + \|\beta\|_{H_{s-2,\delta+5/2}(E,V)}).$$

It is true for $p = 0, 1$. Let us use a reduction argument to prove this for all $p \leq s$. Suppose it is true for $0 \leq q \leq p-1$. Take D_t^{p-2} to the equation $Lu = \beta$, and move all the terms containing t -derivatives of u of order less than p , i.e. $D_t^q u$ with $q < p$, to the right hand side, then we get

$$\begin{aligned} \gamma^{00}\psi^{[p]} &= D_t^{p-2}\beta - \sum_{q=0}^{p-3} \binom{p-2}{q} (D_t^{p-2-q}\gamma^{00})\psi^{[q+2]} - \sum_{q=0}^{p-2} \binom{p-2}{q} \{2(D_t^{p-2-q}\gamma^{0i})D_i\psi^{[q+1]} \\ &\quad + (D_t^{p-2-q}\gamma^{ij})D_iD_j\psi^{[q]} + (D_t^{p-2-q}a_1^0)\psi^{[q+1]} + (D_t^{p-2-q}a_1^i)D_i\psi^{[q]} + (D_t^{p-2-q}a_0)\psi^{[q]}\}. \end{aligned} \quad (5.35)$$

Using the multiplication Lemma 2.2 and Hypothesis (1) in the case:

$$H_{s_2-1-(p-2-q), \delta_2+1/2+(p-2-q)}(E) \times H_{s-(q+2), \delta+1/2+(q+2)}(E) \rightarrow H_{s-p, \delta+1/2+p}(E),$$

we can estimate

$$\|(D_t^{p-2-q} \gamma^{0i}) D_i \psi^{[q+1]}\|_{H_{s-p, \delta+1/2+p}(E)} \leq c_3 \|\gamma - \eta\|_{H_{s_2-1, \delta_2+1/2}(E, V)} \|\psi^{[q+1]}\|_{H_{s-(q+2), \delta+1/2+(q+2)}(E)},$$

where c_3 is a constant depending only on s and δ . Now using similar arguments to evaluate the $H_{s-p, \delta+1/2+p}(E)$ norm of other terms in (5.35), together with our inductive hypothesis, we can get

$$\begin{aligned} \|\gamma^{00} \psi^{[p]}\|_{H_{s-p, \delta+1/2+p}(E)} &\leq \|D_t^{p-2} \beta\|_{H_{s-p, \delta+1/2+p}(E)} + c_4 \mu \sum_{q=0}^{p-1} \|\psi^{[q]}\|_{H_{s-q, \delta+1/2+q}(E)} \\ &\leq c'_p (\|\phi\|_{H_{s, \delta+1/2}(E)} + \|\psi\|_{H_{s-1, \delta+3/2}(E)} + \|\beta\|_{H_{s-2, \delta+5/2}(E, V)}), \end{aligned} \quad (5.36)$$

where μ is defined in (3.17), c_4 is a constant depending only on s , p and δ , while c'_p a constant depending only on μ , s , p and δ .

Here

$$\gamma^{00}|_E = (\gamma^{\mu\nu} D_\mu t D_\nu t)|_{t=0} = \sigma^2 (\gamma^{\mu\nu} D_\mu \tau D_\nu \tau)|_{t=0} = -N^{-2} \sigma^2 \leq -c < 0,$$

where $c > 0$ is a constant depending only on θ and h according to Section 2 in [8]. Now $(\gamma - \eta) \in H_{s_2, \delta_2}(V)$ implies that $(\gamma^{00} + 1)|_E \in H_{s_2-1, \delta_2+1/2}(E)$, hence $((\gamma^{00})^{-1} + 1) \in H_{s_2-1, \delta_2+1/2}(E)$ by Lemma 5.6, and furthermore $\|(\gamma^{00})^{-1} + 1\|_{H_{s_2-1, \delta_2+1/2}(E)}$ is bounded by a constant depending only on n , s_2 , δ_2 and $\|\gamma^{00} + 1\|_{H_{s_2-1, \delta_2+1/2}(E)}$. Now multiply $\gamma^{00} \psi^{[p]}$ by $(\gamma^{00})^{-1}$, and use equation (5.36) and the multiplication Lemma 2.2, then we finish the proof. \square

By combining all the above estimates, we can get the energy estimates in Theorem 3.6.

Theorem 5.8. *Given L a differential operator defined by (3.14) in $V_{\theta, \lambda}$, satisfying hypotheses (1) and (2). Let $\beta \in H_{s-1, \delta+2}(V_{\theta, \lambda})$, $\phi \in H_{s, \delta+\frac{1}{2}}(E)$ and $\psi \in H_{s-1, \delta+\frac{3}{2}}(E)$, with $2 \leq s \leq s'$, $\delta \in \mathbb{R}$. Then every $u \in H_{s+1, \delta}(V)$, which solves $Lu = \beta$, with $u|_E = \phi$, $D_t u|_E = \psi$ satisfies the estimates (3.19).*

Proof. First we can plug in (5.34) to (5.28). Then it follows from an approximation argument similar to the proof of Lemma 4.5 in [8] and an integration of (5.28) w.r.t. τ on $[-\theta, \theta]$. \square

References

- [1] R. Bartnik, Existence of Maximal Surfaces in Asymptotically Flat Spacetimes. Commun. Math. Phys. 94, 155-175(1984).
- [2] R. Bartnik, The Mass of an Asymptotically Flat Manifold. Comm. Pure Appl. Math. 39 661-693(1986).

- [3] R. Bartnik, P. T. Chruściel and N. O'Murchadha, On Maximal Surfaces in Asymptotically Flat Space-Times. *Commun. Math. Phys.* 130, 95-109(1990).
- [4] R. Bartnik and J. Isenberg, The Constraint Equations. Arxiv preprint gr-qc/0405092.
- [5] K. Chang, *Methods in Nonlinear Analysis*(Springer Monographs in Mathematics). Springer, 2005.
- [6] Y. Choquet-bruhat, *General Relativity and the Einstein Equations*. Oxford Science Publications, 2009.
- [7] Y. Choquet-bruhat and D. Christodoulou, Elliptic Systems in $H_{s,\delta}$ Spaces on Manifolds Which Are Euclidean at Infinity. *Acta Math.* 146, 129-150(1981).
- [8] D. Christodoulou, The Boost Problem for Weakly Coupled Quasilinear Hyperbolic Systems of The Second Order. *J. Math. Pur. Appl.* 60, 99-130(1981).
- [9] D. Christodoulou, *Mathematical Problems of General Relativity Theory*. Zurich Lectures in Advanced Mathematics. 2008.
- [10] D. Christodoulou and N. O'Murchadha, The Boost Problem in General Relativity. *Commun. Math. Phys.* 80, 271-300(1981).
- [11] P. T. Chruściel, Mass and Angular-Momentum Inequalities for Axi-Symmetric Initial Data Sets. I. Positivity of Mass, *Ann. Phys.* 323 (2008)2566-2590.
- [12] P. T. Chruściel, Y. Li and G. Weinstein, Mass and Angular-Momentum Inequalities for Axi-Symmetric Initial Data Sets. II. Angular Momentum. *Ann. Phys.* 323 (2008)2591-2613.
- [13] J. Corvino and R. Schoen, On the Asymptotics for the Vacuum Einstein Constraint Equations. *J. Differential Geom.* 73 (2006)185-217.
- [14] S. Dain, Angular Momentum-Mass Inequality for Axisymmetric Black Holes. *Phys. Rev. Lett.* 96(2006) 101101.
- [15] S. Dain, A Variational Principle for Stationary, Axisymmetric Solutions of Einstein's Equations. *Class. Quant. Grav.* 23 (2006)6857-6871.
- [16] S. Dain, Proof of the Angular Momentum-Mass Inequality for Axisymmetric Black Hole. *J. Differential Geom.* 79 (2008)33-67.
- [17] D. Gilbarg and N. Trudinger, *Elliptic Partial Differential Equations of Second Order*. 3rd Edition, Springer. 1998.
- [18] L. Huang, R. Schoen, M. Wang, Specifying Angular Momentum and Center of Mass for Vacuum Initial Data Sets. *Commun. Math. Phys.* 306, 785-803(2011).
- [19] R. Lockhart, Fredholm Properties of A Class of Elliptic Operators on Non-Compact Manifolds. *Duke Math. J.* 48, (1981)289-312.
- [20] W. Ruan, Generalized Special Lagrangian Torus Fibration for Calabi-Yau Hypersurfaces in Toric Varieties I. arXiv:math.DG 0303114.
- [21] R. Schoen and S. T. Yau, On the Proof of the Positive Mass Conjecture in General Relativity. *Comm. Math. Phys.* 65 (1979)45-76.
- [22] R. Schoen and S. T. Yau, Proof of the Positive Mass Theorem, II, *Comm. Math. Phys.* 79 (1981)231-260.
- [23] R. Schoen and X. Zhou, Convexity of Reduced Energy and Mass Angular Momentum Inequalities. arXiv:math.DG 1209.0019.
- [24] R. Wald, *General Relativity*. The University of Chicago Press, Chicago, 1984.
- [25] E. Witten, A New Proof of the Positive Energy Theorem, *Commun. Math. Phys.* 80 (1981) 381-402.

Department of Mathematics, Stanford University

Stanford, CA 94305

E-mail: xzhou08@math.stanford.edu