

Duality and singular continuous spectrum in the almost Mathieu equation

by

A. Y. GORDON

*University of North Carolina
Charlotte, NC, U.S.A.*

S. JITOMIRSKAYA

*University of California
Irvine, CA, U.S.A.*

Y. LAST

and

B. SIMON

*California Institute of Technology
Pasadena, CA, U.S.A.*

*California Institute of Technology
Pasadena, CA, U.S.A.*

1. Introduction

Our main goal in this paper is to study the almost Mathieu operator on $l^2(\mathbf{Z})$ defined by

$$(h_{\lambda,\alpha,\theta}u)(n) = u(n+1) + u(n-1) + \lambda \cos(\pi\alpha n + \theta)u(n). \quad (1.1)$$

Our results in §2 on measurability of (normalizable) eigenfunctions may be of broader applicability. For background on (1.1), see [19], [23].

Our main result here concerns (1.1) at the self-dual point $\lambda=2$.

THEOREM 1. *Let α be an irrational so that there exist $q_n \rightarrow \infty$ and p_n in \mathbf{Z} with*

$$q_n^2 \left| \alpha - \frac{p_n}{q_n} \right| \rightarrow 0 \quad (1.2)$$

as $n \rightarrow \infty$. Then for a.e. θ , $h_{\lambda=2,\alpha,\theta}$ has purely singular continuous spectrum.

Remarks. (1) (1.2) is used because for such α , Last [22] has proven that the spectrum, $\sigma_{\lambda,\alpha}$, of $h_{\lambda,\alpha,\theta}$ (which is θ -independent [5]) has $|\sigma_{2,\alpha}|=0$ (where $|\cdot|$ denotes Lebesgue measure). Our proof is such that for any other α with $|\sigma_{\lambda=2,\alpha}|=0$ (presumably all irrational α), one has that $h_{\lambda=2,\alpha,\theta}$ has purely singular continuous spectrum for a.e. θ .

The first two authors are also affiliated with the Institute of Earthquake Prediction Theory and Mathematical Geophysics, Moscow, Russia. This material is based upon work supported by the National Science Foundation under Grants DMS-9208029, DMS-9501265 and DMS-9401491. The Government has certain rights in this material.

Prior to [22], Helffer–Sjöstrand [17] have shown $|\sigma_{2,\alpha}|=0$ for a set of α 's having all quotients of their continued fraction expansion sufficiently large. While this set is nowhere dense and of zero Lebesgue measure, it includes some α 's for which (1.2) does not hold.

(2) The set of α 's for which (1.2) holds is a dense G_δ whose complement has Lebesgue measure 0, that is, (1.2) is generic in both Baire and Lebesgue sense. Although we do not describe the a.e. θ explicitly, the set with singular continuous spectrum contains a dense G_δ [20], so it is also generic in both Baire and Lebesgue sense.

(3) Delyon [11] proved that $h_{\lambda=2,\alpha,\theta}$ has no eigenfunctions belonging to l^1 (for any α and θ). More recently, Chojnacki [8] has proven that $h_{\lambda=2,\alpha,\theta}$ must have at least some continuous spectrum (for all α 's and a.e. θ). His result does not contradict, however, the possibility of mixed (overlapping continuous and p.p.) spectrum. We note that the absence of absolutely continuous spectrum is obvious whenever the spectrum has zero Lebesgue measure.

While Theorem 1 is our main result, we also prove

THEOREM 2. *If α is irrational and λ is such that $h_{4/\lambda,\alpha,\theta}$ has only p.p. spectrum for a.e. θ , then $h_{\lambda,\alpha,\theta}$ has purely a.c. spectrum for a.e. θ .*

Remarks. (1) It is known that if α has good Diophantine properties, then for $\lambda < \frac{4}{15}$, $h_{4/\lambda,\alpha,\theta}$ has only p.p. spectrum [18] (see [27], [14], [16] for earlier results). For such α, λ , we conclude purely a.c. spectrum of $h_{\lambda,\alpha,\theta}$ for a.e. θ .

(2) Existence of some a.c. spectrum for small λ and Diophantine α has been proven by Bellissard, Lima and Testard [6], who applied ideas earlier developed by Dinaburg–Sinai [12] and Belokolos [7]. Such existence (but not necessarily purely a.c. spectrum) is now known for all $\lambda < 2$ and all α, θ [15], [21]. Purely a.c. spectrum for (unspecified) small λ and Diophantine α has been proven by Chulaevsky and Delyon [9] using duality. Their proof uses detailed information from Sinai's proof of localization [27].

We also provide a new proof of

THEOREM 3. *If α is irrational and $\lambda < 2$, then for a.e. θ , $h_{\lambda,\alpha,\theta}$ has no point spectrum.*

Remark. Delyon [11] has proven that there is no point spectrum for all θ , which is strictly stronger. Moreover, his proof is much simpler. Our proof has a certain methodological advantage in that we don't explicitly use the positivity of the Lyapunov exponent for the dual model.

Our proof of Theorems 1–3 depends on a precise version of Aubry duality [2], [3]. Recall that one way to understand duality is to note the following: Suppose that a_n

solves

$$a_{n+1} + a_{n-1} + \lambda \cos(\pi\alpha n + \theta) a_n = E a_n \quad (1.3)$$

with $a_n \in l^1$. Define

$$\varphi(x) = \sum a_n e^{i(\pi\alpha n + \theta)x}, \quad (1.4)$$

which is continuous on \mathbf{R} with

$$\varphi\left(x + \frac{2}{\alpha}\right) = e^{2i\theta/\alpha} \varphi(x). \quad (1.5)$$

For any η , the sequence

$$u(n) = \varphi\left(n + \frac{\eta}{\pi\alpha}\right) \quad (1.6)$$

is seen to obey

$$u(n+1) + u(n-1) + \frac{4}{\lambda} \cos(\pi\alpha n + \eta) u(n) = \frac{2E}{\lambda} u(n) \quad (1.7)$$

by manipulating (1.4). Thus, nice enough normalizable eigenfunctions at (λ, E) yield Bloch waves at $(4/\lambda, 2E/\lambda)$ for h and, conversely, nice enough Bloch waves (regularity of φ implies decay of the Fourier coefficients a_n in (1.4)) yield normalizable eigenfunctions.

If we slough over what “nice enough” means, we have naive duality:

(D1) point spectrum at $\lambda \Rightarrow$ a.c. spectrum at $4/\lambda$,

(D2) a.c. spectrum at $\lambda \Rightarrow$ point spectrum at $4/\lambda$,

(D3) s.c. spectrum at $\lambda \Rightarrow$ s.c. spectrum at $4/\lambda$,

where the last statement follows from the first two.

The surprise in Last [21] is that this naive expectation is false. There exist α (Liouville numbers) for which the spectrum for $\lambda > 2$ is purely singular continuous, but the spectrum for $\lambda < 2$ has an a.c. component. Thus, (D2) need not be true. In a sense, the main result of the paper is that (D1) is still true. More explicitly, we show that the dual of point spectrum is a.c. spectrum, in the sense that some p.p. spectrum for λ implies some a.c. spectrum for $4/\lambda$, and only p.p. spectrum for λ implies only a.c. spectrum for $4/\lambda$. This strengthens Chojnacki's result [8], which shows (in a more general context, though) that the dual of point spectrum is continuous spectrum (but not necessarily a.c. spectrum).

Thus, there is a kind of more precise duality for the almost Mathieu operator:

(D1') point spectrum at $\lambda \Rightarrow$ a.c. spectrum at $4/\lambda$,

(D2') a.c. spectrum at $\lambda \Rightarrow$ point or s.c. spectrum at $4/\lambda$,

(D3') s.c. spectrum at $\lambda \Rightarrow$ a.c. spectrum (or s.c. spectrum) at $4/\lambda$.

It is unclear if there is any s.c. spectrum for $\lambda < 2$.

Note that while we prove (D1') (and thus (D3')) we do not prove (D2'). It follows, of course, from the known results for the almost Mathieu operator, but we want to point out that this implication fails in certain more general contexts [25].

In §2, we prove that if there is point spectrum, one can always choose the eigenvalues and eigenvectors to be measurable in θ . That allows us to represent the set of all eigenvalues of $h_{\lambda, \alpha, \theta}$ as a union of values of a measurable multivalued function along the trajectory of the rotation by α —an object first introduced by Sinai [27].

In §3, we use this representation and analyze duality to show that point spectrum implies that there are spectral measures for the dual problem (at coupling $4/\lambda$) so that $d\mu_\theta$ is independent of θ . Here we use some important ingredients from [9] and [25].

In §4, we show that results of Deift–Simon [10] imply that the singular components of the spectral measures $d\mu_\theta$ and $d\mu_{\theta'}$ are a.e. disjoint. Thus, the θ -independence of §3 implies that $d\mu_\theta$ is purely absolutely continuous. We make this precise and prove Theorems 1–3 in §5.

A.G. is grateful for the hospitality of the Division of Physics, Mathematics and Astronomy of Caltech and the Department of Mathematics at the University of California at Irvine. S.J. and Y.L. would like to thank J. Avron for the hospitality of the Institute for Theoretical Physics at the Technion where parts of this work were done.

2. Measurability of eigenfunctions

To emphasize that “measurability” here means in Borel sense rather than up to sets of measure zero in the completed measure, we will initially discuss a setup with no measure! At the end of this section, we will link this to the almost periodic situation that is the main focus of this paper.

Let $A < \infty$ be positive and fixed. Let $\Omega = [-A, A]^{\mathbf{Z}}$, that is, $\omega \in \Omega$ is a sequence $\{\omega_n\}_{n=-\infty}^{\infty}$ with $|\omega_n| \leq A$. Ω is a separable compact metric space with Baire sets = Borel sets. We call these sets “measurable”. For each $\omega \in \Omega$, define a self-adjoint operator on $l^2(\mathbf{Z})$ by

$$(h_\omega u)(n) = u(n+1) + u(n-1) + \omega_n u(n).$$

A critical fact we will use below is that (normalizable) eigenvalues are always simple.

Given any normalized eigenvector u for h_ω , we define $j(u)$ to be the leftmost maximum for $|u|$, that is, that j with

$$|u(j)| \begin{cases} \geq |u(k)| & \text{for all } k, \\ > |u(k)| & \text{for all } k < j. \end{cases}$$

We will always fix u by requiring $u(j) > 0$. Since $\sum |u(k)|^2 = 1$, we have $u(k) \rightarrow 0$ as $|k| \rightarrow \infty$, so it has a leftmost maximum. If j is the leftmost maximum for u , we say that u is attached to j .

Let $\{u_n\}$ be the collection of eigenvectors of h_ω . Define $N_j(\omega)$ to be the number of eigenvectors of h_ω attached to j . $N_j(\omega)$ can be finite or infinite. One of our goals below will be to prove

THEOREM 2.1. $N_j(\omega)$ is a measurable function on Ω .

Let $\Omega_{j,k} = \{\omega \mid N_j(\omega) \geq k\}$ for $k = 1, 2, \dots$. On $\Omega_{j,1}$, define $u_1(n; \omega, j)$ to be the eigenfunction attached to j with maximal value of $u_1(j)$. Let $e_1(\omega, j)$ be its eigenvalue. If there are multiple eigenfunctions attached to j with the same value of $u_1(j)$, pick the one with the largest energy. (Note that the Wronskian conservation implies that two eigenfunctions cannot have the same eigenvalue.) Since $\sum_n |u_n(j)|^2 \leq 1$ (by Parseval's inequality for δ_j), there are only finitely many u 's with this maximum value of $u(j)$ and so we can pick the one with largest e .

Similarly, on $\Omega_{j,2}$ we can define $u_2(n; \omega, j)$ by picking the attached eigenvector with second largest $u(j; \omega, j)$, again breaking ties by choosing the largest energy. In this way, we define $u_l(n; \omega, j)$ and $e_l(\omega, j)$ on $\Omega_{j,l}$ so that

- (1) $\{u_l(\cdot; \omega, j)\}_{l=1}^{N_j(\omega)}$ is the set of eigenvectors attached to j ,
- (2) $u_l(j; \omega, j) \geq u_{l+1}(j; \omega, j)$, and if equality holds, then $e_l(\omega, j) > e_{l+1}(\omega, j)$.

Extend u_l and e_l to all of Ω by setting to 0 on $\Omega \setminus \Omega_{j,l}$. Then we will prove

THEOREM 2.2. $e_l(\omega, j)$ and $u_l(n; \omega, j)$ are measurable functions on Ω for each fixed l, j (and n).

Notice

PROPOSITION 2.3. For each ω, l, l' and $j \neq k$,

$$\sum_{n=-\infty}^{\infty} \overline{u_l(n; \omega, j)} u_{l'}(n; \omega, k) = 0. \tag{2.1}$$

This is true because the u 's are distinct eigenfunctions (since they are attached to distinct points) and so orthogonal.

As a first preliminary to the proofs of Theorems 2.1, 2.2, we make several simplifying remarks:

- (i) Without loss, we can take $j = 0$.
- (ii) Instead of looking only at eigenfunctions attached to $j = 0$, we can look at eigenfunctions with $u(0) \neq 0$ normalizing by $u(0) > 0$. If we define $\tilde{N}_0(\omega)$ and $\tilde{u}_l(n; \omega, 0)$ analogously by requiring the analog of (1) and (2), and prove measurability, we recover

Theorems 2.1, 2.2 by noting that since $\tilde{u}_1(n; \omega, 0)$ is measurable, $\{\omega \mid \tilde{u}_1 \text{ is attached to } 0\}$ is measurable, and similarly for \tilde{u}_2, \dots . Then define u_1 in the obvious way and see that it is measurable.

(iii) It will be convenient to deal with that multiple η of u with $\eta(0)=1$. u and η are related, of course, by $\eta(n)=u(n)/u(0)$ and $u(n)=\eta(n)/(\sum_{j=-\infty}^{\infty} |\eta(j)|^2)^{1/2}$. By these relations, if we show that η is measurable, so is u . Notice that since $u(0)=1/\|\eta\|$, ordering by maximal $u(0)$ is the same as ordering by minimal $\|\eta\|$.

Note. Since the passage from η to u involves an infinite sum, weak continuity of η does not imply weak continuity of u , only weak measurability. The use of η below is critical because it, not u , is weakly continuous.

As a second preliminary, we note a few standard facts about Borel functions.

LEMMA 2.4. *Let X be a complete metric space, $Y \subset X$ an arbitrary subspace, and B_X, B_Y their Borel sigma algebras. Then*

$$B_Y = \{Y \cap A \mid A \in B_X\}. \quad (2.2)$$

Proof. Let \tilde{B}_Y be the right side of (2.2). Then $B_Y \subset \tilde{B}_Y$ since \tilde{B}_Y is clearly a sigma algebra containing the closed sets in Y since if C is closed in Y and D is its closure in X , then $D \cap Y = C$. Conversely, let $\tilde{B}_X = \{A \subset X \mid A \cap Y \in B_Y\}$. $\tilde{B}_X \supset B_X$ since \tilde{B}_X is a sigma algebra containing the open sets. Thus, $\tilde{B}_Y \subset B_Y$. \square

LEMMA 2.5. *Let X be a complete metric space and $Y \subset X$ with $Y \in B_X$, the Borel subsets of X . Then any $C \in B_Y$, the Borel subset of Y , is Borel as a subset of X .*

Proof. By Lemma 2.4, $C = Y \cap A$ with $A \in B_X$. Since $Y \in B_X$, so is C . \square

PROPOSITION 2.6. *Let X be a complete metric and B_X its Borel subsets. Let $X = \bigcup_{n=1}^{\infty} X_n$ with $X_n \in B_X$. Let $f: X \rightarrow \mathbf{R}$ be such that for each n , $f_n \equiv f \upharpoonright X_n$ is a Borel function from X_n to \mathbf{R} . Then f is Borel.*

Proof. Let $(a, b) \subset \mathbf{R}$. Then $f^{-1}(a, b) = \bigcup_{n=1}^{\infty} f_n^{-1}(a, b)$ is Borel by Lemma 2.5. \square

Example. Let $X = [0, 1]$ and let $f = \chi_{[1/2, 1]}$ be the characteristic function of $[\frac{1}{2}, 1]$. Let

$$X_n = \left[\frac{1}{2}, 1 \right] \cup \left[0, \frac{1}{2} - \frac{1}{n} \right].$$

Then $f \upharpoonright X_n: X_n \rightarrow \mathbf{R}$ is continuous, so f is Borel. This example shows that Proposition 2.6 is false if a Borel function is replaced by a continuous function. This is useful to keep in mind, given the continuous function argument we use below.

As a final preliminary, we note the following elementary fact:

PROPOSITION 2.7. *Suppose that $\omega^{(m)} \in \Omega$ and $\omega^{(m)} \rightarrow \omega^{(\infty)}$, and let $h_m \equiv h_{\omega^{(m)}}$. Suppose that for each finite m , there is $\eta^{(m)}$ with*

- (i) $h_m \eta^{(m)} = e_m \eta^{(m)}$,
- (ii) $\eta^{(m)}(0) = 1$,
- (iii) $\sup_m \|\eta^{(m)}\| \equiv C < \infty$.

Then there exists a subsequence m_i , $\eta^{(\infty)}$ and e_∞ so that

- (1) $h_\infty \eta^{(\infty)} = e_\infty \eta^{(\infty)}$,
- (2) $\eta^{(\infty)}(0) = 1, \|\eta^{(\infty)}\| \leq C$,
- (3) $e_{m_i} \rightarrow e_\infty, \eta^{(m_i)}(n) \rightarrow \eta^{(\infty)}(n)$ for each n .

Proof. $|e_m| \leq 2 + A$, so by compactness of the unit ball of $l^2(\mathbf{Z})$ in the weak topology, we can pick a subsequence so that (3) holds. (2) is obvious and (1) holds by taking pointwise limits in the equation (i). □

In the proof below, we have to worry about three possibilities that destroy continuity of a function like \tilde{N}_0 . First, $\sup_m \|\eta^{(m)}\| = \infty$. We will avoid this by looking at subsets with $\|\eta\| \leq k$, get measurability, and take k to infinity. Second, as $\omega^{(m)} \rightarrow \omega^{(\infty)}$, two distinct eigenvalues of h_m can converge to a single e so that h_∞ has fewer eigenvalues. We will avoid this by looking at subsets where eigenvalues stay at least a distance 2^{-l} from each other. Then we will take l to infinity. Third, after we restrict to η 's with $\|\eta\| \leq k$, in a limit a bunch of η 's with $\|\eta\| \geq k$ can approach one with $\|\eta\| = k$ increasing N . We will handle this by proving semicontinuity rather than continuity as the starting point of a proof of measurability.

Proof of Theorems 2.1, 2.2. Let, as above,

$$e_m(\omega) = e_m(\omega, 0) \quad \text{and} \quad \eta_m(n; \omega) = \frac{u_m(n; \omega, 0)}{u_m(0; \omega, 0)}.$$

For each pair of positive integers k, p , define $M_{k,p}(\omega)$ to be the maximum number of m 's so that $\|\eta_m\| \leq k$ and $|e_m - e_{m'}| \geq 2^{-p}$, for all $m' \neq m$. Since Parseval's inequality implies that $\sum_m 1/\|\eta_m\|^2 \leq 1$, there are at most k^2 m 's with $\|\eta_m\| \leq k$ and so we can determine the maximum number with $|e_m - e_{m'}| \geq 2^{-p}$.

We claim that $S \equiv \{\omega \mid M_{k,p}(\omega) \geq l\}$ is closed so $M_{k,p}(\cdot)$ is measurable. For if $\omega^{(m)} \in S$ and $\omega^{(m)} \rightarrow \omega^{(\infty)}$, we can use Proposition 2.7 and find η 's and e 's for h_∞ by taking limits. Clearly, the limiting e 's still obey the condition $|e_m - e_{m'}| \geq 2^{-p}$. Thus, S is closed and $M_{k,p}(\cdot)$ is measurable.

Now define

$$M_{k,\infty}(\omega) = \# \text{ of } m\text{'s with } \|\eta_m\| \leq k.$$

Because of simplicity of the spectrum,

$$M_{k,\infty}(\omega) = \max_p M_{k,p}(\omega)$$

so

$$\{\omega \mid M_{k,\infty}(\omega) \geq l\} = \bigcup_{p=1}^{\infty} \{\omega \mid M_{k,p}(\omega) \geq l\}$$

is an F_σ , and so $M_{k,\infty}$ is measurable.

Next define

$$\Sigma_{k,l,p} = \{\omega \mid M_{k,\infty}(\omega) = M_{k,p}(\omega) = l\},$$

the set of ω 's with exactly l eigenvalues with $\|\eta_m\| \leq k$ so that $|e_m - e_{m'}| \geq 2^{-p}$. $\Sigma_{k,l,p}$ is a Borel set. Let $e_1 > \dots > e_l$ be the eigenvalues and η_1, \dots, η_l the eigenvectors (normalized by $\eta_i(0) = 1$). We claim that e_i and η_i are continuous on $\Sigma_{k,l,p}$. For if $\omega^{(m)} \rightarrow \omega^{(\infty)}$, $e_i^{(m)}$ has limit points which are distinct (as i varies) since $|e_i^{(m)} - e_j^{(m)}| \geq 2^{-p}$. By Proposition 2.7, these limit points must be eigenvalues of h_∞ with eigenvectors $\eta_i^{(\infty)}$ obeying $\|\eta_i^{(\infty)}\| \leq k$ and so these limit points must be the $e_i^{(\infty)}$ since h_∞ has only l such eigenvalues. That is, $e_i^{(\infty)}$ are the unique limit points of $e_i^{(m)}$, so $e_i^{(m)} \rightarrow e_i^{(\infty)}$. Similarly, the η 's converge by Proposition 2.7 and the uniqueness of eigenvectors.

Now let

$$\Sigma_{k,l} = \{\omega \mid M_{k,\infty}(\omega) = l\} = \bigcup_p \Sigma_{k,p,l}.$$

By Proposition 2.6, e_i and η_i are measurable on $\Sigma_{k,l}$.

Now change the labeling so that instead of $e_1 > e_2 > \dots > e_l$, we have $\|\eta_i\| \leq \|\eta_{i+1}\|$ with $e_i > e_{i+1}$ if $\|\eta_i\| = \|\eta_{i+1}\|$. This involves a permutation π so that

$$\begin{aligned} e_i^{(\text{new})} &= e_{\pi(i)}^{\text{old}}, \\ \eta_i^{(\text{new})} &= \eta_{\pi(i)}^{\text{old}}. \end{aligned}$$

Because $\eta_i^{(\text{old})}$ and $e_i^{(\text{old})}$ are Borel functions, the set $\Sigma_{k,l}^{(\pi)}$ on which a given permutation π is used is a Borel set. $e_i^{(\text{new})}$ is built out of Borel functions on each $\Sigma^{(\pi)}$ and so we have measurability with the changed labeling.

Once we change labeling, e_i 's and η_i 's are defined consistently on $\Sigma_{k,l}$ as k, l vary, and so, using Proposition 2.6, on all Ω . \square

Note that although the set of all eigenvalues can be naively considered a non-measurable "function" of ω , since it is invariant but nonconstant, we have shown that it admits a measurable selection.

As a final remark on the issue of this section, we want to rewrite (2.1) in a useful way. Taking $\omega_n(\theta) = \lambda \cos(\pi\alpha n + \theta)$ embeds the circle $\theta \in [0, \pi)$ into Ω , so we define $u_l(n; \theta) \equiv u_l(n; \theta, 0)$, the eigenvectors with leftmost maximum at 0. Note that

$$u_l(\cdot - j; \theta + j\alpha\pi) = u_l(\cdot; \theta, j).$$

In particular, (2.1) becomes

$$\sum_n \overline{u_{l'}(n - j; \theta + j\alpha\pi)} u_l(n; \theta) = 0 \tag{2.3}$$

for all l, l' (even with $l = l'$) and $j \neq 0$. If we extend $u_l \equiv 0$ on the set where there are not l eigenvectors with leftmost maximum at 0, then (2.3) holds for all θ .

3. Duality

Fix α irrational throughout this section.

Let $\mathcal{H} = L^2([0, 2\pi), d\theta/2\pi) \times \mathbf{Z}$, that is, functions $\varphi: [0, 2\pi) \times \mathbf{Z} \rightarrow \mathbf{C}$ with

$$\sum_n \int |\varphi(\theta, n)|^2 \frac{d\theta}{2\pi} < \infty.$$

Define Q_λ on \mathcal{H} by

$$(Q_\lambda \varphi)(\theta, n) = \varphi(\theta, n+1) + \varphi(\theta, n-1) + \lambda \cos(\pi\alpha n + \theta) \varphi(\theta, n),$$

that is, Q_λ is the direct integral in θ of $h_{\alpha, \lambda, \theta}$.

Following Chulaevsky–Delyon [9] we define $U: \mathcal{H} \rightarrow \mathcal{H}$ by the formal expression

$$(U\varphi)(\eta, m) = \sum_n \int e^{-i(\eta + \pi\alpha m)n} e^{-im\theta} \varphi(\theta, n) \frac{d\theta}{2\pi}. \tag{3.1}$$

In terms of the Fourier transform $\widehat{\varphi}(m, \eta)$ we have

$$(U\varphi)(\eta, m) = \widehat{\varphi}(m, \eta + \pi\alpha m), \tag{3.2}$$

which gives a precise definition even for cases where the sum in n may not converge absolutely and shows that U is unitary. Here is a precise version of Aubry duality [2], [3]:

THEOREM 3.1.

$$Q_\lambda U = \frac{1}{2}\lambda U Q_{4/\lambda}.$$

Proof. A straightforward calculation. For example, if $(T\varphi)(\theta, n) = \varphi(\theta, n+1)$, then $(U^{-1}TU\varphi)(\theta, n) = e^{-i(\pi\alpha n + \theta)}\varphi(\theta, n)$ so

$$\begin{aligned} (U^{-1}Q_\lambda U\varphi)(\theta, n) &= 2\cos(\pi\alpha n + \theta)\varphi(\theta, n) + \frac{1}{2}\lambda[\varphi(\theta, n+1) + \varphi(\theta, n-1)] \\ &= \frac{1}{2}\lambda(Q_{4/\lambda}\varphi)(\theta, n). \end{aligned} \quad \square$$

Remark. Theorem 3.1 also provides a proof of duality for the integrated density of states first rigorously proven in [4]. For let

$$g(\theta, n) = \delta_{n0}.$$

Then, $Ug = g$. Moreover, if $k_\lambda(E)$ is the integrated density of states, then for any continuous function f ,

$$\langle g, f(Q_\lambda)g \rangle = \int f(E) dk_\lambda(E).$$

Thus Theorem 3.1, which implies

$$\langle g, f(Q_\lambda)g \rangle = \langle Ug, Uf(Q_\lambda)U^{-1}Ug \rangle = \langle g, f(\frac{1}{2}\lambda Q_{4/\lambda})g \rangle,$$

yields the duality of k .

We need one more simple calculation:

PROPOSITION 3.2. For any $\varphi \in \mathcal{H}$, $l \in \mathbf{Z}$, define a unitary operator S_l by

$$(S_l\varphi)(\theta, n) = \varphi(\theta + \pi\alpha l, n - l). \quad (3.3)$$

Then

$$(US_l\varphi)(\eta, m) = e^{-il\eta}(U\varphi)(\eta, m). \quad (3.4)$$

Proof. Let φ be such that there exists N_0 with $\varphi(n, \theta) = 0$ if $|n| > N_0$. Then (3.4) is a simple change of variables in the integral (3.1). Since such φ 's are dense and S_l is bounded, (3.4) holds for all φ . \square

PROPOSITION 3.3. Let $\varphi \in \mathcal{H}$ so that for all $l \neq 0$, $\langle S_l\varphi, \varphi \rangle = 0$. Then

$$\sum_m |(U\varphi)(\eta, m)|^2 = g(\eta)$$

is a.e. independent of η .

Proof. Note that since $U\varphi \in \mathcal{H}$, $g \in L^1(\{0, 2\pi\}, d\theta/2\pi)$. We compute $\int e^{il\eta}g(\eta) d\eta/2\pi = \langle U\varphi, e^{il\eta}U\varphi \rangle = \langle U\varphi, US_{-l}\varphi \rangle = \langle \varphi, S_{-l}\varphi \rangle = 0$, $l \neq 0$, by hypothesis. By the weak-* density of finite linear combinations of $\{e^{il\eta}\}_{l=-\infty}^{\infty}$ in L^∞ , we conclude that $g(\eta)$ is constant. \square

We come now to the main result of this section.

THEOREM 3.4. Fix λ and fix α irrational. Let $u_l(\cdot; \theta, j)$ be the measurable function described in Theorem 2.2 for the Hamiltonian $h_{4/\lambda, \alpha, \theta}$. Let $f(\theta)$ be an arbitrary function in $L^2([0, 2\pi), d\theta/2\pi)$. Let $\varphi(\theta, n) = f(\theta)u_l(n; \theta, j)$ for some fixed l, j . For each η , let

$$\psi_\eta(n) = (U\varphi)(\eta, n)$$

and let $d\mu_\eta(E)$ be the spectral measure for Hamiltonian $h_{\lambda, \alpha, \eta}$ and vector ψ_η . Then $d\mu_\eta$ is a.e. η -independent.

Proof. By (2.3), $S_k\varphi$ is orthogonal (in \mathcal{H}) to φ for any $k \neq 0$. Moreover, since

$$(F(\frac{1}{2}\lambda Q_{4/\lambda})\varphi)(\theta, n) = F(\frac{1}{2}\lambda e_l(\theta, j))\varphi(\theta, n),$$

we have that $S_k\varphi, k \neq 0$, is orthogonal to $F(Q_{4/\lambda})\varphi$ for any continuous function F . As in Proposition 3.3, we conclude that

$$\sum_m \overline{(U\varphi)(\eta, m)} (UF(\frac{1}{2}\lambda Q_{4/\lambda})\varphi)(\eta, m) \tag{3.5}$$

is independent of η . But by Theorem 3.1, $UF(\frac{1}{2}\lambda Q_{4/\lambda}) = F(Q_\lambda)U$. So (3.5) is just $\int F(E) d\mu_\eta(E)$. Since this is a.e. η -independent for each continuous F (and the set of continuous F 's is separable), we conclude that $d\mu_\eta(E)$ is a.e. η -constant. \square

4. A.e. mutual singularity of the singular parts

In this section we want to note a simple consequence of Deift–Simon [10]:

THEOREM 4.1. Let h_ω be an ergodic family of Jacobi matrices and let $d\mu_\omega^s$ be the singular part of a spectral measure for h_ω . Then for a.e. ω and ω' , $d\mu_\omega^s$ and $d\mu_{\omega'}^s$ are mutually singular.

Remark. This is an analog of the celebrated result of Aronszajn [1] and Donoghue [13] of mutual singularity under rank-one perturbations; see [26].

Proof. Let $G_\omega(n, m; z)$ be the Green function for h_ω (matrix elements of $(h_\omega - z)^{-1}$) and let \mathfrak{S}_ω be the set of E_0 in \mathbf{R} so that

$$\overline{\lim}_{\varepsilon \downarrow 0} \{ \text{Im}[G_\omega(0, 0; E_0 + i\varepsilon)] + \text{Im}[G_\omega(1, 1, E_0 + i\varepsilon)] \} = \infty.$$

By the theorem of de la Vallée-Poussin, $d\mu_\omega^s$ is supported by \mathfrak{S}_ω . Deift–Simon [10] prove that for every $E_0 \in \mathbf{R}$, $\{\omega' | E_0 \in \mathfrak{S}_{\omega'}\}$ has measure 0. Thus,

$$\int \text{meas}(\{\omega' | E_0 \in \mathfrak{S}_{\omega'}\}) d\mu_\omega(E_0) = 0,$$

and we see that

$$\mu_\omega(\mathfrak{S}_{\omega'}) = 0$$

for a.e. ω' . Thus for each fixed ω , $d\mu_{\omega'}^s$ is mutually singular to $d\mu_\omega$ for a.e. ω' . \square

Remark. Note that $d\mu_\omega$ in Theorem 4.1 only needs to be “a spectral measure”, namely, the theorem holds regardless of how it is chosen. In particular, $d\mu_\omega$ can be chosen as the spectral measure for some fixed vector, or it can be chosen as the spectral measure for an ω -dependent vector. In the next section we apply Theorem 4.1 to some particular choice of an ω -dependent vector.

5. Putting it all together

THEOREM 5.1. *Fix λ and fix α irrational. Let $u_l(\cdot; \theta, j)$ be the measurable function described in Theorem 2.2 for the Hamiltonian $h_{4/\lambda, \alpha, \theta}$. Let $f(\theta)$ be an arbitrary function in $L^2([0, 2\pi), d\theta/2\pi)$. Let $\varphi(\theta, n) = f(\theta)u_l(n; \theta, j)$ for some fixed l, j . For each η , let $\psi_\eta(n) = (U\varphi)(\eta, n)$ and let $d\mu_\eta(E)$ be the spectral measure for Hamiltonian $h_{\lambda, \alpha, \eta}$ and vector ψ_η . Then $d\mu_\eta$ is purely a.c. for a.e. η .*

Proof. By Theorem 3.4, $d\mu_\eta$ is a.e. constant. By Theorem 4.1, this means that $d\mu_\eta^s$ is a.e. zero. \square

THEOREM 5.2. *Fix λ and fix α irrational. If $h_{4/\lambda, \alpha, \theta}$ has point spectrum for a set of θ 's of positive measure, then $h_{\lambda, \alpha, \theta}$ has some a.c. spectrum for a.e. θ .*

Remark. It then follows by [24] that $h_{\lambda, \alpha, \theta}$ has some a.c. spectrum for all θ .

Proof. $\{u_l(\cdot; \theta, j)\}_{l, j}$ span the point spectrum for $h_{4/\lambda, \alpha, \theta}$, so if there is point spectrum, some $u_1(\cdot; \cdot, j)$ is a nonzero function in \mathcal{H} . Thus by Theorem 5.1, $d\mu_\eta$ is a.e. purely absolutely continuous. Since $\int d\mu_\eta(E) d\eta = \sum_n \int |u_1(n; \theta, j)|^2 d\theta/2\pi > 0$, we conclude that $\int d\mu_\eta(E) \neq 0$ for a set of η 's of positive measure and so for a.e. η since $d\mu_\eta(E)$ is a.e. η -independent. \square

THEOREM 5.3. *Fix λ and fix α irrational. If $h_{4/\lambda, \alpha, \theta}$ has only point spectrum for a.e. θ , then $h_{\lambda, \alpha, \theta}$ has only a.c. spectrum for a.e. θ .*

Proof. Let $f_m(\theta)$ be an orthonormal basis for $L^2([0, 2\pi), d\theta/2\pi)$. By hypothesis for a.e. θ , $\{u_l(\cdot; \theta, j)\}_{l, j}$ is an orthonormal basis for $l^2(\mathbf{Z})$ where we run over those l, j for which $u_l(\cdot; \theta, j) \neq 0$. It follows that if $\varphi_{m, l, j}(\theta, n) \equiv f_m(\theta)u_l(n; \theta, j)$, then $\{\varphi_{m, l, j}\}_{m, l, j}$ is a complete orthogonal set (but not necessarily normalized). By the unitarity of U , $\{U\varphi_{m, l, j}\}_{m, l, j}$ is also a complete orthogonal set. Thus for a.e. η , $\{U\varphi_{m, l, j}(\eta, \cdot)\}$ is a complete set. But these vectors lie in the a.c. spectral subspace by Theorem 5.1. \square

Proof of Theorem 1. Last [22] has shown for such α , the Lebesgue measure of the spectrum $\sigma_{\lambda=2,\alpha}$ is zero. It follows that there is no a.c. spectrum for such α and $\lambda=2$. By Theorem 5.2, there can't be any point spectrum a.e. since such point spectrum would imply a.c. spectrum. Thus for a.e. θ , the spectrum is purely singular continuous. \square

Proof of Theorem 2. This is just a restatement of Theorem 5.3. \square

Proof of Theorem 3. Let $\lambda < 2$. Then $h_{4/\lambda,\alpha,\theta}$ has no a.c. spectrum since $4/\lambda > 2$ (see [3], [5]). Thus by Theorem 5.2, $h_{\lambda,\alpha,\theta}$ can have no point spectrum. \square

References

- [1] ARONSAJN, N., On a problem of Weyl in the theory of Sturm–Liouville equations. *Amer. J. Math.*, 79 (1957), 597–610.
- [2] AUBRY, S., The new concept of transitions by breaking of analyticity in a crystallographic model, in *Solitons and Condensed Matter Physics (Oxford, 1978)*, pp. 264–277. Springer Ser. Solid-State Sci., 8. Springer-Verlag, Berlin–New York, 1978.
- [3] AUBRY, S. & ANDRE, G., Analyticity breaking and Anderson localization in incommensurate lattices. *Ann. Israel Phys. Soc.*, 3 (1980), 133–140.
- [4] AVRON, J. & SIMON, B., Singular continuous spectrum for a class of almost periodic Jacobi matrices. *Bull. Amer. Math. Soc.*, 6 (1982), 81–85.
- [5] — Almost periodic Schrödinger operators, II. The integrated density of states. *Duke Math. J.*, 50 (1983), 369–391.
- [6] BELLISSARD, J., LIMA, R. & TESTARD, D., A metal-insulator transition for the almost Mathieu model. *Comm. Math. Phys.*, 88 (1983), 207–234.
- [7] BELOKOLOS, E. D., A quantum particle in a one-dimensional deformed lattice. Estimates of lacunae dimension in the spectrum. *Teoret. Mat. Fiz.*, 25 (1975), 344–57 (Russian).
- [8] CHOJNACKI, W., A generalized spectral duality theorem. *Comm. Math. Phys.*, 143 (1992), 527–544.
- [9] CHULAEVSKY, V. & DELYON, F., Purely absolutely continuous spectrum for almost Mathieu operators. *J. Statist. Phys.*, 55 (1989), 1279–1284.
- [10] DEIFT, P. & SIMON, B., Almost periodic Schrödinger operators, III. The absolutely continuous spectrum in one dimension. *Comm. Math. Phys.*, 90 (1983), 389–411.
- [11] DELYON, F., Absence of localization for the almost Mathieu equation. *J. Phys. A*, 20 (1987), L21–L23.
- [12] DINABURG, E. & SINAI, YA., The one-dimensional Schrödinger equation with a quasi-periodic potential. *Functional Anal. Appl.*, 9 (1975), 279–289.
- [13] DONOGHUE, W., On the perturbation of the spectra. *Comm. Pure Appl. Math.*, 18 (1965), 559–579.
- [14] FRÖHLICH, J., SPENCER, T. & WITTMER, P., Localization for a class of one-dimensional quasi-periodic Schrödinger operators. *Comm. Math. Phys.*, 132 (1990), 5–25.
- [15] GESZTESY, F. & SIMON, B., The xi function. *Acta Math.*, 176 (1996), 49–71.
- [16] GOLDSTEIN, M., Laplace transform method in perturbation theory of the spectrum of Schrödinger operators, II. One-dimensional quasi-periodic potentials. Preprint, 1992.
- [17] HELFFER, B. & SJÖSTRAND, J., Semi-classical analysis for Harper's equation, III. Cantor structure of the spectrum. *Mém. Soc. Math. France (N.S.)*, 39 (1989), 1–139.

- [18] JITOMIRSKAYA, S., Anderson localization for the almost Mathieu equation: A nonperturbative proof. *Comm. Math. Phys.*, 165 (1994), 49–57.
- [19] — Almost everything about the almost Mathieu operator, II, in *XIth International Congress of Mathematical Physics (Paris, 1994)*, pp. 373–382. Internat. Press, Cambridge, MA, 1995.
- [20] JITOMIRSKAYA, S. & SIMON, B., Operators with singular continuous spectrum, III. Almost periodic Schrödinger operators. *Comm. Math. Phys.*, 165 (1994), 201–205.
- [21] LAST, Y., A relation between a.c. spectrum of ergodic Jacobi matrices and the spectra of periodic approximants. *Comm. Math. Phys.*, 151 (1993), 183–192.
- [22] — Zero measure for the almost Mathieu operator. *Comm. Math. Phys.*, 164 (1994), 421–432.
- [23] — Almost everything about the almost Mathieu operator, I, in *XIth International Congress of Mathematical Physics (Paris, 1994)*, pp. 366–372. Internat. Press, Cambridge, MA, 1995.
- [24] LAST, Y. & SIMON, B., Eigenfunctions, transfer matrices, and a.c. spectrum of one-dimensional Schrödinger operators. Preprint.
- [25] MANDELSHTAM, V. & ZHITOMIRSKAYA, S., 1D-quasiperiodic operators. Latent symmetries. *Comm. Math. Phys.*, 139 (1991), 589–604.
- [26] SIMON, B., Spectral analysis of rank one perturbations and applications, in *Mathematical Quantum Theory, II. Schrödinger Operators (Vancouver, BC, 1993)*, pp. 109–149. CRM Proc. Lecture Notes, 8. Amer. Math. Soc., Providence, RI, 1995.
- [27] SINAI, YA., Anderson localization for one-dimensional Schrödinger operator with quasiperiodic potential. *J. Statist. Phys.*, 46 (1987), 861–909.

A. Y. GORDON
Department of Mathematics
University of North Carolina at Charlotte
Charlotte, NC 28223
U.S.A.
aygordon@uncc.edu

S. JITOMIRSKAYA
Department of Mathematics
University of California
Irvine, CA 92717
U.S.A.
szhitomi@math.uci.edu

Y. LAST
Division of Physics, Mathematics and Astronomy
California Institute of Technology
Pasadena, CA 91125
U.S.A.
ylast@cco.caltech.edu

B. SIMON
Division of Physics, Mathematics and Astronomy
California Institute of Technology
Pasadena, CA 91125
U.S.A.
bsimon@caltech.edu

Received February 7, 1996