

# THE STABILITY OF THE MEAN CURVATURE FLOW IN MANIFOLDS OF SPECIAL HOLONOMY

CHUNG-JUN TSAI AND MU-TAO WANG

ABSTRACT. We study the uniqueness of minimal submanifolds and the stability of the mean curvature flow in several well-known model spaces of manifolds of special holonomy. These include the Stenzel metric on the cotangent bundle of spheres, the Calabi metric on the cotangent bundle of complex projective spaces, and the Bryant-Salamon metrics on vector bundles over certain Einstein manifolds. In particular, we show that the zero sections, as calibrated submanifolds with respect to their respective ambient metrics, are unique among compact minimal submanifolds and are dynamically stable under the mean curvature flow. The proof relies on intricate interconnections of the Ricci flatness of the ambient space and the extrinsic geometry of the calibrated submanifolds.

## 1. INTRODUCTION

Calibrated submanifolds [9] in manifolds of special holonomy are not just minimal submanifolds, they actually minimize the volume functional in their homology classes. Of particular interests are special Lagrangians in Calabi–Yau, associatives and coassociatives in  $G_2$ , and Cayley submanifolds in  $Spin(7)$ . These geometric objects attracted a lot of attentions in recent years. On the one hand, they are natural generalizations of algebraic subvarieties in algebraic manifolds and thus are of immense geometric interest. On the other hand, they appear in various proposals of string theory such as Mirror Symmetry and the M-theory. The most successful construction of metrics of special holonomy is the Calabi–Yau case, where the celebrated theorem of Yau [27] shows the homological condition guarantees the existence of the metric. All other constructions are based on deformation theory, symmetry reductions or gluing constructions [11] (see also a recent flow approach for  $G_2$  construction in [2, 15]). The scenario of the construction of calibrated submanifolds is similar [10, 13]. Even in the Calabi–Yau case, the understanding of special Lagrangians is still rather limited [17, 12].

---

*Date:* May 13, 2016.

Supported in part by Taiwan MOST grants 102-2115-M-002-014-MY2 and 104-2115-M-002-007 (C.-J. Tsai) and NSF grants DMS-1105483 and DMS-1405152 (M.-T. Wang). The second-named author is supported in part by a grant from the Simons Foundation (#305519 to Mu-Tao Wang). Part of this work was carried out when Mu-Tao Wang was visiting the National Center of Theoretical Sciences at National Taiwan University in Taipei, Taiwan.

Among all explicitly constructed manifolds of special holonomy, the most well-known ones seem to be the Stenzel metric [20] on the cotangent bundles of spheres (or the Eguchi–Hanson metric in dimension 4) and the Calabi metric [3] on the cotangent bundles of complex projective spaces. Similar constructions of Bryant–Salamon [1] produce  $G_2$  and  $\text{Spin}(7)$  metrics. All of them are based on bundle constructions and the zero sections are calibrated submanifolds. In the article, we study the uniqueness and the dynamical stability of the zero sections of these manifolds. To be more specific, we consider the following manifolds of special holonomy in this article.

**Assumption 1.1.** Throughout this article,  $M$  is a Riemannian manifold of special holonomy that belongs to one of the followings:

- (i) the total space of the cotangent bundle of  $S^n$  with the Stenzel metric<sup>1</sup> with  $n > 1$ ;
- (ii) the total space of the cotangent bundle of  $\mathbb{C}\mathbb{P}^n$  with the Calabi metric;
- (iii) the total space of  $\mathbb{S}(S^3)$ ,  $\Lambda_-^2(S^4)$ ,  $\Lambda_-^2(\mathbb{C}\mathbb{P}^2)$ , or  $\mathbb{S}_-(S^4)$  with the Ricci flat metric constructed by Bryant–Salamon.

In section 2, 3 and 4, we review the geometry of these metrics in details. In all these examples,  $M$  is the total space of a vector bundle over a base manifold  $B$ . We identify  $B$  with the zero section of the bundle, which is also considered to be an embedded submanifold of  $M$ . Let  $\pi : M \rightarrow B$  be the bundle projection, and let  $\Omega$  be the pull-back of the volume form of  $B$  by the projection  $\pi$ . The smooth differential form  $\Omega$  on  $M$  characterizes  $B$  by the condition that  $\Omega(T_p B) = 1$  for all  $p \in B$  where  $T_p B$  is the (oriented) tangent space of  $B$  as a submanifold of  $M$  at a point  $p$ . Let  $d(\cdot)$  denote the distance function to the zero section  $B$  with respect to the Riemannian metric on  $M$ . For a compact embedded submanifold  $\Sigma$  of  $M$  with  $\dim \Sigma = \dim B$ ,  $\Sigma$  is considered to be  $\mathcal{C}^0$  close to the zero section if  $d(p)$  is close to 0 for all  $p \in \Sigma$ , and  $\Sigma$  is considered to be  $\mathcal{C}^1$  close to the zero section if  $\Omega(T_p \Sigma)$  is close to 1 for all  $p \in \Sigma$ .

Our first result regards the uniqueness property of the zero sections.

**Theorem 1.2.** *In each case considered in Assumption 1.1, the zero section is the unique compact minimal immersed submanifold of the given dimension.*

The mean curvature flow is the parabolic PDE system that deforms a submanifold by its mean curvature vector field, and thus the negative gradient flow of the volume functional. Although a calibrated submanifold represents a local minimum of the volume functional, there is no guarantee that it is stable along the mean curvature flow, due to the nonlinear nature and the degeneracy the PDE system. The nonlinear stability of PDE systems such as the Einstein equation or the Ricci flow is under intense study. Our next result concerns the nonlinear  $\mathcal{C}^1$  stability of the zero sections.

---

<sup>1</sup>When  $n = 1$ , the metric is not only Ricci flat, but flat. We have to exclude this flat case.

**Theorem 1.3.** *In each case considered in Assumption 1.1, there exists an  $\epsilon > 0$  which depends only on the geometry of  $M$  such that if  $\Sigma$  is a compact embedded submanifold of  $M$  and*

$$\sup_{p \in \Sigma} (d(p) + 1 - \Omega(T_p \Sigma)) < \epsilon , \quad (1.1)$$

*then the mean curvature flow of  $\Sigma$  exists for all time and converges to the zero section smoothly.*

**Remark 1.4.** The Ricci flat metric on each manifold  $M$  considered in Assumption 1.1 is constructed under some symmetry Ansatz. From the constructions, one sees that the metric turns out to be uniquely determined by the volume of the zero section  $B$ . That is to say,  $\epsilon$  depends on the volume of the zero section (and also  $n$  in case (i) and (ii)).

Theorem 1.2 is proved in section 5. The main point is to prove the convexity of the distance square to the zero section. It is also the key for the  $\mathcal{C}^0$  convergence in Theorem 1.3, and plays an important role for the  $\mathcal{C}^1$  convergence.

Such a nonlinear  $\mathcal{C}^1$  stability theorem for higher codimensional mean curvature flows has been established for manifolds of reduced holonomy, namely manifolds that are locally Riemannian products, see for examples [19, 21, 22, 23, 25]. Theorem 1.3, to the best of our knowledge, appears to be the first one for manifolds of special holonomy.

Sections 6 and 7 are devoted to the proof of Theorem 1.3. In section 6, we establish estimates on the covariant derivatives of  $\Omega$ , which is needed for proving the  $\mathcal{C}^1$  convergence. In section 7, we put everything together to prove the stability of the zero section under the mean curvature flow.

**Remark 1.5.** A general stability theorem of the mean curvature flow of compact smooth submanifolds can be derived from [18, Theorem 2] under the assumption that the initial data is close enough to a stable minimal submanifold in the  $W^{l+2,2}$  Sobolev norm for large enough  $l$  (which implies at least  $C^2$  smallness). In addition, the smallness constant depends on the functional, in this case the volume of submanifolds. Theorem 1.3 is a  $\mathcal{C}^1$  stability theorem in which the regularity requirement is lower and the dependence of the smallness constant is explicit. The theorem can be turned into a Lipschitz stability theorem (by approximating the initial Lipschitz submanifold by a family of  $C^1$  submanifolds, see [26]), which seems to be the optimal result for the mean curvature flow.

**Acknowledgement.** The first author would like to thank Prof. Shing-Tung Yau for bringing the rigidity question into his attention, and for his generosity in sharing his ideas. The authors would like to thank Mao-Pei Tsui for helpful discussions and his interest in this work.

## 2. GEOMETRY OF THE STENZEL METRIC

**2.1. The Calabi–Yau metric on  $T^*S^n$ .** Consider the  $n$ -dimensional sphere  $S^n$  with the standard metric for  $n > 1$ . Let  $\{\underline{\omega}^\mu\}_{\mu=1}^n$  be a local orthonormal coframe. Denote their connection 1-forms by  $\underline{\omega}^\nu$ . We adopt the Einstein summation convention that repeated indexes are summed. The convention here is that

$$d\underline{\omega}^\mu = -\underline{\omega}^\mu_\nu \wedge \underline{\omega}^\nu, \quad \nabla \underline{\omega}^\mu = -\underline{\omega}^\mu_\nu \otimes \underline{\omega}^\nu.$$

As a space form with curvature equal to 1, its curvature tensor is

$$\underline{\Omega}^\mu_\nu = d\underline{\omega}^\mu_\nu - \underline{\omega}^\mu_\gamma \wedge \underline{\omega}^\gamma_\nu = \underline{\omega}^\mu \wedge \underline{\omega}^\nu.$$

Let  $\{y_\mu\}$  be the coordinate for the fibers of  $T^*S^n$  induced by  $\{\underline{\omega}^\mu\}$ . The standard metric on  $S^n$  induces the following metric on  $T^*S^n$ .

$$\sum_{\mu=1}^n ((\underline{\omega}^\mu)^2 + (dy_\mu - y_\nu \underline{\omega}^\nu_\mu)^2). \quad (2.1)$$

**2.1.1. Spherical coordinate for the fibers.** There are two naturally defined 1-forms outside the zero section:

$$\frac{1}{r} y_\mu \underline{\omega}^\mu \quad \text{and} \quad \frac{1}{r} y_\mu dy_\mu = \frac{1}{r} y_\mu (dy_\mu - y_\nu \underline{\omega}^\nu_\mu)$$

where  $r = (\sum_\mu (y_\mu)^2)^{\frac{1}{2}}$ . The first one is the tautological 1-form rescaled by  $1/r$ ; the second one is the exterior derivative of  $r$ . With respect to (2.1), they are unit-normed and orthogonal to each other.

It is more convenient to consider the metric (2.1) by another coframe, which is an extension of the above two 1-forms. To start, extend the vector  $\frac{1}{r}(y_1, \dots, y_n)$  to an orthonormal frame for  $\mathbb{R}^n$ . This can be done on any simply-connected open subset of  $\mathbb{R}^n \setminus \{0\}$ . To be more precise, choose a smooth map  $T_\nu^\mu$  from a simply-connected open subset of  $\mathbb{R}^n \setminus \{0\}$  to  $O(n)$  such that

$$T_\nu^1(y) = \frac{1}{r} y_\nu \quad \text{for } \nu = 1, 2, \dots, n \quad \text{and} \quad y \in \text{Domain}(T_\nu^\mu) \subset \mathbb{R}^n \setminus \{0\}.$$

For example, the standard spherical coordinate system on  $\mathbb{R}^n \setminus \{0\}$  will do. When  $n = 3$  with  $y_1 = r \sin \theta \sin \phi$ ,  $y_2 = r \sin \theta \cos \phi$ ,  $y_3 = r \cos \theta$ , one can take  $T_\mu^2 = \frac{1}{r} \frac{\partial y_\mu}{\partial \theta}$  and  $T_\mu^3 = \frac{1}{r \sin \theta} \frac{\partial y_\mu}{\partial \phi}$ ,  $\mu = 1, 2, 3$ .

The metric (2.1) has the following orthonormal coframe

$$\begin{aligned} \sigma^\mu &= T_\nu^\mu \underline{\omega}^\nu, \\ \sigma^{n+\mu} &= T_\nu^\mu (dy_\nu - y_\gamma \underline{\omega}^\gamma_\nu) \end{aligned} \quad (2.2)$$

for  $\mu \in \{1, \dots, n\}$ .

2.1.2. *The Stenzel metric.* With this coframe (2.2), the Stenzel metric takes the form

$$(c(r)\sigma^1)^2 + \sum_{j=2}^n (a(r)\sigma^j)^2 + (c(r)\sigma^{n+1})^2 + \sum_{j=2}^n \left(\frac{b(r)}{r}\sigma^{n+j}\right)^2. \quad (2.3)$$

The coefficient functions are defined by

$$\begin{aligned} a^2 &= \frac{1}{4}h'(r)\coth r, \\ b^2 &= \frac{1}{4}h'(r)\tanh r, \\ c^2 &= \frac{1}{4}h''(r) \end{aligned} \quad (2.4)$$

where  $h'(r)$  is the solution of the ODE

$$\frac{d}{dr}(h'(r))^n = 2^{n+1}n(\sinh(2r))^{n-1} \quad \text{with } h'(0) = 0. \quad (2.5)$$

Here, prime ( ' ) denotes the derivative with respect to  $r$ . The function  $h$  is the Kähler potential in the paper of Stenzel [20, section 7]. We remark that the metric here differs from that in [7, section 2] by a factor of  $2^{n-1}/n$  in (2.5). The normalization here is chosen such that the restriction of metric (2.3) to the zero section is the round metric of radius 1, and hence the volume of the zero section is  $2\pi^{\frac{n}{2}}/\Gamma(n/2)$ .

2.1.3. *Connection and Ricci flat equation.* Take the orthonormal coframe:

$$\omega^1 = c(r)\sigma^1, \quad \omega^j = a(r)\sigma^j, \quad \omega^{n+1} = c(r)\sigma^{n+1} = c(r)dr, \quad \omega^{n+j} = \frac{b(r)}{r}\sigma^{n+j} \quad (2.6)$$

for  $j \in \{2, \dots, n\}$ . The indices  $i, j, k, \dots$  will be assumed to belong to  $\{2, \dots, n\}$ .

It is useful to introduce the new radial coordinate  $\rho$  by  $d\rho = c(r)dr = \omega^{n+1}$ , and denote by dot (  $\dot{\phantom{x}}$  ) the derivative with respect to  $\rho$ . This  $\rho$  coordinate is the distance to the zero section with respect to (2.3). Therefore,

$$\dot{f} = \frac{1}{c(r)}f'.$$

The connection 1-forms  $\omega_\nu^\mu$  of (2.6) can be found by a direct computation:

$$\begin{aligned} \omega_{n+1}^1 &= \frac{\dot{c}}{c}\omega^1, & \omega_{n+1}^j &= \frac{\dot{a}}{a}\omega^j, & \omega_{n+1}^{n+j} &= \frac{\dot{b}}{b}\omega^{n+j}, \\ \omega_{n+i}^j &= C\delta_i^j\omega^1, & \omega_1^{n+j} &= A\omega^j, & \omega_j^1 &= B\omega^{n+j} \end{aligned} \quad (2.7)$$

where

$$A = \frac{a^2 - b^2 - c^2}{2abc}, \quad B = \frac{b^2 - a^2 - c^2}{2abc}, \quad C = \frac{c^2 - a^2 - b^2}{2abc}. \quad (2.8)$$

The rest of the components  $\omega_i^j$  and  $\omega_{n+i}^{n+j}$  satisfy

$$\omega_i^j = \omega_{n+i}^{n+j} = T_\mu^j \omega_\nu^\mu T_\nu^i - (dT_\mu^j) T_\mu^i. \quad (2.9)$$

In [7, section 2], Cvetic et al. showed that the Ricci flat equations admit first integrals which are given by

$$\frac{\dot{a}}{a} + A = 0, \quad \frac{\dot{b}}{b} + B = 0, \quad \frac{\dot{c}}{c} + (n-1)C = 0. \quad (2.10)$$

They solved the equations (2.4) and reconstructed the Stenzel metric in the above form (2.3). The expressions of the Kähler form and the holomorphic volume form are quite simple in terms of the coframe (2.6). The Kähler form is  $\sum_{\mu=1}^n \omega^\mu \wedge \omega^{n+\mu}$ , and the holomorphic volume form is  $(\omega^1 + i\omega^{n+1}) \wedge (\omega^2 + i\omega^{n+2}) \wedge \cdots (\omega^n + i\omega^{2n})$ . With the above relations, one can check that these two differential forms are parallel. With this understanding, the complex structure  $I$  in terms of the dual frame  $\{\bar{e}_1, \dots, \bar{e}_{2n}\}$  of (2.6) sends  $\bar{e}_\mu$  to  $\bar{e}_{n+\mu}$  and sends  $\bar{e}_{n+\mu}$  to  $-\bar{e}_\mu$  for any  $\mu \in \{1, \dots, n\}$ .

**2.2. Coefficient functions and curvature.** The geometry of the Stenzel metric is encoded in the functions  $A$ ,  $B$  and  $C$ . In this subsection, we summarize the properties that will be used later. Note that (2.5) implies that  $h' > 0$  and  $h'' > 0$  when  $r > 0$ . By (2.4) and (2.10),

$$\begin{aligned} A &= -\frac{1}{\sqrt{h''}} \left( \frac{h''}{h'} - \frac{2}{\sinh(2r)} \right), \\ B &= -\frac{1}{\sqrt{h''}} \left( \frac{h''}{h'} + \frac{2}{\sinh(2r)} \right), \\ C &= -\frac{1}{n-1} \frac{1}{\sqrt{h''}} \frac{h'''}{h''}. \end{aligned} \quad (2.11)$$

Instead of  $r$ , we state the estimates in terms of  $\rho = \int_0^r c(u)du$ .

**Lemma 2.1.** *The functions  $A$ ,  $B$  and  $C$  are negative when  $\rho > 0$ . Moreover, there exists a constant  $K > 1$  which depends only on  $n$  such that  $|A|/\rho$ ,  $|C|/\rho$  and  $|B|/\rho$  are all bounded between  $1/K$  and  $K$  for any point  $p$  with  $0 < \rho(p) < 1$ .*

*Proof.* It follows from (2.5) that

$$\begin{aligned} \frac{h''}{h'} &= \frac{(\sinh(2r))^{n-1}}{n \int_0^r (\sinh(2u))^{n-1} du}, \\ \frac{1}{n-1} \frac{h'''}{h''} &= \frac{2n \cosh(2r) \left( \int_0^r (\sinh(2u))^{n-1} du \right) - (\sinh(2r))^n}{n \sinh(2r) \left( \int_0^r (\sinh(2u))^{n-1} du \right)}. \end{aligned} \quad (2.12)$$

To prove that  $C < 0$ , we estimate

$$2n \cosh(2r) \left( \int_0^r (\sinh(2u))^{n-1} du \right) > 2n \int_0^r \cosh(2u) (\sinh(2u))^{n-1} du = (\sinh(2r))^n$$

for any  $r > 0$ . It follows that  $C$  is negative. For  $A < 0$ ,

$$\begin{aligned} \frac{h''}{h'} - \frac{2}{\sinh(2r)} &= \frac{(\sinh(2r))^n - 2n \int_0^r (\sinh(2u))^{n-1} du}{n \sinh(2r) \int_0^r (\sinh(2u))^{n-1} du} \\ &= \frac{2n \int_0^r (\sinh(2u))^{n-1} (\cosh(2u) - 1) ds}{n \sinh(2r) \int_0^r (\sinh(2u))^{n-1} du} > 0 \end{aligned}$$

for any  $r > 0$ . It follows that  $A < 0$ . Since  $B \leq A$ ,  $B$  is also negative.

The second assertion is a direct consequence of the power series expansion at  $r = 0$ . It follows from (2.5) that the Taylor series expansion of  $h'(r)$  near  $r = 0$  is

$$h' = 4r \left( 1 + \frac{2(n-1)}{3(n+2)} r^2 + \mathcal{O}(r^3) \right) .$$

Thus,  $\rho = r + \mathcal{O}(r^2)$ ,  $A = -\frac{n}{n+2}r + \mathcal{O}(r^2)$ ,  $C = -\frac{2}{n+2}r + \mathcal{O}(r^2)$  and  $B = -r^{-1} + \mathcal{O}(1)$ . This finishes the proof of the lemma.  $\square$

We remark that (2.10) and the negativity of  $A$ ,  $B$  and  $C$  imply that  $a$ ,  $b$  and  $c$  are increasing functions. Other quantities we will encounter are the derivatives of  $A$ ,  $B$  and  $C$  with respect to  $\rho$ . It turns out that they are degree two polynomials in  $A$ ,  $B$  and  $C$ .

**Lemma 2.2.** *The functions  $\dot{A}$ ,  $\dot{B}$  and  $\dot{C}$  obey:*

$$\begin{aligned} \dot{A} &= -nBC + A(A + B + C) , \\ \dot{B} &= -nAC + B(A + B + C) , \\ \dot{C} &= -2AB + (n-1)C(A + B + C) . \end{aligned}$$

*Proof.* The equation (2.8) can be rewritten as

$$B + C = -\frac{a}{bc} , \quad C + A = -\frac{b}{ac} , \quad A + B = -\frac{c}{ab} . \quad (2.13)$$

By using (2.10),

$$\begin{aligned} \dot{B} + \dot{C} &= (B + C)(-A + B + (n-1)C) , \\ \dot{C} + \dot{A} &= (C + A)(-B + A + (n-1)C) , \\ \dot{A} + \dot{B} &= (A + B)(-(n-1)C + A + B) . \end{aligned}$$

The lemma follows from these formulae.  $\square$

2.2.1. *The Riemann curvature tensor.* In [7, section 2], Cvetič et al. also computed the components of the Riemann curvature tensor of the metric (2.3).

Note that (2.13) implies that

$$(A + B)(A + C) = \frac{1}{a^2} , \quad (B + A)(B + C) = \frac{1}{b^2} , \quad (C + A)(C + B) = \frac{1}{c^2} .$$

By applying (2.10), (2.13) and Lemma 2.2 to [7, (2.14)], we find that the components of the Riemann curvature tensor are all degree two polynomials in  $A$ ,  $B$  and  $C$ :

$$\begin{aligned}
R(\bar{e}_1, \bar{e}_j, \bar{e}_1, \bar{e}_j) &= AB + BC - nCA , \\
R(\bar{e}_1, \bar{e}_{n+j}, \bar{e}_1, \bar{e}_{n+j}) &= AB + CA - nBC , \\
\frac{1}{n-1}R(\bar{e}_1, \bar{e}_{n+1}, \bar{e}_1, \bar{e}_{n+1}) &= R(\bar{e}_1, \bar{e}_{n+1}, \bar{e}_{n+j}, \bar{e}_j) \\
&= (n-1)(CA + BC) - 2AB , \\
R(\bar{e}_i, \bar{e}_{n+k}, \bar{e}_j, \bar{e}_{n+l}) &= -AB(\delta_{ij}\delta_{kl} + \delta_{il}\delta_{jk}) + (BC + AC)\delta_{ik}\delta_{jl} , \\
R(\bar{e}_i, \bar{e}_k, \bar{e}_j, \bar{e}_l) &= (AB + BC + CA)(\delta_{ij}\delta_{kl} - \delta_{il}\delta_{jk})
\end{aligned}$$

and other inequivalent components vanish. Here,  $\bar{e}_1, \dots, \bar{e}_{2n}$  is the dual frame of (2.6), and the convention of the Riemann curvature tensor is

$$R(X, Y, Z, W) = \langle \nabla_Z \nabla_W Y - \nabla_W \nabla_Z Y - \nabla_{[Z, W]} Y, X \rangle .$$

By equivalence we mean that the curvature is a quadrilinear map  $R$  satisfying the condition

$$R(X, Y, Z, W) = -R(X, Y, W, Z) = R(Z, W, X, Y) = R(X, Y, IZ, IW) . \quad (2.14)$$

The most important property of the Riemann curvature tensor of the Stenzel metric that will be used later is  $R(\bar{e}_\mu, \bar{e}_{n+\nu}, \bar{e}_{n+\delta}, \bar{e}_{n+\epsilon}) = 0 = R(\bar{e}_{n+\mu}, \bar{e}_\nu, \bar{e}_\delta, \bar{e}_\epsilon)$  for any  $\mu, \nu, \delta, \epsilon \in \{1, \dots, n\}$ .

### 3. GEOMETRY OF THE CALABI METRIC

The Eguchi–Hanson metric has another higher dimensional generalization. There exists a hyper-Kähler metric on the cotangent bundle of the complex projective space,  $T^*\mathbb{C}\mathbb{P}^n$ . The metric is thus Ricci flat. It was constructed by Calabi in [3, section 5] by solving the Kähler potential under an ansatz. Since we are going to study the Riemannian geometric properties of the metric, it is more convenient to describe the metric in terms of a moving frame.

**3.1. The hyper-Kähler metric on  $T^*\mathbb{C}\mathbb{P}^n$ .** Consider the  $n$ -dimensional complex projective space  $\mathbb{C}\mathbb{P}^n$  with the Fubini–Study metric. Let  $\{\theta^\mu\}$  be a local unitary coframe of type  $(1, 0)$ . That is to say, the Fubini–Study metric is

$$\sum_{\mu=1}^n |\theta^\mu|^2 .$$

Denote by  $\theta_\nu^\mu$  the corresponding connection 1-forms. They are determined uniquely by the relations:

$$d\theta^\mu = -\theta_\nu^\mu \wedge \theta^\nu \quad \text{and} \quad \theta_\mu^\nu + \overline{\theta_\nu^\mu} = 0 . \quad (3.1)$$



The curvature of the Fubini–Study metric is

$$\Theta_\nu^\mu = d\theta_\nu^\mu - \theta_\nu^\gamma \wedge \theta_\gamma^\mu = \theta^\mu \wedge \overline{\theta^\nu} + \delta_\nu^\mu \theta^\gamma \wedge \overline{\theta^\gamma} .$$

The curvature formula implies that its sectional curvature lies between 1 and 4, and is equal to 4 if and only if the 2-plane is complex. The detailed discussion of the Fubini–Study metric in terms of the moving frame can be found in [5, section 8].

Let  $\{z_\mu\}$  be the complex coordinate for the fibers of  $(T^*\mathbb{C}\mathbb{P}^n)^{(1,0)}$ . Then, the Fubini–Study metric induces the following metric on  $T^*\mathbb{C}\mathbb{P}^n$ .

$$\sum_{\mu=1}^n (|\theta^\mu|^2 + |dz_\mu - z_\nu \theta_\mu^\nu|^2) . \quad (3.2)$$

The complex structure of  $\mathbb{C}\mathbb{P}^n$  induces a complex structure on  $T^*\mathbb{C}\mathbb{P}^n$ , with respect to which  $\theta^\mu$  and  $dz_\mu - z_\mu \theta_\mu^\nu$  are  $(1,0)$ -forms.

**Remark 3.1.** We briefly explain the convention of the correspondence between real and complex moving frames. Write  $\theta^\mu$  as  $\underline{\omega}^{2\mu-1} + i\underline{\omega}^{2\mu}$ . Then,  $\{\underline{\omega}^{2\mu-1}\}_{\mu=1}^n \cup \{\underline{\omega}^{2\mu}\}_{\mu=1}^n$  constitutes an orthonormal coframe. Let  $\{\underline{e}_{2\mu-1}\}_{\mu=1}^n \cup \{\underline{e}_{2\mu}\}_{\mu=1}^n$  be the dual orthonormal frame. They satisfy  $J(\underline{e}_{2\mu-1}) = \underline{e}_{2\mu}$ , where  $J$  is the complex structure as an endomorphism on the (real) tangent bundle. Denote by  $\underline{\omega}_A^B$  the connection 1-forms, where  $1 \leq A, B \leq 2n$ . Namely,  $\nabla \underline{e}_A = \underline{\omega}_A^B \otimes \underline{e}_B$ . Since  $J$  is parallel,  $\underline{\omega}_{2\nu}^{2\mu} = \underline{\omega}_{2\nu-1}^{2\mu-1}$  and  $\underline{\omega}_{2\nu}^{2\mu-1} = -\underline{\omega}_{2\nu-1}^{2\mu}$ . The Hermitian connection  $\theta_\nu^\mu$  is equal to

$$\underline{\omega}_{2\nu-1}^{2\mu-1} + i\underline{\omega}_{2\nu-1}^{2\mu} = \underline{\omega}_{2\nu}^{2\mu} - i\underline{\omega}_{2\nu}^{2\mu-1} . \quad (3.3)$$

For the total space of the cotangent bundle, write  $z_\mu$  as  $x_\mu - iy_\mu$ . Under the (real) isomorphism

$$\begin{aligned} (T^*\mathbb{C}\mathbb{P})^{(1,0)} &\cong T^*\mathbb{C}\mathbb{P}^n \quad (\text{real cotangent bundle}) \\ z_\mu \theta^\mu &\leftrightarrow \text{Re}(z_\mu \theta^\mu) = x_\mu \underline{\omega}^{2\mu-1} + y_\mu \underline{\omega}^{2\mu} , \end{aligned}$$

the metric (3.2) is equal to

$$\sum_{\mu=1}^n \left( (\underline{\omega}^{2\mu-1})^2 + (\underline{\omega}^{2\mu})^2 + (dx_\mu - x_\nu \underline{\omega}_{2\mu-1}^{2\nu-1} - y_\nu \underline{\omega}_{2\mu-1}^{2\nu})^2 + (dy_\mu - x_\nu \underline{\omega}_{2\mu}^{2\nu-1} - y_\nu \underline{\omega}_{2\mu}^{2\nu})^2 \right) .$$

3.1.1. *Spherical coordinate for the fibers.* Let  $r = \sqrt{\sum_\mu |z_\mu|^2}$  be the distance to the zero section with respect to the metric (3.2). Again, the exterior derivative of  $r$  and the tautological 1-form are two naturally defined 1-forms on  $T^*\mathbb{C}\mathbb{P}^n$ . In terms of the complex coordinate, they read

$$\text{Re} \left( \frac{1}{r} \bar{z}_\mu (dz_\mu - z_\nu \theta_\mu^\nu) \right) \quad \text{and} \quad \text{Re}(z_\mu \theta^\mu) ,$$

respectively. Their images under the complex structure give another two 1-forms, which are the imaginary part of the above two  $(1,0)$ -forms multiplied by  $-1$ .

With this understood, consider the complex version of the spherical change of gauge. It means an extension of  $\frac{1}{r}(z_1, \dots, z_n)$  to a unitary frame for  $\mathbb{C}^n$ , which can be done on any simply connected open subset of  $\mathbb{C}^n \setminus \{0\}$ . More precisely, choose a smooth map  $T_\nu^\mu$  from a simply connected open subset of  $\mathbb{C}^n \setminus \{0\}$  to  $U(n)$  such that

$$T_\mu^1(z) = \frac{1}{r} z_\mu \quad \text{for } \mu = 1, 2, \dots, n \quad \text{and } z \in \text{Domain}(T_\mu^\nu) \subset \mathbb{C}^n \setminus \{0\}.$$

It follows that the following 1-forms also constitute a unitary coframe of type (1, 0) for (3.2)

$$\begin{aligned} \sigma^\mu &= T_\nu^\mu \theta^\nu, \\ \sigma^{n+\mu} &= \overline{T_\nu^\mu} (dz_\nu - z_\nu \theta_\nu^\gamma) \end{aligned} \quad (3.4)$$

where  $\mu \in \{1, \dots, n\}$ .

3.1.2. *The Calabi metric.* In terms of this coframe, the Calabi metric is of the following form

$$|c(r) \sigma^1|^2 + \sum_{j=2}^n |b(r) \sigma^j|^2 + (h(r) dr)^2 + \left( \frac{f(r)}{r} \text{Im } \sigma^{n+1} \right)^2 + \sum_{j=2}^n \left| \frac{a(r)}{r} \sigma^{n+j} \right|^2, \quad (3.5)$$

where

$$\begin{aligned} a &= \sinh(r), & c &= h = \sqrt{\cosh(2r)}, \\ b &= \cosh(r), & f &= \frac{1}{2} \frac{\sinh(2r)}{\sqrt{\cosh(2r)}}. \end{aligned} \quad (3.6)$$

In [8], Dancer and Swann found an easy way to construct the hyper-Kähler metric. They wrote down the ansätze for the three hyper-Kähler forms, and imposed the d-closed condition. The three Kähler forms are

$$\frac{hf}{r} dr \wedge \text{Im } \sigma^{n+1} + \frac{i}{2} c^2 \sigma^1 \wedge \overline{\sigma^1} + \frac{i}{2} b^2 \sum_{j=2}^n \sigma^j \wedge \overline{\sigma^j} + \frac{i}{2} \frac{a^2}{r^2} \sum_{j=2}^n \sigma^{n+j} \wedge \overline{\sigma^{n+j}}, \quad (3.7)$$

and the imaginary and real part of

$$hc dr \wedge \sigma^1 + \frac{ifc}{r} (\text{Im } \sigma^{n+1}) \wedge \sigma^1 - \frac{ab}{r} \sum_{j=2}^n \sigma^j \wedge \sigma^{n+j}. \quad (3.8)$$

If (3.7) and (3.8) are annihilated by the exterior derivative, the coefficient functions must obey

$$\begin{aligned} \frac{da^2}{dr} &= 2hf = \frac{db^2}{dr}, & \frac{dc^2}{dr} &= 4hf, & a^2 + b^2 &= c^2, \\ \frac{d(ab)}{dr} &= hc, & \frac{d(fc)}{dr} &= hc, & \text{and } ab &= fc. \end{aligned} \quad (3.9)$$

It is a straightforward computation to check that (3.6) does solve (3.9). One can consult [6, section 4] for the discussion on solving (3.9).

For the connection and curvature computation in the following subsections, it is more transparent to simplify the expressions by using the hyper-Kähler equation (3.9) than by plugging in the explicit solution (3.6). Note that (3.9) implies that  $b^2 - a^2$  is a constant, which is 1 for the explicit solution (3.6).

3.1.3. *Connection 1-forms.* To compute the connection 1-forms of the metric, it is easier to choose a complex structure and take a unitary frame. With respect to the complex structure corresponding to the Kähler form (3.7), we have the unitary coframe:

$$\xi^1 = c \sigma^1, \quad \xi^j = b \sigma^j, \quad \xi^{n+1} = h dr + \frac{if}{r} \operatorname{Im} \sigma^{n+1}, \quad \xi^{n+j} = \frac{a}{r} \sigma^{n+j} \quad (3.10)$$

for  $j \in \{2, \dots, n\}$ . Denote by  $\xi_\nu^\mu$  the Hermitian connection 1-forms. They can be found by the structure equation (3.1):

$$\begin{aligned} \xi_1^1 &= i \left( \frac{2f}{c^2} - \frac{1}{f} \right) \operatorname{Im} \xi^{n+1} = -\xi_{n+1}^{n+1}, & \xi_{n+1}^j &= \frac{f}{b^2} \xi^j = \xi_{n+j}^1, \\ \xi_{n+1}^1 &= \frac{2f}{c^2} \xi^1, & \xi_{n+k}^j &= 0, & \xi_{n+1}^{n+j} &= \frac{f}{a^2} \xi^{n+j} = -\xi_j^1. \end{aligned} \quad (3.11)$$

The hyper-Kähler equation (3.9) is used to simplify the above expressions. The components  $\xi_j^k$  and  $\xi_{n+j}^{n+k}$  are related to the connection of the Fubini–Study metric as follows:

$$\xi_j^k = -\xi_{n+k}^{n+j} = T_\mu^k \theta_\nu^\mu \bar{T}_\nu^j - (dT_\mu^k) \bar{T}_\mu^j + i \frac{f}{b^2} \delta_j^k \operatorname{Im} \xi^{n+1}. \quad (3.12)$$

3.1.4. *Riemann curvature tensor.* In terms of the unitary coframe (3.10), the curvatures are as follows:

$$\begin{aligned} \Omega_1^1 &= \frac{2}{c^6} \left( \xi^1 \wedge \bar{\xi}^1 - \xi^{n+1} \wedge \bar{\xi}^{n+1} \right) + \frac{1}{c^4} \left( \xi^j \wedge \bar{\xi}^j - \xi^{n+j} \wedge \bar{\xi}^{n+j} \right) = -\Omega_{n+1}^{n+1}, \\ \Omega_1^j &= \frac{1}{c^4} \left( \xi^j \wedge \bar{\xi}^1 - \xi^{n+1} \wedge \bar{\xi}^{n+j} \right) = -\Omega_{n+j}^{n+1}, \\ \Omega_1^{n+j} &= -\frac{1}{c^4} \left( \xi^{n+j} \wedge \bar{\xi}^1 + \xi^{n+1} \wedge \bar{\xi}^j \right) = \Omega_j^{n+1}, \\ \Omega_1^{n+1} &= -\frac{2}{c^6} \xi^{n+1} \wedge \bar{\xi}^1, \\ \Omega_j^{n+k} &= -\frac{1}{c^2} \left( \xi^{n+j} \wedge \bar{\xi}^k + \xi^{n+k} \wedge \bar{\xi}^j \right), \\ \Omega_j^k &= \frac{1}{c^2} \left( \xi^k \wedge \bar{\xi}^j - \xi^{n+j} \wedge \bar{\xi}^{n+k} \right) \\ &\quad + \delta_j^k \left( \frac{1}{c^4} \left( \xi^1 \wedge \bar{\xi}^1 - \xi^{n+1} \wedge \bar{\xi}^{n+1} \right) + \frac{1}{c^2} \left( \xi^i \wedge \bar{\xi}^i - \xi^{n+i} \wedge \bar{\xi}^{n+i} \right) \right) = -\Omega_{n+k}^{n+j}. \end{aligned}$$

The equality between different curvature components is a consequence of the hyper-Kähler geometry.

## 4. GEOMETRY OF LOCAL $G_2$ AND LOCAL $\text{Spin}(7)$

In [1], Bryant and Salamon constructed complete manifolds with special holonomy. They constructed three examples with holonomy  $G_2$ , and one example with holonomy  $\text{Spin}(7)$ , each of which is the total space of a vector bundle.

**4.1. Bryant–Salamon manifolds.** This subsection is a brief review on the construction of Bryant and Salamon. We first review the general framework of the metric construction on a vector bundle, and then specialize in their examples.

**4.1.1. Bundle construction.** Let  $(B^n, \underline{g})$  be a Riemannian manifold, and  $E \xrightarrow{\pi} B$  be a rank  $m$  vector bundle. Suppose that  $E$  carries a bundle metric and a metric connection. Then, these datum naturally induce a Riemannian metric on the total space  $E$ . The construction goes as follows. With the connection, the tangent space of  $E$  decomposes into vertical and horizontal subspaces. These two subspaces are defined to be orthogonal to each other. The metric on the vertical subspace is given by the original bundle metric; the metric on the horizontal subspace is the pull-back of the metric  $\underline{g}$ .

This metric can be seen explicitly in terms of the moving frame. Take a local orthonormal coframe  $\{\underline{\omega}^j\}_{j=1}^n$  on an open subset  $U \subset B$ , and a local orthonormal basis of sections  $\{\mathfrak{s}_\nu\}_{\nu=1}^m$  that trivializes  $E|_U$ . Denote by  $\nabla_A$  a metric connection for  $E$ . Let  $A_\nu^\mu$  be the the connection 1-forms of  $\nabla_A$  with respect to  $\mathfrak{s}_\nu$ , namely,  $\nabla_A \mathfrak{s}_\nu = \sum_{\mu=1}^m A_\nu^\mu \mathfrak{s}_\mu$ . Thus,  $[A_\nu^\mu]$  is an  $\mathfrak{o}(m)$ -valued 1-forms on  $U$ . Let  $\{y^\mu\}_{\mu=1}^m$  be the coordinate for the fibers of  $E|_U$  induced by  $\{\mathfrak{s}_\mu\}_{\mu=1}^m$ . The Riemannian metric  $g_b$  on  $E$  induced by the bundle metric is

$$g_b = \sum_{j=1}^n (\underline{\omega}^j)^2 + \sum_{\mu=1}^m (dy^\mu + A_\nu^\mu y^\nu)^2. \quad (4.1)$$

In our discussion on the bundle construction, the indices  $i, j, k$  are assumed to belong to  $\{1, 2, \dots, n = \dim B\}$ , and the indices  $\mu, \nu, \gamma, \sigma$  are assumed to belong to  $\{1, 2, \dots, m = \text{rank } E\}$ .

Here is a relation that will be used later. The exterior derivative of  $dy^\mu + A_\nu^\mu y^\nu$  can be written as  $F_\nu^\mu y^\nu - A_\nu^\mu \wedge (dy^\nu + A_\gamma^\nu y^\gamma)$ , where

$$F_\nu^\mu = dA_\nu^\mu + A_\gamma^\mu \wedge A_\nu^\gamma = \frac{1}{2} F_{\nu ij}^\mu \underline{\omega}^i \wedge \underline{\omega}^j \quad (4.2)$$

is the curvature of  $\nabla_A$ .

**4.1.2. Rescaling the metric.** Let  $s = \sum_{\mu} (y^\mu)^2$  be the distance square to the zero section with respect to the Riemannian metric  $g_b$  in (4.1). For any two smooth, positive functions  $\alpha(s), \beta(s)$  defined for  $s \geq 0$ ,

$$g_{\alpha, \beta} = \sum_j (\alpha \underline{\omega}^j)^2 + \sum_{\mu} (\beta (dy^\mu + A_\nu^\mu y^\nu))^2 \quad (4.3)$$

also defines a Riemannian metric on  $E$ . Let

$$\omega^j = \alpha \underline{\omega}^j \quad \text{and} \quad \omega^{n+\mu} = \beta (dy^\mu + A_\nu^\mu y^\nu) . \quad (4.4)$$

They form an orthonormal coframe of the metric (4.3). Their exterior derivatives read

$$\begin{aligned} d\omega^j &= -\underline{\omega}_i^j \wedge \omega^i - \frac{2\alpha'}{\alpha\beta} y^\mu \omega^j \wedge \omega^{n+\mu} , \\ d\omega^{n+\mu} &= \beta F_\nu^\mu y^\nu - A_\nu^\mu \wedge \omega^{n+\nu} - \frac{2\beta'}{\beta^2} y^\nu \omega^{n+\mu} \wedge \omega^{n+\nu} \end{aligned} \quad (4.5)$$

where  $\underline{\omega}_i^j$  is the connection 1-form of the Levi-Civita connection of  $(B, \underline{g})$ . By a direct computation, we find that the connection 1-forms of the Levi-Civita connection of  $(E, g_{\alpha,\beta})$  are

$$\omega_i^j = \underline{\omega}_i^j + \frac{\beta}{2\alpha^2} F_{\nu ij}^\mu y^\nu \omega^{n+\mu} , \quad (4.6)$$

$$\omega_i^{n+\mu} = \frac{\beta}{2\alpha^2} F_{\nu ij}^\mu y^\nu \omega^j - \frac{2\alpha'}{\alpha\beta} y^\mu \omega^i , \quad (4.7)$$

$$\omega_{n+\nu}^{n+\mu} = A_\nu^\mu + \frac{2\beta'}{\beta^2} (y^\nu \omega^{n+\mu} - y^\mu \omega^{n+\nu}) . \quad (4.8)$$

The mixed component  $\omega_i^{n+\mu}$  is different from the other two components; it involves only the curvature but not the connection. That is to say,  $\omega_i^{n+\mu}$  is a tensor:

$$\omega^i \otimes \omega_i^{n+\mu} \otimes \mathfrak{s}_\mu = \beta F(\cdot, \cdot)(y^\mu \mathfrak{s}_\mu) - \frac{2\alpha\alpha'}{\beta} \underline{g}(\cdot, \cdot)(y^\mu \mathfrak{s}_\mu) : \mathcal{H} \times \mathcal{H} \rightarrow \mathcal{V}$$

where  $\mathcal{H} \cong \pi^*TB$  is the horizontal subspace, and  $\mathcal{V} \equiv \pi^*E$  is the vertical subspace over the total space of  $E$ .

**4.1.3. Examples of Bryant and Salamon.** For the examples of Bryant and Salamon, the base manifold  $B$  is either a sphere or a complex projective space. The metric  $\underline{g}$  is the standard metric. The vector bundle  $E$  is constructed from the tangent bundle or the spinor bundle, and the metric connection is induced from the Levi-Civita connection. In what follows,  $\kappa$  is the sectional curvature of the round metric when the base is the sphere, and is half the holomorphic sectional curvature of the Fubini-Study metric when the base is  $\mathbb{C}\mathbb{P}^2$ .

The first example [1, p.840] is the spinor bundle over the 3-sphere,  $\mathbb{S}(S^3)$ . The  $G_2$  metric has

$$\alpha(s) = (3\kappa)^{\frac{1}{2}}(1+s)^{\frac{1}{3}} \quad \text{and} \quad \beta(s) = 2(1+s)^{-\frac{1}{6}} . \quad (4.9)$$

The next two examples [1, p.844] are the bundle of anti-self-dual 2-forms over the 4-sphere and the 2-dimensional complex projective space,  $\Lambda_-^2(S^4)$  and  $\Lambda_-^2(\mathbb{C}\mathbb{P}^2)$ . They have

$$\alpha(s) = (2\kappa)^{\frac{1}{2}}(1+s)^{\frac{1}{4}} \quad \text{and} \quad \beta(s) = (1+s)^{-\frac{1}{4}} . \quad (4.10)$$

The last example [1, p.847] is the spinor bundle of negative chirality over the 4-sphere,  $\mathbb{S}_-(S^4)$ . The  $\text{Spin}(7)$  metric has

$$\alpha(s) = (5\kappa)^{\frac{1}{2}}(1+s)^{\frac{3}{10}} \quad \text{and} \quad \beta(s) = 2(1+s)^{-\frac{1}{5}}. \quad (4.11)$$

We will refer  $\mathbb{S}(S^3)$ ,  $\Lambda_-^2(S^4)$ ,  $\Lambda_-^2(\mathbb{CP}^2)$  and  $\mathbb{S}_-(S^4)$  with the Ricci flat metric as the *Bryant–Salamon manifolds*.

The above coefficient functions are derived from the special holonomy equation, which is a first order elliptic system. According to their paper, the special holonomy equation reduces to the following equation:

$$\alpha' = \kappa_1 \frac{\beta^2}{\alpha} \quad \text{and} \quad \beta' = -\kappa_2 \frac{\beta^3}{\alpha^2} \quad (4.12)$$

for some *positive* constants  $\kappa_1$  and  $\kappa_2$ . For  $\mathbb{S}(S^3)$ ,  $\kappa_1 = \kappa/4$  and  $\kappa_2 = \kappa/8$ . For  $\Lambda_-^2(S^4)$  and  $\Lambda_-^2(\mathbb{CP}^2)$ ,  $\kappa_1 = \kappa_2 = \kappa/2$ . For  $\mathbb{S}_-(S^4)$ ,  $\kappa_1 = 3\kappa/8$  and  $\kappa_2 = \kappa/4$ .

One can easily construct some functional equations from (4.12). Here are two relations that will be used later:

$$\left(\frac{\alpha^2}{\beta^2}\right)' = 2(\kappa_1 + \kappa_2) \quad \Rightarrow \quad \frac{\alpha^2}{\beta^2} = \left(\frac{\alpha(0)}{\beta(0)}\right)^2 + 2(\kappa_1 + \kappa_2)s; \quad (4.13)$$

$$\left(\frac{\beta}{\alpha^2}\right)' = -(2\kappa_1 + \kappa_2) \frac{\beta^3}{\alpha^4}. \quad (4.14)$$

## 5. THE UNIQUENESS OF THE ZERO SECTION

We first prove a general lemma regarding the location of a compact minimal submanifold.

**Lemma 5.1.** *Let  $(M, g)$  be a Riemannian manifold. Suppose that  $\psi$  is a smooth function on  $M$  whose Hessian is non-negative definite. Then, any compact, minimal submanifold of  $M$  must be contained in where  $\text{Hess}_M \psi$  degenerates. In addition,  $\psi$  takes constant value on the submanifold if it is connected.*

*Proof.* Let  $\Sigma \subset M$  be a compact submanifold, and  $p$  be a point in  $\Sigma$ . Choose an orthonormal frame  $\{e_j\}$  for  $T\Sigma$  on some neighborhood of  $p$  in  $\Sigma$ . Consider the trace of the Hessian of  $\psi$  on  $T\Sigma$ :

$$\begin{aligned} 0 \leq \text{tr}_\Sigma \text{Hess } \psi &= \sum_j \text{Hess } \psi(e_j, e_j) = \sum_j (e_j(e_j(\psi)) - (\nabla_{e_j} e_j)(\psi)) \\ &= \Delta^\Sigma \psi - H(\psi) \end{aligned}$$

where  $\nabla$  is the covariant derivative of  $(M, g)$ ,  $\Delta^\Sigma$  is the Laplace–Beltrami operator of the induced metric on  $\Sigma$ , and  $H = \sum_j (\nabla_{e_j} e_j)^\perp$  is the mean curvature vector of  $\Sigma$ . When  $\Sigma$  is minimal, it implies that  $\Delta^\Sigma \psi \geq 0$ . The lemma follows from the compactness of  $\Sigma$  and the maximum principle.  $\square$

**5.1. The Stenzel metric case.** In this subsection, we apply Lemma 5.1 to show that the zero section is the only compact, special Lagrangian submanifold in  $T^*S^n$ .

**Theorem 5.2.** *When  $n > 1$ , any compact, minimal submanifold in  $T^*S^n$  with the Stenzel metric must belong to the zero section.*

*Proof.* Consider the smooth function  $\psi = \rho^2$ , which is the square of the distance to the zero section (with respect to the Stenzel metric). Due to Lemma 5.1, it suffices to show that the Hessian of  $\psi$  is positive definite outside the zero section. Since  $\text{Hess}(\rho) = \nabla d\rho$  and  $d\rho = \omega^{n+1}$ , we compute

$$\text{Hess}(\psi) = 2\omega^{n+1} \otimes \omega^{n+1} + 2\rho \nabla \omega^{n+1} .$$

By the third equation in (6.6), we obtain

$$\text{Hess}(\psi) = 2\omega^{n+1} \otimes \omega^{n+1} - 2\rho(n-1)C\omega^1 \otimes \omega^1 - 2\rho A \sum_{j=2}^n \omega^j \otimes \omega^j - 2\rho B \sum_{j=2}^n \omega^{n+j} \otimes \omega^{n+j} . \quad (5.1)$$

According to Lemma 2.1,  $\text{Hess}(\psi)$  is positive definite when  $r > 0$  (equivalently, when  $\rho > 0$ ). This finishes the proof of the theorem.  $\square$

**5.2. The Calabi metric case.** In this subsection, we prove that the zero section is the only compact, special Lagrangian<sup>2</sup> submanifold in  $T^*\mathbb{C}\mathbb{P}^n$ .

**Theorem 5.3.** *Any compact, minimal submanifold in  $T^*\mathbb{C}\mathbb{P}^n$  with the Calabi metric must belong to the zero section.*

*Proof.* The function  $\rho = \int_0^r \sqrt{\cosh(2u)} du$  is the distance function to the zero section with respect to the Calabi metric and  $d\rho = \text{Re} \xi^{n+1}$  by (3.6) and (3.10). Consider the smooth function  $\psi = \rho^2$  and compute as in the last theorem,

$$\text{Hess}(\psi) = 2(\text{Re} \xi^{n+1}) \otimes (\text{Re} \xi^{n+1}) + 2\rho \text{Re}(\nabla \xi^{n+1}) .$$

By the third equation in (6.12),

$$\text{Re}(\nabla \xi^{n+1}) = \frac{2f}{c^2} |\xi^1|^2 + \frac{f}{b^2} \sum_{j=2}^n |\xi^j|^2 + \left(\frac{1}{f} - \frac{2f}{c^2}\right) (\text{Im} \xi^{n+1}) \otimes (\text{Im} \xi^{n+1}) + \frac{f}{a^2} \sum_{j=2}^n |\xi^{n+j}|^2 .$$

According (3.6), it is not hard to see that  $\text{Hess}(\psi)$  is positive definite when  $r > 0$ , and the theorem follows from Lemma 5.1.  $\square$

---

<sup>2</sup>The zero section is Lagrangian with respect to any Kähler structure orthogonal to (3.7). Equivalently, it is holomorphic with respect to the complex structure corresponding to the Kähler form (3.7).

**5.3. The Bryant–Salamon metric case.** In this subsection, we examine the uniqueness of the zero section as a minimal submanifold.

**Lemma 5.4.** *Let  $(B^n, g)$  be a Riemannian manifold. Let  $E \rightarrow B$  be a rank  $m$  vector bundle with a bundle metric and a metric connection. Denote by  $s$  the square of the distance to the zero section with respect to the metric  $g_b$  (4.1) on  $E$ . For any two smooth positive functions  $\alpha(s)$  and  $\beta(s)$ , endow  $E$  the Riemannian metric  $g_{\alpha, \beta}$  defined by (4.3). Then, the Hessian of  $s$  with respect to  $g_{\alpha, \beta}$  is positive definite outside the zero section if and only if  $\alpha' > 0$  and  $\beta > 2s|\beta'|$  for  $s > 0$ .*

*Proof.* Suppose that  $\psi$  is a smooth function on  $E$  depending only on  $s = \sum_{\mu} (y^{\mu})^2$ . Its exterior derivative is

$$d\psi = \psi' ds = 2\psi' y^{\mu} dy^{\mu} = \frac{2\psi'}{\beta} y^{\mu} \omega^{n+\mu} . \quad (5.2)$$

Let  $\{\bar{e}_j\}_{j=1}^n \cup \{\bar{e}_{n+\mu}\}_{\mu=1}^m$  be the frame dual to the coframe (4.4). The Hessian of  $\psi$  along  $(\bar{e}_i, \bar{e}_j)$  is

$$\begin{aligned} \text{Hess}(\psi)(\bar{e}_i, \bar{e}_j) &= \bar{e}_i(\bar{e}_j(\psi)) - (\nabla_{\bar{e}_i} \bar{e}_j)(\psi) \\ &= \frac{4\alpha'\psi'}{\alpha^2\beta} s \delta_i^j + \frac{\psi'}{\alpha^2} y^{\mu} y^{\nu} F_{\nu}^{\mu}{}_{ij} = \frac{4\alpha'\psi'}{\alpha^2\beta} s \delta_i^j \end{aligned} \quad (5.3)$$

where equality uses the fact that  $[F_{\nu}^{\mu}]$  is skew-symmetric. It is not hard to see that the Hessian of  $\psi$  along  $(\bar{e}_{n+\mu}, \bar{e}_j)$  vanishes. Along  $(\bar{e}_{n+\mu}, \bar{e}_{n+\nu})$ ,

$$\begin{aligned} \text{Hess}(\psi)(\bar{e}_{n+\mu}, \bar{e}_{n+\nu}) &= \bar{e}_{n+\mu}(\bar{e}_{n+\nu}(\psi)) - (\nabla_{\bar{e}_{n+\mu}} \bar{e}_{n+\nu})(\psi) \\ &= \left( \frac{2\psi'}{\beta^2} \delta_{\mu}^{\nu} + y^{\mu} y^{\nu} \frac{2}{\beta} \left( \frac{2\psi'}{\beta} \right)' \right) - \left( y^{\mu} y^{\nu} \frac{4\beta'\psi'}{\beta^3} - \frac{4\beta'\psi'}{\beta^3} \rho \delta_{\mu}^{\nu} \right) \\ &= \left( \frac{2}{\beta^2} + \frac{4\beta'}{\beta^3} s \right) \psi' \delta_{\mu}^{\nu} + \left( \frac{4\psi'}{\beta^2} \right)' y^{\mu} y^{\nu} . \end{aligned} \quad (5.4)$$

Substituting  $\psi = s$ , the lemma follows from (5.3), (5.4) and the fact that the eigenvalues of  $[y^{\mu} y^{\nu}]$  are  $s$  and 0, where 0 has geometric multiplicity  $k - 1$ .  $\square$

Applying this lemma to the Bryant–Salamon manifolds leads to the following theorem.

**Theorem 5.5.** *Any compact minimal submanifold of the Bryant–Salamon manifolds must be contained in the zero section.*

*Proof.* This follows directly from (4.12), (4.13) and Lemma 5.4 with  $\psi = s$ . Moreover, by (4.13), we have

$$\text{Hess}(\psi) \geq 4\kappa_1 \frac{\beta}{\alpha^3} s \sum_{i=1}^n \omega^i \otimes \omega^i + \frac{2}{\alpha^2} \left( \left( \frac{\alpha(0)}{\beta(0)} \right)^2 + 2\kappa_1 s \right) \sum_{\mu=1}^m \omega^{n+\mu} \otimes \omega^{n+\mu} . \quad (5.5)$$



This Hessian estimate will be needed later. □

## 6. FURTHER ESTIMATES NEEDED FOR THE STABILITY THEOREM

In this section, we begin preparations for the proof of Theorem 1.3. Each manifold in Assumption 1.1 is the total space of a vector bundle  $\pi : E \rightarrow B$ . The base  $B$  naturally sits inside  $E$  as the zero section. The pull-back of the volume form on the zero section defines a differential form  $\Omega$  on  $E$ . We are going to calculate the covariant derivatives of  $\Omega$ , which will be applied to derive  $\mathcal{C}^1$  estimate of the mean curvature flow.

**6.1. Estimates from linear-algebraic decomposition.** Each metric in Assumption 1.1 admits a local orthonormal coframe  $\{\omega^j\}_{j=1}^n \cup \{\omega^{n+\mu}\}_{\mu=1}^m$  such that  $\bigcap_{j=1}^n \ker \omega^j = \pi^*E$ . As in section 4.1.1, The subbundle  $\pi^*E \subset TE$  will be called the *vertical subspace*, and will be denoted by  $\mathcal{V}$ . The orthogonal subbundle  $\mathcal{H} \subset TE$  is given by  $\bigcap_{\mu=1}^m \ker \omega^{n+\mu}$ , and is isomorphic to  $\pi^*TB$ . This bundle  $\mathcal{H}$  will be referred as the *horizontal subspace*. For each manifold in Assumption 1.1, the dimension of the base manifold,  $n$ , is no less than the rank of the vector bundle,  $m$ .

In terms of the frame, the  $n$ -form  $\Omega$  is  $\omega^1 \wedge \omega^2 \wedge \cdots \wedge \omega^n$ . Let  $p \in E$ , and suppose that  $L \subset T_pE$  is an oriented  $n$ -dimensional subspace with  $\Omega(L) > 0$ . Then,  $L$  can be regarded as the graph of a linear map from  $\mathcal{H}_p$  to  $\mathcal{V}_p$ . By the singular value decomposition, there exist orthonormal bases  $\{u_j\}_{j=1}^n$  for  $\mathcal{H}_p$ ,  $\{v_\mu\}_{\mu=1}^m$  for  $\mathcal{V}_p$ , and angles  $\theta_j \in [0, \pi)$  such that

$$\{e_j = \cos \theta_j u_j + \sin \theta_j v_j\}_{j=1}^n \quad \text{and} \quad \{e_{n+\mu} = -\sin \theta_\mu u_\mu + \cos \theta_\mu v_\mu\}_{\mu=1}^m \quad (6.1)$$

constitute an orthonormal basis for  $L$  and  $L^\perp$ , respectively. For  $j > m$ ,  $v_j$  is set to be the zero vector, and  $\theta_j$  is set to be zero.

Note that neither the frame  $\{e_j\} \cup \{e_{n+\mu}\}$  nor  $\{u_j\} \cup \{v_\mu\}$  is necessarily dual to  $\{\omega^j\} \cup \{\omega^{n+\mu}\}$ . In any event,  $[\omega^j(u_i)]_{i,j}$  is an  $n \times n$  (special) orthogonal matrix, and  $[\omega^{n+\mu}(v_\nu)]_{\nu,\mu}$  is a  $m \times m$  orthogonal matrix. Denote by

$$\mathfrak{s} = \max_j |\sin \theta_j|. \quad (6.2)$$

The following estimates are straightforward to come by:

$$\sum_{i=1}^n |(\omega^j \otimes \omega^k)(e_i, e_i)| \leq n, \quad \sum_{i=1}^n |(\omega^{n+\mu} \wedge \omega^j)(e_i, e_i)| \leq 2n\mathfrak{s} \quad (6.3)$$

and

$$\begin{aligned}
& \left| (\omega^1 \wedge \cdots \wedge \omega^n)(e_{n+\mu}, e_1, \dots, \widehat{e}_i, \dots, e_n) \right| \leq \mathfrak{s} , \\
& \left| (\omega^1 \wedge \cdots \wedge \omega^n)(e_{n+\mu}, e_{n+\nu}, e_1, \dots, \widehat{e}_i, \dots, \widehat{e}_j, \dots, e_n) \right| \leq \mathfrak{s}^2 , \\
& \left| (\omega^{n+\mu} \wedge \omega^1 \wedge \cdots \wedge \widehat{\omega}^i \wedge \cdots \wedge \omega^n)(e_1, \dots, e_n) \right| \leq n\mathfrak{s} , \\
& \left| (\omega^{n+\mu} \wedge \omega^1 \wedge \cdots \wedge \widehat{\omega}^i \wedge \cdots \wedge \omega^n)(e_{n+\nu}, e_1, \dots, \widehat{e}_j, \dots, e_n) \right| \leq 1 , \\
& \left| (\omega^{n+\mu} \wedge \omega^{n+\nu} \wedge \omega^1 \wedge \cdots \wedge \widehat{\omega}^i \wedge \cdots \wedge \widehat{\omega}^j \wedge \cdots \wedge \omega^n)(e_1, \dots, e_n) \right| \leq n(n-1)\mathfrak{s}^2
\end{aligned} \tag{6.4}$$

for any  $i, j, k \in \{1, \dots, n\}$  and  $\mu, \nu \in \{1, \dots, m\}$ .

Suppose that  $\Sigma \subset E$  is an oriented,  $n$ -dimensional submanifold with  $\Omega(T_p\Sigma) > 0$ . Applying the above construction to  $T_p\Sigma$  gives a continuous function  $\mathfrak{s}$  on  $\Sigma$ . With this understanding, the remainder of this section is devoted to estimating

$$\nabla_{e_j}\Omega \quad \text{and} \quad (\text{tr}_{T_p\Sigma} \nabla^2\Omega)(T_p\Sigma) = \sum_{j=1}^n (\nabla_{e_j}^2\Omega)(e_1, \dots, e_n)$$

in terms of  $\mathfrak{s}$  and the distance to the zero section. These estimates will be used in the proof of Theorem 1.3

**6.2. The Stenzel metric case.** Consider the Stenzel metric on  $T^*S^n$ , and adopt the notations introduced in section 2. The  $n$ -form  $\Omega = \omega^1 \wedge \omega^2 \wedge \cdots \wedge \omega^n$  is not parallel. In order to establish the estimates on  $\nabla\Omega$  and  $\nabla^2\Omega$ , it is convenient to introduce the following notations:

$$\begin{aligned}
\Phi &= \omega^2 \wedge \cdots \wedge \omega^n , \\
\Phi^j &= \iota(e_j)\Phi = (-1)^j \omega^2 \wedge \cdots \wedge \widehat{\omega}^j \wedge \cdots \wedge \omega^n , \\
\Phi^{jk} &= \iota(e_k)\iota(e_j)\Phi = \begin{cases} (-1)^{j+k} \omega^2 \wedge \cdots \wedge \widehat{\omega}^k \wedge \cdots \wedge \widehat{\omega}^j \wedge \cdots \wedge \omega^n & \text{if } k < j , \\ (-1)^{j+k+1} \omega^2 \wedge \cdots \wedge \widehat{\omega}^j \wedge \cdots \wedge \widehat{\omega}^k \wedge \cdots \wedge \omega^n & \text{if } k > j \end{cases}
\end{aligned} \tag{6.5}$$

for any  $j, k \in \{2, \dots, n\}$ .

By (2.7), (2.9) and (2.10), the covariant derivatives of the coframe 1-forms are as follows.

$$\begin{aligned}
\nabla\omega^1 &= -B\omega^{n+j} \otimes \omega^j + (n-1)C\omega^1 \otimes \omega^{n+1} + A\omega^j \otimes \omega^{n+j} , \\
\nabla\omega^j &= -\theta_k^j \otimes \omega^k + B\omega^{n+j} \otimes \omega^1 + A\omega^j \otimes \omega^{n+1} - C\omega^1 \otimes \omega^{n+j} , \\
\nabla\omega^{n+1} &= -(n-1)C\omega^1 \otimes \omega^1 - A\omega^j \otimes \omega^j - B\omega^{n+j} \otimes \omega^{n+j} , \\
\nabla\omega^{n+j} &= -A\omega^j \otimes \omega^1 + C\omega^1 \otimes \omega^j + B\omega^{n+j} \otimes \omega^{n+1} - \theta_k^j \otimes \omega^{n+k} .
\end{aligned} \tag{6.6}$$

We compute the covariant derivative of  $\Phi$  and  $\Phi^j$ .

$$\begin{aligned}\nabla\Phi &= (\nabla\omega^j) \wedge \Phi^j \\ &= (B\omega^{n+j}) \otimes (\omega^1 \wedge \Phi^j) - (C\omega^1) \otimes (\omega^{n+j} \wedge \Phi^j) + (A\omega^j) \otimes (\omega^{n+1} \wedge \Phi^j),\end{aligned}\quad (6.7)$$

$$\begin{aligned}\nabla\Phi^j &= (\nabla\omega^k) \wedge \Phi^{jk} \\ &= (B\omega^{n+k}) \otimes (\omega^1 \wedge \Phi^{jk}) + (A\omega^k) \otimes (\omega^{n+1} \wedge \Phi^{jk}) \\ &\quad - (C\omega^1) \otimes (\omega^{n+k} \wedge \Phi^{jk}) + \theta_j^k \otimes \Phi^k.\end{aligned}\quad (6.8)$$

Putting (6.6) and (6.7) together gives the covariant derivative of  $\Omega$ .

$$\begin{aligned}\nabla\Omega &= (n-1)(C\omega^1) \otimes (\omega^{n+1} \wedge \Phi) + (A\omega^j) \otimes (\omega^{n+j} \wedge \Phi) \\ &\quad - (C\omega^1) \otimes (\omega^1 \wedge \omega^{n+j} \wedge \Phi^j) + (A\omega^j) \otimes (\omega^1 \wedge \omega^{n+1} \wedge \Phi^j).\end{aligned}\quad (6.9)$$

We also compute the second covariant derivative of  $\Omega$  by computing the covariant derivative of the four terms on the right hand side of (6.9). By (6.6), (6.7) and (6.8):

$$\nabla^2\Omega = (n-1)\text{I} + \text{II} + \text{III} + \text{IV}$$

where

$$\begin{aligned}\text{I} &= \nabla((C\omega^1) \otimes (\omega^{n+1} \wedge \Phi)) \\ &= -(n-1)(C^2\omega^1 \otimes \omega^1) \otimes \Omega - (BC\omega^{n+j} \otimes \omega^1) \otimes (\omega^{n+j} \wedge \Phi + \omega^1 \wedge \omega^{n+1} \wedge \Phi^j) \\ &\quad + \left(\dot{C}\omega^{n+1} \otimes \omega^1 - BC\omega^{n+j} \otimes \omega^j + (n-1)C^2\omega^1 \otimes \omega^{n+1} + AC\omega^j \otimes \omega^{n+j}\right) \otimes (\omega^{n+1} \wedge \Phi) \\ &\quad - (C^2\omega^1 \otimes \omega^1) \otimes (\omega^{n+1} \wedge \omega^{n+j} \wedge \Phi^j),\end{aligned}$$

$$\begin{aligned}\text{II} &= \nabla((A\omega^j) \otimes (\omega^{n+j} \wedge \Phi)) \\ &= -(A^2\omega^j \otimes \omega^j) \otimes \Omega + (AB\omega^{n+j} \otimes \omega^j) \otimes (\omega^{n+1} \wedge \Phi) + (AB\omega^{n+k} \otimes \omega^j) \otimes (\omega^{n+j} \wedge \omega^1 \wedge \Phi^k) \\ &\quad + \left(\dot{A}\omega^{n+1} \otimes \omega^j + A^2\omega^j \otimes \omega^{n+1} + AB\omega^{n+j} \otimes \omega^1 - AC\omega^1 \otimes \omega^{n+j}\right) \otimes (\omega^{n+j} \wedge \Phi) \\ &\quad - (AC\omega^1 \otimes \omega^j) \otimes (\omega^{n+j} \wedge \omega^{n+k} \wedge \Phi^k) + (A^2\omega^k \otimes \omega^j) \otimes (\omega^{n+j} \wedge \omega^{n+1} \wedge \Phi^k),\end{aligned}$$

$$\begin{aligned}\text{III} &= -\nabla((C\omega^1) \otimes (\omega^1 \wedge \omega^{n+j} \wedge \Phi^j)) \\ &= -(C^2\omega^1 \otimes \omega^1) \otimes \Omega - (C^2\omega^1 \otimes \omega^1) \otimes \left((n-1)\omega^{n+1} \wedge \omega^{n+j} \wedge \Phi^j + \omega^1 \wedge \omega^{n+j} \wedge \omega^{n+k} \wedge \Phi^{jk}\right) \\ &\quad - \left(\dot{C}\omega^{n+1} \otimes \omega^1 - BC\omega^{n+j} \otimes \omega^j + (n-1)C^2\omega^1 \otimes \omega^{n+1} + AC\omega^j \otimes \omega^{n+j}\right) \otimes (\omega^1 \wedge \omega^{n+k} \wedge \Phi^k) \\ &\quad - (AC\omega^k \otimes \omega^1) \otimes (\omega^{n+k} \wedge \omega^{n+j} \wedge \Phi^j + \omega^1 \wedge \omega^{n+j} \wedge \omega^{n+1} \wedge \Phi^{jk}) \\ &\quad - (BC\omega^{n+j} \otimes \omega^1) \otimes (\omega^{n+j} \wedge \Phi + \omega^1 \wedge \omega^{n+1} \wedge \Phi^j),\end{aligned}$$

$$\begin{aligned}
\text{IV} &= \nabla \left( (A\omega^j) \otimes (\omega^1 \wedge \omega^{n+1} \wedge \Phi^j) \right) \\
&= -(A^2 \omega^j \otimes \omega^j) \otimes \Omega + (A^2 \omega^k \otimes \omega^j) \otimes (\omega^{n+k} \wedge \omega^{n+1} \wedge \Phi^j) \\
&\quad + \left( \dot{A} \omega^{n+1} \otimes \omega^j + A^2 \omega^j \otimes \omega^{n+1} + AB \omega^{n+j} \otimes \omega^1 - AC \omega^1 \otimes \omega^{n+j} \right) \otimes (\omega^1 \wedge \omega^{n+1} \wedge \Phi^j) \\
&\quad + (AB \omega^{n+j} \otimes \omega^j) \otimes (\omega^{n+1} \wedge \Phi) - (AB \omega^{n+k} \otimes \omega^j) \otimes (\omega^1 \wedge \omega^{n+k} \wedge \Phi^j) \\
&\quad - (AC \omega^1 \otimes \omega^{n+j}) \otimes (\omega^1 \wedge \omega^{n+1} \wedge \omega^{n+k} \wedge \Phi^{jk}) .
\end{aligned}$$

By examining the coefficient functions carefully, we conclude the following lemma. Recall that the  $\rho$  coordinate is the distance to the zero section.

**Lemma 6.1.** *Consider  $M = T^*S^n$  with the Stenzel metric (2.3), and consider the  $n$ -form  $\Omega = \omega^1 \wedge \cdots \wedge \omega^n$ . There exists a constant  $K > 0$  depending only on  $n$  with the following property. Suppose that  $\Sigma$  is an oriented  $n$ -dimensional submanifold of  $M$  such that  $\rho(p) < 1$  and  $\Omega(T_p\Sigma) > 0$  for any  $p \in \Sigma$ . Then,*

$$\begin{aligned}
|\nabla_X \Omega|(p) &< K\rho(p) |X| \quad \text{for any } X \in T_p M, \text{ and} \\
-K(\rho^2(p) + \mathfrak{s}^2(p)) &< (\text{tr}_{T_p\Sigma} \nabla^2 \Omega)(T_p\Sigma) < K\mathfrak{s}^2(p) - \frac{1}{K}\rho^2(p) .
\end{aligned}$$

Here,  $|\nabla_X \Omega|$  means the metric norm of  $\nabla_X \Omega$  as a section of  $(\Lambda^n T^*M)|_\Sigma$  with respect to the metric induced by the Stenzel metric.

*Proof.* In (6.9), the coefficient functions consist of multiples of  $A$  and  $C$ , which are of order  $\rho$  by the second assertion of Lemma 2.1.

There are three types of terms in the formula of  $\nabla^2 \Omega$ .

- (i) The first type is in the direction of  $\Omega$ . The first term of each of I, II, III, IV is of this type and their sum is

$$\left[ -(n^2 - 2n + 2)C^2 \omega^1 \otimes \omega^1 - 2A^2 \sum_{j=2}^n \omega^j \otimes \omega^j \right] \otimes \Omega .$$

The two-tensor in the bracket is clearly semi-negative definite, and is of order  $\rho^2$  by Lemma 2.1.

- (ii) The second one is the linear combination of

$$\eta = (\omega^{n+i} \otimes \omega^j) \otimes (\omega^{n+k} \wedge \omega^1 \wedge \cdots \wedge \widehat{\omega^l} \wedge \cdots \wedge \omega^n) \quad \text{and} \quad (\omega^i \otimes \omega^{n+j}) \otimes (\text{same})$$

for  $i, j, k, l \in \{1, \dots, n\}$ . By (6.3) and the first line of (6.4),  $|(\text{tr}_{T_p\Sigma} \eta)(T_p\Sigma)| \leq 2n\mathfrak{s}^2$ . On the other hand, the coefficient functions are constant multiples of  $AB, BC, AC, C^2, \dot{A}$  or  $\dot{C}$ . They are at most of order 1 by Lemma 2.1 and 2.2.

- (iii) The third one is the linear combination of

$$\eta = (\omega^p \otimes \omega^q) \otimes (\omega^{n+i} \wedge \omega^{n+j} \wedge \omega^1 \wedge \cdots \wedge \widehat{\omega^k} \wedge \cdots \wedge \widehat{\omega^l} \wedge \cdots \wedge \omega^n)$$

for  $i, j, k, l, p, q \in \{1, \dots, n\}$ . By (6.3) and the last line of (6.4),  $|\langle \text{tr}_{T_p \Sigma} \eta \rangle(T_p \Sigma)| \leq n^3 \mathfrak{s}^2$ . In these terms, the coefficient functions are multiples of  $A^2, AC$  and  $C^2$ . They are of order  $\rho^2$  by Lemma 2.1.

By the triangle inequality, it finishes the proof of this lemma.  $\square$

**6.3. The Calabi metric case.** For the Calabi metric, we follow the notations introduced in section 3 and consider the  $2n$ -form

$$\Omega = \left(\frac{i}{2}\right)^n (\xi^1 \wedge \xi^2 \wedge \dots \wedge \xi^n) \wedge \overline{(\xi^1 \wedge \xi^2 \wedge \dots \wedge \xi^n)}. \quad (6.10)$$

Its restriction to the zero section coincides with the volume form of the zero section. Let  $\mathbf{u}_\nu$  be the complexified tangent vector defined by  $\xi^\mu(\mathbf{u}_\nu) = \delta_\nu^\mu$  and  $\overline{\xi^\mu}(\mathbf{u}_\nu) = 0$ . Similar to the case of the Stenzel metric, let

$$\begin{aligned} \Xi &= \xi^2 \wedge \dots \wedge \xi^n, \\ \Xi^j &= \iota(\mathbf{u}_j)\Xi = (-1)^j \xi^2 \wedge \dots \wedge \widehat{\xi^j} \wedge \dots \wedge \xi^n, \\ \Xi^{jk} &= \iota(\mathbf{u}_k)\iota(\mathbf{u}_j)\Xi = \begin{cases} (-1)^{j+k} \xi^2 \wedge \dots \wedge \widehat{\xi^k} \wedge \dots \wedge \widehat{\xi^j} \wedge \dots \wedge \xi^n & \text{if } k < j, \\ (-1)^{j+k+1} \xi^2 \wedge \dots \wedge \widehat{\xi^j} \wedge \dots \wedge \widehat{\xi^k} \wedge \dots \wedge \xi^n & \text{if } k > j \end{cases} \end{aligned} \quad (6.11)$$

for any  $j, k \in \{2, \dots, n\}$ .

By (3.11) and (3.12), the covariant derivative of the unitary coframe reads:

$$\begin{aligned} \nabla \xi^1 &= -i\left(\frac{2f}{c^2} - \frac{1}{f}\right) (\text{Im } \xi^{n+1}) \otimes \xi^1 + \frac{b}{ac} \xi^{n+j} \otimes \xi^j - \frac{2f}{c^2} \xi^1 \otimes \xi^{n+1} - \frac{a}{bc} \xi^j \otimes \xi^{n+j}, \\ \nabla \xi^j &= -\frac{b}{ac} \overline{\xi^{n+j}} \otimes \xi^1 - \xi_k^j \otimes \xi^k - \frac{f}{b^2} \xi^j \otimes \xi^{n+1}, \\ \nabla \xi^{n+1} &= \frac{2f}{c^2} \overline{\xi^1} \otimes \xi^1 + \frac{f}{b^2} \overline{\xi^j} \otimes \xi^j + i\left(\frac{2f}{c^2} - \frac{1}{f}\right) (\text{Im } \xi^{n+1}) \otimes \xi^{n+1} + \frac{f}{a^2} \overline{\xi^{n+j}} \otimes \xi^{n+j}, \\ \nabla \xi^{n+j} &= \frac{a}{bc} \overline{\xi^j} \otimes \xi^1 - \frac{f}{a^2} \xi^{n+j} \otimes \xi^{n+1} + \xi_j^k \otimes \xi^{n+k}. \end{aligned} \quad (6.12)$$

It follows that the covariant derivative of  $\Xi$  and  $\Xi^j$  are as follows:

$$\begin{aligned} \nabla \Xi &= (\nabla \xi^j) \wedge \Xi^j \\ &= -\frac{b}{ac} \overline{\xi^{n+j}} \otimes (\xi^1 \wedge \Xi^j) - \xi_j^k \otimes \Xi - \frac{f}{b^2} \xi^j \otimes (\xi^{n+1} \wedge \Xi^j), \end{aligned} \quad (6.13)$$

$$\begin{aligned} \nabla \Xi^j &= (\nabla \xi^k) \wedge \Xi^{jk} \\ &= -\frac{b}{ac} \overline{\xi^{n+k}} \otimes (\xi^1 \wedge \Xi^{jk}) - \xi_k^k \otimes \Xi^j + \xi_j^k \otimes \Xi^k - \frac{f}{b^2} \xi^k \otimes (\xi^{n+1} \wedge \Xi^{jk}). \end{aligned} \quad (6.14)$$

By combining (6.12) and (6.13), the covariant derivative of  $\xi^1 \wedge \Xi = \xi^1 \wedge \xi^2 \wedge \cdots \wedge \xi^n$  is

$$\begin{aligned} \nabla(\xi^1 \wedge \Xi) &= i\left(\frac{1}{f} - \frac{2f}{c^2}\right) (\text{Im } \xi^{n+1}) \otimes (\xi^1 \wedge \Xi) - \frac{2f}{c^2} \xi^1 \otimes (\xi^{n+1} \wedge \Xi) \\ &\quad - \frac{a}{bc} \xi^j \otimes (\xi^{n+j} \wedge \Xi) - \xi_j^j \otimes (\xi^1 \wedge \Xi) - \frac{f}{b^2} \xi^j \otimes (\xi^1 \wedge \xi^{n+1} \wedge \Xi^j). \end{aligned} \quad (6.15)$$

Since  $\xi_j^k$  is skew-Hermitian,

$$\begin{aligned} \nabla\Omega &= -\frac{2f}{c^2} \xi^1 \otimes \left( \left(\frac{i}{2}\right)^n \xi^{n+1} \wedge \Xi \wedge \bar{\xi}^1 \wedge \bar{\Xi} \right) - \frac{a}{bc} \xi^j \otimes \left( \left(\frac{i}{2}\right)^n \xi^{n+j} \wedge \Xi \wedge \bar{\xi}^1 \wedge \bar{\Xi} \right) \\ &\quad - \frac{f}{b^2} \xi^j \otimes \left( \left(\frac{i}{2}\right)^n \xi^1 \wedge \xi^{n+1} \wedge \Xi^j \wedge \bar{\xi}^1 \wedge \bar{\Xi} \right) + (\text{their conjugates}). \end{aligned} \quad (6.16)$$

Note that by (3.9), the second coefficient,  $a/(bc)$ , is equal to  $f/b^2$ . The next step is to calculate the second order derivative of  $\Omega$ , which is a sum of the covariant derivative of the six terms on the right hand side of (6.16). Due to (6.12), (6.13), (6.14), (6.15) and the relations (3.9):

$$\nabla^2\Omega = -\left(\frac{i}{2}\right)^n ((\text{I} + \text{II} + \text{III}) + (-1)^n (\bar{\text{I}} + \bar{\text{II}} + \bar{\text{III}}))$$

where

$$\begin{aligned} \text{I} &= \nabla \left( \frac{2f}{c^2} \xi^1 \otimes \left( \xi^{n+1} \wedge \Xi \wedge \bar{\xi}^1 \wedge \bar{\Xi} \right) \right) \\ &= \frac{4f^2}{c^4} (\bar{\xi}^1 \otimes \xi^1) \otimes \left( \xi^1 \wedge \Xi \wedge \bar{\xi}^1 \wedge \bar{\Xi} + \xi^{n+1} \wedge \Xi \wedge \bar{\xi}^{n+1} \wedge \bar{\Xi} \right) \\ &\quad + \left( \left(\frac{2f}{c^2}\right)' \frac{1}{h} (\text{Re } \xi^{n+1}) \otimes \xi^1 + i\frac{2f}{c^2} \left(\frac{2f}{c^2} - \frac{1}{f}\right) (\text{Im } \xi^{n+1}) \otimes \xi^1 - \frac{4f^2}{c^4} \xi^1 \otimes \xi^{n+1} \right. \\ &\quad \left. + \frac{2b^2}{c^4} \xi^{n+j} \otimes \xi^j - \frac{2a^2}{c^4} \xi^j \otimes \xi^{n+j} \right) \otimes \left( \xi^{n+1} \wedge \Xi \wedge \bar{\xi}^1 \wedge \bar{\Xi} \right) \\ &\quad + \frac{2b^2}{c^4} (\bar{\xi}^{n+j} \otimes \xi^1) \otimes \left( \xi^{n+j} \wedge \Xi \wedge \bar{\xi}^1 \wedge \bar{\Xi} \right) - \frac{2a^2}{c^4} (\bar{\xi}^j \otimes \xi^1) \otimes \left( \xi^{n+1} \wedge \Xi \wedge \bar{\xi}^{n+j} \wedge \bar{\Xi} \right) \\ &\quad + \frac{2b^2}{c^4} (\bar{\xi}^{n+j} \otimes \xi^1) \otimes \left( \xi^1 \wedge \xi^{n+1} \wedge \Xi^j \wedge \bar{\xi}^1 \wedge \bar{\Xi} \right) - \frac{2a^2}{c^4} (\bar{\xi}^j \otimes \xi^1) \otimes \left( \xi^{n+1} \wedge \Xi \wedge \bar{\xi}^1 \wedge \bar{\xi}^{n+1} \wedge \bar{\Xi}^j \right), \end{aligned}$$

$$\begin{aligned}
\text{II} &= \nabla \left( \frac{f}{b^2} \xi^j \otimes \left( \xi^{n+j} \wedge \Xi \wedge \bar{\xi}^1 \wedge \bar{\Xi} \right) \right) \\
&= \frac{f^2}{b^4} (\bar{\xi}^j \otimes \xi^j) \otimes \left( \xi^1 \wedge \Xi \wedge \bar{\xi}^1 \wedge \bar{\Xi} \right) - \frac{1}{c^2} (\xi^{n+j} \otimes \xi^j) \otimes \left( \xi^{n+1} \wedge \Xi \wedge \bar{\xi}^1 \wedge \bar{\Xi} \right) \\
&\quad + \left( \left( \frac{f}{b^2} \right)' \frac{1}{h} (\text{Re } \xi^{n+1}) \otimes \xi^j + i \frac{f}{b^2} \left( \frac{2f}{c^2} - \frac{1}{f} \right) (\text{Im } \xi^{n+1}) \otimes \xi^j - \frac{1}{c^2} \overline{\xi^{n+j}} \otimes \xi^1 \right. \\
&\quad \left. - \frac{f^2}{b^4} \xi^j \otimes \xi^{n+1} \right) \otimes \left( \xi^{n+j} \wedge \Xi \wedge \bar{\xi}^1 \wedge \bar{\Xi} \right) \\
&\quad - \frac{2a^2}{c^4} (\bar{\xi}^1 \otimes \xi^j) \otimes \left( \xi^{n+j} \wedge \Xi \wedge \bar{\xi}^{n+1} \wedge \bar{\Xi} \right) - \frac{f^2}{b^4} (\bar{\xi}^k \otimes \xi^j) \otimes \left( \xi^{n+j} \wedge \Xi \wedge \bar{\xi}^{n+k} \wedge \bar{\Xi} \right) \\
&\quad + \frac{1}{c^2} (\overline{\xi^{n+k}} \otimes \xi^j) \otimes \left( \xi^1 \wedge \xi^{n+j} \wedge \Xi^k \wedge \bar{\xi}^1 \wedge \bar{\Xi} \right) + \frac{f^2}{b^4} (\xi^k \otimes \xi^j) \otimes \left( \xi^{n+1} \wedge \xi^{n+j} \wedge \Xi^k \wedge \bar{\xi}^1 \wedge \bar{\Xi} \right) \\
&\quad - \frac{f^2}{b^4} (\bar{\xi}^k \otimes \xi^j) \otimes \left( \xi^{n+j} \wedge \Xi \wedge \bar{\xi}^1 \wedge \bar{\xi}^{n+1} \wedge \bar{\Xi}^k \right) ,
\end{aligned}$$

$$\begin{aligned}
\text{III} &= \nabla \left( \frac{f}{b^2} \xi^j \otimes \left( \xi^1 \wedge \xi^{n+1} \wedge \Xi^j \wedge \bar{\xi}^1 \wedge \bar{\Xi} \right) \right) \\
&= \frac{f^2}{b^4} (\bar{\xi}^j \otimes \xi^j) \otimes \left( \xi^1 \wedge \Xi \wedge \bar{\xi}^1 \wedge \bar{\Xi} \right) - \frac{1}{c^2} (\xi^{n+j} \otimes \xi^j) \otimes \left( \xi^{n+1} \wedge \Xi \wedge \bar{\xi}^1 \wedge \bar{\Xi} \right) \\
&\quad + \left( \left( \frac{f}{b^2} \right)' \frac{1}{h} (\text{Re } \xi^{n+1}) \otimes \xi^j + i \frac{f}{b^2} \left( \frac{2f}{c^2} - \frac{1}{f} \right) (\text{Im } \xi^{n+1}) \otimes \xi^j - \frac{1}{c^2} \overline{\xi^{n+j}} \otimes \xi^1 \right. \\
&\quad \left. - \frac{f^2}{b^4} \xi^j \otimes \xi^{n+1} \right) \otimes \left( \xi^1 \wedge \xi^{n+1} \wedge \Xi^j \wedge \bar{\xi}^1 \wedge \bar{\Xi} \right) \\
&\quad + \frac{f^2}{b^4} (\xi^k \otimes \xi^j) \otimes \left( \xi^{n+1} \wedge \xi^{n+k} \wedge \Xi^j \wedge \bar{\xi}^1 \wedge \bar{\Xi} \right) + \frac{1}{c^2} (\overline{\xi^{n+k}} \otimes \xi^j) \otimes \left( \xi^1 \wedge \xi^{n+k} \wedge \Xi^j \wedge \bar{\xi}^1 \wedge \bar{\Xi} \right) \\
&\quad - \frac{2a^2}{c^4} (\bar{\xi}^1 \otimes \xi^j) \otimes \left( \xi^1 \wedge \xi^{n+1} \wedge \Xi^j \wedge \bar{\xi}^{n+1} \wedge \bar{\Xi} \right) - \frac{f^2}{b^4} (\bar{\xi}^k \otimes \xi^j) \otimes \left( \xi^1 \wedge \xi^{n+1} \wedge \Xi^j \wedge \bar{\xi}^{n+k} \wedge \bar{\Xi} \right) \\
&\quad - \frac{f^2}{b^4} (\bar{\xi}^k \otimes \xi^j) \otimes \left( \xi^1 \wedge \xi^{n+1} \wedge \Xi^j \wedge \bar{\xi}^1 \wedge \bar{\xi}^{n+1} \wedge \bar{\Xi}^k \right) .
\end{aligned}$$

Recall that the distance to the zero section with respect to the Calabi metric is  $\rho = \int_0^r \sqrt{\cosh(2u)} du$ . With the help of (6.3), (6.4) and the coefficients (3.6), a completely parallel argument as that in the proof of Lemma 6.1 leads to the following lemma.

**Lemma 6.2.** *Consider  $M = T^*\mathbb{C}\mathbb{P}^n$  with the Calabi metric (3.5), and consider the 2n-form  $\Omega$  defined by (6.10). There exists a constant  $K > 0$  depending only on  $n$  which has the following property. Suppose that  $\Sigma$  is an oriented 2n-dimensional submanifold of  $M$  such that  $\rho(p) < 1$  and  $\Omega(T_p\Sigma) > 0$  for any  $p \in \Sigma$ . Then,*

$$\begin{aligned}
|\nabla_X \Omega| &< K \rho(p) |X| \quad \text{for any } X \in T_p M , \text{ and} \\
-K(\rho^2(p) + \mathfrak{s}^2(p)) &< (\text{tr}_{T_p\Sigma} \nabla^2 \Omega)(T_p\Sigma) < K \mathfrak{s}^2(p) - \frac{1}{K} \rho^2(p) .
\end{aligned}$$

**6.4. The Bryant–Salamon metric case.** Consider the  $n$ -form  $\Omega = \omega^1 \wedge \omega^2 \wedge \cdots \wedge \omega^n$ . The notations in this subsection follow those in section 4. Similar to the case of the Stenzel metric, it is convenient to introduce the following shorthand notations:

$$\Omega^j = \iota(e_j)\Omega = (-1)^{j+1}\omega^1 \wedge \cdots \wedge \widehat{\omega^j} \wedge \cdots \wedge \omega^n, \quad (6.17)$$

$$\Omega^{jk} = \iota(e_k)\iota(e_j)\Omega = \begin{cases} (-1)^{j+k}\omega^1 \wedge \cdots \wedge \widehat{\omega^k} \wedge \cdots \wedge \widehat{\omega^j} \wedge \cdots \wedge \omega^n & \text{if } k < j, \\ (-1)^{j+k+1}\omega^1 \wedge \cdots \wedge \widehat{\omega^j} \wedge \cdots \wedge \widehat{\omega^k} \wedge \cdots \wedge \omega^n & \text{if } k > j. \end{cases} \quad (6.18)$$

The covariant derivative of  $\Omega$  is

$$\nabla\Omega = (\nabla\omega^j) \wedge \Omega^j = \omega_j^{n+\mu} \otimes (\omega^{n+\mu} \wedge \Omega^j). \quad (6.19)$$

For  $\nabla^2\Omega$ , one expects that the covariant derivative of the curvature,  $\nabla_A F \in \mathcal{C}^\infty(B; T^*B \otimes (\Lambda^2 T^*B \otimes \text{End } E))$ , will show up. The coefficient 1-form of  $\nabla_A F$  is

$$(\nabla_A F)_{\nu jk}^\mu = dF_{\nu jk}^\mu + A_\gamma^\mu F_{\nu jk}^\gamma - F_{\gamma jk}^\mu A_\nu^\gamma - F_{\nu ik}^\mu \underline{\omega}_j^i - F_{\nu ji}^\mu \underline{\omega}_k^i. \quad (6.20)$$

We are now ready to calculate the second order derivative of  $\Omega$ . Since

$$\nabla\omega^{n+\mu} = -\omega_i^{n+\mu} \otimes \omega^i - \omega_{n+\nu}^{n+\mu} \otimes \omega^{n+\nu} \quad \text{and} \quad (6.21)$$

$$\nabla\Omega^j = (\nabla\omega^k) \wedge \Omega^{jk} = \omega_j^k \otimes \Omega^k + \omega_k^{n+\nu} \otimes (\omega^{n+\nu} \wedge \Omega^{jk}), \quad (6.22)$$

the covariant derivative of (6.19) is

$$\begin{aligned} \nabla^2\Omega &= -(\omega_i^{n+\mu} \otimes \omega_i^{n+\mu}) \otimes \Omega + (\omega_k^{n+\nu} \otimes \omega_j^{n+\mu}) \otimes (\omega^{n+\mu} \wedge \omega^{n+\nu} \wedge \Omega^{jk}) \\ &\quad + \left( \nabla\omega_j^{n+\mu} + \omega_{n+\nu}^{n+\mu} \otimes \omega_j^{n+\nu} + \omega_k^j \otimes \omega_k^{n+\mu} \right) \otimes (\omega^{n+\mu} \wedge \Omega^j). \end{aligned} \quad (6.23)$$

The first two coefficients on the right hand side of (6.23) can be substituted by (4.7). We compute the coefficient in the second line of (6.23). By (4.7),

$$\nabla\omega_j^{n+\mu} = \nabla \left( \frac{\beta}{2\alpha^2} F_{\nu jk}^\mu y^\nu \omega^k \right) - \nabla \left( \frac{\alpha'}{r\alpha\beta} y^\mu \omega^j \right).$$

With the help of (4.4), (4.6), (5.2) and (6.21), we have

$$\begin{aligned} \nabla \left( \frac{\beta}{2\alpha^2} F_{\nu jk}^\mu y^\nu \omega^k \right) &= \frac{1}{r\beta} \left( \frac{\beta}{2\alpha^2} \right)' F_{\nu jk}^\mu y^\nu y^\gamma \omega^{n+\gamma} \otimes \omega^k + \frac{1}{2\alpha^2} F_{\nu jk}^\mu \omega^{n+\nu} \otimes \omega^k \\ &\quad + \frac{\beta}{2\alpha^2} F_{\nu jk}^\mu y^\nu \omega_k^{n+\gamma} \otimes \omega^{n+\gamma} + \frac{\beta^2}{4\alpha^2} (F_{\nu jk}^\mu y^\nu) (F_{\gamma ik}^\sigma y^\gamma) \omega^{n+\sigma} \otimes \omega^i \\ &\quad + \frac{\beta}{2\alpha^2} y^\nu \left( dF_{\nu jk}^\mu - F_{\gamma jk}^\mu A_\nu^\gamma - F_{\nu ji}^\mu \underline{\omega}_k^i \right) \otimes \omega^k, \end{aligned} \quad (6.24)$$



and

$$\begin{aligned}
-\nabla \left( \frac{\alpha'}{r\alpha\beta} y^\mu \omega^j \right) &= -\frac{1}{r\beta} \left( \frac{\alpha'}{r\alpha\beta} \right)' y^\mu y^\nu \omega^{n+\nu} \otimes \omega^j - \frac{\alpha'}{r\alpha\beta^2} \omega^{n+\mu} \otimes \omega^j \\
&+ \frac{\alpha'}{2r\alpha^2} y^\mu F_\nu^\gamma y^\nu \omega^{n+\gamma} \otimes \omega^k - \frac{\alpha'}{r\alpha\beta} y^\mu \omega_j^{n+\nu} \otimes \omega^{n+\nu} \\
&+ \frac{\alpha'}{r\alpha\beta} y^\gamma A_\gamma^\mu \otimes \omega^j + \frac{\alpha'}{r\alpha\beta} y^\mu \underline{\omega}_k^j \otimes \omega^k .
\end{aligned} \tag{6.25}$$

Due to (4.6) (4.7) and (4.8),

$$\begin{aligned}
\omega_{n+\nu}^{n+\mu} \otimes \omega_j^{n+\nu} &= \frac{\beta'}{r\beta^2} (y^\nu \omega^{n+\mu} - y^\mu \omega^{n+\nu}) \otimes \omega_j^{n+\nu} \\
&- \frac{\alpha'}{r\alpha\beta} y^\nu A_\nu^\mu \otimes \omega^j + \frac{\beta}{2\alpha^2} y^\nu (A_\gamma^\mu F_\nu^\gamma y_{jk}) \otimes \omega^k ,
\end{aligned} \tag{6.26}$$

and

$$\begin{aligned}
-\omega_j^k \otimes \omega_k^{n+\mu} &= -\frac{\beta}{2\alpha^2} F_\nu^\gamma y^\nu \omega^{n+\gamma} \otimes \omega_k^{n+\mu} \\
&+ \frac{\alpha'}{r\alpha\beta} y^\mu \underline{\omega}_j^k \otimes \omega^k - \frac{\beta^2}{2\alpha^2} y^\nu (F_\nu^\mu \underline{\omega}_j^i) \otimes \omega^k .
\end{aligned} \tag{6.27}$$

To summarize the above computations, note that

- the sum of the last terms of (6.24), (6.26) and (6.27) is a multiple of  $\nabla_A F$  by (6.20);
- the last term of (6.25) cancels with the second-last term of (6.27);
- the second-last term of (6.25) cancels with that of (6.26).

The above computation is for a general bundle construction. We now examine the expressions for the Bryant-Salamon metric. It is more convenient to use the function  $s = \sum_\mu (y^\mu)^2$ . On any compact region, it is equivalent to the distance square to the zero section.

**Lemma 6.3.** *Consider the  $n$ -form  $\Omega = \omega^1 \wedge \dots \wedge \omega^n$  on the Bryant-Salamon manifolds. There exists a constant  $K > 0$  which has the following property. Suppose that  $\Sigma$  is an oriented  $n$ -dimensional submanifold of  $M$  such that  $s(p) < 1$  and  $\Omega(T_p\Sigma) > 0$  for any  $p \in \Sigma$ . Then,*

$$\begin{aligned}
|\nabla_X \Omega| &< K \sqrt{s(p)} |X| \quad \text{for any } X \in T_p M , \text{ and} \\
-K (s(p) + \mathfrak{s}^2(p)) &< (\text{tr}_{T_p\Sigma} \nabla^2 \Omega)(T_p\Sigma) < K \mathfrak{s}^2(p) - \frac{1}{K} s(p) .
\end{aligned}$$

*Proof.* The coefficient functions  $\alpha$  and  $\beta$  have explicit expressions, (4.9), (4.10) and (4.11). The estimate on  $\nabla_X \Omega$  follows directly from (4.7) and (6.19).

For  $\nabla^2 \Omega$ , observe that:

- The first coefficient 2-tensor on the right hand side of (6.23) is non-positive definite, and in the order of  $s$  when  $s(p) < 1$ .
- The second term on the right hand side of (6.23) carries  $\omega^{n+\mu} \wedge \omega^{n+\nu} \wedge \dots$ .

- As explained in Appendix A.1,  $\nabla_A F \equiv 0$ . Thus, each term of the third coefficient 2-tensor on the right hand side of (6.23) carries at least one  $\omega^{n+\mu}$ -codirection.

Then, the lemma follows from (6.3) and (6.4).  $\square$

## 7. THE STABILITY OF ZERO SECTIONS

Suppose  $\Sigma_t$  is a mean curvature flow of  $n$ -dimensional compact submanifolds in an ambient Riemannian manifold  $M$  and  $\Omega$  is an  $n$ -form on  $M$ . For any point  $p \in \Sigma_t$ , let  $\{e_1, \dots, e_n\}$  be an orthonormal frame of  $T\Sigma_t$  near  $p$  and  $\{e_{n+1}, \dots, e_{n+m}\}$  be an orthonormal frame of the normal bundle of  $\Sigma_t$  near  $p$ . In the following, the indexes  $i, j, k$  range from 1 to  $n$ , the indexes  $\alpha, \beta, \gamma$  range from  $n+1$  to  $n+m$ , and repeated indexes are summed.  $h_{\alpha ij} = \langle \nabla_{e_i} e_j, e_\alpha \rangle$  denote the coefficients of the second fundamental form of  $\Sigma_t$ . Here,  $\nabla$  is the Levi-Civita connection of the ambient manifold  $M$ .

We first recall the following proposition from [25, Proposition 3.1]

**Proposition 7.1.** *Along the mean curvature flow  $\Sigma_t$  in  $M$ ,  $*\Omega = \Omega(e_1, \dots, e_n)$  satisfies*

$$\begin{aligned} \frac{d}{dt} * \Omega &= \Delta^{\Sigma_t} * \Omega + * \Omega \left( \sum_{\alpha, i, k} h_{\alpha ik}^2 \right) \\ &\quad - 2 \sum_{\alpha, \beta, k} [\Omega_{\alpha\beta 3 \dots n} h_{\alpha 1k} h_{\beta 2k} + \Omega_{\alpha 2\beta \dots n} h_{\alpha 1k} h_{\beta 3k} + \dots + \Omega_{1 \dots (n-2)\alpha\beta} h_{\alpha(n-1)k} h_{\beta nk}] \\ &\quad - 2(\nabla_{e_k} \Omega)(e_\alpha, \dots, e_n) h_{\alpha 1k} - \dots - 2(\nabla_{e_k} \Omega)(e_1, \dots, e_\alpha) h_{\alpha nk} \\ &\quad - \sum_{\alpha, k} [\Omega_{\alpha 2 \dots n} R_{\alpha k k 1} + \dots + \Omega_{1 \dots (n-1)\alpha} R_{\alpha k k n}] - (\nabla_{e_k, e_k}^2 \Omega)(e_1, \dots, e_n) \end{aligned} \quad (7.1)$$

where  $\Delta^{\Sigma_t}$  denotes the time-dependent Laplacian on  $\Sigma_t$ ,  $\Omega_{\alpha\beta 3 \dots n} = \Omega(e_\alpha, e_\beta, e_3, \dots, e_n)$  etc., and  $R_{\alpha k k 1} = R(e_\alpha, e_k, e_k, e_1)$ , etc. are the coefficients of the curvature operators of  $M$ .

When  $\Omega$  is a parallel form in  $M$ ,  $\nabla \Omega \equiv 0$ , this recovers an important formula in proving the long time existence result of the graphical mean curvature flow in [24].

**Remark 7.2.** In [25], the frame  $\{e_k\}_{k=1}^n$  is the geodesic frame at some  $p \in M$ , i.e.  $\nabla_{e_j}^\Sigma e_i$  vanishes at  $p$ . Thus, the last term of (7.1) is

$$(\nabla_{e_k, e_k}^2 \Omega)(e_1, \dots, e_n) = -(\nabla_{e_k} \nabla_{e_k} \Omega)(e_1, \dots, e_n) + (\nabla_H \Omega)(e_1, \dots, e_n)$$

at  $p \in M$ . This is exactly the formula in [25, Proposition 3.1].

### 7.1. Proof of Theorem 1.3 for the Stenzel metric.

*Proof.* We deal with the Stenzel metric first. Let  $\epsilon$  be the constant to be determined and  $\Sigma$  be a compact submanifold of  $M$  that satisfies the assumption (1.1). Throughout the proof,

$K_i, i = 0, 1, 2, \dots$  denotes a positive constant that depends only on the dimension  $n$ . Denote by  $\Sigma_t$  the mean curvature flow in  $M$  with  $\Sigma$  as the initial data.

We first prove the  $\mathcal{C}^0$  estimate. As in the proof of Theorem 5.2, let  $\psi$  be the distance square to the zero section with respect to the Stenzel metric, or the square of the  $\rho$  coordinate. Its evolution equation along the mean curvature flow  $\Sigma_t$  reads (see the proof of Theorem C in [23])

$$\frac{d}{dt}\psi = \Delta^{\Sigma_t}\psi - \text{tr}_{\Sigma_t} \text{Hess } \psi, \quad (7.2)$$

where  $\text{tr}_{\Sigma_t} \text{Hess } \psi$  is the trace of the Hessian of  $\psi$  over  $\Sigma_t$  and is always non-negative by Theorem 5.2. By the maximum principle, the maximum of  $\psi$  on  $\Sigma_t$  is non-increasing, and thus  $\Sigma_t$  remains close to the zero section along the flow.

Next, we derive the  $\mathcal{C}^1$  estimate which amounts to showing that  $*\Omega = \Omega(T_p\Sigma_t)$  remains close to one. We claim that there exists a small enough  $\epsilon \in (0, \frac{1}{2})$  and a large enough  $K_0 > 1$ , both depending only on the dimension  $n$ , such that if the inequality  $*\Omega - K_0\psi > 1 - \epsilon$  holds on the initial data, then it remains true on  $\Sigma_t$  at any subsequent time as long as the flow exists smoothly.

For the argument of contradiction, suppose  $*\Omega - K_0\psi > 1 - \epsilon$  holds initially, and  $*\Omega - K_0\psi = 1 - \epsilon$  for the first time at  $T_0$ . Our goal is to apply equation (7.1) and the estimates in section 6.2 to show that for certain  $\epsilon$  and  $K_0$ ,

$$\frac{d}{dt}(*\Omega - K_0\psi) \geq \Delta^{\Sigma_t}(*\Omega - K_0\psi) + \frac{1}{2}(*\Omega - K_0\psi)|\mathcal{A}|^2. \quad (7.3)$$

where  $|\mathcal{A}|^2 = \sum_{\alpha,i,k} h_{\alpha ik}^2$ . Therefore, the minimum of  $*\Omega - K_0\psi$  is non-decreasing in the interval  $[0, T_0)$  and  $*\Omega - K_0\psi$  is indeed strictly greater than  $1 - \epsilon$  at  $T_0$ .

By (7.1) and (7.2), the evolution equation of  $*\Omega - K_0\psi$  is:

$$\begin{aligned} & \frac{d}{dt}(*\Omega - K_0\psi) - \Delta^{\Sigma_t}(*\Omega - K_0\psi) \\ &= *\Omega \left( \sum_{\alpha,i,k} h_{\alpha ik}^2 \right) - 2 \sum_{\alpha,\beta,k} [\Omega_{\alpha\beta 3\dots n} h_{\alpha 1k} h_{\beta 2k} + \Omega_{\alpha 2\beta\dots n} h_{\alpha 1k} h_{\beta 3k} + \dots + \Omega_{1\dots(n-2)\alpha\beta} h_{\alpha(n-1)k} h_{\beta nk}] \\ & \quad - 2(\nabla_{e_k}\Omega)(e_\alpha, \dots, e_n)h_{\alpha 1k} - \dots - 2(\nabla_{e_k}\Omega)(e_1, \dots, e_\alpha)h_{\alpha nk} \\ & \quad - \sum_{\alpha,k} [\Omega_{\alpha 2\dots n} R_{\alpha k k 1} + \dots + \Omega_{1\dots(n-1)\alpha} R_{\alpha k k n}] - (\nabla_{e_k, e_k}^2 \Omega)(e_1, \dots, e_n) + K_0 \text{tr}_{\Sigma_t} \text{Hess } \psi. \end{aligned} \quad (7.4)$$

We aim at using the first and last term on the right hand side of (7.4) to control the rest terms. At any  $p \in \Sigma_t$ ,  $T_p\Sigma$  and  $(T_p\Sigma)^\perp$  have the following orthonormal basis constructed in section 6.1:

$$\{e_j = \cos \theta_j u_j + \sin \theta_j v_j\} \quad \text{and} \quad \{e_{n+j} = -\sin \theta_j u_j + \cos \theta_j v_j\},$$

respectively. Recall that  $\{u_j, v_j\}_{j=1, \dots, n}$ , is an orthonormal basis for  $T_p M$  such that  $\omega^i(v_j) = \omega^{n+i}(u_j) = 0$  for any  $i, j$ , and  $[\omega^i(u_j)]_{i,j}$  and  $[\omega^{n+i}(v_j)]_{i,j}$  are both  $n \times n$  orthogonal matrices. As in (6.2), set  $\mathfrak{s}$  to be  $\max_j |\sin \theta_j|$ .

It follows from  $*\Omega - K_0\psi > 1 - \epsilon$  that  $\psi = \rho^2 < \epsilon/K_0 < 1$ . In what follows, the point  $p$  is always assumed to have distance less than 1 from the zero section. It also follows from  $*\Omega - K_0\psi > 1 - \epsilon$  that  $*\Omega = \Omega(e_1, \dots, e_n) = \prod_{j=1}^n \cos \theta_j > 1 - \epsilon$ . Hence,  $\cos \theta_j > 1 - \epsilon > \frac{1}{2}$  and  $\sin^2 \theta_j < 2\epsilon$  for any  $j \in \{1, \dots, n\}$ , and  $\mathfrak{s} < 2\epsilon^{\frac{1}{2}}$ .

We now analyze the terms on the right hand side of (7.4)

(i) By (5.1) and Lemma 2.1,  $\text{Hess } \psi \geq \frac{1}{K_1} \sum_{j=1}^n (\rho^2 \omega^j \otimes \omega^j + \omega^{n+j} \otimes \omega^{n+j})$  for some constant  $K_1 > 0$ . It follows that

$$\text{tr}_{\Sigma_t} \text{Hess } \psi \geq \frac{1}{K_1} (\rho^2 \sum_{j=1}^n \cos^2 \theta_j + \sum_{j=1}^n \sin^2 \theta_j) > \frac{1}{K_1} (\frac{1}{4} \rho^2 + \mathfrak{s}^2). \quad (7.5)$$

(ii) With the second line of (6.4) and the Cauchy–Schwarz inequality,  $|\Omega_{\alpha\beta 3 \dots n} h_{\alpha 1k} h_{\beta 2k}|$  is bounded by  $\mathfrak{s}^2 |\mathcal{A}|^2$ . Thus,

$$\left| 2 \sum_{\alpha, \beta, k} (\Omega_{\alpha\beta 3 \dots n} h_{\alpha 1k} h_{\beta 2k} + \dots + \Omega_{1 \dots (n-2) \alpha \beta} h_{\alpha (n-1)k} h_{\beta nk}) \right| \leq K_2 \mathfrak{s}^2 |\mathcal{A}|^2$$

for some constant  $K_2 > 0$ .

(iii) Due to the first assertion of Lemma 6.1,

$$2 |(\nabla_{e_k} \Omega)(e_\alpha, \dots, e_n) h_{\alpha 1k} + \dots + (\nabla_{e_k} \Omega)(e_1, \dots, e_\alpha) h_{\alpha nk}| < K_3 \rho |\mathcal{A}| \leq K_3^2 \rho^2 + \frac{1}{4} |\mathcal{A}|^2$$

for some constant  $K_3 > 0$ .

(iv) According to the curvature computation in section 2.2.1, the Riemann curvature tensor of the Stenzel metric satisfies  $R(u_i, v_j, v_k, v_l) = 0 = R(v_i, u_j, u_k, u_l)$  for any  $i, j, k, l \in \{1, \dots, n\}$ . It follows that

$$|R(e_{n+j}, e_k, e_k, e_i)| \leq K_4 (|\sin \theta_j| + |\sin \theta_k| + |\sin \theta_i|) \leq 3K_4 \mathfrak{s}$$

for some constant  $K_4 > 0$ . This together with the first line of (6.4) implies that

$$\left| \sum_{\alpha, k} (\Omega_{\alpha 2 \dots n} R_{\alpha k k 1} + \dots + \Omega_{1 \dots (n-1) \alpha} R_{\alpha k k n}) \right| \leq 3n^3 K_4 \mathfrak{s}^2.$$

(v) By Lemma 6.1, there exists a constant  $K_5 > 0$  such that

$$-\sum_{k=1}^n (\nabla_{e_k, e_k}^2 \Omega)(e_1, \dots, e_n) \geq -K_5 \mathfrak{s}^2.$$

It follows that the right hand side of (7.4) is greater than

$$*\Omega|\mathcal{A}|^2 + \frac{K_0}{K_1}\left(\frac{1}{4}\rho^2 + \mathfrak{s}^2\right) - K_2\mathfrak{s}^2|\mathcal{A}|^2 - K_3^2\rho^2 - \frac{1}{4}|\mathcal{A}|^2 - (3n^3K_4 + K_5)\mathfrak{s}^2 .$$

It is clear that by taking  $\epsilon$  to be sufficiently small and  $K_0$  to be sufficiently large, the expression is greater than  $\frac{1}{2}*\Omega|\mathcal{A}|^2 \geq \frac{1}{2}(*\Omega - K_0\psi)|\mathcal{A}|^2$ . This proves the differential inequality (7.3).

Finally, we prove the  $\mathcal{C}^2$  estimate, which amounts to bounding the norm of the second fundamental form from above. The evolution equation for the norm of the second fundamental form for a mean curvature flow in a general Riemannian manifold is derived in [22, Proposition 7.1]. In particular,  $|\mathcal{A}|^2 = \sum_{\alpha,i,k} h_{\alpha ik}^2$  satisfies the following equation along the flow:

$$\begin{aligned} \frac{d}{dt}|\mathcal{A}|^2 &= \Delta_{\Sigma_t}|\mathcal{A}|^2 - 2|\nabla^{\Sigma_t}\mathcal{A}|^2 + 2[(\nabla_{e_k}R)_{\alpha ijk} + (\nabla_{e_j}R)_{\alpha kik}]h_{\alpha ij} \\ &\quad - 4R_{lij k}h_{\alpha lk}h_{\alpha ij} + 8R_{\alpha\beta jk}h_{\beta ik}h_{\alpha ij} - 4R_{lkik}h_{\alpha lj}h_{\alpha ij} + 2R_{\alpha k\beta k}h_{\beta ij}h_{\alpha ij} \\ &\quad + 2 \sum_{\alpha,\gamma,i,m} \left( \sum_k h_{\alpha ik}h_{\gamma mk} - h_{\alpha mk}h_{\gamma ik} \right)^2 + 2 \sum_{i,j,m,k} \left( \sum_{\alpha} h_{\alpha ij}h_{\alpha mk} \right)^2 . \end{aligned} \quad (7.6)$$

In particular, we have

$$\frac{d}{dt}|\mathcal{A}|^2 \leq \Delta^{\Sigma_t}|\mathcal{A}|^2 - 2|\nabla^{\Sigma_t}\mathcal{A}|^2 + K_6|\mathcal{A}|^4 + K_7|\mathcal{A}|^2 ,$$

where  $K_6, K_7$  are positive constant that only depend on the geometry of  $M$ . Combining this with equation (7.4) and applying the method in [24, p.534–542] yield the boundedness of the second fundamental form. Standard estimates for second order quasilinear parabolic system imply all higher derivatives are bounded, and we can apply Simon's convergence theorem [18] to conclude the smooth convergence as  $t \rightarrow \infty$ . It follows from (7.5) and (7.2) that  $\psi|_{\Sigma_t}$  converges exponentially to zero. Similarly, it follows from (7.3) that  $(1 - *\Omega + K_0\psi)|_{\Sigma_t}$  converges exponentially to zero, and thus  $*\Omega$  converges exponentially to 1.  $\square$

**7.2. Proof of Theorem 1.3 for the other cases.** The proof for the Calabi metric and the Bryant–Salamon metric is almost the same as above. We just highlight where needs to be modified. For the Calabi metric, the function  $\psi$  is taken to be the distance square to the zero section,  $(\int_0^r \sqrt{\cosh(2u)}du)^2$ . For the Bryant–Salamon metric, the function  $\psi$  is taken to be  $s = \sum_{\mu}(y^{\mu})^2$ .

First of all, due to Theorem 5.3 and 5.5, the  $\mathcal{C}^0$  estimate follows from the same argument. Moreover, as can be seen in their proofs, there exists a constant  $K_8 > 0$  such that

$$\text{Hess}(\psi) > \frac{1}{K_8}(\psi + \mathfrak{s}^2)$$

provided  $\psi \leq 1$ . This is item (i) in the proof of the  $\mathcal{C}^1$  estimate. For the rest items,

- one simply has to replace Lemma 6.1 by Lemma 6.2 and 6.3, respectively;
- according to the curvature computation in section 3.1.4 and appendix A.2, their Riemann curvatures also admit the property that  $R(\mathcal{H}, \mathcal{V}, \mathcal{V}, \mathcal{V}) = 0 = R(\mathcal{V}, \mathcal{H}, \mathcal{H}, \mathcal{H})$ .

For  $\mathcal{C}^2$  and all higher derivative estimates, the argument is completely the same.

## APPENDIX A. CURVATURE OF BRYANT–SALAMON MANIFOLDS

This appendix is a brief summary on the curvature properties of the Bryant–Salamon manifolds. For these bundle manifolds, the isometry group of the base acts transitively, and the action lifts the total space of the vector bundle isometrically. Hence, it suffices to examine the curvature at a particular fiber.

Fix a point  $p$  in the base, and let  $\underline{\omega}^j$  be the geodesic frame at  $p$ . Thus, the Levi-Civita connection of the base metric vanishes,  $\underline{\omega}_i^j|_p = 0$ . Since the bundle is either the spinor bundle or the bundle of anti-self-dual 2-forms,  $\{\underline{\omega}^j\}$  induces an orthonormal trivialization  $\{\mathfrak{s}_\nu\}$  of the bundle by representation theory. In other words, the bundle connection  $A_\nu^\mu$  is a linear combination of  $\underline{\omega}_i^j$ , and also vanishes at  $p$ . It follows that  $(d\underline{\omega}_i^j)|_p = \underline{\Omega}_i^j|_p$  and  $(dA_\nu^\mu)|_p = F_\nu^\mu|_p$ .

**A.1. The bundle curvature of the connection.** In order to use (4.6), (4.7) and (4.8) to compute the curvature of (4.3) over the fiber at  $p$ , we also need to know the exterior derivative of  $F_\nu^\mu$ . They are locally defined functions on the base. Since the metric here is the round metric or the Fubini-Study metric, these locally defined functions are actually locally constants. The readers are directed to [1] for the detail of the curvature computation. We simply write down the answer.

For  $\mathbb{S}(S^3)$ ,

$$F = \frac{\kappa}{2} \begin{bmatrix} 0 & -\underline{\omega}^2 \wedge \underline{\omega}^3 & \underline{\omega}^1 \wedge \underline{\omega}^3 & -\underline{\omega}^1 \wedge \underline{\omega}^2 \\ \underline{\omega}^2 \wedge \underline{\omega}^3 & 0 & \underline{\omega}^1 \wedge \underline{\omega}^2 & \underline{\omega}^1 \wedge \underline{\omega}^3 \\ -\underline{\omega}^1 \wedge \underline{\omega}^3 & -\underline{\omega}^1 \wedge \underline{\omega}^2 & 0 & \underline{\omega}^2 \wedge \underline{\omega}^3 \\ \underline{\omega}^1 \wedge \underline{\omega}^2 & -\underline{\omega}^1 \wedge \underline{\omega}^3 & -\underline{\omega}^2 \wedge \underline{\omega}^3 & 0 \end{bmatrix}. \quad (\text{A.1})$$

It is understood as an endomorphism-valued 2-form with respect to the trivialization  $\{\mathfrak{s}_\nu\}$ . The components of the curvature can be read from the matrix. For instance,  $F_{413}^2 = \kappa/2$ .

For  $\Lambda_-^2(S^4)$  and  $\Lambda_-^2(\mathbb{C}\mathbb{P}^2)$ ,

$$F = \kappa \begin{bmatrix} 0 & -\underline{\omega}^1 \wedge \underline{\omega}^4 + \underline{\omega}^2 \wedge \underline{\omega}^3 & \underline{\omega}^1 \wedge \underline{\omega}^3 + \underline{\omega}^2 \wedge \underline{\omega}^4 \\ \underline{\omega}^1 \wedge \underline{\omega}^4 - \underline{\omega}^2 \wedge \underline{\omega}^3 & 0 & -\underline{\omega}^1 \wedge \underline{\omega}^2 + \underline{\omega}^3 \wedge \underline{\omega}^4 \\ -\underline{\omega}^1 \wedge \underline{\omega}^3 - \underline{\omega}^2 \wedge \underline{\omega}^4 & \underline{\omega}^1 \wedge \underline{\omega}^2 - \underline{\omega}^3 \wedge \underline{\omega}^4 & 0 \end{bmatrix}. \quad (\text{A.2})$$

For  $\mathbb{S}_-(S^4)$ ,

$$F = \frac{\kappa}{2} \begin{bmatrix} 0 & \underline{\omega}^1 \wedge \underline{\omega}^2 - \underline{\omega}^3 \wedge \underline{\omega}^4 & \underline{\omega}^1 \wedge \underline{\omega}^3 + \underline{\omega}^2 \wedge \underline{\omega}^4 & \underline{\omega}^1 \wedge \underline{\omega}^4 - \underline{\omega}^2 \wedge \underline{\omega}^3 \\ -\underline{\omega}^1 \wedge \underline{\omega}^2 + \underline{\omega}^3 \wedge \underline{\omega}^4 & 0 & -\underline{\omega}^1 \wedge \underline{\omega}^4 + \underline{\omega}^2 \wedge \underline{\omega}^3 & \underline{\omega}^1 \wedge \underline{\omega}^3 + \underline{\omega}^2 \wedge \underline{\omega}^4 \\ -\underline{\omega}^1 \wedge \underline{\omega}^3 - \underline{\omega}^2 \wedge \underline{\omega}^4 & \underline{\omega}^1 \wedge \underline{\omega}^4 - \underline{\omega}^2 \wedge \underline{\omega}^3 & 0 & -\underline{\omega}^1 \wedge \underline{\omega}^2 + \underline{\omega}^3 \wedge \underline{\omega}^4 \\ -\underline{\omega}^1 \wedge \underline{\omega}^4 + \underline{\omega}^2 \wedge \underline{\omega}^3 & -\underline{\omega}^1 \wedge \underline{\omega}^3 - \underline{\omega}^2 \wedge \underline{\omega}^4 & \underline{\omega}^1 \wedge \underline{\omega}^2 - \underline{\omega}^3 \wedge \underline{\omega}^4 & 0 \end{bmatrix}. \quad (\text{A.3})$$

By (6.20), it is clear that  $\nabla_A F$  vanishes at  $p$ . Since the argument applies to any  $p \in B$ ,  $\nabla_A F \equiv 0$ .

**A.2. The Riemann curvature tensor of the bundle metric.** We are now ready to compute the curvature of Bryant–Salamon metric (4.3) over the fiber at  $p$ . Abuse the notation by still denoting  $|_p$ . The Levi-Civita connection of (4.3) is discussed in section 4.

By (4.6), (4.5), (4.12) and (4.14),

$$\begin{aligned} (d\omega_i^j)|_p &= \underline{\Omega}_i^j - \frac{\beta^2}{4\alpha^4} y^\nu F_{\mu ij}^\nu F_{\gamma kl}^\mu y^\gamma \omega^k \wedge \omega^l + \frac{1}{2\alpha^2} F_{\mu ij}^\mu \omega^{n+\nu} \wedge \omega^{n+\mu} \\ &\quad - \frac{2\beta^2}{4\alpha^4} (\kappa_1 + \kappa_2) F_{\nu ij}^\mu y^\nu y^\gamma \omega^{n+\gamma} \wedge \omega^{n+\mu}, \\ (-\omega_i^k \wedge \omega_k^j)|_p &= \frac{\beta^2}{4\alpha^4} F_{\nu ik}^\mu y^\nu \omega^{n+\mu} \wedge F_{\gamma jk}^\eta y^\gamma \omega^{n+\eta}, \\ (-\omega_i^{n+\mu} \wedge \omega_{n+\mu}^j)|_p &= \frac{4\beta^2}{\alpha^4} \kappa_1^2 \rho \omega^i \wedge \omega^j - \frac{\beta^2}{4\alpha^4} y^\nu F_{\mu ik}^\nu F_{\gamma jl}^\mu y^\gamma \omega^k \wedge \omega^l, \end{aligned}$$

and thus

$$\begin{aligned} R_{jjkl} &= \frac{1}{\alpha^2} \underline{R}_{jjkl} - \frac{4\beta^2}{\alpha^4} \kappa_1^2 \rho (\delta_{jk} \delta_{il} - \delta_{jl} \delta_{ik}) \\ &\quad - \frac{\beta^2}{4\alpha^4} y^\nu (2F_{\mu ij}^\nu F_{\gamma kl}^\mu + F_{\mu ik}^\nu F_{\gamma jl}^\mu - F_{\mu il}^\nu F_{\gamma jk}^\mu) y^\gamma, \end{aligned} \quad (\text{A.4})$$

$$\begin{aligned} R_{ji(n+\mu)(n+\nu)} &= -\frac{1}{\alpha^2} F_{\nu ij}^\mu + \frac{2\beta^2}{\alpha^4} (\kappa_1 + \kappa_2) (y^\nu F_{\gamma ij}^\mu - y^\mu F_{\gamma ij}^\nu) y^\gamma \\ &\quad + \frac{\beta^2}{4\alpha^4} y^\gamma (F_{\gamma ik}^\mu F_{\eta jk}^\nu - F_{\gamma ik}^\nu F_{\eta jk}^\mu) y^\eta. \end{aligned} \quad (\text{A.5})$$

In the above expression,  $\underline{R}_{jjkl}$  is the Riemann curvature tensor of  $(B, \underline{g})$  along  $(\underline{e}_j, \underline{e}_i, \underline{e}_k, \underline{e}_l)$ , where  $\{\underline{e}_j\}$  is the dual frame of  $\{\underline{\omega}^j\}$ .

For the curvature component  $\Omega_i^{n+\mu}$ ,

$$\begin{aligned}
(d\omega_i^{n+\mu})|_p &= -\frac{2}{\alpha^2}\kappa_1\omega^{n+\mu}\wedge\omega^i + \frac{1}{2\alpha^2}F_{\nu ij}^\mu\omega^{n+\nu}\wedge\omega^j + \frac{4\beta^2}{\alpha^4}\kappa_1(\kappa_1+\kappa_2)y^\mu y^\gamma\omega^{n+\gamma}\wedge\omega^i \\
&\quad - \frac{\beta^2}{\alpha^4}(\kappa_1+\kappa_2)F_{\gamma ij}^\mu y^\gamma y^\nu\omega^{n+\nu}\wedge\omega^j, \\
(-\omega_i^j\wedge\omega_j^{n+\mu})|_p &= \frac{\beta^2}{\alpha^4}\kappa_1 F_{\gamma ij}^\nu y^\gamma y^\mu\omega^{n+\nu}\wedge\omega^j + \frac{\beta^2}{\alpha^4}F_{\gamma ik}^\nu y^\gamma F_{\eta jk}^\mu y^\eta\omega^{n+\nu}\wedge\omega^j, \\
(-\omega_i^{n+\nu}\wedge\omega_{n+\nu}^i)|_p &= \frac{4\beta^2}{\alpha^4}\kappa_1\kappa_2\rho\omega^{n+\mu}\wedge\omega^i - \frac{4\beta^2}{\alpha^4}\kappa_1\kappa_2 y^\mu y^\nu\omega^{n+\nu}\wedge\omega^i + \frac{\beta^2}{\alpha^4}\kappa_2 F_{\gamma ij}^\nu y^\gamma y^\mu\omega^{n+\nu}\wedge\omega^j,
\end{aligned}$$

and then

$$\begin{aligned}
R_{(n+\mu)i(n+\nu)j} &= -\left(\frac{2\kappa_1}{\alpha^2} - \frac{4\kappa_1\kappa_2\beta^2}{\alpha^4}\rho\right)\delta_{\mu\nu}\delta_{ij} + \frac{1}{2\alpha^2}F_{\nu ij}^\mu + \frac{4\kappa_1^2\beta^2}{\alpha^4}y^\mu y^\nu\delta_{ij} \\
&\quad + (\kappa_1+\kappa_2)\frac{\beta^2}{\alpha^4}(y^\mu F_{\gamma ij}^\nu - y^\nu F_{\gamma ij}^\mu)y^\gamma + \frac{\beta^2}{4\alpha^4}y^\gamma F_{\gamma ik}^\nu F_{\eta jk}^\mu y^\eta.
\end{aligned} \tag{A.6}$$

For the curvature component  $\Omega_{n+\nu}^{n+\mu}$ ,

$$\begin{aligned}
(d\omega_{n+\nu}^{n+\mu})|_p &= \frac{1}{2\alpha^2}F_{\nu ij}^\mu\omega^i\wedge\omega^j + \kappa_2\frac{\beta^2}{\alpha^4}(y^\mu F_{\gamma ij}^\nu - y^\nu F_{\gamma ij}^\mu)y^\gamma\omega^i\wedge\omega^j \\
&\quad + \frac{4\kappa_2}{\alpha^2}\omega^{n+\mu}\wedge\omega^{n+\nu} - 8\kappa_2(\kappa_1+\kappa_2)\frac{\beta^2}{\alpha^4}y^\gamma\omega^{n+\gamma}\wedge(y^\mu\omega^{n+\nu} - y^\nu\omega^{n+\mu}), \\
(-\omega_{n+\nu}^i\wedge\omega_i^{n+\mu})|_p &= \frac{\beta^2}{4\alpha^4}y^\gamma F_{\gamma ik}^\nu F_{\eta jk}^\mu y^\eta\omega^i\wedge\omega^j - \kappa_1\frac{\beta^2}{\alpha^4}(y^\nu F_{\gamma ij}^\mu - y^\mu F_{\gamma ij}^\nu)y^\gamma\omega^i\wedge\omega^j, \\
(-\omega_{n+\nu}^{n+\gamma}\wedge\omega_{n+\nu}^{n+\mu})|_p &= \kappa_2^2\frac{4\beta^2}{\alpha^4}\rho\omega^{n+\nu}\wedge\omega^{n+\mu} - \kappa_2^2\frac{4\beta^2}{\alpha^4}y^\nu y^\gamma\omega^{n+\gamma}\wedge\omega^{n+\mu} - 4\kappa_2^2 y^\mu y^\gamma\omega^{n+\nu}\wedge\omega^{n+\gamma},
\end{aligned}$$

and

$$\begin{aligned}
R_{(n+\mu)(n+\nu)(n+\gamma)(n+\eta)} &= \left(\frac{4\kappa_2}{\alpha^2} - \frac{4\kappa_2^2\beta^2}{\alpha^4}\rho\right)(\delta_{\mu\gamma}\delta_{\nu\eta} - \delta_{\mu\eta}\delta_{\nu\gamma}) \\
&\quad + 4\kappa_2(2\kappa_1+\kappa_2)\frac{\beta^2}{\alpha^4}(y^\nu y^\gamma\delta_{\mu\eta} - y^\nu y^\eta\delta_{\mu\gamma} + y^\mu y^\eta\delta_{\nu\gamma} - y^\mu y^\gamma\delta_{\eta\nu}).
\end{aligned} \tag{A.7}$$

## REFERENCES

- [1] R. L. Bryant and S. M. Salamon, *On the construction of some complete metrics with exceptional holonomy*, Duke Math. J. **58** (1989), no. 3, 829–850.
- [2] R. L. Bryant and F. Xu, *Laplacian flow for closed  $G_2$  structures: short time behavior*, arXiv:1101.2004.
- [3] E. Calabi, *Métriques kählériennes et fibrés holomorphes*, Ann. Sci. École Norm. Sup. (4) **12** (1979), no. 2, 269–294 (French).
- [4] P. Candelas and X. C. de la Ossa, *Comments on conifolds*, Nuclear Phys. B **342** (1990), no. 1, 246–268.
- [5] S.-S. Chern, *Complex manifolds without potential theory*, 2nd ed., Springer-Verlag, New York-Heidelberg, 1979.
- [6] M. Cvetič, G. W. Gibbons, H. Lü, and C. N. Pope, *Hyper-Kähler Calabi metrics,  $L^2$  harmonic forms, resolved M2-branes, and AdS<sub>4</sub>/CFT<sub>3</sub> correspondence*, Nuclear Phys. B **617** (2001), no. 1-3, 151–197.



- [7] ———, *Ricci-flat metrics, harmonic forms and brane resolutions*, Comm. Math. Phys. **232** (2003), no. 3, 457–500.
- [8] A. Dancer and A. Swann, *Hyper-Kähler metrics of cohomogeneity one*, J. Geom. Phys. **21** (1997), no. 3, 218–230.
- [9] R. Harvey and H. B. Lawson Jr., *Calibrated geometries*, Acta Math. **148** (1982), 47–157.
- [10] M. Ionel, S. Karigiannis, and M. Min-Oo, *Bundle constructions of calibrated submanifolds in  $\mathbb{R}^7$  and  $\mathbb{R}^8$* , Math. Res. Lett. **12** (2005), no. 4, 493–512.
- [11] D. D. Joyce, *Compact manifolds with special holonomy*, Oxford University Press, 2000.
- [12] ———, *Lectures on special Lagrangian geometry*, Clay Math. Proc. **2** (2005), 667–695.
- [13] S. Karigiannis and M. Min-Oo, *Calibrated subbundles in noncompact manifolds of special holonomy*, Ann. Global Anal. Geom. **28** (2005), no. 4, 371–394.
- [14] H. B. Lawson Jr. and R. Osserman, *Non-existence, non-uniqueness and irregularity of solutions to the minimal surface system*, Acta Math. **139** (1977), no. no. 1-2, 1–17.
- [15] J. D. Lotay and Y. Wei, *Stability of torsion-free  $G_2$  structures along the Laplacian flow*, arXiv:1504.07771.
- [16] R. C. McLean, *Deformations of calibrated submanifolds*, Comm. Anal. Geom. **6** (1998), no. 4, 705–747.
- [17] R. Schoen and J. Wolfson, *Minimizing area among Lagrangian surfaces: the mapping problem*, J. Differential Geom. **58** (2001), no. 1, 1–86.
- [18] L. Simon, *Asymptotics for a class of nonlinear evolution equations, with applications to geometric problems*, Ann. of Math. (2) **118** (1983), no. 3, 525–571.
- [19] K. Smoczyk and M.-T. Wang, *Mean curvature flows of Lagrangian submanifolds with convex potentials*, J. Differential Geom. **62** (2002), no. 2, 243–257.
- [20] M. B. Stenzel, *Ricci-flat metrics on the complexification of a compact rank one symmetric space*, Manuscripta Math. **80** (1993), no. 2, 151–163.
- [21] M.-P. Tsui and M.-T. Wang, *Mean curvature flows and isotopy of maps between spheres*, Comm. Pure Appl. Math. **57** (2004), no. 8, 1110–1126.
- [22] M.-T. Wang, *Mean curvature flow of surfaces in Einstein four-manifolds*, J. Differential Geom. **57** (2001), no. 2, 301–338.
- [23] ———, *Deforming area preserving diffeomorphism of surfaces by mean curvature flow*, Math. Res. Lett. **8** (2001), no. 5-6, 651–661.
- [24] ———, *Long-time existence and convergence of graphic mean curvature flow in arbitrary codimension*, Invent. Math. **148** (2002), no. 3, 525–543.
- [25] ———, *Subsets of Grassmannians preserved by mean curvature flows*, Comm. Anal. Geom. **13** (2005), no. 5, 981–998.
- [26] ———, *The mean curvature flow smoothes Lipschitz submanifolds*, Comm. Anal. Geom. **12** (2004), no. 3, 581–599.
- [27] S.-T. Yau, *On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation. I*, Comm. Pure Appl. Math. **31** (1978), no. 3, 339–411.

DEPARTMENT OF MATHEMATICS, NATIONAL TAIWAN UNIVERSITY, TAIPEI 10617, TAIWAN

*E-mail address:* [cjtsai@ntu.edu.tw](mailto:cjtsai@ntu.edu.tw)

DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, NEW YORK, NY 10027, USA

*E-mail address:* [mtwang@math.columbia.edu](mailto:mtwang@math.columbia.edu)