

# Global Solutions to the Gas-Vacuum Interface Problem of Isentropic Compressible Inviscid Flows with Damping in Spherically Symmetric Motions and Physical Vacuum

Huihui Zeng<sup>a,b</sup>

<sup>a</sup> *Mathematical Sciences Center, Tsinghua University, Beijing, 100084, China*<sup>1</sup>

E-mail: hhzeng@mail.tsinghua.edu.cn

<sup>b</sup> *Center of Mathematical Sciences and Applications, Harvard University, 1 Oxford Street, Cambridge, MA 02318, USA*

MSC : 35B35, 35B 65, 76N10

*Keywords* : Physical vacuum, Compressible Euler equations, Damping, Barenblatt solutions, Spherically symmetric motions.

## Abstract

For the physical vacuum free boundary problem with the sound speed being  $C^{1/2}$ -Hölder continuous near vacuum boundaries of the compressible Euler equations with damping, the global existence of solutions and convergence to Barenblatt self-similar solutions of the porous media equation was recently proved in [34] for 1-d motions by Luo and the author. This paper generalizes the results for 1-d motions to 3-d spherically symmetric motions. Compared with the 1-d theory, besides the high degeneracy of the equations near the physical vacuum boundary, the analytical difficulties lie in the complexity of equations and the coordinates singularity in the center of symmetry which is resolved by constructing suitable weights. The results obtained in this work contribute to the theory of global solutions to free boundary problems of compressible inviscid fluids in the presence of vacuum states, for which the currently available results are mainly for the local-in-time well-posedness theory, also to the theory of global smooth solutions of dissipative hyperbolic systems which fail to be strictly hyperbolic.

## 1 Introduction

Due to its great physical importance and mathematical challenges, the motion of physical vacuum in compressible fluids has received much attention recently (cf. [7], [9]-[13], [20]-[23], [31]-[35], [42, 43]), and significant progress has been made on the local well-posedness theory (cf. [7, 9, 10, 20, 21, 35]). However, much less is known on the global existence and long time dynamics of solutions, which are of fundamental importance in both physics and nonlinear partial differential equations. This is the main issue we address in this work for the spherically symmetric motions of three-dimensional isentropic compressible inviscid flows with frictional damping. Physical vacuum problems arise in many physical situations naturally, for example, in the study of the evolution and structure of gaseous stars (cf. [5, 11]) for which vacuum boundaries are natural boundaries. Another situation in which the physical vacuum plays an important role is the gas-vacuum interface problem of compressible

---

<sup>1</sup>Permanent address

isentropic Euler equations with damping (cf. [31]-[33], [42, 43]). In three dimensions, this problem is given as follows:

$$\begin{aligned}
\rho_t + \operatorname{div}(\rho \mathbf{u}) &= 0 && \text{in } \Omega(t), \\
(\rho \mathbf{u})_t + \operatorname{div}(\rho \mathbf{u} \otimes \mathbf{u}) + \nabla_{\mathbf{x}} p(\rho) &= -\rho \mathbf{u} && \text{in } \Omega(t), \\
\rho > 0 &&& \text{in } \Omega(t), \\
\rho = 0 &&& \text{on } \Gamma(t) := \partial\Omega(t), \\
\mathcal{V}(\Gamma(t)) &= \mathbf{u} \cdot \mathbf{n}, \\
(\rho, \mathbf{u}) &= (\rho_0, \mathbf{u}_0) && \text{on } \Omega := \Omega(0).
\end{aligned} \tag{1.1}$$

Here  $(\mathbf{x}, t) \in \mathbb{R}^3 \times [0, \infty)$ ,  $\rho$ ,  $\mathbf{u}$ , and  $p$  denote, respectively, the space and time variable, density, velocity and pressure;  $\Omega(t) \subset \mathbb{R}^3$ ,  $\Gamma(t)$ ,  $\mathcal{V}(\Gamma(t))$  and  $\mathbf{n}$  represent, respectively, the changing volume occupied by a gas at time  $t$ , moving interface of fluids and vacuum states, normal velocity of  $\Gamma(t)$  and exterior unit normal vector to  $\Gamma(t)$ . We consider a polytropic gas: the equation of state is given by

$$p(\rho) = \rho^\gamma \quad \text{for } \gamma > 1 \tag{1.2}$$

with the adiabatic constant set to be unity for convenience. Equations (1.1)<sub>1,2</sub> describe the balance laws of mass and momentum, respectively; conditions (1.1)<sub>3,4</sub> state that  $\Gamma(t)$  is the interface to be investigated; (1.1)<sub>5</sub> indicates that the interface moves with the normal component of the fluid velocity; and (1.1)<sub>6</sub> is the initial conditions for the density, velocity and domain. Let  $c = \sqrt{p'(\rho)} = \sqrt{\gamma \rho^{\gamma-1}}$  be the sound speed, the condition

$$-\infty < \nabla_{\mathbf{n}}(c^2) < 0 \quad \text{on } \Gamma(t) \tag{1.3}$$

defines a physical boundary (cf. [7, 10, 21], [31]-[33]).

The compressible Euler equations of isentropic flow with damping are closely related to the porous media equation (cf. [15, 16, 17, 31]):

$$\rho_t = \Delta p(\rho), \tag{1.4}$$

when (1.1)<sub>2</sub> is simplified to Darcy's law:

$$\nabla_{\mathbf{x}} p(\rho) = -\rho \mathbf{u}. \tag{1.5}$$

For (1.4), basic understanding of the solution with finite mass  $M > 0$  is provided by Barenblatt (cf. [4]), which is spherically symmetric and given by

$$\bar{\rho}(\mathbf{x}, t) = \bar{\rho}(r, t) = (1+t)^{-\frac{3}{3\gamma-1}} \left( A - B(1+t)^{-\frac{2}{3\gamma-1}} r^2 \right)^{\frac{1}{\gamma-1}} \quad \text{with } r = |\mathbf{x}|, \tag{1.6}$$

where

$$B = \frac{\gamma-1}{2\gamma(3\gamma-1)} \quad \text{and} \quad (\gamma A)^{\frac{3\gamma-1}{2(\gamma-1)}} = M\gamma^{\frac{1}{\gamma-1}} (\gamma B)^{\frac{3}{2}} \left( \int_0^1 y^2 (1-y^2)^{\frac{1}{\gamma-1}} dy \right)^{-1}. \tag{1.7}$$

Clearly,

$$\int_0^{\bar{R}(t)} r^2 \bar{\rho}(r, t) dr = M \quad \text{for } t \geq 0, \quad \text{where } \bar{R}(t) = \sqrt{A/B}(1+t)^{1/(3\gamma-1)}. \quad (1.8)$$

The corresponding Barenblatt velocity  $\bar{\mathbf{u}}$  is defined by

$$\bar{\mathbf{u}}(\mathbf{x}, t) = (\mathbf{x}/r)\bar{u}(r, t) \quad \text{in the region } \{(r, t) : 0 \leq r \leq \bar{R}(t), t > 0\},$$

where

$$\bar{u}(r, t) = -\frac{p(\bar{\rho})_r}{\bar{\rho}} = \frac{r}{(3\gamma-1)(1+t)} \quad \text{satisfying } \bar{u}(0, t) = 0 \quad \text{and } \dot{\bar{R}}(t) = \bar{u}(\bar{R}(t), t).$$

So,  $(\bar{\rho}, \bar{\mathbf{u}})$  defined in the region  $\{(r, t) : 0 \leq r \leq \bar{R}(t), t > 0\}$  solves (1.4) and (1.5).

The vacuum boundary  $r = \bar{R}(t)$  of Barenblatt's solution is physical. This is the major motivation to study the physical vacuum free boundary problem of compressible Euler equations with damping. To this end, a class of explicit spherically symmetric solutions to problem (1.1) was constructed in [31], which are of the following form:

$$\Omega(t) = B_{R(t)}(\mathbf{0}), \quad c^2(\mathbf{x}, t) = c^2(r, t) = e(t) - b(t)r^2 \quad \text{and} \quad \mathbf{u}(\mathbf{x}, t) = (\mathbf{x}/r)u(r, t), \quad (1.9)$$

where

$$R(t) = \sqrt{e(t)/b(t)} \quad \text{and} \quad u(r, t) = a(t)r.$$

In [31], a system of ordinary differential equations for  $(e, b, a)(t)$  was derived with  $e(t), b(t) > 0$  for  $t \geq 0$  by substituting (1.9) into (1.1)<sub>1,2</sub> and the time-asymptotic equivalence of this explicit solution and Barenblatt's solution with the same total mass was shown. Indeed, the Barenblatt solution of (1.4) and (1.5) can be obtained by the same ansatz as (1.9):

$$\bar{c}^2(\mathbf{x}, t) = \bar{e}(t) - \bar{b}(t)r^2 \quad \text{and} \quad \mathbf{u}(\mathbf{x}, t) = \bar{a}(t)\mathbf{x}.$$

Substituting this into (1.4), (1.5) and (1.8) with  $\bar{R}(t) = \sqrt{\bar{e}(t)/\bar{b}(t)}$  gives

$$\bar{e}(t) = \gamma A(1+t)^{-3(\gamma-1)/(3\gamma-1)}, \quad \bar{b}(t) = \gamma B(1+t)^{-1} \quad \text{and} \quad \bar{a}(t) = (3\gamma-1)^{-1}(1+t)^{-1},$$

where  $A$  and  $B$  are determined by (1.7). Precisely, it was proved in [31] the following time-asymptotic equivalence:

$$(a, b, e)(t) = (\bar{a}, \bar{b}, \bar{e})(t) + O(1)(1+t)^{-1} \ln(1+t) \quad \text{as } t \rightarrow \infty.$$

A question was raised in [31] whether this equivalence is still true for general solutions to problem (1.1). For this purpose, we seek solutions to problem (1.1) of the form:

$$\Omega(t) = B_{R(t)}(\mathbf{0}), \quad \rho(\mathbf{x}, t) = \rho(r, t), \quad \mathbf{u}(\mathbf{x}, t) = (\mathbf{x}/r)u(r, t) \quad \text{with } r = |\mathbf{x}|.$$

Then problem (1.1) reduces to

$$\begin{aligned} (r^2 \rho)_t + (r^2 \rho u)_r &= 0 && \text{in } (0, R(t)), \\ \rho(u_t + uu_r) + p_r &= -\rho u && \text{in } (0, R(t)), \\ \rho &> 0 && \text{in } [0, R(t)), \\ \rho(R(t), t) &= 0, \quad u(0, t) = 0, && \\ \dot{R}(t) &= u(R(t), t) \quad \text{with } R(0) = R_0, && \\ (\rho, u)(r, t=0) &= (\rho_0, u_0)(r) && \text{on } (0, R_0), \end{aligned} \quad (1.10)$$

so that  $R(t)$  is the radius of the domain occupied by the gas at time  $t$  and  $r = R(t)$  represents the vacuum free boundary which issues from  $r = R_0$  and moves with the fluid velocity. One of motivations to study the spherically symmetric solution is that the Barenblatt solution possesses the same symmetry, and it is expected that spherically symmetric solutions will provide insights on the local and long time behavior of solutions to the general three-dimensional problem (1.1). Locally, at each point  $\mathbf{x}$  in  $\Omega(t)$ , it might be plausible to rotate a solution in all possible ways about  $\mathbf{x}$  and average all rotations in the spirit of spherical mean. In long time, for a general three-dimensional problem, it is expected that the geometry of the boundary becomes more and more symmetric due to the dissipation of damping which dissipates the total energy.

In the spherically symmetric setting, the physical vacuum boundary condition (1.3) reduces to

$$-\infty < (c^2)_r < 0$$

in a small neighborhood of the boundary. To capture this singularity, the initial domain is taken to be a ball  $\{0 \leq r \leq R_0\}$  and the initial density is assumed to satisfy

$$\rho_0(r) > 0 \text{ for } 0 \leq r < R_0, \quad \rho_0(R_0) = 0 \text{ and } -\infty < (\rho_0^{\gamma-1})_r < 0 \text{ at } r = R_0. \quad (1.11)$$

We require that the initial total mass is the same as that of the Barenblatt solution, that is,

$$\int_0^{R_0} r^2 \rho_0(r) dr = \int_0^{\bar{R}(0)} r^2 \bar{\rho}_0(r) dr = M. \quad (1.12)$$

The conservation law of mass, (1.10)<sub>1</sub>, and (1.8) give

$$\int_0^{R(t)} r^2 \rho(r, t) dr = \int_0^{R_0} r^2 \rho_0(r) dr = M = \int_0^{\bar{R}(t)} r^2 \bar{\rho}(r, t) dr \text{ for } t \geq 0.$$

In the present work, we prove the global existence of smooth solutions to problem (1.10) when initial data are small spherically symmetric perturbations of Barenblatt solutions and they have the same total masses. Moreover, we obtain the pointwise convergence with a rate of density which gives the detailed behavior of the density, the convergence rate of velocity in supreme norm and the precise expanding rate of physical vacuum boundaries. The results obtained in this article also prove the nonlinear asymptotic stability of Barenblatt solutions in the setting of physical vacuum free boundary problems.

The physical vacuum with the sound speed being  $C^{1/2}$ -Hölder continuous across vacuum boundaries makes it challenging and interesting in the study of free boundary problems in compressible fluids, even for the local-in-time existence theory, since standard methods of symmetric hyperbolic systems (cf. [24]) do not apply. Indeed, characteristic speeds of the compressible isentropic Euler equations become singular with infinite spatial derivatives at vacuum boundaries which creates much severe difficulties in analyzing the regularity near boundaries. Recently, important progress has been made in the local-in-time well-posedness theory for the compressible Euler equations (cf. [7, 9, 10, 20, 21]). On the other hand, it poses a great challenge to extend the local-in-time existence theory to the global one of smooth solutions, due to the strong degeneracy near vacuum states caused by the singular behavior of physical vacuum. Obtaining the global-in-time regularity of solutions

near vacuum boundaries by establishing the uniform-in-time higher-order estimates is the key to analyses. This is nontrivial due to the strong degenerate nonlinear hyperbolic nature. To obtain global-in-time estimates, it is essential to show decay estimates, which are achieved in the present work by introducing time weights to quantify the long time behavior of solutions. This is in sharp contrast to the weighted estimates used in establishing the local-in-time well-posedness theory (cf. [7, 9, 10, 20, 21, 35]), where only spatial weights are involved.

It should be noted that, as the first step to understand global solutions and their long time behavior for physical vacuum boundary problems of the isentropic compressible Euler equations with frictional damping, Luo and the author (cf. [34]) proved the global smooth solutions and convergence to Barenblatt solutions as time goes to infinity in one-dimensional case, based on a construction of higher-order weighted functionals with both space and time weights capturing the behavior of solutions both near vacuum states and in large time, an introduction of a new ansatz, higher-order nonlinear energy estimates and elliptic estimates. In general, much more obstacles appear in the study of multi-dimensional problems of compressible Euler equations as a prototype. Compared with the one-dimensional case studied in [34], it is much more difficult and involved to solve the three-dimensional spherically symmetric problem, (1.10), in the construction of the nonlinear weighted functionals, nonlinear weighted estimates and elliptic estimates. Besides the difficulty of degeneracy of the equations at vacuum states, one of the difficulties in solving (1.10) is the coordinates singularity at the origin, the center of symmetry, which carries the true three-dimensional nature. We succeed in constructing suitable weights to resolve the coordinates singularity in this paper. As an intermediate step passing from one-dimensional case in [34] to the general three-dimensional problem, (1.1), we believe the ideas and estimates including the nonlinear weighted functionals and pointwise decay estimates developed in this paper will contribute to a understanding of the behavior of solutions to problem (1.1).

It should be pointed that the  $L^p$ -convergence of  $L^\infty$ -weak solutions for the Cauchy problem of the one-dimensional compressible Euler equations with damping to Barenblatt solutions of the porous media equations was given in [16] with  $p = 2$  if  $1 < \gamma \leq 2$  and  $p = \gamma$  if  $\gamma > 2$  and in [17] with  $p = 1$ , respectively, using entropy-type estimates for the solution itself without deriving estimates for derivatives. However, the interfaces separating gases and vacuum cannot be traced in the framework of  $L^\infty$ -weak solutions. The aim of this work is to understand the behavior and long time dynamics of physical vacuum boundaries, for which obtaining the global-in-time regularity of solutions is essential.

There has been a recent explosion of interests in the analysis of free boundary problems for both compressible and incompressible inviscid flows. (As for viscous flows, there have been many results on the free boundary Navier-Stokes equations which cause quite different difficulties in analyses from those for inviscid flows, so we do not discuss the works in that regime here.) For incompressible inviscid flows, one may refer to [2, 3, 6, 8, 26, 28, 36, 38, 39, 47] for the local-in-time theory; while the global-in-time theory is rather recent which is for both 2-d and 3-d water wave problems of irrotational flows (cf. [18, 19, 40, 41]). For compressible inviscid flows, besides the aforementioned results on vacuum boundary problems, the local-in-time existence and uniqueness for the 3-d compressible Euler equations modeling a liquid rather than a gas were established in [29] by using Lagrangian variables combined with Nash-Moser iteration to construct solutions. (For a compressible liquid, the density is assumed to be a strictly positive constant on the moving boundary. As such,

the compressible liquid provides a uniformly hyperbolic, but characteristic, system.) An alternative proof for the existence of a compressible liquid was given in [37], employing a strategy based on symmetric hyperbolic systems combined with Nash-Moser iteration. From the above discussions, one may see that the current available theories of free boundary problems for inviscid flows, in particular for compressible inviscid flows, are mainly on local-in-time solutions. The results obtained in this paper are among the first ones on the global solutions of free boundary problems for compressible inviscid fluids in the presence of vacuum states.

In a broader context, the equation we consider in this work fall into the class of hyperbolic systems with dissipation for which most available results on the global existence of smooth solutions are for *strictly* hyperbolic systems or systems endowed with *strict* convex entropy (cf. [14, 27, 30, 44, 45, 46]). Indeed, the isentropic compressible Euler equations with frictional damping fail to be strictly hyperbolic at the vacuum state  $\rho = 0$ , and the standard mechanic entropy  $\eta(\rho, m) = p(\rho)/(\gamma - 1) + m^2/(2\rho)$  with  $m = \rho u$  being the momentum fails to be strictly convex. The results obtained in this paper, together with those in [34], contribute therefore to the global existence theory of smooth solutions of hyperbolic systems which are not strictly hyperbolic.

## 2 Reformulation of the problem and main results

### 2.1 Fix the domain and Lagrangian variables

We make the initial domain of the Barenblatt solution,  $(0, \bar{R}(0))$ , as the reference domain and define a diffeomorphism  $\eta_0 : (0, \bar{R}(0)) \rightarrow (0, R_0)$  by

$$\int_0^{\eta_0(r)} r^2 \rho_0(r) dr = \int_0^r r^2 \bar{\rho}_0(r) dr \quad \text{for } r \in (0, \bar{R}(0)),$$

where  $\bar{\rho}_0(r) := \bar{\rho}(r, 0)$  is the initial density of the Barenblatt solution. Clearly,

$$\eta_0^2(r) \rho_0(\eta_0(r)) \eta_{0r}(r) = r^2 \bar{\rho}_0(r) \quad \text{for } r \in (0, \bar{R}(0)). \quad (2.1)$$

Due to (1.11), (1.6) and the fact that the total mass of the Barenblatt solution is the same as that of  $\rho_0$ , (1.12), the diffeomorphism  $\eta_0$  is well defined. For simplicity of presentation, set

$$\mathcal{I} := (0, \bar{R}(0)) = \left(0, \sqrt{A/B}\right).$$

To fix the boundary, we transform system (1.10) into Lagrangian variables. For  $r \in \mathcal{I}$ , we define the Lagrangian variable  $\eta(r, t)$  by

$$\eta_t(r, t) = u(\eta(r, t), t) \quad \text{for } t > 0 \quad \text{and} \quad \eta(r, 0) = \eta_0(r), \quad (2.2)$$

and set the Lagrangian density and velocity by

$$f(r, t) = \rho(\eta(r, t), t) \quad \text{and} \quad v(r, t) = u(\eta(r, t), t). \quad (2.3)$$

Then the Lagrangian version of system (1.10) can be written on the reference domain  $\mathcal{I}$  as

$$\begin{aligned}
(\eta^2 f)_t + r^2 f v_r / \eta_r &= 0 & \text{in } \mathcal{I} \times (0, \infty), \\
f v_t + (f^\gamma)_r / \eta_r &= -f v & \text{in } \mathcal{I} \times (0, \infty), \\
v(0, t) &= 0 & \text{on } (0, \infty), \\
(f, v) &= (\rho_0(\eta_0), u_0(\eta_0)) & \text{on } \mathcal{I} \times \{t = 0\}.
\end{aligned} \tag{2.4}$$

It should be noted that we need  $\eta_r(r, t) > 0$  for  $r \in \mathcal{I}$  and  $t \geq 0$  to make the Lagrangian transformation sensible, which will be verified in (3.3). Indeed,  $\eta_r > 0$  implies  $\eta(r, t) > 0$  for  $r \in \mathcal{I}$  and  $t \geq 0$ , due to the boundary condition that the center of the symmetry does not move,  $v(0, t) = 0$ . The map  $\eta(\cdot, t)$  defined above can be extended to  $\bar{\mathcal{I}} = [0, \sqrt{A/B}]$ . In the setting, the vacuum free boundaries for problem (1.10) are given by

$$R(t) = \eta(\bar{R}(0), t) = \eta\left(\sqrt{A/B}, t\right) \quad \text{for } t \geq 0. \tag{2.5}$$

It follows from solving (2.4)<sub>1</sub> and using (2.1) that

$$f(r, t) \eta^2(r, t) \eta_r(r, t) = \rho_0(\eta_0(r)) \eta_0^2(r) \eta_{0r}(r) = r^2 \bar{\rho}_0(r), \quad r \in \mathcal{I}. \tag{2.6}$$

So, the initial density of the Barenblatt solution,  $\bar{\rho}_0$ , can be viewed as a parameter and system (2.4) can be rewritten as

$$\begin{aligned}
\bar{\rho}_0 \eta_{tt} + \bar{\rho}_0 \eta_t + \left(\frac{\eta}{r}\right)^2 \left[ \left(\frac{r^2 \bar{\rho}_0}{\eta^2 \eta_r}\right)^\gamma \right]_r &= 0 & \text{in } \mathcal{I} \times (0, \infty), \\
\eta(0, t) &= 0, & \text{on } (0, \infty), \\
(\eta, \eta_t) &= (\eta_0, u_0(\eta_0)) & \text{on } \mathcal{I} \times (0, \infty).
\end{aligned} \tag{2.7}$$

## 2.2 Ansatz

Define the Lagrangian variable  $\bar{\eta}(r, t)$  for the Barenblatt flow in  $\bar{\mathcal{I}}$  by

$$\partial_t \bar{\eta}(r, t) = \bar{u}(\bar{\eta}(r, t), t) = \frac{\bar{\eta}(r, t)}{(3\gamma - 1)(1 + t)} \quad \text{for } t > 0 \quad \text{and} \quad \bar{\eta}(r, 0) = r, \tag{2.8}$$

so that

$$\bar{\eta}(r, t) = r (1 + t)^{1/(3\gamma - 1)} \quad \text{for } (r, t) \in \bar{\mathcal{I}} \times [0, \infty) \tag{2.9}$$

and

$$\bar{\rho}_0 \bar{\eta}_t + \left(\frac{\bar{\eta}}{r}\right)^2 \left[ \left(\frac{r^2 \bar{\rho}_0}{\bar{\eta}^2 \bar{\eta}_r}\right)^\gamma \right]_r = 0 \quad \text{in } \mathcal{I} \times (0, \infty).$$

Since  $\bar{\eta}$  does not solve (2.7)<sub>1</sub> exactly, we introduce a correction  $h(t)$  which is a solution of the following initial value problem of ordinary differential equations:

$$\begin{aligned}
h_{tt} + h_t - (\bar{\eta}_r + h)^{2-3\gamma} / (3\gamma - 1) + \bar{\eta}_{rtt} + \bar{\eta}_{rt} &= 0, \quad t > 0, \\
h(t = 0) = h_t(t = 0) &= 0.
\end{aligned} \tag{2.10}$$

(Notice that  $\bar{\eta}_r$ ,  $\bar{\eta}_{rt}$  and  $\bar{\eta}_{rtt}$  are independent of  $r$ .) The new ansatz is then given by

$$\tilde{\eta}(r, t) := \bar{\eta}(r, t) + rh(t), \quad (2.11)$$

so that

$$\bar{\rho}_0 \tilde{\eta}_{tt} + \bar{\rho}_0 \tilde{\eta}_t + \left( \frac{\tilde{\eta}}{r} \right)^2 \left[ \left( \frac{r^2 \bar{\rho}_0}{\tilde{\eta}^2 \tilde{\eta}_r} \right)^\gamma \right]_r = 0 \quad \text{in } \mathcal{I} \times (0, \infty). \quad (2.12)$$

It should be noted that  $\tilde{\eta}_r$  is independent of  $r$ . We will prove in the Appendix that  $\tilde{\eta}$  behaves similar to  $\bar{\eta}$ , that is, there exist positive constants  $K$  and  $C(n)$  independent of time  $t$  such that for all  $t \geq 0$ ,

$$\begin{aligned} (1+t)^{1/(3\gamma-1)} \leq \tilde{\eta}_r(t) \leq K(1+t)^{1/(3\gamma-1)}, \quad \tilde{\eta}_{rt} \geq 0, \\ \left| \frac{d^k \tilde{\eta}_r(t)}{dt^k} \right| \leq C(n) (1+t)^{\frac{1}{3\gamma-1}-k}, \quad k = 1, 2, \dots, n. \end{aligned} \quad (2.13)$$

Moreover, there exists a certain constant  $C$  independent of  $t$  such that

$$0 \leq h(t) \leq C(1+t)^{\frac{1}{3\gamma-1}-1} \ln(1+t) \quad \text{and} \quad |h_t(t)| \leq C(1+t)^{\frac{1}{3\gamma-1}-2} \ln(1+t), \quad t \geq 0. \quad (2.14)$$

The proof of (2.14) will also be given in the Appendix.

## 2.3 Main results

To state the main theorem, we write equation (2.7)<sub>1</sub> in a perturbation form around the Barenblatt solution. Let

$$\zeta(r, t) := \eta(r, t)/r - \tilde{\eta}(r, t)/r.$$

Thus,

$$\eta(r, t) = \tilde{\eta}(r, t) + r\zeta(r, t) \quad \text{and} \quad \eta_r(r, t) = \tilde{\eta}_r(t) + \zeta(r, t) + r\zeta_r(r, t). \quad (2.15)$$

It follows from (2.7)<sub>1</sub> and (2.12) that

$$r\bar{\rho}_0 \zeta_{tt} + r\bar{\rho}_0 \zeta_t + (\tilde{\eta}_r + \zeta)^2 [\bar{\rho}_0^\gamma (\tilde{\eta}_r + \zeta)^{-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma}]_r - \tilde{\eta}_r^{2-3\gamma} (\bar{\rho}_0^\gamma)_r = 0, \quad (2.16)$$

Denote

$$\alpha := 1/(\gamma - 1), \quad l := 3 + \min \{m \in \mathbb{N} : m > \alpha\} = 4 + [\alpha].$$

For  $j = 0, \dots, l$  and  $i = 0, \dots, l - j$ , we set

$$\begin{aligned} \mathcal{E}_j(t) &:= (1+t)^{2j} \int_{\mathcal{I}} \left[ r^4 \bar{\rho}_0 (\partial_t^j \zeta)^2 + r^2 \bar{\rho}_0^\gamma |\partial_t^j (\zeta, r\zeta_r)|^2 + (1+t)r^4 \bar{\rho}_0 (\partial_t^j \zeta_t)^2 \right] dr, \\ \mathcal{E}_{j,i}(t) &:= (1+t)^{2j} \int_{\mathcal{I}} \left[ r^2 \bar{\rho}_0^{1+(i-1)(\gamma-1)} (\partial_t^j \partial_r^i \zeta)^2 + r^4 \bar{\rho}_0^{1+(i+1)(\gamma-1)} (\partial_t^j \partial_r^{i+1} \zeta)^2 \right] dr. \end{aligned}$$

The higher-order norm is defined by

$$\mathcal{E}(t) := \sum_{j=0}^l \left( \mathcal{E}_j(t) + \sum_{i=1}^{l-j} \mathcal{E}_{j,i}(t) \right).$$



It will be proved in Lemma 3.7 that

$$\sup_{r \in \mathcal{I}} \left\{ \sum_{j=0}^2 (1+t)^{2j} |\partial_t^j \zeta(r, t)|^2 + \sum_{j=0}^1 (1+t)^{2j} |\partial_t^j \zeta_r(r, t)|^2 \right\} \leq C\mathcal{E}(t)$$

for some constant  $C$  independent of  $t$ . So the boundedness of  $\mathcal{E}(t)$  gives the uniform boundedness and decay of the perturbation  $\zeta$  and its derivatives. In what follows, we state our main result.

**Theorem 2.1** *Suppose that (1.12) holds. There exists a constant  $\bar{\delta} > 0$  such that if  $\mathcal{E}(0) \leq \bar{\delta}$ , then the problem (2.7) admits a global unique smooth solution in  $\mathcal{I} \times [0, \infty)$  satisfying for all  $t \geq 0$ ,*

$$\mathcal{E}(t) \leq C\mathcal{E}(0)$$

and

$$\begin{aligned} & \sup_{r \in \mathcal{I}} \left\{ \sum_{j=0}^2 (1+t)^{2j} |\partial_t^j \zeta(r, t)|^2 + \sum_{j=0}^1 (1+t)^{2j} |\partial_t^j \zeta_r(r, t)|^2 + \sum_{i+j \leq l-2, 2i+j \geq 3} (1+t)^{2j} \right. \\ & \quad \times \left| \bar{\rho}_0^{\frac{(\gamma-1)(2i+j-3)}{2}} \partial_t^j \partial_r^i \zeta(r, t) \right|^2 + \sum_{i+j=l-1} (1+t)^{2j} \left| r \bar{\rho}_0^{\frac{(\gamma-1)(2i+j-3)}{2}} \partial_t^j \partial_r^i \zeta(r, t) \right|^2 \\ & \quad \left. + \sum_{i+j=l} (1+t)^{2j} \left| r^2 \bar{\rho}_0^{\frac{(\gamma-1)(2i+j-3)}{2}} \partial_t^j \partial_r^i \zeta(r, t) \right|^2 \right\} \leq C\mathcal{E}(0), \end{aligned} \quad (2.17)$$

where  $C$  is a positive constant independent of  $t$ .

It should be noticed that the time derivatives involved in the initial higher-order energy norm,  $\mathcal{E}(0)$ , can be determined via the equation by the initial data  $\rho_0$  and  $u_0$  (see [9, 35] for instance).

As a corollary of Theorem 2.1, we have the following theorem for solutions to the original vacuum free boundary problem (1.10).

**Theorem 2.2** *Suppose that (1.12) holds. There exists a constant  $\bar{\delta} > 0$  such that if  $\mathcal{E}(0) \leq \bar{\delta}$ , then the problem (1.10) admits a global unique smooth solution  $(\rho(\eta, t), u(\eta, t), R(t))$  for  $t \in [0, \infty)$  satisfying*

$$\begin{aligned} |\rho(\eta(r, t), t) - \bar{\rho}(\bar{\eta}(r, t), t)| & \leq C \left( A - Br^2 \right)^{\frac{1}{\gamma-1}} (1+t)^{-\frac{4}{3\gamma-1}} \\ & \quad \times \left( \sqrt{\mathcal{E}(0)} + (1+t)^{-\frac{3\gamma-2}{3\gamma-1}} \ln(1+t) \right), \end{aligned} \quad (2.18)$$

$$|u(\eta(r, t), t) - \bar{u}(\bar{\eta}(r, t), t)| \leq Cr(1+t)^{-1} \left( \sqrt{\mathcal{E}(0)} + (1+t)^{-\frac{3\gamma-2}{3\gamma-1}} \ln(1+t) \right), \quad (2.19)$$

for all  $r \in \mathcal{I}$  and  $t \geq 0$ ; and for all  $t \geq 0$ ,

$$c_1(1+t)^{\frac{1}{3\gamma-1}} \leq R(t) \leq c_2(1+t)^{\frac{1}{3\gamma-1}}, \quad (2.20)$$

$$\left| \frac{d^k R(t)}{dt^k} \right| \leq C(1+t)^{\frac{1}{3\gamma-1}-k}, \quad k = 1, 2, 3, \quad (2.21)$$

$$c_3(1+t)^{-\frac{3\gamma-2}{3\gamma-1}} \leq \left| (\rho^{\gamma-1})_\eta(\eta, t) \right| \leq c_4(1+t)^{-\frac{3\gamma-2}{3\gamma-1}} \quad \text{when} \quad \frac{1}{2}R(t) \leq \eta \leq R(t). \quad (2.22)$$

Here  $C$ ,  $c_1$ ,  $c_2$ ,  $c_3$  and  $c_4$  are positive constants independent of  $t$ .

The pointwise behavior of the density and velocity for the vacuum free boundary problem (1.10) to that of the Barenblatt solution are given by (2.18) and (2.19), respectively. It is also shown in (2.18) that the difference of density to problem (1.10) and the corresponding Barenblatt density decays at the rate of  $(1+t)^{-4/(\gamma+1)}$  in  $L^\infty$ , while the density of the Barenblatt solution,  $\bar{\rho}$ , decays at the rate of  $(1+t)^{-3/(\gamma+1)}$  in  $L^\infty$  (see (1.6)). (2.20) gives the precise expanding rate of the vacuum boundaries of the problem (1.10), which is the same as that for the Barenblatt solution shown in (1.8). Furthermore, it verifies in (2.22) that the vacuum boundary  $R(t)$  is physical at any finite time.

### 3 Proof of Theorem 2.1

The proof is based on the local existence of smooth solutions (cf. [35, 9, 20]) and continuation arguments. The uniqueness of the smooth solutions can be obtained as in section 11 of [35]. In order to prove the global existence of smooth solutions, we need to obtain the uniform-in-time *a priori* estimates on any given time interval  $[0, T]$  satisfying  $\sup_{t \in [0, T]} \mathcal{E}(t) < \infty$ . For this purpose, we use a bootstrap argument by making the following *a priori* assumption: Let  $\zeta$  be a smooth solution to (2.16) on  $[0, T]$ , there exists a suitably small fixed positive number  $\epsilon_0 \in (0, 1)$  independent of  $t$  such that for  $t \in [0, T]$ ,

$$\begin{aligned} & \sum_{j=0}^2 (1+t)^{2j} \left\| \partial_t^j \zeta(\cdot, t) \right\|_{L^\infty}^2 + \sum_{j=0}^1 (1+t)^{2j} \left\| \partial_t^j \zeta_r(\cdot, t) \right\|_{L^\infty}^2 + \sum_{i+j \leq l-2, 2i+j \geq 3} (1+t)^{2j} \\ & \times \left\| \bar{\rho}_0^{\frac{(\gamma-1)(2i+j-3)}{2}} \partial_t^j \partial_r^i \zeta(\cdot, t) \right\|_{L^\infty}^2 + \sum_{i+j=l-1} (1+t)^{2j} \left\| r \bar{\rho}_0^{\frac{(\gamma-1)(2i+j-3)}{2}} \partial_t^j \partial_r^i \zeta(\cdot, t) \right\|_{L^\infty}^2 \leq \epsilon_0^2. \end{aligned} \quad (3.1)$$

This in particular implies, noting (2.13), that for  $0 \leq \theta_1, \theta_2 \leq 1$ ,

$$\frac{1}{2}(1+t)^{\frac{1}{3\gamma-1}} \leq (\tilde{\eta}_r + \theta_1 \zeta + \theta_2 r \zeta_r)(r, t) \leq 2K(1+t)^{\frac{1}{3\gamma-1}}, \quad (r, t) \in \mathcal{I} \times [0, T]. \quad (3.2)$$

Moreover, it follows from (2.15) and (3.2) that

$$\frac{1}{2}(1+t)^{\frac{1}{3\gamma-1}} \leq \eta_r(r, t), \quad r^{-1} \eta(r, t) \leq 2K(1+t)^{\frac{1}{3\gamma-1}}, \quad (r, t) \in \mathcal{I} \times [0, T]. \quad (3.3)$$

Here  $K$  is the positive constant appearing in (2.13)<sub>1</sub>.

Under this *a priori* assumption, we prove in section 3.2 the following elliptic estimates:

$$\mathcal{E}_{j,i}(t) \leq C \sum_{\iota=0}^{i+j} \mathcal{E}_\iota(t), \quad \text{when } j \geq 0, \quad i \geq 1, \quad i+j \leq l,$$

where  $C$  is a positive constant independent of  $t$ . With the *a priori* assumption and elliptic estimates, we show in section 3.3 the following nonlinear weighted energy estimate: for some positive constant  $C$  independent of  $t$ ,

$$\mathcal{E}_j(t) \leq C \sum_{\iota=0}^j \mathcal{E}_\iota(0), \quad j = 0, 1, \dots, l.$$

Finally, the *a priori* assumption (3.1) can be verified in section 3.4 by proving

$$\begin{aligned} & \sum_{j=0}^2 (1+t)^{2j} \left\| \partial_t^j \zeta(\cdot, t) \right\|_{L^\infty}^2 + \sum_{j=0}^1 (1+t)^{2j} \left\| \partial_t^j \zeta_r(\cdot, t) \right\|_{L^\infty}^2 + \sum_{i+j \leq l-2, 2i+j \geq 3} (1+t)^{2j} \\ & \times \left\| \bar{\rho}_0^{\frac{(\gamma-1)(2i+j-3)}{2}} \partial_t^j \partial_r^i \zeta(\cdot, t) \right\|_{L^\infty}^2 + \sum_{i+j=l-1} (1+t)^{2j} \left\| r \bar{\rho}_0^{\frac{(\gamma-1)(2i+j-3)}{2}} \partial_t^j \partial_r^i \zeta(\cdot, t) \right\|_{L^\infty}^2 \\ & + \sum_{i+j=l} (1+t)^{2j} \left\| r^2 \bar{\rho}_0^{\frac{(\gamma-1)(2i+j-3)}{2}} \partial_t^j \partial_r^i \zeta(\cdot, t) \right\|_{L^\infty}^2 \leq C \mathcal{E}(t). \end{aligned}$$

for some positive constant  $C$  independent of  $t$ . This closes the whole bootstrap argument for small initial perturbations and completes the proof of Theorem 2.1.

### 3.1 Preliminaries

In this subsection, we list some embedding estimates for weighted Sobolev spaces which will be used later and introduce some notations to simplify the presentation. For any bounded interval  $I$ , set  $d(r) = \text{dist}(r, \partial I)$ . For any  $a > 0$  and nonnegative integer  $b$ , the weighted Sobolev space  $H^{a,b}(I)$  is given by

$$H^{a,b}(I) := \left\{ d^{a/2} F \in L^2(I) : \int_I d^a |\partial_r^k F|^2 dr < \infty, \quad 0 \leq k \leq b \right\}$$

with the norm

$$\|F\|_{H^{a,b}(I)}^2 := \sum_{k=0}^b \int_I d^a |\partial_r^k F|^2 dr.$$

Then for  $b \geq a/2$ , it holds the following *embedding of weighted Sobolev spaces* (cf. [25]):

$$H^{a,b}(I) \hookrightarrow H^{b-a/2}(I)$$

with the estimate

$$\|F\|_{H^{b-a/2}(I)} \leq C \|F\|_{H^{a,b}(I)} \tag{3.4}$$

for some positive constant  $C$  depending on  $a$ ,  $b$  and  $I$ .

The following general version of the *Hardy inequality* whose proof can be found in [25] will also be used often in this paper. Let  $k > 1$  be a given real number and  $F$  be a function satisfying

$$\int_0^\delta r^k (F^2 + F_r^2) dr < \infty,$$

where  $\delta$  is a positive constant; then it holds that

$$\int_0^\delta r^{k-2} F^2 dr \leq C(\delta, k) \int_0^\delta r^k (F^2 + F_r^2) dr,$$

where  $C(\delta, k)$  is a constant depending only on  $\delta$  and  $k$ . As a consequence, one has

$$\int_{\sqrt{A/(4B)}}^{\sqrt{A/B}} \left( \sqrt{A/B} - r \right)^{k-2} F^2 dr \leq C \int_{\sqrt{A/(4B)}}^{\sqrt{A/B}} \left( \sqrt{A/B} - r \right)^k (F^2 + F_r^2) dr, \quad (3.5)$$

where  $C$  is a constant depending on  $A$ ,  $B$  and  $k$ .

**Notations:**

1) Throughout the rest of paper,  $C$  will denote a positive constant which only depend on the parameters of the problem,  $\gamma$  and  $M$ , but does not depend on the data. They are referred as universal and can change from one inequality to another one. Also we use  $C(\beta)$  to denote a certain positive constant depending on quantity  $\beta$ .

2) We will employ the notation  $a \lesssim b$  to denote  $a \leq Cb$ ,  $a \sim b$  to denote  $C^{-1}b \leq a \leq Cb$  and  $a \gtrsim b$  to denote  $a \geq C^{-1}b$ , where  $C$  is the universal constant as defined above.

3) In the rest of the paper, we will use the notations

$$\int =: \int_{\mathcal{I}}, \quad \|\cdot\| =: \|\cdot\|_{L^2(\mathcal{I})} \quad \text{and} \quad \|\cdot\|_{L^\infty} =: \|\cdot\|_{L^\infty(\mathcal{I})}.$$

4) We set

$$\sigma(r) := \bar{\rho}_0^{\gamma-1}(r) = A - Br^2, \quad r \in \mathcal{I}.$$

Then  $\mathcal{E}_j$  and  $\mathcal{E}_{j,i}$  can be rewritten as

$$\begin{aligned} \mathcal{E}_j(t) &= (1+t)^{2j} \int \left[ r^4 \sigma^\alpha (\partial_t^j \zeta)^2 + r^2 \sigma^{\alpha+1} |\partial_t^j (\zeta, r\zeta_r)|^2 + (1+t)r^4 \sigma^\alpha (\partial_t^j \zeta_t)^2 \right] (r, t) dr, \\ \mathcal{E}_{j,i}(t) &= (1+t)^{2j} \int \left[ r^2 \sigma^{\alpha+i-1} (\partial_t^j \partial_r^i \zeta)^2 + r^4 \sigma^{\alpha+i+1} (\partial_t^j \partial_r^{i+1} \zeta)^2 \right] (r, t) dr. \end{aligned}$$

5) We set

$$\mathcal{I}_o := \left( 0, \sqrt{A/(4B)} \right) \quad \text{and} \quad \mathcal{I}_b := \left( \sqrt{A/(4B)}, \sqrt{A/B} \right).$$

Then

$$\mathcal{I} = \mathcal{I}_o \cup \mathcal{I}_b.$$

Moreover, it gives from the Hardy inequality (3.5) that for  $k > 1$ ,

$$\int_{\mathcal{I}_b} \sigma^{k-2}(r) F^2 dr \leq C(A, B, k) \int_{\mathcal{I}_b} \sigma^k(r) (F^2 + F_r^2) dr, \quad (3.6)$$

provided that the right-hand side of (3.6) is finite.

## 3.2 Elliptic estimates

In this subsection, we prove the following elliptic estimates.

**Proposition 3.1** *Suppose that (3.1) holds for suitably small positive number  $\epsilon_0 \in (0, 1)$ . Then it holds that for  $t \in [0, T]$ ,*

$$\mathcal{E}_{j,i}(t) \lesssim \sum_{\iota=0}^{i+j} \mathcal{E}_\iota(t) \quad \text{when } j \geq 0, \quad i \geq 1, \quad i+j \leq l. \quad (3.7)$$

The proof of this proposition consists of Lemma 3.2 and Lemma 3.3.

### 3.2.1 Lower-order elliptic estimates

Dividing equation (2.16) by  $\bar{\rho}_0$ , one has

$$\begin{aligned} & r\zeta_{tt} + r\zeta_t + \sigma (\tilde{\eta}_r + \zeta)^2 [(\tilde{\eta}_r + \zeta)^{-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma}]_r \\ & + \frac{\gamma}{\gamma-1} \sigma_r [(\tilde{\eta}_r + \zeta)^{2-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma} - \tilde{\eta}_r^{2-3\gamma}] = 0. \end{aligned}$$

Note that

$$\begin{aligned} & (\tilde{\eta}_r + \zeta)^2 [(\tilde{\eta}_r + \zeta)^{-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma}]_r \\ & = -2\gamma (\tilde{\eta}_r + \zeta)^{1-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma} \zeta_r - \gamma (\tilde{\eta}_r + \zeta)^{2-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma-1} (2\zeta_r + r\zeta_{rr}) \\ & = -\gamma \tilde{\eta}_r^{1-3\gamma} (4\zeta_r + r\zeta_{rr}) + \mathfrak{J}_1, \\ & (\tilde{\eta}_r + \zeta)^{2-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma} - \tilde{\eta}_r^{2-3\gamma} = -\gamma \tilde{\eta}_r^{1-3\gamma} (r\zeta_r) + (2-3\gamma) \tilde{\eta}_r^{1-3\gamma} \zeta + \mathfrak{J}_2, \end{aligned}$$

where

$$\begin{aligned} \mathfrak{J}_1 & := -2\gamma [(\tilde{\eta}_r + \zeta)^{1-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma} - \tilde{\eta}_r^{1-3\gamma}] \zeta_r \\ & \quad - \gamma [(\tilde{\eta}_r + \zeta)^{2-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma-1} - \tilde{\eta}_r^{1-3\gamma}] (2\zeta_r + r\zeta_{rr}), \\ \mathfrak{J}_2 & := (\tilde{\eta}_r + \zeta)^{2-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma} - \tilde{\eta}_r^{2-3\gamma} + \gamma \tilde{\eta}_r^{1-3\gamma} (r\zeta_r) - (2-3\gamma) \tilde{\eta}_r^{1-3\gamma} \zeta. \end{aligned} \quad (3.8)$$

Then,

$$\begin{aligned} & \gamma \tilde{\eta}_r^{1-3\gamma} \left[ r\sigma \zeta_{rr} + 4\sigma \zeta_r + \frac{\gamma}{\gamma-1} r\sigma_r \zeta_r \right] \\ & = r\zeta_{tt} + r\zeta_t + \frac{\gamma(2-3\gamma)}{\gamma-1} \sigma_r \tilde{\eta}_r^{1-3\gamma} \zeta + \sigma \mathfrak{J}_1 + \frac{\gamma}{\gamma-1} \sigma_r \mathfrak{J}_2. \end{aligned} \quad (3.9)$$

**Lemma 3.2** *Assume that (3.1) holds for suitably small positive number  $\epsilon_0 \in (0, 1)$ . Then,*

$$\mathcal{E}_{0,1}(t) \lesssim \mathcal{E}_0(t) + \mathcal{E}_1(t), \quad 0 \leq t \leq T.$$

**Proof.** Multiply equation (3.9) by  $\tilde{\eta}_r^{3\gamma-1} r \sigma^{\alpha/2}$  and square the spatial  $L^2$ -norm of the product to obtain

$$\begin{aligned} & \left\| r^2 \sigma^{1+\frac{\alpha}{2}} \zeta_{rr} + 4r \sigma^{1+\frac{\alpha}{2}} \zeta_r + (1+\alpha) r^2 \sigma^{\frac{\alpha}{2}} \sigma_r \zeta_r \right\|^2 \\ & \lesssim (1+t)^2 \left( \left\| r^2 \sigma^{\frac{\alpha}{2}} \zeta_{tt} \right\|^2 + \left\| r^2 \sigma^{\frac{\alpha}{2}} \zeta_t \right\|^2 + \left\| r \sigma^{1+\frac{\alpha}{2}} \mathfrak{J}_1 \right\|^2 + \left\| r \sigma^{\frac{\alpha}{2}} \sigma_r \mathfrak{J}_2 \right\|^2 \right) + \left\| r \sigma^{\frac{\alpha}{2}} \zeta \right\|^2 \quad (3.10) \\ & \lesssim \mathcal{E}_1 + (1+t)^2 \left( \left\| r \sigma^{1+\frac{\alpha}{2}} \mathfrak{J}_1 \right\|^2 + \left\| r \sigma^{\frac{\alpha}{2}} \sigma_r \mathfrak{J}_2 \right\|^2 \right) + \left\| r \sigma^{\frac{\alpha}{2}} \zeta \right\|^2 \end{aligned}$$

where we have used (2.13) and the definition of  $\mathcal{E}_1$ . It follows from the Taylor expansion, (3.2) and (3.1) that

$$\begin{aligned} |\mathfrak{J}_1| & \lesssim (1+t)^{-\frac{3\gamma}{3\gamma-1}} (|r\zeta_r| + |\zeta|) (|r\zeta_{rr}| + |\zeta_r|) \lesssim (1+t)^{-\frac{3\gamma}{3\gamma-1}} \epsilon_0 (|r\zeta_{rr}| + |\zeta_r|), \\ |\mathfrak{J}_2| & \lesssim (1+t)^{-\frac{3\gamma}{3\gamma-1}} (|r\zeta_r|^2 + |\zeta|^2) \lesssim (1+t)^{-\frac{3\gamma}{3\gamma-1}} \epsilon_0 (|r\zeta_r| + |\zeta|). \end{aligned}$$

Thus,

$$\begin{aligned} & (1+t)^2 \left( \left\| r \sigma^{1+\frac{\alpha}{2}} \mathfrak{J}_1 \right\|^2 + \left\| r \sigma^{\frac{\alpha}{2}} \sigma_r \mathfrak{J}_2 \right\|^2 \right) \\ & \lesssim \epsilon_0^2 \left( \left\| r^2 \sigma^{1+\frac{\alpha}{2}} \zeta_{rr} \right\|^2 + \left\| r \sigma^{1+\frac{\alpha}{2}} \zeta_r \right\|^2 + \left\| r^2 \sigma^{\frac{\alpha}{2}} \sigma_r \zeta_r \right\|^2 + \left\| r \sigma^{\frac{\alpha}{2}} \sigma_r \zeta \right\|^2 \right). \quad (3.11) \end{aligned}$$

Note that

$$\left\| r \sigma^{\frac{\alpha}{2}} \zeta \right\|^2 = \int_{\mathcal{I}_o} r^2 \sigma^\alpha \zeta^2 dr + \int_{\mathcal{I}_b} r^2 \sigma^\alpha \zeta^2 dr \lesssim \int_{\mathcal{I}_o} r^2 \sigma^{1+\alpha} \zeta^2 dr + \int_{\mathcal{I}_b} r^4 \sigma^\alpha \zeta^2 dr \lesssim \mathcal{E}_0. \quad (3.12)$$

Then, it yields from (3.10), (3.11) and (3.12) that

$$\begin{aligned} & \left\| r^2 \sigma^{1+\frac{\alpha}{2}} \zeta_{rr} + 4r \sigma^{1+\frac{\alpha}{2}} \zeta_r + (1+\alpha) r^2 \sigma^{\frac{\alpha}{2}} \sigma_r \zeta_r \right\|^2 \\ & \lesssim \mathcal{E}_0 + \mathcal{E}_1 + \epsilon_0^2 \left( \left\| r^2 \sigma^{1+\frac{\alpha}{2}} \zeta_{rr} \right\|^2 + \left\| r \sigma^{1+\frac{\alpha}{2}} \zeta_r \right\|^2 + \left\| r^2 \sigma^{\frac{\alpha}{2}} \sigma_r \zeta_r \right\|^2 \right). \quad (3.13) \end{aligned}$$

In what follows, we analyze the left-hand side of (3.13), which can be expanded as

$$\begin{aligned} & \left\| r^2 \sigma^{1+\frac{\alpha}{2}} \zeta_{rr} + 4r \sigma^{1+\frac{\alpha}{2}} \zeta_r + (1+\alpha) r^2 \sigma^{\frac{\alpha}{2}} \sigma_r \zeta_r \right\|^2 \\ & = \left\| r^2 \sigma^{1+\frac{\alpha}{2}} \zeta_{rr} \right\|^2 + 16 \left\| r \sigma^{1+\frac{\alpha}{2}} \zeta_r \right\|^2 + (1+\alpha)^2 \left\| r^2 \sigma^{\frac{\alpha}{2}} \sigma_r \zeta_r \right\|^2 \\ & \quad + \int \left[ 4r^3 \sigma^{2+\alpha} + (1+\alpha) r^4 \sigma^{1+\alpha} \sigma_r \right] (\zeta_r^2)_r dr + 8(1+\alpha) \int r^3 \sigma^{1+\alpha} \sigma_r \zeta_r^2 dr. \quad (3.14) \end{aligned}$$

With the help of the integration by parts and the fact  $\sigma_r = -2Br$ , one has

$$\begin{aligned} & \int \left[ 4r^3 \sigma^{2+\alpha} + (1+\alpha) r^4 \sigma^{1+\alpha} \sigma_r \right] (\zeta_r^2)_r dr \\ & = - \int \left[ 4r^3 \sigma^{2+\alpha} + (1+\alpha) r^4 \sigma^{1+\alpha} \sigma_r \right]_r \zeta_r^2 dr \\ & \geq -12 \int r^2 \sigma^{2+\alpha} \zeta_r^2 dr - (1+\alpha)^2 \int r^4 \sigma^\alpha \sigma_r^2 \zeta_r^2 dr - C \int r^4 \sigma^{1+\alpha} \zeta_r^2 dr. \end{aligned}$$

Substitute this into (3.14) and use  $\sigma_r = -2Br$  to give

$$\begin{aligned} & \left\| r^2 \sigma^{1+\frac{\alpha}{2}} \zeta_{rr} + 4r \sigma^{1+\frac{\alpha}{2}} \zeta_r + (1+\alpha) r^2 \sigma^{\frac{\alpha}{2}} \sigma_r \zeta_r \right\|^2 \\ & \geq \left\| r^2 \sigma^{1+\frac{\alpha}{2}} \zeta_{rr} \right\|^2 + 4 \left\| r \sigma^{1+\frac{\alpha}{2}} \zeta_r \right\|^2 - C \int r^4 \sigma^{1+\alpha} \zeta_r^2 dr. \end{aligned}$$

In view of (3.13), we then see that

$$\begin{aligned} & \left\| r^2 \sigma^{1+\frac{\alpha}{2}} \zeta_{rr} \right\|^2 + 4 \left\| r \sigma^{1+\frac{\alpha}{2}} \zeta_r \right\|^2 \\ & \lesssim \mathcal{E}_0 + \mathcal{E}_1 + \epsilon_0^2 \left( \left\| r^2 \sigma^{1+\frac{\alpha}{2}} \zeta_{rr} \right\|^2 + \left\| r \sigma^{1+\frac{\alpha}{2}} \zeta_r \right\|^2 + \left\| r^2 \sigma^{\frac{\alpha}{2}} \sigma_r \zeta_r \right\|^2 \right). \end{aligned} \quad (3.15)$$

On the other hand, it follows from (3.13) and (3.15) that

$$\begin{aligned} & \left\| (1+\alpha) r^2 \sigma^{\frac{\alpha}{2}} \sigma_r \zeta_r \right\|^2 \\ & \leq 2 \left\| r^2 \sigma^{1+\frac{\alpha}{2}} \zeta_{rr} + 4r \sigma^{1+\frac{\alpha}{2}} \zeta_r + (1+\alpha) r^2 \sigma^{\frac{\alpha}{2}} \sigma_r \zeta_r \right\|^2 + 2 \left\| r^2 \sigma^{1+\frac{\alpha}{2}} \zeta_{rr} + 4r \sigma^{1+\frac{\alpha}{2}} \zeta_r \right\|^2 \\ & \lesssim \mathcal{E}_0 + \mathcal{E}_1 + \epsilon_0^2 \left( \left\| r^2 \sigma^{1+\frac{\alpha}{2}} \zeta_{rr} \right\|^2 + \left\| r \sigma^{1+\frac{\alpha}{2}} \zeta_r \right\|^2 + \left\| r^2 \sigma^{\frac{\alpha}{2}} \sigma_r \zeta_r \right\|^2 \right). \end{aligned}$$

This, together with (3.15), gives

$$\begin{aligned} & \left\| r^2 \sigma^{1+\frac{\alpha}{2}} \zeta_{rr} \right\|^2 + \left\| r \sigma^{1+\frac{\alpha}{2}} \zeta_r \right\|^2 + \left\| r^2 \sigma^{\frac{\alpha}{2}} \sigma_r \zeta_r \right\|^2 \\ & \lesssim \mathcal{E}_0 + \mathcal{E}_1 + \epsilon_0^2 \left( \left\| r^2 \sigma^{1+\frac{\alpha}{2}} \zeta_{rr} \right\|^2 + \left\| r \sigma^{1+\frac{\alpha}{2}} \zeta_r \right\|^2 + \left\| r^2 \sigma^{\frac{\alpha}{2}} \sigma_r \zeta_r \right\|^2 \right), \end{aligned} \quad (3.16)$$

which implies, with the aid of the smallness of  $\epsilon_0$ , that

$$\left\| r^2 \sigma^{1+\frac{\alpha}{2}} \zeta_{rr} \right\|^2 + \left\| r \sigma^{1+\frac{\alpha}{2}} \zeta_r \right\|^2 + \left\| r^2 \sigma^{\frac{\alpha}{2}} \sigma_r \zeta_r \right\|^2 \lesssim \mathcal{E}_0 + \mathcal{E}_1.$$

In view of  $\sigma_r = -2Br$ , we then see that

$$\left\| r^2 \sigma^{1+\frac{\alpha}{2}} \zeta_{rr} \right\|^2 + \left\| r \sigma^{1+\frac{\alpha}{2}} \zeta_r \right\|^2 + \left\| r^3 \sigma^{\frac{\alpha}{2}} \zeta_r \right\|^2 \lesssim \mathcal{E}_0 + \mathcal{E}_1,$$

which implies

$$\begin{aligned} \left\| r \sigma^{\frac{\alpha}{2}} \zeta_r \right\|^2 &= \int_{\mathcal{I}_o} r^2 \sigma^\alpha \zeta_r^2 dr + \int_{\mathcal{I}_b} r^2 \sigma^\alpha \zeta_r^2 dr \\ &\lesssim \int_{\mathcal{I}_o} r^2 \sigma^{2+\alpha} \zeta_r^2 dr + \int_{\mathcal{I}_b} r^6 \sigma^\alpha \zeta_r^2 dr \lesssim \mathcal{E}_0 + \mathcal{E}_1. \end{aligned} \quad (3.17)$$

This finishes the proof of Lemma 3.2.  $\square$

### 3.2.2 Higher-order elliptic estimates

For  $i \geq 1$  and  $j \geq 0$ , it yields from  $\partial_t^j \partial_r^{i-1}$ (3.9) and  $\sigma_r = -2Br$  that

$$\begin{aligned} & \gamma \tilde{\eta}_r^{1-3\gamma} \left[ r \sigma \partial_t^j \partial_r^{i+1} \zeta + (i+3) \sigma \partial_t^j \partial_r^i \zeta + (\alpha+i) r \sigma_r \partial_t^j \partial_r^i \zeta \right] \\ & = r \partial_t^{j+2} \partial_r^{i-1} \zeta + r \partial_t^{j+1} \partial_r^{i-1} \zeta + \mathfrak{P}_1 + \mathfrak{P}_2. \end{aligned} \quad (3.18)$$

where

$$\begin{aligned}
\mathfrak{P}_1 := & -\gamma \sum_{\iota=1}^j [\partial_t^\iota (\tilde{\eta}_r^{1-3\gamma})] \partial_t^{j-\iota} [r\sigma \partial_r^{i+1} \zeta + (i+3)\sigma \partial_r^i \zeta + (\alpha+i)r\sigma_r \partial_r^i \zeta] \\
& - \gamma \partial_t^j \left\{ \tilde{\eta}_r^{1-3\gamma} \left[ \sum_{\iota=2}^{i-1} C_{i-1}^\iota [\partial_r^\iota (r\sigma)] \partial_r^{i+1-\iota} \zeta + 4 \sum_{\iota=1}^{i-1} C_{i-1}^\iota (\partial_r^\iota \sigma) \partial_r^{i-\iota} \zeta \right. \right. \\
& \left. \left. + (\alpha+1) \sum_{\iota=1}^{i-1} C_{i-1}^\iota [\partial_r^\iota (r\sigma_r)] \partial_r^{i-\iota} \zeta \right] \right\} + (i-1) \partial_r^{i-2} (\partial_t^{j+2} \zeta + \partial_t^{j+1} \zeta) \\
& - \frac{2\gamma(2-3\gamma)B}{\gamma-1} \partial_t^j [\tilde{\eta}_r^{1-3\gamma} (r\partial_r^{i-1} \zeta + (i-1)\partial_r^{i-2} \zeta)],
\end{aligned} \tag{3.19}$$

$$\mathfrak{P}_2 := \partial_r^{i-1} (\sigma \partial_t^j \mathfrak{J}_1) + (1+\alpha) \partial_r^{i-1} (\sigma_r \partial_t^j \mathfrak{J}_2). \tag{3.20}$$

(Recall that  $\mathfrak{J}_1$  and  $\mathfrak{J}_2$  are defined in (3.8).) Here and thereafter  $C_m^j$  is used to denote the binomial coefficients for  $0 \leq j \leq m$ ,

$$C_m^j = \frac{m!}{j!(m-j)!}$$

and summations  $\sum_{\iota=1}^{i-1}$  and  $\sum_{\iota=2}^{i-1}$  should be understood as zero when  $i = 1$  and  $i = 1, 2$ , respectively. Multiply equation (3.18) by  $\tilde{\eta}_r^{3\gamma-1} r\sigma^{(\alpha+i-1)/2}$ , square the spatial  $L^2$ -norm of the product and use (2.13) to give

$$\begin{aligned}
& \left\| r^2 \sigma^{\frac{\alpha+i+1}{2}} \partial_t^j \partial_r^{i+1} \zeta + (i+3) r \sigma^{\frac{\alpha+i+1}{2}} \partial_t^j \partial_r^i \zeta + (\alpha+i) r^2 \sigma^{\frac{\alpha+i-1}{2}} \sigma_r \partial_t^j \partial_r^i \zeta \right\|^2 \\
& \lesssim (1+t)^2 \left( \left\| r^2 \sigma^{\frac{\alpha+i-1}{2}} \partial_t^{j+2} \partial_r^{i-1} \zeta \right\|^2 + \left\| r^2 \sigma^{\frac{\alpha+i-1}{2}} \partial_t^{j+1} \partial_r^{i-1} \zeta \right\|^2 \right) \\
& + (1+t)^2 \left( \left\| r \sigma^{\frac{\alpha+i-1}{2}} \mathfrak{P}_1 \right\|^2 + \left\| r \sigma^{\frac{\alpha+i-1}{2}} \mathfrak{P}_2 \right\|^2 \right).
\end{aligned}$$

Similar to the derivation of (3.16) and (3.17), we can then obtain

$$\begin{aligned}
(1+t)^{-2j} \mathcal{E}_{j,i}(t) & = \left\| r^2 \sigma^{\frac{\alpha+i+1}{2}} \partial_t^j \partial_r^{i+1} \zeta \right\|^2 + \left\| r \sigma^{\frac{\alpha+i-1}{2}} \partial_t^j \partial_r^i \zeta \right\|^2 \\
& \lesssim \left\| r^2 \sigma^{\frac{\alpha+i}{2}} \partial_t^j \partial_r^i \zeta \right\|^2 + (1+t)^2 \left( \left\| r^2 \sigma^{\frac{\alpha+i-1}{2}} \partial_t^{j+2} \partial_r^{i-1} \zeta \right\|^2 + \left\| r^2 \sigma^{\frac{\alpha+i-1}{2}} \partial_t^{j+1} \partial_r^{i-1} \zeta \right\|^2 \right) \\
& + (1+t)^2 \left( \left\| r \sigma^{\frac{\alpha+i-1}{2}} \mathfrak{P}_1 \right\|^2 + \left\| r \sigma^{\frac{\alpha+i-1}{2}} \mathfrak{P}_2 \right\|^2 \right).
\end{aligned} \tag{3.21}$$

We will use this estimate to prove the following lemma by the mathematical induction.

**Lemma 3.3** *Assume that (3.1) holds for suitably small positive number  $\epsilon_0 \in (0, 1)$ . Then for  $j \geq 0$ ,  $i \geq 1$  and  $2 \leq i+j \leq l$ ,*

$$\mathcal{E}_{j,i}(t) \lesssim \sum_{\iota=0}^{i+j} \mathcal{E}_\iota(t), \quad t \in [0, T]. \tag{3.22}$$



**Proof.** We use the induction for  $i + j$  to prove this lemma. As shown in Lemma 3.2 we know that (3.22) holds for  $i + j = 1$ . For  $1 \leq k \leq l - 1$ , we make the induction hypothesis that (3.22) holds for all  $j \geq 0$ ,  $i \geq 1$  and  $i + j \leq k$ , that is,

$$\mathcal{E}_{j,i}(t) \lesssim \sum_{\iota=0}^{i+j} \mathcal{E}_{\iota}(t), \quad j \geq 0, \quad i \geq 1, \quad i + j \leq k. \quad (3.23)$$

It then suffices to prove (3.22) for  $j \geq 0$ ,  $i \geq 1$  and  $i + j = k + 1$ . (Indeed, there exists an order of  $(i, j)$  for the proof. For example, when  $i + j = k + 1$  we will bound  $\mathcal{E}_{k+1-\iota, \iota}$  from  $\iota = 1$  to  $k + 1$  step by step.)

Before going to the estimate, we notice a fact that  $\mathcal{E}_{j,0} \lesssim \mathcal{E}_j$  for  $j = 0, \dots, l$ . Indeed, it follows from (3.6) that

$$\begin{aligned} \int_{\mathcal{I}_b} \sigma^{\alpha-1} (\partial_t^j \zeta)^2 dr &\lesssim \int_{\mathcal{I}_b} \sigma^{\alpha+1} \left[ (\partial_t^j \zeta)^2 + (\partial_t^j \zeta_r)^2 \right] dr \\ &\lesssim \int_{\mathcal{I}_b} \sigma^{\alpha+1} \left[ r^2 (\partial_t^j \zeta)^2 + r^4 (\partial_t^j \zeta_r)^2 \right] dr \leq (1+t)^{-2j} \mathcal{E}_j(t), \end{aligned}$$

which implies

$$\begin{aligned} \mathcal{E}_{j,0}(t) &= (1+t)^{2j} \int \left[ r^2 \sigma^{\alpha-1} (\partial_t^j \zeta)^2 + r^4 \sigma^{\alpha+1} (\partial_t^j \zeta_r)^2 \right] (r, t) dr \\ &\leq (1+t)^{2j} \left[ \int_{\mathcal{I}_o} r^2 \sigma^{\alpha-1} (\partial_t^j \zeta)^2 (r, t) dr + \int_{\mathcal{I}_b} r^2 \sigma^{\alpha-1} (\partial_t^j \zeta)^2 (r, t) dr \right] + \mathcal{E}_j(t) \\ &\lesssim (1+t)^{2j} \left[ \int_{\mathcal{I}_o} r^2 \sigma^{\alpha+1} (\partial_t^j \zeta)^2 (r, t) dr + \int_{\mathcal{I}_b} \sigma^{\alpha-1} (\partial_t^j \zeta)^2 (r, t) dr \right] + \mathcal{E}_j(t) \\ &\lesssim \mathcal{E}_j(t), \quad j = 0, 1, \dots, l. \end{aligned} \quad (3.24)$$

This, together with the induction hypothesis (3.23), gives

$$\mathcal{E}_{j,i}(t) \lesssim \sum_{\iota=0}^{i+j} \mathcal{E}_{\iota}(t), \quad j \geq 0, \quad i \geq 0, \quad i + j \leq k. \quad (3.25)$$

In what follows, we assume  $j \geq 0$ ,  $i \geq 1$  and  $i + j = k + 1 \leq l$ . First, We estimate  $\mathfrak{P}_1$  and  $\mathfrak{P}_2$  given by (3.19) and (3.20), respectively. For  $\mathfrak{P}_1$ , it follows from (2.13) and  $\sigma_r = -2Br$  that

$$\begin{aligned} |\mathfrak{P}_1| &\lesssim \sum_{\iota=1}^j (1+t)^{-1-\iota} \left( |rw \partial_t^{j-\iota} \partial_r^{i+1} \zeta| + |\partial_t^{j-\iota} \partial_r^i \zeta| \right) + \sum_{\iota=0}^j \sum_{m=1}^{i-1} (1+t)^{-1-\iota} |\partial_t^{j-\iota} \partial_r^m \zeta| \\ &\quad + (i-1) \left( |\partial_t^{j+2} \partial_r^{i-2} \zeta| + |\partial_t^{j+1} \partial_r^{i-2} \zeta| + \sum_{\iota=0}^j (1+t)^{-1-\iota} |\partial_t^{j-\iota} \partial_r^{i-2} \zeta| \right) \\ &\quad + \sum_{\iota=0}^j (1+t)^{-1-\iota} |r \partial_t^{j-\iota} \partial_r^{i-1} \zeta|, \end{aligned}$$

which implies

$$\begin{aligned} \left\| r\sigma^{\frac{\alpha+i-1}{2}} \mathfrak{P}_1 \right\|^2 &\lesssim \sum_{\iota=1}^j (1+t)^{-2-2\iota} \left( \left\| r^2\sigma^{\frac{\alpha+i+1}{2}} \partial_t^{j-\iota} \partial_r^{i+1} \zeta \right\|^2 + \left\| r\sigma^{\frac{\alpha+i-1}{2}} \partial_t^{j-\iota} \partial_r^i \zeta \right\|^2 \right) \\ &\quad + \sum_{\iota=0}^j (1+t)^{-2-2\iota} \left( \sum_{m=1}^{i-1} \left\| r\sigma^{\frac{\alpha+i-1}{2}} \partial_t^{j-\iota} \partial_r^m \zeta \right\|^2 + \left\| r^2\sigma^{\frac{\alpha+i-1}{2}} \partial_t^{j-\iota} \partial_r^{i-1} \zeta \right\|^2 \right) \\ &\quad + (i-1)^2 \left( \sum_{\iota=j+1}^{j+2} \left\| r\sigma^{\frac{\alpha+i-1}{2}} \partial_t^\iota \partial_r^{i-2} \zeta \right\|^2 + \sum_{\iota=0}^j (1+t)^{-2-2\iota} \left\| r\sigma^{\frac{\alpha+i-1}{2}} \partial_t^{j-\iota} \partial_r^{i-2} \zeta \right\|^2 \right). \end{aligned}$$

So,

$$\left\| r\sigma^{\frac{\alpha+i-1}{2}} \mathfrak{P}_1 \right\|^2 \lesssim \begin{cases} (1+t)^{-2-2j} \left( \sum_{\iota=0}^{j-1} \mathcal{E}_{\iota,1} + \sum_{\iota=0}^j \mathcal{E}_\iota \right) (t), & i = 1, \\ (1+t)^{-2-2j} \left( \sum_{\iota=0}^{j-1} \mathcal{E}_{\iota,i} + \sum_{\iota=0}^j \sum_{m=1}^{i-1} \mathcal{E}_{\iota,m} + \sum_{\iota=0}^{j+2} \mathcal{E}_{\iota,i-2} \right) (t), & i \geq 2. \end{cases} \quad (3.26)$$

For  $\mathfrak{P}_2$ , it follows from (2.13), (3.1), (3.2) and  $\sigma_r = -2Br$  that

$$\begin{aligned} |\mathfrak{P}_2| &\lesssim \sum_{n=0}^j \sum_{m=0}^{i-1} K_{nm} \left( \left| \partial_t^{j-n} \partial_r^{i-1-m} (\sigma_r \zeta_{rr}) \right| + \left| \partial_t^{j-n} \partial_r^{i-1-m} (\sigma \zeta_r) \right| \right. \\ &\quad \left. + \left| \partial_t^{j-n} \partial_r^{i-1-m} (\sigma_r r \zeta_r) \right| + \left| \partial_t^{j-n} \partial_r^{i-1-m} (\sigma_r \zeta) \right| \right) \\ &\lesssim \sum_{n=0}^j \sum_{m=0}^{i-1} K_{nm} \left( \left| \sigma_r \partial_t^{j-n} \partial_r^{i-m+1} \zeta \right| + \sum_{\iota=0}^{i-m} \left| \partial_t^{j-n} \partial_r^\iota \zeta \right| \right) =: \sum_{n=0}^j \sum_{m=0}^{i-1} \mathfrak{P}_{2nm}, \end{aligned}$$

where

$$\begin{aligned} K_{00} &= \epsilon_0 (1+t)^{-1-\frac{1}{3\gamma-1}}; \\ K_{10} &= \epsilon_0 (1+t)^{-2-\frac{1}{3\gamma-1}}, \quad K_{01} = (1+t)^{-1-\frac{1}{3\gamma-1}} (\epsilon_0 + |r \partial_r^2 \zeta|); \\ K_{20} &= \epsilon_0 (1+t)^{-3-\frac{1}{3\gamma-1}} + (1+t)^{-1-\frac{1}{3\gamma-1}} |r \partial_t^2 \partial_r \zeta|, \\ K_{11} &= (1+t)^{-2-\frac{1}{3\gamma-1}} (\epsilon_0 + |r \partial_r^2 \zeta|) + (1+t)^{-1-\frac{1}{3\gamma-1}} |r \partial_t \partial_r^2 \zeta|, \\ K_{02} &= (1+t)^{-1-\frac{1}{3\gamma-1}} (|\partial_r^2 \zeta| + |r \partial_r^3 \zeta|) + (1+t)^{-1-\frac{2}{3\gamma-1}} (\epsilon_0^2 + |r \partial_r^2 \zeta|^2). \end{aligned}$$

We do not list here  $K_{nm}$  for  $n+m \geq 3$  since we can use the same method to estimate  $\mathfrak{P}_{2nm}$  for  $n+m \geq 3$  as that for  $n+m \leq 2$ . Easily,  $\mathfrak{P}_{200}$  and  $\mathfrak{P}_{210}$  can be bounded by

$$\begin{aligned} \left\| r\sigma^{\frac{\alpha+i-1}{2}} \mathfrak{P}_{200} \right\|^2 &\lesssim \epsilon_0^2 (1+t)^{-2} \left( \left\| r^2\sigma^{\frac{\alpha+i+1}{2}} \partial_t^j \partial_r^{i+1} \zeta \right\|^2 + \sum_{\iota=0}^i \left\| r\sigma^{\frac{\alpha+i-1}{2}} \partial_t^j \partial_r^\iota \zeta \right\|^2 \right) \\ &\lesssim \epsilon_0^2 (1+t)^{-2-2j} \left( \mathcal{E}_{j,i} + \sum_{\iota=0}^{i-1} \mathcal{E}_{j,\iota} \right) (t), \end{aligned}$$

$$\begin{aligned} \left\| r\sigma^{\frac{\alpha+i-1}{2}} \mathfrak{P}_{210} \right\|^2 &\lesssim \epsilon_0^2 (1+t)^{-4} \left( \left\| r^2 \sigma^{\frac{\alpha+i+1}{2}} \partial_t^{j-1} \partial_r^{i+1} \zeta \right\|^2 + \sum_{\iota=0}^i \left\| r\sigma^{\frac{\alpha+i-1}{2}} \partial_t^{j-1} \partial_r^\iota \zeta \right\|^2 \right) \\ &\lesssim \epsilon_0^2 (1+t)^{-2-2j} \sum_{\iota=0}^i \mathcal{E}_{j-1,\iota}(t). \end{aligned}$$

For  $\mathfrak{P}_{201}$ , we use (3.1) to get  $|\sigma^{1/2} \partial_r^2 \zeta| \lesssim \epsilon_0$  and then obtain

$$\begin{aligned} \left\| r\sigma^{\frac{\alpha+i-1}{2}} \mathfrak{P}_{201} \right\|^2 &\lesssim \epsilon_0^2 (1+t)^{-2} \left( \left\| r^2 \sigma^{\frac{\alpha+i}{2}} \partial_t^j \partial_r^i \zeta \right\|^2 + \sum_{\iota=0}^{i-1} \left\| r\sigma^{\frac{\alpha+i-2}{2}} \partial_t^j \partial_r^\iota \zeta \right\|^2 \right) \\ &\lesssim \epsilon_0^2 (1+t)^{-2-2j} \sum_{\iota=0}^{i-1} \mathcal{E}_{j,\iota}(t), \end{aligned}$$

For  $\mathfrak{P}_{220}$ , we use (3.1) again to get  $|r\sigma^{1/2} \partial_t^2 \partial_r \zeta| \lesssim \epsilon_0 (1+t)^{-2}$  and then achieve

$$\begin{aligned} \left\| r\sigma^{\frac{\alpha+i-1}{2}} \mathfrak{P}_{220} \right\|^2 &\lesssim \epsilon_0^2 (1+t)^{-6} \left( \left\| r^2 \sigma^{\frac{\alpha+i}{2}} \partial_t^{j-2} \partial_r^{i+1} \zeta \right\|^2 + \sum_{\iota=0}^i \left\| r\sigma^{\frac{\alpha+i-2}{2}} \partial_t^{j-2} \partial_r^\iota \zeta \right\|^2 \right) \\ &\lesssim \epsilon_0^2 (1+t)^{-2-2j} \sum_{\iota=0}^{i+1} \mathcal{E}_{j-2,\iota}(t), \end{aligned}$$

because it can be derived from (3.6) that

$$\begin{aligned} \left\| r\sigma^{\frac{\alpha+i-2}{2}} \partial_t^{j-2} \partial_r^i \zeta \right\|^2 &= \int_{\mathcal{I}_o} r^2 \sigma^{\alpha+i-2} |\partial_t^{j-2} \partial_r^i \zeta|^2 dr + \int_{\mathcal{I}_b} r^2 \sigma^{\alpha+i-2} |\partial_t^{j-2} \partial_r^i \zeta|^2 dr \\ &\lesssim \int_{\mathcal{I}_o} r^2 \sigma^{\alpha+i-1} |\partial_t^{j-2} \partial_r^i \zeta|^2 dr + \int_{\mathcal{I}_b} \sigma^{\alpha+i-2} |\partial_t^{j-2} \partial_r^i \zeta|^2 dr \\ &\lesssim \int_{\mathcal{I}_o} r^2 \sigma^{\alpha+i-1} |\partial_t^{j-2} \partial_r^i \zeta|^2 dr + \int_{\mathcal{I}_b} \sigma^{\alpha+i} \left( |\partial_t^{j-2} \partial_r^i \zeta|^2 + |\partial_t^{j-2} \partial_r^{i+1} \zeta|^2 \right) dr \\ &\lesssim \int_{\mathcal{I}_o} r^2 \sigma^{\alpha+i-1} |\partial_t^{j-2} \partial_r^i \zeta|^2 dr + \int_{\mathcal{I}_b} \sigma^{\alpha+i} \left( r^2 |\partial_t^{j-2} \partial_r^i \zeta|^2 + r^2 |\partial_t^{j-2} \partial_r^{i+1} \zeta|^2 \right) dr \\ &\lesssim \int_{\mathcal{I}_o} r^2 \sigma^{\alpha+i-1} |\partial_t^{j-2} \partial_r^i \zeta|^2 dr + \int_{\mathcal{I}_b} r^2 \sigma^{\alpha+i} |\partial_t^{j-2} \partial_r^{i+1} \zeta|^2 dr. \end{aligned}$$

Similar to the estimate for  $\mathfrak{P}_{220}$ , we can obtain

$$\begin{aligned} \left\| r\sigma^{\frac{\alpha+i-1}{2}} \mathfrak{P}_{211} \right\|^2 &\lesssim \epsilon_0^2 (1+t)^{-4} \left( \left\| r^2 \sigma^{\frac{\alpha+i-1}{2}} \partial_t^{j-1} \partial_r^i \zeta \right\|^2 + \sum_{\iota=0}^{i-1} \left\| r\sigma^{\frac{\alpha+i-3}{2}} \partial_t^{j-1} \partial_r^\iota \zeta \right\|^2 \right) \\ &\lesssim \epsilon_0^2 (1+t)^{-2-2j} \sum_{\iota=0}^i \mathcal{E}_{j-1,\iota}(t), \\ \left\| r\sigma^{\frac{\alpha+i-1}{2}} \mathfrak{P}_{202} \right\|^2 &\lesssim \epsilon_0^2 (1+t)^{-2} \left( \left\| r^2 \sigma^{\frac{\alpha+i-2}{2}} \partial_t^j \partial_r^{i-1} \zeta \right\|^2 + \sum_{\iota=0}^{i-2} \left\| r\sigma^{\frac{\alpha+i-4}{2}} \partial_t^j \partial_r^\iota \zeta \right\|^2 \right) \\ &\lesssim \epsilon_0^2 (1+t)^{-2-2j} \sum_{\iota=0}^{i-1} \mathcal{E}_{j,\iota}(t). \end{aligned}$$

It should be noted that  $\mathfrak{P}_{211}$  and  $\mathfrak{P}_{202}$  appear when  $i \geq 2$  and  $i \geq 3$ , respectively. This ensures the application of the Hardy inequality (3.6). Other cases can be done similarly, since the leading term of  $K_{nm}$  is

$$\sum_{q=0}^n (1+t)^{-1-\frac{1}{3\gamma-1}-q} (|r\partial_t^{n-q}\partial_r^{m+1}\zeta| + |\partial_t^{n-q}\partial_r^m\zeta|)$$

and

$$\begin{aligned} & \sum_{n=0}^j \sum_{m=0}^{i-1} \sum_{q=0}^n (1+t)^{-2-2q} \left\| r\sigma^{\frac{\alpha+i-1}{2}} (|r\partial_t^{n-q}\partial_r^{m+1}\zeta| + |\partial_t^{n-q}\partial_r^m\zeta|) (|\sigma r\partial_t^{j-n}\partial_r^{i-m+1}\zeta| \right. \\ & \left. + \sum_{\iota=0}^{i-m} |\partial_t^{j-n}\partial_r^\iota\zeta|) \right\|^2 \lesssim \epsilon_0^2 (1+t)^{-2-2j} \left( \mathcal{E}_{j,i} + \sum_{0 \leq \iota \leq j, p \geq 0, \iota+p \leq i+j-1} \mathcal{E}_{\iota,p} \right) (t). \end{aligned} \quad (3.27)$$

(Estimate (3.27) will be verified in the Appendix.) Now, we may conclude that

$$\left\| r\sigma^{\frac{\alpha+i-1}{2}} \mathfrak{P}_2 \right\|^2 \lesssim \epsilon_0^2 (1+t)^{-2-2j} \left( \mathcal{E}_{j,i} + \sum_{0 \leq \iota \leq j, p \geq 0, \iota+p \leq i+j-1} \mathcal{E}_{\iota,p} \right) (t). \quad (3.28)$$

Substitute (3.26) and (3.28) into (3.21) gives, for suitably small  $\epsilon_0$ , that

$$\mathcal{E}_{j,i}(t) \lesssim \begin{cases} \mathcal{E}_j(t) + \mathcal{E}_{j+1}(t) + \sum_{\iota \geq 0, p \geq 0, \iota+p \leq j} \mathcal{E}_{\iota,p}(t) + \sum_{\iota=0}^j \mathcal{E}_\iota(t), & i = 1, \\ \mathcal{E}_{j,i-1}(t) + \mathcal{E}_{j+2,i-2}(t) + \mathcal{E}_{j+1,i-2}(t) + \sum_{0 \leq \iota \leq j, p \geq 0, \iota+p \leq i+j-1} \mathcal{E}_{\iota,p}(t), & i \geq 2. \end{cases} \quad (3.29)$$

Now, we use estimate (3.25), derived from the induction hypothesis (3.23), and (3.29) to show that (3.22) holds for  $i+j = k+1$ . First, choosing  $j = k$  and  $i = 1$  in (3.29) gives

$$\mathcal{E}_{k,1}(t) \lesssim \sum_{\iota=0}^{k+1} \mathcal{E}_\iota(t) + \sum_{\iota \geq 0, p \geq 0, \iota+p \leq k} \mathcal{E}_{\iota,p}(t) \lesssim \sum_{\iota=0}^{k+1} \mathcal{E}_\iota(t). \quad (3.30)$$

We choose  $j = k-1$  and  $i = 2$  in (3.29) and use (3.24)-(3.25) to show

$$\mathcal{E}_{k-1,2}(t) \lesssim \mathcal{E}_{k-1,1}(t) + \mathcal{E}_{k+1,0}(t) + \mathcal{E}_{k,0}(t) + \sum_{0 \leq \iota \leq k-1, p \geq 0, \iota+p \leq k} \mathcal{E}_{\iota,p}(t) \lesssim \sum_{\iota=0}^{k+1} \mathcal{E}_\iota(t).$$

For  $\mathcal{E}_{k-2,3}$ , it follows from (3.29), (3.25) and (3.30) to obtain

$$\mathcal{E}_{k-2,3}(t) \lesssim \mathcal{E}_{k-2,2}(t) + \mathcal{E}_{k,1}(t) + \mathcal{E}_{k-1,1}(t) + \sum_{0 \leq \iota \leq k-2, p \geq 0, \iota+p \leq k} \mathcal{E}_{\iota,p}(t) \lesssim \sum_{\iota=0}^{k+1} \mathcal{E}_\iota(t).$$

The other cases can be handled similarly. So we have proved (3.22) when  $i+j = k+1$ . This finishes the proof of Lemma 3.3.  $\square$

### 3.3 Nonlinear weighted energy estimates

In this section, we show that the weighted energy  $\mathcal{E}_j(t)$  can be bounded by the initial data for all  $t \in [0, T]$ .

**Proposition 3.4** *Suppose that (3.1) holds for suitably small positive number  $\epsilon_0 \in (0, 1)$ . Then it holds that for  $t \in [0, T]$ ,*

$$\mathcal{E}_j(t) \lesssim \sum_{\iota=0}^j \mathcal{E}_\iota(0), \quad j = 0, 1, \dots, l. \quad (3.31)$$

The proof of this proposition consists of Lemma 3.5 and Lemma 3.6.

#### 3.3.1 Basic energy estimates

**Lemma 3.5** *Assume that (3.1) holds for suitably small positive number  $\epsilon_0 \in (0, 1)$ . Then,*

$$\mathcal{E}_0(t) + \int_0^t \int [(1+s)^{-1} r^2 \bar{\rho}_0^\gamma (\zeta^2 + (r\zeta_r)^2) + (1+s)r^4 \bar{\rho}_0 \zeta_s^2] dr ds \lesssim \mathcal{E}_0(0), \quad t \in [0, T]. \quad (3.32)$$

**Proof.** Multiplying (2.16) by  $r^3 \zeta_t$ , and integrating the product with respect to the spatial variable, we obtain, using the integration by parts, that

$$\frac{d}{dt} \int \frac{1}{2} r^4 \bar{\rho}_0 \zeta_t^2 dr + \int r^4 \bar{\rho}_0 \zeta_t^2 dr + \int \bar{\rho}_0^\gamma \mathfrak{L}_1 dr = 0 \quad (3.33)$$

where

$$\mathfrak{L}_1 := -(\tilde{\eta}_r + \zeta)^{-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma} [r^3 (\tilde{\eta}_r + \zeta)^2 \zeta_t]_r + \tilde{\eta}_r^{2-3\gamma} (r^3 \zeta_t)_r =: -\mathfrak{L}_{11} + \mathfrak{L}_{12}.$$

For  $\mathfrak{L}_{11}$ , note that

$$\begin{aligned} & [r^3 (\tilde{\eta}_r + \zeta)^2 \zeta_t]_r \\ &= 3r^2 (\tilde{\eta}_r + \zeta)^2 \zeta_t + 2r^2 (\tilde{\eta}_r + \zeta) (r\zeta_r) \zeta_t + r^2 (\tilde{\eta}_r + \zeta)^2 (r\zeta_{rt}) \\ &= 2r^2 (\tilde{\eta}_r + \zeta) (\tilde{\eta}_r + \zeta + r\zeta_r) \zeta_t + r^2 (\tilde{\eta}_r + \zeta)^2 (\zeta + r\zeta_r)_t, \end{aligned}$$

thus,

$$\begin{aligned} \mathfrak{L}_{11} &= 2r^2 (\tilde{\eta}_r + \zeta)^{1-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{1-\gamma} \zeta_t + r^2 (\tilde{\eta}_r + \zeta)^{2-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma} (\zeta + r\zeta_r)_t \\ &= \frac{r^2}{1-\gamma} [(\tilde{\eta}_r + \zeta)^{2-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{1-\gamma}]_t \\ &\quad - r^2 [2(\tilde{\eta}_r + \zeta)^{1-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{1-\gamma} + (\tilde{\eta}_r + \zeta)^{2-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma}] \tilde{\eta}_{rt}. \end{aligned}$$

Clearly,  $\mathfrak{L}_{12}$  can be rewritten as

$$\mathfrak{L}_{12} = r^2 (3\zeta + r\zeta_r)_t \tilde{\eta}_r^{2-3\gamma} = r^2 [(3\zeta + r\zeta_r) \tilde{\eta}_r^{2-3\gamma}]_t - (2-3\gamma)r^2 (3\zeta + r\zeta_r) \tilde{\eta}_r^{1-3\gamma} \tilde{\eta}_{rt}.$$

Substitute these calculations into (3.33) to give

$$\frac{d}{dt} \int \left( \frac{1}{2} r^4 \bar{\rho}_0 \zeta_t^2 + r^2 \bar{\rho}_0^\gamma \tilde{\mathfrak{E}}_0 \right) dr + \int r^4 \bar{\rho}_0 \zeta_t^2 dr + \int r^2 \bar{\rho}_0^\gamma \tilde{\eta}_{rt} \mathfrak{F} dr = 0, \quad (3.34)$$

where

$$\begin{aligned} \tilde{\mathfrak{E}}_0 &:= \frac{1}{\gamma - 1} \left[ (\tilde{\eta}_r + \zeta)^{2-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{1-\gamma} - \tilde{\eta}_r^{3-3\gamma} + (\gamma - 1) (3\zeta + r\zeta_r) \tilde{\eta}_r^{2-3\gamma} \right], \\ \mathfrak{F} &:= 2 (\tilde{\eta}_r + \zeta)^{1-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{1-\gamma} + (\tilde{\eta}_r + \zeta)^{2-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma} \\ &\quad - 3\tilde{\eta}_r^{2-3\gamma} - (2 - 3\gamma) (3\zeta + r\zeta_r) \tilde{\eta}_r^{1-3\gamma}. \end{aligned}$$

It follows from the Taylor expansion, the smallness of  $\zeta$  and  $r\zeta_r$  which is a consequence of (3.1), and (2.13) that

$$\begin{aligned} \tilde{\mathfrak{E}}_0 &= \tilde{\eta}_r^{1-3\gamma} \left[ \frac{3}{2} (3\gamma - 2) \zeta^2 + (3\gamma - 2) \zeta r\zeta_r + \frac{\gamma}{2} (r\zeta_r)^2 \right] \\ &\quad + O(1) \tilde{\eta}_r^{-3\gamma} (|\zeta| + |r\zeta_r|) (\zeta^2 + (r\zeta_r)^2) \\ &\sim \tilde{\eta}_r^{1-3\gamma} (\zeta^2 + (r\zeta_r)^2) \sim (1+t)^{-1} (\zeta^2 + (r\zeta_r)^2), \\ \mathfrak{F} &\geq (3\gamma - 1) \tilde{\eta}_r^{-3\gamma} \left[ \frac{3}{2} (3\gamma - 2) \zeta^2 + (3\gamma - 2) \zeta r\zeta_r + \frac{\gamma}{2} (r\zeta_r)^2 \right] \\ &\quad - C \tilde{\eta}_r^{-3\gamma-1} (|\zeta| + |r\zeta_r|) (\zeta^2 + (r\zeta_r)^2) \\ &\geq (1+t)^{-\frac{3\gamma}{3\gamma-1}} (\zeta^2 + (r\zeta_r)^2) \geq 0. \end{aligned}$$

Here and thereafter the notation  $O(1)$  represents a finite number could be positive or negative. We then have, by integrating (3.34) with respect to the temporal variable, that

$$\int \left( \frac{1}{2} r^4 \bar{\rho}_0 \zeta_t^2 + r^2 \bar{\rho}_0^\gamma \tilde{\mathfrak{E}}_0 \right) (r, s) dr \Big|_{s=0}^t + \int_0^t \int r^4 \bar{\rho}_0 \zeta_s^2 dr ds \leq 0$$

and

$$\begin{aligned} &\int [r^4 \bar{\rho}_0 \zeta_t^2 + (1+t)^{-1} r^2 \bar{\rho}_0^\gamma (\zeta^2 + (r\zeta_r)^2)] (r, t) dr + \int_0^t \int r^4 \bar{\rho}_0 \zeta_s^2 dr ds \\ &\lesssim \int [r^4 \bar{\rho}_0 \zeta_t^2 + r^2 \bar{\rho}_0^\gamma (\zeta^2 + (r\zeta_r)^2)] (r, 0) dr. \end{aligned} \quad (3.35)$$

Multiplying (2.16) by  $r^3 \zeta$ , and integrating the product with respect to the spatial variable, we have, using the integration by parts, that

$$\frac{d}{dt} \int r^4 \bar{\rho}_0 \left( \frac{1}{2} \zeta^2 + \zeta \zeta_t \right) dr + \int \bar{\rho}_0^\gamma \mathfrak{L}_2 dr = \int r^4 \bar{\rho}_0 \zeta_t^2 dr, \quad (3.36)$$

where

$$\mathfrak{L}_2 := - (\tilde{\eta}_r + \zeta)^{-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma} [r^3 (\tilde{\eta}_r + \zeta)^2 \zeta]_r + \tilde{\eta}_r^{2-3\gamma} (r^3 \zeta)_r,$$

which can be rewritten as

$$\begin{aligned}
\mathfrak{L}_2 &= -2r^2 (\tilde{\eta}_r + \zeta)^{1-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{1-\gamma} \zeta - r^2 (\tilde{\eta}_r + \zeta)^{2-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma} \\
&\quad \times (\zeta + r\zeta_r) + r^2 \tilde{\eta}_r^{2-3\gamma} (3\zeta + r\zeta_r) \\
&= r^2 \left[ 3\tilde{\eta}_r^{2-3\gamma} - 2(\tilde{\eta}_r + \zeta)^{1-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{1-\gamma} - (\tilde{\eta}_r + \zeta)^{2-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma} \right] \zeta \\
&\quad + r^2 \left[ \tilde{\eta}_r^{2-3\gamma} - (\tilde{\eta}_r + \zeta)^{2-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma} \right] r\zeta_r.
\end{aligned}$$

Again, we use the Taylor expansion, (3.2) and (3.1) to obtain

$$\begin{aligned}
\mathfrak{L}_2 &\gtrsim r^2 (1+t)^{-1} \left[ 3(3\gamma - 2)\zeta^2 + 2(3\gamma - 2)\zeta r\zeta_r + \gamma (r\zeta_r)^2 \right] \\
&\quad - Cr^2 (1+t)^{-\frac{3\gamma}{3\gamma-1}} (|\zeta| + |r\zeta_r|) (\zeta^2 + (r\zeta_r)^2) \\
&\gtrsim r^2 (1+t)^{-1} \left[ 3(3\gamma - 2)\zeta^2 + 2(3\gamma - 2)\zeta r\zeta_r + \gamma (r\zeta_r)^2 - C\epsilon_0 (\zeta^2 + (r\zeta_r)^2) \right] \\
&\gtrsim r^2 (1+t)^{-1} (\zeta^2 + (r\zeta_r)^2),
\end{aligned}$$

provide that  $\epsilon_0$  is suitably small. It then follows from (3.36), the Cauchy inequality and (3.35) that

$$\begin{aligned}
&\int (r^4 \bar{\rho}_0 \zeta^2) (r, t) dr + \int_0^t \int (1+s)^{-1} r^2 \bar{\rho}_0^\gamma (\zeta^2 + (r\zeta_r)^2) dr ds \\
&\lesssim \int (r^4 \bar{\rho}_0 (\zeta^2 + \zeta_t^2)) (r, 0) dr + \int (r^4 \bar{\rho}_0 \zeta_t^2) (r, t) dr + \int_0^t \int r^4 \bar{\rho}_0 \zeta_s^2 dr ds \quad (3.37) \\
&\lesssim \int [r^4 \bar{\rho}_0 (\zeta^2 + \zeta_t^2) + r^2 \bar{\rho}_0^\gamma (\zeta^2 + (r\zeta_r)^2)] (r, 0) dr = \mathcal{E}_0(0).
\end{aligned}$$

Next, we show the time decay of the energy norm. Multiply equation (3.34) by  $(1+t)$  and integrate the product with respect to the temporal variable to get

$$\begin{aligned}
&(1+t) \int \left( \frac{1}{2} r^4 \bar{\rho}_0 \zeta_t^2 + r^2 \bar{\rho}_0^\gamma \tilde{\mathfrak{E}}_0 \right) (r, t) dr + \int_0^t (1+s) \int r^4 \bar{\rho}_0 \zeta_s^2 dr ds \\
&\leq \int \left( \frac{1}{2} r^4 \bar{\rho}_0 \zeta_t^2 + r^2 \bar{\rho}_0^\gamma \tilde{\mathfrak{E}}_0 \right) (r, 0) dr + \int_0^t \int \left( \frac{1}{2} r^4 \bar{\rho}_0 \zeta_s^2 + \tilde{\mathfrak{E}}_0 \right) dr ds \\
&\lesssim \int \left( \frac{1}{2} r^4 \bar{\rho}_0 \zeta_t^2 + r^2 \bar{\rho}_0^\gamma \tilde{\mathfrak{E}}_0 \right) (r, 0) dr + \int_0^t \int [r^4 \bar{\rho}_0 \zeta_s^2 + (1+s)^{-1} r^2 \bar{\rho}_0^\gamma (\zeta^2 + (r\zeta_r)^2)] dr ds \\
&\lesssim \int [r^4 \bar{\rho}_0 (\zeta^2 + \zeta_t^2) + r^2 \bar{\rho}_0^\gamma (\zeta^2 + (r\zeta_r)^2)] (r, 0) dr = \mathcal{E}_0(0),
\end{aligned}$$

where estimates (3.35) and (3.37) have been used to derive the last inequality. This means

$$\int [(1+t)r^4 \bar{\rho}_0 \zeta_t^2 + r^2 \bar{\rho}_0^\gamma (\zeta^2 + (r\zeta_r)^2)] (r, t) dr + \int_0^t (1+s) \int r^4 \bar{\rho}_0 \zeta_s^2 dr ds \lesssim \mathcal{E}_0(0),$$

which, together with (3.37), gives (3.32). This finishes the proof of Lemma 3.5.  $\square$

### 3.3.2 Higher-order energy estimates

Equation (2.16) reads

$$\begin{aligned} & r\bar{\rho}_0\zeta_{tt} + r\bar{\rho}_0\zeta_t + [\bar{\rho}_0^\gamma (\tilde{\eta}_r + \zeta)^{2-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma}]_r - \tilde{\eta}_r^{2-3\gamma} (\bar{\rho}_0^\gamma)_r \\ & - 2\bar{\rho}_0^\gamma (\tilde{\eta}_r + \zeta)^{1-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma} \zeta_r = 0. \end{aligned}$$

Let  $k \geq 1$  be an integer and take the  $k$ -th time derivative of the equation above. One has

$$\begin{aligned} & r\bar{\rho}_0\partial_t^k\zeta_{tt} + r\bar{\rho}_0\partial_t^k\zeta_t + [\bar{\rho}_0^\gamma (w_1\partial_t^k\zeta + w_2r\partial_t^k\zeta_r + K_1)]_r + \bar{\rho}_0^\gamma [(3w_2 - w_1)\partial_t^k\zeta_r + K_2] \\ & - 2\bar{\rho}_0^\gamma (w_3\zeta_r\partial_t^k\zeta + K_3) + \partial_t^{k-1} \{ \bar{\rho}_0^\gamma \tilde{\eta}_{rt} [w_1 - (2 - 3\gamma)\tilde{\eta}_r^{1-3\gamma}] \}_r - 2\bar{\rho}_0^\gamma \partial_t^{k-1} (\tilde{\eta}_{rt}w_3\zeta_r) = 0. \end{aligned} \quad (3.38)$$

Here

$$\begin{aligned} w_1 &= (2 - 2\gamma) (\tilde{\eta}_r + \zeta)^{1-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma} - \gamma (\tilde{\eta}_r + \zeta)^{2-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma-1}, \\ w_2 &= -\gamma (\tilde{\eta}_r + \zeta)^{2-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma-1}, \\ w_3 &= (1 - 2\gamma) (\tilde{\eta}_r + \zeta)^{-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma} - \gamma (\tilde{\eta}_r + \zeta)^{1-2\gamma} (\tilde{\eta}_r + \zeta + r\zeta_r)^{-\gamma-1}; \end{aligned}$$

and

$$\begin{aligned} K_1 &= \partial_t^{k-1} (w_1\zeta_t + w_2r\zeta_{tr}) - (w_1\partial_t^k\zeta + w_2r\partial_t^k\zeta_r), \\ K_2 &= \partial_t^{k-1} [(3w_2 - w_1)\zeta_{tr}] - (3w_2 - w_1)\partial_t^k\zeta_r, \\ K_3 &= \partial_t^{k-1} (w_3\zeta_r\zeta_t) - w_3\zeta_r\partial_t^k\zeta. \end{aligned}$$

It should be noted that  $K_1$ ,  $K_2$  and  $K_3$  contain lower-order terms involving  $\partial_t^\iota(\zeta, \zeta_r)$  with  $\iota = 0, \dots, k-1$ ; and  $w_1$ ,  $w_2$  and  $w_3$  can be expanded, according to the Taylor expansion and the smallness of  $\zeta$  and  $r\zeta_r$  which is a consequence of (3.1), as follows

$$\begin{aligned} w_1 &= (2 - 3\gamma)\tilde{\eta}_r^{1-3\gamma} + (3\gamma - 1)\tilde{\eta}_r^{-3\gamma} [(3\gamma - 2)\zeta + \gamma r\zeta_r] + \bar{w}_1 \\ w_2 &= -\gamma\tilde{\eta}_r^{1-3\gamma} + \gamma\tilde{\eta}_r^{-3\gamma} [(3\gamma - 1)\zeta + (\gamma + 1)r\zeta_r] + \bar{w}_2, \\ w_3 &= (1 - 3\gamma)\tilde{\eta}_r^{-3\gamma} + \bar{w}_3. \end{aligned} \quad (3.39)$$

Here  $\bar{w}_i$  satisfies

$$|\bar{w}_1| + |\bar{w}_2| \lesssim \tilde{\eta}_r^{-3\gamma-1} (|\zeta|^2 + |r\zeta_r|^2), \quad \text{and} \quad |\bar{w}_3| \lesssim \tilde{\eta}_r^{-3\gamma-1} (|\zeta| + |r\zeta_r|). \quad (3.40)$$

In particular,  $K_1 = K_2 = K_3 = 0$  when  $k = 1$ .

**Lemma 3.6** *Assume that (3.1) holds for suitably small positive number  $\epsilon_0 \in (0, 1)$ . Then for all  $j = 1, \dots, l$ ,*

$$\begin{aligned} & \mathcal{E}_j(t) + \int_0^t \int \left[ (1+s)^{2j-1} r^2 \bar{\rho}_0^\gamma |\partial_s^j(\zeta, r\zeta_r)|^2 + (1+s)^{2j+1} r^4 \bar{\rho}_0 (\partial_s^j \zeta_s)^2 \right] dr ds \\ & \lesssim \sum_{\iota=0}^j \mathcal{E}_\iota(0), \quad t \in [0, T]. \end{aligned} \quad (3.41)$$



**Proof.** We use induction to prove (3.41). As shown in Lemma 3.5, we know that (3.41) holds for  $j = 0$ . For  $1 \leq k \leq l$ , we make the induction hypothesis that (3.41) holds for all  $j = 0, \dots, k-1$ , that is,

$$\begin{aligned} & \mathcal{E}_j(t) + \int_0^t \int \left[ (1+s)^{2j-1} r^2 \bar{\rho}_0^\gamma |\partial_s^j(\zeta, r\zeta_r)|^2 + (1+s)^{2j+1} r^4 \bar{\rho}_0 (\partial_s^j \zeta_s)^2 \right] dr ds \\ & \lesssim \sum_{\iota=0}^j \mathcal{E}_\iota(0) \quad \text{for all } j = 0, 1, \dots, k-1. \end{aligned} \quad (3.42)$$

It suffices to prove (3.41) holds for  $j = k$  under the induction hypothesis (3.42).

**Step 1.** In this step, we prove that

$$\begin{aligned} & \frac{d}{dt} \left[ \int \frac{1}{2} r^4 \bar{\rho}_0 (\partial_t^k \zeta_t)^2 dr + \bar{\mathfrak{E}}_k \right] + \int r^4 \bar{\rho}_0 (\partial_t^k \zeta_t)^2 dr \\ & \lesssim (\epsilon_0 + \delta)(1+t)^{-2k-2} \mathcal{E}_k(t) + (\epsilon_0 + \delta^{-1})(1+t)^{-2k-2} \sum_{\iota=0}^{k-1} \mathcal{E}_\iota(t), \end{aligned} \quad (3.43)$$

for any positive number  $\delta > 0$  which will be specified later, where  $\bar{\mathfrak{E}}_k := \int r^2 \bar{\rho}_0^\gamma \tilde{\mathfrak{E}}_k dr + M_2$ . Here  $\tilde{\mathfrak{E}}_k$  and  $M_2$  are defined by (3.49) and (3.47), respectively. Moreover, we show that  $\bar{\mathfrak{E}}_k$  satisfies the following estimates:

$$\bar{\mathfrak{E}}_k \geq C^{-1}(1+t)^{-1} \int r^2 \bar{\rho}_0^\gamma |\partial_t^k(\zeta, r\zeta_r)|^2 dr - C(1+t)^{-2k-1} \sum_{\iota=0}^{k-1} \mathcal{E}_\iota(t), \quad (3.44)$$

$$\bar{\mathfrak{E}}_k \lesssim (1+t)^{-1} \int r^2 \bar{\rho}_0^\gamma |\partial_t^k(\zeta, r\zeta_r)|^2 dr + (1+t)^{-2k-1} \sum_{\iota=0}^{k-1} \mathcal{E}_\iota(t). \quad (3.45)$$

We start with integrating the production of (3.38) and  $r^3 \partial_t^k \zeta_t$  with respect to the spatial variable which gives

$$\frac{d}{dt} \int \frac{1}{2} r^4 \bar{\rho}_0 (\partial_t^k \zeta_t)^2 dr + \int r^4 \bar{\rho}_0 (\partial_t^k \zeta_t)^2 dr + N_1 + N_2 = 0, \quad (3.46)$$

where

$$\begin{aligned} N_1 := & - \int \bar{\rho}_0^\gamma (w_1 \partial_t^k \zeta + w_2 r \partial_t^k \zeta_r) (r^3 \partial_t^k \zeta_t)_r dr + \int r^2 \bar{\rho}_0^\gamma (3w_2 - w_1) (r \partial_t^k \zeta_r) \partial_t^k \zeta_t dr \\ & - 2 \int r^2 \bar{\rho}_0^\gamma w_3 (r \zeta_r) (\partial_t^k \zeta) \partial_t^k \zeta_t dr, \end{aligned}$$

$$\begin{aligned} N_2 := & - \int \bar{\rho}_0^\gamma \{ K_1 + \partial_t^{k-1} [\tilde{\eta}_{rt} (w_1 - (2-3\gamma)\tilde{\eta}_r^{1-3\gamma})] \} (r^3 \partial_t^k \zeta_t)_r dr \\ & + \int r^2 \bar{\rho}_0^\gamma (\partial_t^k \zeta_t) [r(K_2 - 2K_3) - 2\partial_t^{k-1} (\tilde{\eta}_{rt} w_3 r \zeta_r)] dr. \end{aligned}$$

Note that  $N_1$  and  $N_2$  can be rewritten as

$$\begin{aligned}
N_1 &= -\frac{1}{2} \int r^2 \bar{\rho}_0^\gamma \left[ (3w_1 + 2w_3 r \zeta_r) \left[ (\partial_t^k \zeta)^2 \right]_t + 2w_1 \left[ (\partial_t^k \zeta) r \partial_t^k \zeta_r \right]_t + w_2 \left[ (r \partial_t^k \zeta_r)^2 \right]_t \right] dr \\
&= -\frac{1}{2} \frac{d}{dt} \int r^2 \bar{\rho}_0^\gamma \left[ (3w_1 + 2w_3 r \zeta_r) (\partial_t^k \zeta)^2 + 2w_1 (\partial_t^k \zeta) r \partial_t^k \zeta_r + w_2 (r \partial_t^k \zeta_r)^2 \right] dr + \tilde{N}_1, \\
N_2 &= \frac{d}{dt} M_2 + \int \bar{\rho}_0^\gamma \left\{ K_{1t} + \partial_t^k \left[ \tilde{\eta}_{rt} (w_1 - (2 - 3\gamma) \tilde{\eta}_r^{1-3\gamma}) \right] \right\} (r^3 \partial_t^k \zeta)_r dr \\
&\quad - \int r^2 \bar{\rho}_0^\gamma (\partial_t^k \zeta) \left[ r(K_2 - 2K_3)_t - 2\partial_t^k (\tilde{\eta}_{rt} w_3 r \zeta_r) \right] dr =: \frac{d}{dt} M_2 + \tilde{N}_2,
\end{aligned}$$

where

$$\begin{aligned}
\tilde{N}_1 &:= \frac{1}{2} \int r^2 \bar{\rho}_0^\gamma \left[ (3w_1 + 2w_3 r \zeta_r)_t (\partial_t^k \zeta)^2 + 2w_{1t} (\partial_t^k \zeta) r \partial_t^k \zeta_r + w_{2t} (r \partial_t^k \zeta_r)^2 \right] dr, \\
M_2 &:= - \int \bar{\rho}_0^\gamma \left\{ K_1 + \partial_t^{k-1} \left[ \tilde{\eta}_{rt} (w_1 - (2 - 3\gamma) \tilde{\eta}_r^{1-3\gamma}) \right] \right\} (r^3 \partial_t^k \zeta)_r dr \\
&\quad + \int r^2 \bar{\rho}_0^\gamma (\partial_t^k \zeta) \left[ r(K_2 - 2K_3) - 2\partial_t^{k-1} (\tilde{\eta}_{rt} w_3 r \zeta_r) \right] dr.
\end{aligned} \tag{3.47}$$

It then follows from equation (3.46) that

$$\frac{d}{dt} \left[ \int \left( \frac{1}{2} r^4 \bar{\rho}_0 (\partial_t^k \zeta_t)^2 + r^2 \bar{\rho}_0^\gamma \tilde{\mathfrak{E}}_k \right) dr + M_2 \right] + \int r^4 \bar{\rho}_0 (\partial_t^k \zeta_t)^2 dr = -\tilde{N}_1 - \tilde{N}_2, \tag{3.48}$$

where

$$\tilde{\mathfrak{E}}_k := -\frac{1}{2} \left[ (3w_1 + 2w_3 r \zeta_r) (\partial_t^k \zeta)^2 + 2w_1 (\partial_t^k \zeta) r \partial_t^k \zeta_r + w_2 (r \partial_t^k \zeta_r)^2 \right] \tag{3.49}$$

which satisfies

$$\begin{aligned}
\tilde{\mathfrak{E}}_k &= \tilde{\eta}_r^{1-3\gamma} \left[ \frac{3}{2} (3\gamma - 2) (\partial_t^k \zeta)^2 + (3\gamma - 2) (\partial_t^k \zeta) r \partial_t^k \zeta_r + \frac{\gamma}{2} (r \partial_t^k \zeta_r)^2 \right] \\
&\quad + O(1) \tilde{\eta}_r^{-3\gamma} (|\zeta| + |r \zeta_r|) \left( (\partial_t^k \zeta)^2 + (r \partial_t^k \zeta_r)^2 \right) \\
&\sim \tilde{\eta}_r^{1-3\gamma} \left[ (\partial_t^k \zeta)^2 + (r \partial_t^k \zeta_r)^2 \right] \sim (1+t)^{-1} \left[ (\partial_t^k \zeta)^2 + (r \partial_t^k \zeta_r)^2 \right].
\end{aligned} \tag{3.50}$$

Here we have used (3.39), (2.13) and the smallness of  $\zeta$  and  $r \zeta_r$  which is a consequence of (3.1) to derive the above equivalence. We will show later that  $M_2$  can be bounded by the integral of  $\tilde{\mathfrak{E}}_k$  and lower-order terms, see (3.65).

In what follows, we analyze the terms on the right-hand side of (3.48). Clearly,  $-\tilde{N}_1$  can be bounded by

$$\begin{aligned}
-\tilde{N}_1 &\leq (1 - 3\gamma) \int r^2 \bar{\rho}_0^\gamma \tilde{\eta}_r^{-3\gamma} \tilde{\eta}_{rt} \left[ \left( \frac{9}{2} \gamma - 3 \right) (\partial_t^k \zeta)^2 + (3\gamma - 2) (\partial_t^k \zeta) (r \partial_t^k \zeta_r) \right. \\
&\quad \left. + \frac{\gamma}{2} (r \partial_t^k \zeta_r)^2 \right] dr + C \int r^2 \bar{\rho}_0^\gamma \tilde{\eta}_r^{-3\gamma} \left[ \tilde{\eta}_r^{-1} \tilde{\eta}_{rt} (|\zeta| + |r \zeta_r|) \right. \\
&\quad \left. + (1 + \tilde{\eta}_r^{-1} (|\zeta| + |r \zeta_r|)) (|\zeta_t| + |r \zeta_{rt}|) \right] \left( (\partial_t^k \zeta)^2 + (r \partial_t^k \zeta_r)^2 \right) dr.
\end{aligned}$$

It should be noted that the first integral on the right-hand side of the inequality above is non-positive due to  $\tilde{\eta}_{rt} \geq 0$ . Thus, we have by use of (2.13) and (3.1) that

$$-\tilde{N}_1 \lesssim \epsilon_0(1+t)^{-2-\frac{1}{3\gamma-1}} \int r^2 \bar{\rho}_0^\gamma \left( (\partial_t^k \zeta)^2 + (r\partial_t^k \zeta_r)^2 \right) dr. \quad (3.51)$$

To control  $\tilde{N}_2$ , we may rewrite it as

$$\begin{aligned} \tilde{N}_2 &= \int r^2 \bar{\rho}_0^\gamma \left\{ (3\partial_t^k \zeta + r\partial_t^k \zeta_r) \partial_t^k [\tilde{\eta}_{rt} (w_1 - (2-3\gamma)\tilde{\eta}_r^{1-3\gamma})] + 2(\partial_t^k \zeta) \partial_t^k (\tilde{\eta}_{rt} w_3 r \zeta_r) \right\} dr \\ &\quad + \int r^2 \bar{\rho}_0^\gamma [K_{1t} (3\partial_t^k \zeta + r\partial_t^k \zeta_r) - r(K_2 - 2K_3)_t (\partial_t^k \zeta)] dr =: \tilde{N}_{21} + \tilde{N}_{22}. \end{aligned}$$

For  $\tilde{N}_{21}$ , note that

$$\begin{aligned} &\partial_t^k [\tilde{\eta}_{rt} (w_1 - (2-3\gamma)\tilde{\eta}_r^{1-3\gamma})] \\ &= (3\gamma-1) \partial_t^k [\tilde{\eta}_{rt} \tilde{\eta}_r^{-3\gamma} ((3\gamma-2)\zeta + \gamma r \zeta_r)] + \partial_t^k (\tilde{\eta}_{rt} \bar{w}_1) \\ &= (3\gamma-1) \tilde{\eta}_{rt} \tilde{\eta}_r^{-3\gamma} ((3\gamma-2) \partial_t^k \zeta + \gamma r \partial_t^k \zeta_r) \\ &\quad + O(1) \sum_{\iota=1}^k |\partial_t^\iota (\tilde{\eta}_{rt} \tilde{\eta}_r^{-3\gamma})| |\partial_t^{k-\iota} (\zeta, r\zeta_r)| + \partial_t^k (\tilde{\eta}_{rt} \bar{w}_1). \end{aligned}$$

and

$$\begin{aligned} \partial_t^k (\tilde{\eta}_{rt} w_3 r \zeta_r) &= (1-3\gamma) \partial_t^k (\tilde{\eta}_{rt} \tilde{\eta}_r^{-3\gamma} r \zeta_r) + \partial_t^k (\tilde{\eta}_{rt} \bar{w}_3 r \zeta_r) \\ &= (1-3\gamma) \tilde{\eta}_{rt} \tilde{\eta}_r^{-3\gamma} (r \partial_t^k \zeta_r) + O(1) \sum_{\iota=1}^k |\partial_t^\iota (\tilde{\eta}_{rt} \tilde{\eta}_r^{-3\gamma})| |r \partial_t^{k-\iota} \zeta_r| + \partial_t^k (\tilde{\eta}_{rt} \bar{w}_3 r \zeta_r). \end{aligned}$$

Thus,

$$\begin{aligned} -\tilde{N}_{21} &\leq (1-3\gamma) \int r^2 \bar{\rho}_0^\gamma \tilde{\eta}_r^{-3\gamma} \tilde{\eta}_{rt} \left[ (9\gamma-6) (\partial_t^k \zeta)^2 + (6\gamma-4) (\partial_t^k \zeta) (r \partial_t^k \zeta_r) \right. \\ &\quad \left. + \gamma (r \partial_t^k \zeta_r)^2 \right] dr + C \int r^2 \bar{\rho}_0^\gamma (|\partial_t^k \zeta| + |r \partial_t^k \zeta_r|) \\ &\quad \times \left[ \sum_{\iota=1}^k |\partial_t^\iota (\tilde{\eta}_{rt} \tilde{\eta}_r^{-3\gamma})| |\partial_t^{k-\iota} (\zeta, r\zeta_r)| + |\partial_t^k (\tilde{\eta}_{rt} \bar{w}_1, \tilde{\eta}_{rt} \bar{w}_3 r \zeta_r)| \right] dr. \end{aligned} \quad (3.52)$$

For  $\tilde{N}_{22}$ , note that

$$\begin{aligned} K_{1t} &= (k-1) (w_{1t} \partial_t^k \zeta + w_{2t} r \partial_t^k \zeta_r) + O(1) \sum_{\iota=2}^k |\partial_t^\iota (w_1, w_2)| |\partial_t^{k+1-\iota} (\zeta, r\zeta_r)|, \\ rK_{2t} &= (k-1) (3w_2 - w_1)_t (r \partial_t^k \zeta_r) + O(1) \sum_{\iota=2}^k |\partial_t^\iota (w_1, w_2)| |\partial_t^{k+1-\iota} (r\zeta_r)|, \\ rK_{3t} &= (k-1) (w_3 r \zeta_r)_t (\partial_t^k \zeta) + O(1) \sum_{\iota=2}^k |\partial_t^\iota (w_3 r \zeta_r)| |\partial_t^{k+1-\iota} \zeta|. \end{aligned}$$

Thus,

$$\begin{aligned} \tilde{N}_{22} &\geq 2(k-1)\tilde{N}_1 - C \int r^2 \bar{\rho}_0^\gamma (|\partial_t^k \zeta| + |r\partial_t^k \zeta_r|) \\ &\quad \times \sum_{\iota=2}^k |\partial_t^\iota (w_1, w_2, w_3 r \zeta_r)| |\partial_t^{k+1-\iota} (\zeta, r\zeta_r)| dr. \end{aligned} \quad (3.53)$$

In a similar way to dealing with  $\tilde{N}_1$  shown in (3.51), we have, with the aid of (3.52) and (3.53), that

$$\begin{aligned} -\tilde{N}_2 &\lesssim \epsilon_0 (1+t)^{-2-\frac{1}{3\gamma-1}} \int r^2 \bar{\rho}_0^\gamma \left( (\partial_t^k \zeta)^2 + (r\partial_t^k \zeta_r)^2 \right) dr \\ &\quad + \int r^2 \bar{\rho}_0^\gamma (|\partial_t^k \zeta| + |r\partial_t^k \zeta_r|) Q dr. \end{aligned} \quad (3.54)$$

where

$$\begin{aligned} Q &= \sum_{\iota=1}^k |\partial_t^\iota (\tilde{\eta}_{rt} \tilde{\eta}_r^{-3\gamma})| |\partial_t^{k-\iota} (\zeta, r\zeta_r)| + |\partial_t^k (\tilde{\eta}_{rt} \bar{w}_1, \tilde{\eta}_{rt} \bar{w}_3 r \zeta_r)| \\ &\quad + \sum_{\iota=2}^k |\partial_t^\iota (w_1, w_2, w_3 r \zeta_r)| |\partial_t^{k+1-\iota} (\zeta, r\zeta_r)|. \end{aligned} \quad (3.55)$$

Therefore, it produces from (3.48), (3.51) and (3.54) that

$$\begin{aligned} &\frac{d}{dt} \left[ \int \left( \frac{1}{2} r^4 \bar{\rho}_0 (\partial_t^k \zeta_t)^2 + r^2 \bar{\rho}_0^\gamma \tilde{\mathfrak{E}}_k \right) dr + M_2 \right] + \int r^4 \bar{\rho}_0 (\partial_t^k \zeta_t)^2 dr \\ &\lesssim \epsilon_0 (1+t)^{-2-\frac{1}{3\gamma-1}} \int r^2 \bar{\rho}_0^\gamma |\partial_t^k (\zeta, r\zeta_r)|^2 dr + \int r^2 \bar{\rho}_0^\gamma |\partial_t^k (\zeta, r\zeta_r)| Q dr. \end{aligned} \quad (3.56)$$

We are to bound the last term on the right-hand side of (3.56). It follows from (2.13) and (3.1) that

$$Q \lesssim \epsilon_0 (1+t)^{-2-\frac{1}{3\gamma-1}} |\partial_t^k (\zeta, r\zeta_r)| + \tilde{Q}, \quad (3.57)$$

where

$$\begin{aligned} \tilde{Q} &:= \sum_{\iota=1}^k (1+t)^{-2-\iota} |\partial_t^{k-\iota} (\zeta, r\zeta_r)| + (1+t)^{-1-\frac{1}{3\gamma-1}} |\partial_t^2 (\zeta, r\zeta_r)| |\partial_t^{k-1} (\zeta, r\zeta_r)| \\ &\quad + (1+t)^{-2-\frac{1}{3\gamma-1}} |\partial_t^2 (\zeta, r\zeta_r)| |\partial_t^{k-2} (\zeta, r\zeta_r)| + (1+t)^{-1-\frac{1}{3\gamma-1}} |\partial_t^3 (\zeta, r\zeta_r)| |\partial_t^{k-2} (\zeta, r\zeta_r)| \\ &\quad + \left[ (1+t)^{-1-\frac{1}{3\gamma-1}} |\partial_t^4 (\zeta, r\zeta_r)| + (1+t)^{-2-\frac{1}{3\gamma-1}} |\partial_t^3 (\zeta, r\zeta_r)| + (1+t)^{-3-\frac{1}{3\gamma-1}} \right. \\ &\quad \left. \times |\partial_t^2 (\zeta, r\zeta_r)| + (1+t)^{-1-\frac{2}{3\gamma-1}} |\partial_t^2 (\zeta, r\zeta_r)|^2 \right] |\partial_t^{k-3} (\zeta, r\zeta_r)| + \text{l.o.t.} \end{aligned} \quad (3.58)$$

Here and thereafter the notation l.o.t. is used to represent the lower-order terms involving  $\partial_t^\iota (\zeta, r\zeta_r)$  with  $\iota = 2, \dots, k-4$ . It should be noticed that the second term on the right-hand

side of (3.58) only appears as  $k - 1 \geq 2$ , the third term as  $k - 2 \geq 2$ , the fourth term as  $k - 2 \geq 3$ , and so on. Clearly, we use (3.1) again to obtain

$$\begin{aligned} \tilde{Q} &\lesssim \sum_{\iota=1}^k (1+t)^{-2-\iota} |\partial_t^{k-\iota}(\zeta, r\zeta_r)| + \epsilon_0 \sigma^{-\frac{1}{2}} (1+t)^{-3-\frac{1}{3\gamma-1}} |\partial_t^{k-1}(\zeta, r\zeta_r)| \\ &\quad + \epsilon_0 \sigma^{-1} (1+t)^{-4-\frac{1}{3\gamma-1}} |\partial_t^{k-2}(\zeta, r\zeta_r)| + \epsilon_0 \sigma^{-\frac{3}{2}} (1+t)^{-5-\frac{1}{3\gamma-1}} |\partial_t^{k-3}(\zeta, r\zeta_r)| + \text{l.o.t.}, \end{aligned}$$

if  $k \geq 7$ . Similarly, we can bound l.o.t. and achieve

$$\tilde{Q} \lesssim \sum_{\iota=1}^k (1+t)^{-2-\iota} |\partial_t^{k-\iota}(\zeta, r\zeta_r)| + \epsilon_0 \sum_{\iota=1}^{[(k-1)/2]} \sigma^{-\frac{\iota}{2}} (1+t)^{-2-\iota-\frac{1}{3\gamma-1}} |\partial_t^{k-\iota}(\zeta, r\zeta_r)|,$$

which implies

$$\begin{aligned} \int r^2 \bar{\rho}_0^\gamma |\partial_t^k(\zeta, r\zeta_r)| \tilde{Q} dr &\lesssim \sum_{\iota=1}^k (1+t)^{-2-\iota} \int r^2 \bar{\rho}_0^\gamma |\partial_t^k(\zeta, r\zeta_r)| |\partial_t^{k-\iota}(\zeta, r\zeta_r)| dr \\ &\quad + \epsilon_0 \sum_{\iota=1}^{[(k-1)/2]} (1+t)^{-2-\iota} \int r^2 \bar{\rho}_0^\gamma \sigma^{-\frac{\iota}{2}} |\partial_t^k(\zeta, r\zeta_r)| |\partial_t^{k-\iota}(\zeta, r\zeta_r)| dr =: \tilde{Q}_1 + \tilde{Q}_2. \end{aligned} \quad (3.59)$$

Easily, it follows from the Cauchy-Schwarz inequality that for any  $\delta > 0$ ,

$$\begin{aligned} \tilde{Q}_1 &\lesssim \delta (1+t)^{-2} \int r^2 \bar{\rho}_0^\gamma |\partial_t^k(\zeta, r\zeta_r)|^2 dr + \delta^{-1} \sum_{\iota=1}^k (1+t)^{-2-2\iota} \int r^2 \bar{\rho}_0^\gamma |\partial_t^{k-\iota}(\zeta, r\zeta_r)|^2 dr \\ &\lesssim \delta (1+t)^{-2-2k} \mathcal{E}_k(t) + \delta^{-1} (1+t)^{-2-2k} \sum_{\iota=0}^{k-1} \mathcal{E}_\iota(t) \end{aligned} \quad (3.60)$$

and

$$\begin{aligned} \tilde{Q}_2 &\lesssim \epsilon_0 (1+t)^{-2} \int r^2 \bar{\rho}_0^\gamma |\partial_t^k(\zeta, r\zeta_r)|^2 dr \\ &\quad + \epsilon_0 \sum_{\iota=1}^{[(k-1)/2]} (1+t)^{-2-2\iota} \int r^2 \bar{\rho}_0^\gamma \sigma^{-\iota} |\partial_t^{k-\iota}(\zeta, r\zeta_r)|^2 dr. \end{aligned} \quad (3.61)$$

In view of the Hardy inequality (3.6), we see that for  $\iota = 1, \dots, [(k-1)/2]$ ,

$$\begin{aligned} \int_{\mathcal{I}_b} \sigma^{\alpha+1-\iota} |\partial_t^{k-\iota}(\zeta, \zeta_r)|^2 dr &\lesssim \int_{\mathcal{I}_b} \sigma^{\alpha+3-\iota} |\partial_t^{k-\iota}(\zeta, \zeta_r, \zeta_{rr})|^2 dr \lesssim \dots \\ &\lesssim \sum_{i=0}^{\iota+1} \int_{\mathcal{I}_b} \sigma^{\alpha+1+i} |\partial_t^{k-\iota} \partial_r^i \zeta|^2 dr \lesssim \sum_{i=0}^{\iota+1} \int_{\mathcal{I}_b} r^4 \sigma^{\alpha+1+i} |\partial_t^{k-\iota} \partial_r^i \zeta|^2 dr \\ &\lesssim (1+t)^{2\iota-2k} \left( \mathcal{E}_{k-\iota} + \sum_{i=1}^{\iota} \mathcal{E}_{k-\iota, i} \right) (t) \lesssim (1+t)^{2\iota-2k} \sum_{\iota=0}^k \mathcal{E}_\iota(t), \end{aligned}$$

due to  $\alpha + 1 - \iota \geq \alpha - [([\alpha] + 1)/2] \geq 0$  for  $k \leq l$ , which ensures the application of the Hardy inequality. Here the last inequality follows from the elliptic estimate (3.7). Thus, we can obtain for  $\iota = 1, \dots, [(k-1)/2]$ ,

$$\begin{aligned}
& \int r^2 \bar{\rho}_0^\gamma \sigma^{-\iota} |\partial_t^{k-\iota}(\zeta, r\zeta_r)|^2 dr = \int r^2 \sigma^{\alpha+1-\iota} |\partial_t^{k-\iota}(\zeta, r\zeta_r)|^2 dr \\
& = \int_{\mathcal{I}_o} r^2 \sigma^{\alpha+1-\iota} |\partial_t^{k-\iota}(\zeta, r\zeta_r)|^2 dr + \int_{\mathcal{I}_b} r^2 \sigma^{\alpha+1-\iota} |\partial_t^{k-\iota}(\zeta, r\zeta_r)|^2 dr \\
& \lesssim \int_{\mathcal{I}_o} r^2 \sigma^{\alpha+1} |\partial_t^{k-\iota}(\zeta, r\zeta_r)|^2 dr + \int_{\mathcal{I}_b} \sigma^{\alpha+1-\iota} |\partial_t^{k-\iota}(\zeta, \zeta_r)|^2 dr \\
& \lesssim (1+t)^{2\iota-2k} \sum_{\iota=0}^k \mathcal{E}_\iota(t). \tag{3.62}
\end{aligned}$$

This, together with (3.61), implies

$$\tilde{Q}_2 \lesssim \epsilon_0 (1+t)^{-2-2k} \sum_{\iota=0}^k \mathcal{E}_\iota(t). \tag{3.63}$$

So, it yields from (3.57), (3.59), (3.60) and (3.63) that for  $\delta > 0$ ,

$$\int r^2 \bar{\rho}_0^\gamma |\partial_t^k(\zeta, r\zeta_r)| Q dr \lesssim (1+t)^{-2k-2} \left[ (\epsilon_0 + \delta) \mathcal{E}_k(t) + (\epsilon_0 + \delta^{-1}) \sum_{\iota=0}^{k-1} \mathcal{E}_\iota(t) \right].$$

Substitute this into (3.56) to give (3.43).

To prove (3.44) and (3.45), we adopt a similar but much easier way to dealing with  $\tilde{N}_2$  as shown in (3.54) to show

$$|M_2| \lesssim \int r^2 \bar{\rho}_0^\gamma (|\partial_t^k \zeta| + |r \partial_t^k \zeta_r|) P dr, \tag{3.64}$$

where

$$\begin{aligned}
P & = \sum_{\iota=0}^{k-1} |\partial_t^\iota (\tilde{\eta}_{rt} \tilde{\eta}_r^{-3\gamma})| |\partial_t^{k-1-\iota}(\zeta, r\zeta_r)| + |\partial_t^{k-1}(\tilde{\eta}_{rt} \bar{w}_1, \tilde{\eta}_{rt} \bar{w}_3 r \zeta_r)| \\
& \quad + \sum_{\iota=1}^{k-1} |\partial_t^\iota(w_1, w_2, w_3 r \zeta_r)| |\partial_t^{k-\iota}(\zeta, r\zeta_r)|.
\end{aligned}$$

In view of (2.13) and (3.1), we have

$$\begin{aligned}
P & \lesssim \sum_{\iota=0}^{k-1} (1+t)^{-2-\iota} |\partial_t^{k-1-\iota}(\zeta, r\zeta_r)| + |\partial_t^{k-2}(\zeta, r\zeta_r)| (1+t)^{-1-\frac{1}{3\gamma-1}} |\partial_t^2(\zeta, r\zeta_r)| \\
& \quad + |\partial_t^{k-3}(\zeta, r\zeta_r)| \left[ (1+t)^{-1-\frac{1}{3\gamma-1}} |\partial_t^3(\zeta, r\zeta_r)| + (1+t)^{-2-\frac{1}{3\gamma-1}} |\partial_t^2(\zeta, r\zeta_r)| \right] + \text{l.o.t.},
\end{aligned}$$

which implies

$$P \lesssim \sum_{\iota=0}^{k-1} (1+t)^{-2-\iota} |\partial_t^{k-1-\iota}(\zeta, r\zeta_r)| + \epsilon_0 \sum_{\iota=2}^{[k/2]} \sigma^{\frac{1-\iota}{2}} (1+t)^{-1-\iota-\frac{1}{3\gamma-1}} |\partial_t^{k-\iota}(\zeta, r\zeta_r)|.$$

Similar to the derivation of (3.62), we can use the Hardy inequality (3.6) and elliptic estimate (3.7) to obtain

$$\begin{aligned} \int r^2 \bar{\rho}_0^\gamma P^2 dr &\lesssim (1+t)^{-2-2k} \sum_{\iota=0}^{k-1} \mathcal{E}_\iota(t) + \epsilon_0^2 \sum_{\iota=2}^{[k/2]} (1+t)^{-2-2\iota} \int r^2 \bar{\rho}_0^\gamma \sigma^{1-\iota} |\partial_t^{k-\iota}(\zeta, r\zeta_r)|^2 dr \\ &\lesssim (1+t)^{-2-2k} \sum_{\iota=0}^{k-1} \mathcal{E}_\iota(t). \end{aligned}$$

It then gives from the Cauchy inequality and (3.64) that for any  $\delta > 0$ ,

$$\begin{aligned} |M_2| &\lesssim \delta (1+t)^{-1} \int r^2 \bar{\rho}_0^\gamma |\partial_t^k(\zeta, r\zeta_r)|^2 dr + \delta^{-1} (1+t) \int r^2 \bar{\rho}_0^\gamma P^2 dr \\ &\lesssim \delta (1+t)^{-1} \int r^2 \bar{\rho}_0^\gamma |\partial_t^k(\zeta, r\zeta_r)|^2 dr + \delta^{-1} (1+t)^{-1-2k} \sum_{\iota=0}^{k-1} \mathcal{E}_\iota(t). \end{aligned} \quad (3.65)$$

This, together with (3.50), proves (3.44) (by choosing suitably small  $\delta$ ) and (3.45).

**Step 2.** To control the first term on the right-hand side of (3.43), we will prove that

$$\begin{aligned} &\frac{d}{dt} \mathfrak{E}_k + \int \left[ (1+t)^{-1} r^2 \bar{\rho}_0^\gamma |\partial_t^k(\zeta, r\zeta_r)|^2 + r^4 \bar{\rho}_0 (\partial_t^k \zeta_t)^2 \right] dr \\ &\lesssim (1+t)^{-1-2k} \sum_{\iota=0}^{k-1} (1+t)^{2\iota} \int \left[ r^2 \bar{\rho}_0^\gamma |\partial_t^\iota(\zeta, r\zeta_r)|^2 + (1+t) r^4 \bar{\rho}_0 (\partial_t^\iota \zeta_t)^2 \right] dr, \end{aligned} \quad (3.66)$$

where

$$\mathfrak{E}_k := \int r^4 \bar{\rho}_0 \left[ (\partial_t^k \zeta_t)^2 + (\partial_t^k \zeta) \partial_t^k \zeta_t + \frac{1}{2} (\partial_t^k \zeta)^2 \right] dr + 2\bar{\mathfrak{E}}_k.$$

We start with integrating the product of (3.38) and  $r^3 \partial_t^k \zeta$  with respect to  $r$  to give

$$\frac{d}{dt} \int r^4 \bar{\rho}_0 \left( (\partial_t^k \zeta) \partial_t^k \zeta_t + \frac{1}{2} (\partial_t^k \zeta)^2 \right) dr - \int r^4 \bar{\rho}_0 (\partial_t^k \zeta_t)^2 dr + M_1 + M_2 = 0, \quad (3.67)$$

where

$$\begin{aligned} M_1 &= - \int \bar{\rho}_0^\gamma (w_1 \partial_t^k \zeta + w_2 r \partial_t^k \zeta_r) (r^3 \partial_t^k \zeta)_r dr + \int r^2 \bar{\rho}_0^\gamma (3w_2 - w_1) (r \partial_t^k \zeta_r) \partial_t^k \zeta dr \\ &\quad - 2 \int r^2 \bar{\rho}_0^\gamma w_3 (r \zeta_r) (\partial_t^k \zeta)^2 dr \end{aligned}$$

and  $M_2$  is defined in (3.47). A direct calculation shows that  $M_1$  is positive and can be bounded from below as follows

$$\begin{aligned}
M_1 &= - \int r^2 \bar{\rho}_0^\gamma \left[ (3w_1 + 2w_3 r \zeta_r) (\partial_t^k \zeta)^2 + 2w_1 (\partial_t^k \zeta) (r \partial_t^k \zeta_r) + w_2 (r \partial_t^k \zeta_r)^2 \right] dr \\
&\geq \int r^2 \bar{\rho}_0^\gamma \tilde{\eta}_r^{1-3\gamma} \left\{ (9\gamma - 6) (\partial_t^k \zeta)^2 + (6\gamma - 4) (\partial_t^k \zeta) (r \partial_t^k \zeta_r) + \gamma (r \partial_t^k \zeta_r)^2 \right. \\
&\quad \left. - C (|\zeta| + |r \zeta_r|) \left[ (\partial_t^k \zeta)^2 + (r \partial_t^k \zeta_r)^2 \right] \right\} dr \\
&\gtrsim \int r^2 \bar{\rho}_0^\gamma \tilde{\eta}_r^{1-3\gamma} \left[ (\partial_t^k \zeta)^2 + (r \partial_t^k \zeta_r)^2 \right] dr \gtrsim (1+t)^{-1} \int r^2 \bar{\rho}_0^\gamma \left[ (\partial_t^k \zeta)^2 + (r \partial_t^k \zeta_r)^2 \right] dr,
\end{aligned}$$

due to (3.39), the smallness of  $\zeta_r$  and  $r \zeta_r$  and (2.13). We then obtain, by making a summation of  $2 \times$  (3.43) and (3.67), that

$$\begin{aligned}
&\frac{d}{dt} \mathfrak{E}_k + \int r^4 \bar{\rho}_0 (\partial_t^k \zeta_t)^2 dr + (1+t)^{-1} \int r^2 \bar{\rho}_0^\gamma \left[ (\partial_t^k \zeta)^2 + (r \partial_t^k \zeta_r)^2 \right] dr \\
&\lesssim \delta (1+t)^{-1} \int r^2 \bar{\rho}_0^\gamma |\partial_t^k (\zeta, r \zeta_r)|^2 dr + (\epsilon_0 + \delta) (1+t)^{-2k-2} \mathcal{E}_k(t) \\
&\quad + \delta^{-1} (1+t)^{-1-2k} \sum_{\iota=0}^{k-1} \mathcal{E}_\iota(t) + (\epsilon_0 + \delta^{-1}) (1+t)^{-2k-2} \sum_{\iota=0}^{k-1} \mathcal{E}_\iota(t),
\end{aligned} \tag{3.68}$$

because of (3.65). Notice from the Hardy inequality (3.6) that for  $j = 0, 1, \dots, l$ ,

$$\begin{aligned}
\int r^4 \bar{\rho}_0 (\partial_t^j \zeta)^2 dr &= \int_{\mathcal{I}_o} r^4 \sigma^\alpha (\partial_t^j \zeta)^2 dr + \int_{\mathcal{I}_b} r^4 \sigma^\alpha (\partial_t^j \zeta)^2 dr \lesssim \int_{\mathcal{I}_o} r^2 \sigma^{\alpha+1} (\partial_t^j \zeta)^2 dr \\
&\quad + \int_{\mathcal{I}_b} \sigma^\alpha (\partial_t^j \zeta)^2 dr \lesssim \int_{\mathcal{I}_o} r^2 \sigma^{\alpha+1} (\partial_t^j \zeta)^2 dr + \int_{\mathcal{I}_b} \sigma^{\alpha+2} \left[ (\partial_t^j \zeta)^2 + (\partial_t^j \zeta_r)^2 \right] dr \\
&\lesssim \int r^2 \sigma^{\alpha+1} (\partial_t^j \zeta)^2 dr + \int_{\mathcal{I}_b} \sigma^{\alpha+1} \left[ r^2 (\partial_t^j \zeta)^2 + r^4 (\partial_t^j \zeta_r)^2 \right] dr \lesssim \int r^2 \bar{\rho}_0^\gamma |\partial_t^j (\zeta, r \zeta_r)|^2 dr.
\end{aligned}$$

Thus,

$$\mathcal{E}_j(t) \lesssim (1+t)^{2j} \int \left[ r^2 \bar{\rho}_0^\gamma |\partial_t^j (\zeta, r \zeta_r)|^2 + (1+t) r^4 \bar{\rho}_0 (\partial_t^j \zeta_t)^2 \right] dr, \quad j = 0, \dots, l. \tag{3.69}$$

This finishes the proof of (3.66), by using (3.68) and (3.69), choosing suitably small  $\delta$  and noting the smallness of  $\epsilon_0$ . Moreover, it follows from (3.44) and (3.45) that

$$\begin{aligned}
\mathfrak{E}_k &\geq C^{-1} \int r^4 \bar{\rho}_0 |\partial_t^k (\zeta, \zeta_t)|^2 dr + C^{-1} (1+t)^{-1} \int r^2 \bar{\rho}_0^\gamma |\partial_t^k (\zeta, r \zeta_r)|^2 dr \\
&\quad - C (1+t)^{-2k-1} \sum_{\iota=0}^{k-1} \mathcal{E}_\iota(t),
\end{aligned} \tag{3.70}$$

$$\begin{aligned}
\mathfrak{E}_k &\lesssim \int r^4 \bar{\rho}_0 |\partial_t^k (\zeta, \zeta_t)|^2 dr + (1+t)^{-1} \int r^2 \bar{\rho}_0^\gamma |\partial_t^k (\zeta, r \zeta_r)|^2 dr \\
&\quad + (1+t)^{-2k-1} \sum_{\iota=0}^{k-1} \mathcal{E}_\iota(t).
\end{aligned} \tag{3.71}$$



**Step 3.** In this step, we show the time decay of the norm. We integrate (3.66) and use the induction hypothesis (3.42) to show, noting (3.70) and (3.71), that

$$\begin{aligned}
& \int \left[ r^4 \bar{\rho}_0 |\partial_t^k (\zeta, \zeta_t)|^2 + (1+t)^{-1} r^2 \bar{\rho}_0^\gamma |\partial_t^k (\zeta, r\zeta_r)|^2 \right] (r, t) dr \\
& + \int_0^t \int \left[ (1+s)^{-1} r^2 \bar{\rho}_0^\gamma |\partial_s^k (\zeta, r\zeta_r)|^2 + r^4 \bar{\rho}_0 (\partial_s^k \zeta_s)^2 \right] dr ds \\
& \lesssim \sum_{\iota=0}^k \mathcal{E}_\iota(0) + \sum_{\iota=0}^{k-1} \int_0^t (1+s)^{2\iota-1-2k} \int \left[ r^2 \bar{\rho}_0^\gamma |\partial_s^\iota (\zeta, r\zeta_r)|^2 + (1+s) r^4 \bar{\rho}_0 (\partial_s^\iota \zeta_s)^2 \right] dr \\
& \lesssim \sum_{\iota=0}^k \mathcal{E}_\iota(0).
\end{aligned}$$

Multiply (3.66) by  $(1+t)^p$  and integrate the product with respect to the temporal variable from  $p=1$  to  $p=2k$  step by step to get

$$\begin{aligned}
& (1+t)^{2k} \int \left[ r^4 \bar{\rho}_0 |\partial_t^k (\zeta, \zeta_t)|^2 + (1+t)^{-1} r^2 \bar{\rho}_0^\gamma |\partial_t^k (\zeta, r\zeta_r)|^2 \right] (r, t) dr \\
& + \int_0^t (1+s)^{2k} \int \left[ (1+s)^{-1} r^2 \bar{\rho}_0^\gamma |\partial_s^k (\zeta, r\zeta_r)|^2 + r^4 \bar{\rho}_0 (\partial_s^k \zeta_s)^2 \right] dr ds \\
& \lesssim \sum_{\iota=0}^k \mathcal{E}_\iota(0) + \sum_{\iota=0}^{k-1} \int_0^t (1+s)^{2\iota-1} \int \left[ r^2 \bar{\rho}_0^\gamma |\partial_s^\iota (\zeta, r\zeta_r)|^2 + (1+s) r^4 \bar{\rho}_0 (\partial_s^\iota \zeta_s)^2 \right] dr \\
& \lesssim \sum_{\iota=0}^k \mathcal{E}_\iota(0).
\end{aligned} \tag{3.72}$$

With this estimate at hand, we finally integrate  $(1+t)^{2k+1}$ (3.43) with respect to the temporal variable and use (3.69), (3.42) and (3.72) to show

$$\begin{aligned}
& (1+t)^{2k} \int \left[ (1+t) r^4 \bar{\rho}_0 |\partial_t^k \zeta_t|^2 + r^2 \bar{\rho}_0^\gamma |\partial_t^k (\zeta, r\zeta_r)|^2 \right] (r, t) dr \\
& + \int_0^t (1+s)^{2k+1} \int r^4 \bar{\rho}_0 (\partial_s^k \zeta_s)^2 dr ds \\
& \lesssim \sum_{\iota=0}^k \mathcal{E}_\iota(0) + \sum_{\iota=0}^k \int_0^t (1+s)^{2\iota-1} \int \left[ r^2 \bar{\rho}_0^\gamma |\partial_s^\iota (\zeta, r\zeta_r)|^2 + (1+s) r^4 \bar{\rho}_0 (\partial_s^\iota \zeta_s)^2 \right] dr \\
& \lesssim \sum_{\iota=0}^k \mathcal{E}_\iota(0).
\end{aligned} \tag{3.73}$$

It finally follows from (3.72) and (3.73) that

$$\mathcal{E}_k(t) + \int_0^t \int (1+s)^{2k-1} \left[ r^2 \bar{\rho}_0^\gamma |\partial_s^k (\zeta, r\zeta_r)|^2 + (1+s)^2 r^4 \bar{\rho}_0 (\partial_s^k \zeta_s)^2 \right] dr ds \lesssim \sum_{\iota=0}^k \mathcal{E}_\iota(0).$$

This completes the proof of Lemma 3.6.  $\square$

### 3.4 Verification of the a priori assumption

In this subsection, we prove the following lemma.

**Lemma 3.7** *Suppose that  $\mathcal{E}(t)$  is finite, then it holds that*

$$\begin{aligned}
& \sum_{j=0}^2 (1+t)^{2j} \left\| \partial_t^j \zeta(\cdot, t) \right\|_{L^\infty}^2 + \sum_{j=0}^1 (1+t)^{2j} \left\| \partial_t^j \zeta_r(\cdot, t) \right\|_{L^\infty}^2 + \sum_{i+j \leq l-2, 2i+j \geq 3} (1+t)^{2j} \\
& \times \left\| \sigma^{\frac{2i+j-3}{2}} \partial_t^j \partial_r^i \zeta(\cdot, t) \right\|_{L^\infty}^2 + \sum_{i+j=l-1} (1+t)^{2j} \left\| r \sigma^{\frac{2i+j-3}{2}} \partial_t^j \partial_r^i \zeta(\cdot, t) \right\|_{L^\infty}^2 \\
& + \sum_{i+j=l} (1+t)^{2j} \left\| r^2 \sigma^{\frac{2i+j-3}{2}} \partial_t^j \partial_r^i \zeta(\cdot, t) \right\|_{L^\infty}^2 \lesssim \mathcal{E}(t). \tag{3.74}
\end{aligned}$$

Once this lemma is proved, the a priori assumption (3.1) is then verified and the proof of Theorem 2.1 is finished, since it follows from the elliptic estimate (3.7) and the nonlinear weighted energy estimate (3.31) that

$$\mathcal{E}(t) \lesssim \mathcal{E}(0), \quad t \in [0, T].$$

**Proof.** The proof consists of two steps. In Step 1, we derive the  $L^\infty$ -bounds away from the boundary, that is,

$$\begin{aligned}
& \sum_{i+j \leq l-2} \left\| \partial_t^j \partial_r^i \zeta \right\|_{L^\infty(\mathcal{I}_o)}^2 + \sum_{i+j=l-1} \left\| r \partial_t^j \partial_r^i \zeta \right\|_{L^\infty(\mathcal{I}_o)}^2 + \sum_{i+j=l-2} \left\| r^2 \partial_t^j \partial_r^i \zeta \right\|_{L^\infty(\mathcal{I}_o)}^2 \\
& \lesssim (1+t)^{-2j} \mathcal{E}(t). \tag{3.75}
\end{aligned}$$

Away from the origin, we show in Step 2 the following  $L^\infty$ -estimates:

$$\sum_{j=0}^3 (1+t)^{2j} \left\| \partial_t^j \zeta \right\|_{L^\infty(\mathcal{I}_b)}^2 + \sum_{j=0}^1 (1+t)^{2j} \left\| \partial_t^j \zeta_r \right\|_{L^\infty(\mathcal{I}_b)}^2 \lesssim \mathcal{E}(t), \tag{3.76}$$

$$\left\| \sigma^{\frac{2i+j-3}{2}} \partial_t^j \partial_r^i \zeta \right\|_{L^\infty(\mathcal{I}_b)}^2 \lesssim (1+t)^{-2j} \mathcal{E}(t) \quad \text{when } 2i+j \geq 4. \tag{3.77}$$

We obtain (3.74) by using (3.75)-(3.77) and noting the facts  $l \geq 4$  and  $\mathcal{I} = \mathcal{I}_o \cup \mathcal{I}_b$ . It suffices to show (3.75)-(3.77).

To this end, we first notice some facts. It follows from (3.24) that  $\mathcal{E}_{j,0} \lesssim \mathcal{E}_j$  for  $j = 0, \dots, l$ , which implies

$$\sum_{j=0}^l \left( \mathcal{E}_j(t) + \sum_{i=0}^{l-j} \mathcal{E}_{j,i}(t) \right) \lesssim \mathcal{E}(t). \tag{3.78}$$

The following embedding (cf. [1]):  $H^{1/2+\delta}(\mathcal{I}) \hookrightarrow L^\infty(\mathcal{I})$  with the estimate

$$\|F\|_{L^\infty(\mathcal{I})} \leq C(\delta) \|F\|_{H^{1/2+\delta}(\mathcal{I})}, \tag{3.79}$$

for  $\delta > 0$  will be used in the rest of the proof.

**Step 1** (away from the boundary). It follows from (3.78) that for  $j = 0, 1, \dots, l$ ,

$$(1+t)^{2j} \left[ \sum_{i=0}^{l-j} \int_{\mathcal{I}_o} r^2 (\partial_t^j \partial_r^i \zeta)^2 dr + \int_{\mathcal{I}_o} r^4 (\partial_t^j \partial_r^{l-j+1} \zeta)^2 dr \right] \lesssim \sum_{i=0}^{l-j} \mathcal{E}_{j,i}(t) \leq \mathcal{E}(t), \quad (3.80)$$

which implies, using (3.4), that for  $j = 0, 1, \dots, l-1$ ,

$$\begin{aligned} \|\partial_t^j \zeta\|_{H^{l-j-1}(\mathcal{I}_o)}^2 &\lesssim \|\partial_t^j \zeta\|_{H^{2,l-j}(\mathcal{I}_o)}^2 = \sum_{i=0}^{l-j} \int_{\mathcal{I}_o} \text{dist}^2(r, \partial \mathcal{I}_o) (\partial_t^j \partial_r^i \zeta)^2 dr \\ &\leq \sum_{i=0}^{l-j} \int_{\mathcal{I}_o} r^2 (\partial_t^j \partial_r^i \zeta)^2 dr \leq (1+t)^{-2j} \mathcal{E}(t). \end{aligned} \quad (3.81)$$

In view of (3.79) and (3.81), we see that for  $j = 0, 1, \dots, l-2$ ,

$$\sum_{i=0}^{l-j-2} \|\partial_t^j \partial_r^i \zeta\|_{L^\infty(\mathcal{I}_o)}^2 \lesssim \sum_{i=0}^{l-j-2} \|\partial_t^j \partial_r^i \zeta\|_{H^1(\mathcal{I}_o)}^2 \lesssim \|\partial_t^j \zeta\|_{H^{l-j-1}(\mathcal{I}_o)}^2 \lesssim (1+t)^{-2j} \mathcal{E}(t). \quad (3.82)$$

It gives from (3.79), (3.80) and (3.81) that for  $j = 0, 1, \dots, l-1$ ,

$$\begin{aligned} \|r \partial_t^j \partial_r^{l-j-1} \zeta\|_{L^\infty(\mathcal{I}_o)}^2 &\lesssim \|r \partial_t^j \partial_r^{l-j-1} \zeta\|_{H^1(\mathcal{I}_o)}^2 \lesssim \|\partial_t^j \partial_r^{l-j-1} \zeta\|_{L^2(\mathcal{I}_o)}^2 + \|r \partial_t^j \partial_r^{l-j} \zeta\|_{L^2(\mathcal{I}_o)}^2 \\ &\lesssim (1+t)^{-2j} \mathcal{E}(t) \end{aligned} \quad (3.83)$$

and for  $j = 0, 1, \dots, l$ ,

$$\begin{aligned} \|r^2 \partial_t^j \partial_r^{l-j} \zeta\|_{L^\infty(\mathcal{I}_o)}^2 &\lesssim \|r^2 \partial_t^j \partial_r^{l-j} \zeta\|_{H^1(\mathcal{I}_o)}^2 \lesssim \|r \partial_t^j \partial_r^{l-j} \zeta\|_{L^2(\mathcal{I}_o)}^2 + \|r^2 \partial_t^j \partial_r^{l-j+1} \zeta\|_{L^2(\mathcal{I}_o)}^2 \\ &\lesssim (1+t)^{-2j} \mathcal{E}(t). \end{aligned} \quad (3.84)$$

So that we can derive (3.75) from (3.82)-(3.84).

**Step 2** (away from the origin). We set

$$d_b(r) := \text{dist}(r, \partial \mathcal{I}_b) \leq \sqrt{A/B} - r \lesssim \sigma(r), \quad r \in \mathcal{I}_b. \quad (3.85)$$

It follows from (3.4) and (3.85) that for  $j \leq 5 + [\alpha] - \alpha$ ,

$$\begin{aligned} \|\partial_t^j \zeta\|_{H^{\frac{5-j+[\alpha]-\alpha}{2}}(\mathcal{I}_b)}^2 &= \|\partial_t^j \zeta\|_{H^{l-j+1-\frac{l-j+1+\alpha}{2}}(\mathcal{I}_b)}^2 \lesssim \|\partial_t^j \zeta\|_{H^{l-j+1+\alpha, l-j+1}(\mathcal{I}_b)}^2 \\ &= \sum_{k=0}^{l-j+1} \int_{\mathcal{I}_b} d_b^{\alpha+1+l-j}(r) |\partial_r^k \partial_t^j \zeta|^2 dr \lesssim \sum_{k=0}^{l-j+1} \int_{\mathcal{I}_b} \sigma^{\alpha+1+l-j} |\partial_r^k \partial_t^j \zeta|^2 dr \\ &\lesssim \sum_{k=0}^{l-j+1} \int_{\mathcal{I}_b} \sigma^{\alpha+k} |\partial_r^k \partial_t^j \zeta|^2 dr \lesssim \sum_{k=0}^{l-j+1} \int_{\mathcal{I}_b} r^4 \sigma^{\alpha+k} |\partial_r^k \partial_t^j \zeta|^2 dr \\ &\leq (1+t)^{-2j} \left( \mathcal{E}_j(t) + \sum_{k=1}^{l-j} \mathcal{E}_{j,k}(t) \right) \leq (1+t)^{-2j} \mathcal{E}(t). \end{aligned}$$

This, together with (3.79), gives (3.76).

To prove (3.77), we denote

$$\psi := \sigma^{\frac{2i+j-3}{2}} \partial_t^j \partial_r^i \zeta.$$

In what follows, we assume  $2i + j \geq 4$  and  $i + j \leq l$  and show that

$$\|\psi\|_{L^\infty(\mathcal{I}_b)}^2 \lesssim (1+t)^{-2j} \mathcal{E}(t). \quad (3.86)$$

The estimate (3.86) will be proved by separating the cases when  $\alpha$  is or is not an integer.

**Case 1** ( $\alpha \neq [\alpha]$ ). When  $\alpha$  is not an integer, we choose  $\sigma^{2(l-i-j)+\alpha-[\alpha]}$  as the spatial weight. A simple calculation yields

$$\begin{aligned} |\partial_r \psi| &\lesssim \left| \sigma^{\frac{2i+j-3}{2}} \partial_t^j \partial_r^{i+1} \zeta \right| + \left| \sigma^{\frac{2i+j-3}{2}-1} \partial_t^j \partial_r^i \zeta \right|, \\ |\partial_r^2 \psi| &\lesssim \left| \sigma^{\frac{2i+j-3}{2}} \partial_t^j \partial_r^{i+2} \zeta \right| + \left| \sigma^{\frac{2i+j-3}{2}-1} \partial_t^j \partial_r^{i+1} \zeta \right| + \left| \sigma^{\frac{2i+j-3}{2}-2} \partial_t^j \partial_r^i \zeta \right|, \\ &\dots\dots\dots \\ |\partial_r^k \psi| &\lesssim \sum_{p=0}^k \left| \sigma^{\frac{2i+j-3}{2}-p} \partial_t^j \partial_r^{i+k-p} \zeta \right| \quad \text{for } k = 1, 2, \dots, l+1-j-i. \end{aligned} \quad (3.87)$$

It follows from (3.87) that for  $1 \leq k \leq l+1-i-j$ ,

$$\begin{aligned} \int_{\mathcal{I}_b} \sigma^{2(l-i-j)+\alpha-[\alpha]} |\partial_r^k \psi|^2 dr &\lesssim \sum_{p=0}^k \int_{\mathcal{I}_b} \sigma^{\alpha+l-j+1-2p} |\partial_t^j \partial_r^{i+k-p} \zeta|^2 dr \\ &\lesssim \int_{\mathcal{I}_b} \sigma^{l-i-j+1-k} \sum_{p=0}^1 \sigma^{\alpha+i+k-2p} |\partial_t^j \partial_r^{i+k-p} \zeta|^2 dr + \sum_{p=2}^k \int_{\mathcal{I}_b} \sigma^{\alpha+l-j+1-2p} |\partial_t^j \partial_r^{i+k-p} \zeta|^2 dr \\ &\lesssim \sum_{p=0}^1 \int_{\mathcal{I}_b} r^4 \sigma^{\alpha+i+k-2p} |\partial_t^j \partial_r^{i+k-p} \zeta|^2 dr + \sum_{p=2}^k \int_{\mathcal{I}_b} \sigma^{\alpha+l-j+1-2p} |\partial_t^j \partial_r^{i+k-p} \zeta|^2 dr \\ &\lesssim (1+t)^{-2j} \mathcal{E}_{j,i+k-1} + \sum_{p=2}^k \int_{\mathcal{I}_b} \sigma^{\alpha+l-j+1-2p} |\partial_t^j \partial_r^{i+k-p} \zeta|^2 dr. \end{aligned}$$

To bound the 2nd term on the right-hand side of the inequality above, notice that

$$\begin{aligned} &\alpha + l - j + 1 - 2p \\ &= 2(l+1-i-j-k) + 2(k-p) + (\alpha - [\alpha]) + (2i+j-4) - 1 > -1 \end{aligned} \quad (3.88)$$

for  $p \in [2, k]$ , due to  $\alpha > [\alpha]$  and  $2i + j \geq 4$ . We then have, with the aid of the Hardy inequality (3.6), that for  $p \in [2, k]$ ,

$$\begin{aligned} \int_{\mathcal{I}_b} \sigma^{\alpha+l-j+1-2p} |\partial_t^j \partial_r^{i+k-p} \zeta|^2 dr &\lesssim \int_{\mathcal{I}_b} \sigma^{\alpha+l-j+1-2p+2} \sum_{\iota=0}^1 |\partial_t^j \partial_r^{i+k-p+\iota} \zeta|^2 dr \lesssim \dots \\ &\lesssim \int_{\mathcal{I}_b} \sigma^{\alpha+l-j+1} \sum_{\iota=0}^p |\partial_t^j \partial_r^{i+k-p+\iota} \zeta|^2 dr = \sum_{\iota=0}^p \int_{\mathcal{I}_b} \sigma^{(l+1-i-j-k)+(p-\iota)} \sigma^{\alpha+i+k-p+\iota} |\partial_t^j \partial_r^{i+k-p+\iota} \zeta|^2 dr \\ &\lesssim \sum_{\iota=0}^p \int_{\mathcal{I}_b} r^4 \sigma^{\alpha+i+k-p+\iota} |\partial_t^j \partial_r^{i+k-p+\iota} \zeta|^2 dr \leq \sum_{\iota=i+k-p}^{i+k-1} (1+t)^{-2j} \mathcal{E}_{j,\iota}. \end{aligned}$$

That yields, for  $1 \leq k \leq l+1-i-j$ ,

$$\begin{aligned} \int_{\mathcal{I}_b} \sigma^{2(l-i-j)+\alpha-[\alpha]} |\partial_r^k \psi|^2 dr &\lesssim (1+t)^{-2j} \mathcal{E}_{j,i+k-1} + \sum_{p=2}^k \sum_{\iota=i+k-p}^{i+k-1} (1+t)^{-2j} \mathcal{E}_{j,\iota} \\ &\lesssim (1+t)^{-2j} \sum_{\iota=i}^{i+k-1} \mathcal{E}_{j,\iota}. \end{aligned}$$

Therefore, it follows from (3.85) and (3.78) that

$$\begin{aligned} \|\psi\|_{H^{2(l-i-j)+\alpha-[\alpha], l+1-i-j}(\mathcal{I}_b)}^2 &= \sum_{k=0}^{l+1-i-j} \int_{\mathcal{I}_b} d_b^{2(l-i-j)+\alpha-[\alpha]} |\partial_r^k \psi|^2 dr \\ &\lesssim \sum_{k=0}^{l+1-i-j} \int_{\mathcal{I}_b} \sigma^{2(l-i-j)+\alpha-[\alpha]} |\partial_r^k \psi|^2 dr \lesssim \int_{\mathcal{I}_b} \sigma^{\alpha+l-j+1} |\partial_t^j \partial_r^i \zeta|^2 dr + (1+t)^{-2j} \sum_{\iota=i}^{l-j} \mathcal{E}_{j,\iota} \\ &\lesssim \int_{\mathcal{I}_b} r^4 \sigma^{\alpha+i+1} |\partial_t^j \partial_r^i \zeta|^2 dr + (1+t)^{-2j} \sum_{\iota=i}^{l-j} \mathcal{E}_{j,\iota} \lesssim (1+t)^{-2j} \sum_{\iota=i}^{l-j} \mathcal{E}_{j,\iota} \leq (1+t)^{-2j} \mathcal{E}(t). \end{aligned}$$

When  $\alpha$  is not an integer,  $\alpha - [\alpha] \in (0, 1)$ . So, it follows from (3.79) and (3.4) that

$$\|\psi\|_{L^\infty(\mathcal{I}_b)}^2 \lesssim \|\psi\|_{H^{1-\frac{\alpha-[\alpha]}{2}}(\mathcal{I}_b)}^2 \lesssim \|\psi\|_{H^{2(l-i-j)+\alpha-[\alpha], l+1-i-j}(\mathcal{I}_b)}^2 \lesssim (1+t)^{-2j} \mathcal{E}(t). \quad (3.89)$$

**Case 2** ( $\alpha = [\alpha]$ ). In this case  $\alpha$  is an integer, we choose  $\sigma^{2(l-i-j)+1/2}$  as the spatial weight. As shown in Case 1, we have for  $1 \leq k \leq l+1-i-j$ ,

$$\int_{\mathcal{I}_b} \sigma^{2(l-i-j)+1/2} |\partial_r^k \psi|^2 dr \lesssim (1+t)^{-2j} \mathcal{E}_{j,i+k-1} + \sum_{p=2}^k \int_{\mathcal{I}_b} \sigma^{\alpha+l-j+1-2p+1/2} |\partial_t^j \partial_r^{i+k-p} \zeta|^2 dr.$$

Note that for  $1 \leq k \leq l+1-i-j$  and  $2 \leq p \leq k$ ,

$$\alpha + l - j + 1 - 2p + \frac{1}{2} = 2(l+1-i-j-k) + 2(k-p) + (2i+j-4) - \frac{1}{2} \geq -\frac{1}{2}.$$

We can then use the Hardy inequality (3.6) to obtain

$$\int_{\mathcal{I}_b} \sigma^{2(l-i-j)+1/2} |\partial_r^k \psi|^2 dr \lesssim (1+t)^{-2j} \sum_{\iota=i}^{i+k-1} \mathcal{E}_{j,\iota}, \quad k = 1, 2, \dots, l-j+1-i,$$

which, together with (3.85) and (3.78), implies that

$$\|\psi\|_{H^{2(l-i-j)+1/2, l+1-i-j}(\mathcal{I}_b)}^2 \lesssim (1+t)^{-2j} \sum_{\iota=i}^{l-j} \mathcal{E}_{j,\iota} \leq (1+t)^{-2j} \mathcal{E}(t).$$

Therefore, it follows from (3.79) and (3.4) that

$$\|\psi\|_{L^\infty(\mathcal{I}_b)}^2 \lesssim \|\psi\|_{H^{3/4}(\mathcal{I}_b)}^2 \lesssim \|\psi\|_{H^{2(l-i-j)+1/2, l+1-i-j}(\mathcal{I}_b)}^2 \lesssim (1+t)^{-2j} \mathcal{E}(t). \quad (3.90)$$

In view of (3.89) and (3.90), we obtain (3.86) or equivalently (3.77). This completes the proof of Lemma 3.7.  $\square$

## 4 Proof of Theorem 2.2

In this section, we prove Theorem 2.2. First, it follows from (2.3), (2.6), (1.6), (2.9), (2.2) and (2.8) that for  $(r, t) \in \mathcal{I} \times [0, \infty)$ ,

$$\rho(\eta(r, t), t) - \bar{\rho}(\bar{\eta}(r, t), t) = \frac{r^2 \bar{\rho}_0(r)}{\eta^2(r, t) \eta_r(r, t)} - \frac{r^2 \bar{\rho}_0(r)}{\bar{\eta}^2(r, t) \bar{\eta}_r(r, t)}$$

and

$$u(\eta(r, t), t) - \bar{u}(\bar{\eta}(r, t), t) = \eta_t(r, t) - \bar{\eta}_t(r, t).$$

Then, we have, using (2.15), (2.11), (2.9), (3.2), (2.14) and (2.17) that

$$|\rho(\eta(r, t), t) - \bar{\rho}(\bar{\eta}(r, t), t)| \lesssim (A - Br^2)^{\frac{1}{\gamma-1}} (1+t)^{-\frac{4}{3\gamma-1}} \left[ \sqrt{\mathcal{E}(0)} + (1+t)^{-\frac{3\gamma-2}{3\gamma-1}} \ln(1+t) \right]$$

and

$$|u(\eta(r, t), t) - \bar{u}(\bar{\eta}(r, t), t)| \lesssim r(1+t)^{-1} \left[ \sqrt{\mathcal{E}(0)} + (1+t)^{-\frac{3\gamma-2}{3\gamma-1}} \ln(1+t) \right].$$

This gives the proof of (2.18) and (2.19).

For the boundary behavior, it follows from (2.5), (2.15), (2.11) and (2.9) that

$$\begin{aligned} R(t) &= \eta \left( \sqrt{A/B}, t \right) = (\bar{\eta} + r\zeta) \left( \sqrt{A/B}, t \right) = (\bar{\eta} + rh + r\zeta) \left( \sqrt{A/B}, t \right) \\ &= [r(\bar{\eta}_r + h + \zeta)] \left( \sqrt{A/B}, t \right) = \sqrt{A/B} \left[ (1+t)^{1/(3\gamma-1)} + h(t) + \zeta(t) \right] \end{aligned}$$

which, together with (2.17) and (2.14) that

$$R(t) \geq \sqrt{A/B} \left[ (1+t)^{\frac{1}{3\gamma-1}} - C\sqrt{\mathcal{E}(0)} \right]$$

and

$$R(t) \leq \sqrt{A/B} \left[ (1+t)^{\frac{1}{3\gamma-1}} + C(1+t)^{-\frac{3\gamma-2}{3\gamma-1}} \ln(1+t) + C\sqrt{\mathcal{E}(0)} \right].$$

Thus, (2.20) follows from the smallness of  $\mathcal{E}(0)$ . Notice that for  $k = 1, 2, 3$ ,

$$\frac{d^k R(t)}{dt^k} = \partial_t^k \bar{\eta} \left( \sqrt{A/B}, t \right) + (r \partial_t^k \zeta) \left( \sqrt{A/B}, t \right).$$

Therefore, (2.21) follows from (2.13) and (2.17).

We are to verify the physical vacuum condition, (2.22). It follows from (2.3), (2.6), (2.15) that

$$\begin{aligned} (\rho^{\gamma-1})_\eta(\eta, t) &= \frac{(f^{\gamma-1})_r(r, t)}{\eta_r(r, t)} = \frac{1}{\eta_r} \left[ \bar{\rho}_0^{\gamma-1} \left( \frac{r^2}{\eta^2 \eta_r} \right)^{\gamma-1} \right]_r \\ &= \frac{1}{\eta_r} \left\{ \bar{\rho}_0^{\gamma-1} \left[ \bar{\rho}_0^{\gamma-1} \left( \frac{r^2}{\eta^2 \eta_r} \right)^{\gamma-1} \right]_r - 2Br \left( \frac{r^2}{\eta^2 \eta_r} \right)^{\gamma-1} \right\} \\ &= (1-\gamma) \bar{\rho}_0^{\gamma-1} \left[ 2 \left( \frac{\eta}{r} \right)^{1-2\gamma} \eta_r^{-\gamma} \zeta_r + \left( \frac{\eta}{r} \right)^{2-2\gamma} \eta_r^{-\gamma-1} (2\zeta_r + r\zeta_{rr}) \right] - 2Br \left( \frac{\eta}{r} \right)^{2-2\gamma} \eta_r^{-\gamma}, \end{aligned}$$

which implies, with the aid of (3.3) and (2.17), that

$$\left| (\rho^{\gamma-1})_{\eta}(\eta, t) \right| \lesssim (1+t)^{-1} \sqrt{\mathcal{E}(0)} + r(1+t)^{-1+\frac{1}{3\gamma-1}} \quad (4.1)$$

and

$$\begin{aligned} \left| (\rho^{\gamma-1})_{\eta}(\eta, t) \right| &\geq 2Br \left( \frac{\eta}{r} \right)^{2-2\gamma} \eta_r^{-\gamma} - C(1+t)^{-1} \sqrt{\mathcal{E}(0)} \\ &\geq C^{-1} r (1+t)^{-1+\frac{1}{3\gamma-1}} - C(1+t)^{-1} \sqrt{\mathcal{E}(0)}. \end{aligned} \quad (4.2)$$

In view of (3.3), we see that

$$\eta(r, t) \sim (1+t)^{\frac{1}{3\gamma-1}} r,$$

which, together with (2.20), (4.1) and (4.2), gives for  $R(t)/2 \leq \eta \leq R(t)$ ,

$$C^{-1}(1+t)^{-\frac{3\gamma-2}{3\gamma-1}} - C(1+t)^{-1} \sqrt{\mathcal{E}(0)} \leq \left| (\rho^{\gamma-1})_{\eta}(\eta, t) \right| \lesssim (1+t)^{-1} \sqrt{\mathcal{E}(0)} + (1+t)^{-\frac{3\gamma-2}{3\gamma-1}}.$$

Thus, (2.22) follows from the smallness of  $\mathcal{E}(0)$ . This finishes the proof of Theorem 2.2.  $\square$

## 5 Appendix

**Proof of (2.13)<sub>1</sub>.** We may write (2.10) as the following system

$$\begin{cases} h_t = z, \\ z_t = -z - [\bar{\eta}_r^{2-3\gamma} - (\bar{\eta}_r + h)^{2-3\gamma}] / (3\gamma - 1) - \bar{\eta}_{rtt} \\ (h, z)(t=0) = (0, 0). \end{cases} \quad (5.1)$$

Recalling that  $\bar{\eta}_r(t) = (1+t)^{1/(3\gamma-1)}$ , thus  $\bar{\eta}_{rtt} < 0$ . A simple phase plane analysis shows that there exist  $0 < t_0 < t_1 < t_2$  such that, starting from  $(h, z) = (0, 0)$  at  $t = 0$ ,  $h$  and  $z$  increases in the interval  $[0, t_0]$  and  $z$  reaches its positive maxima at  $t_0$ ; in the interval  $[t_0, t_1]$ ,  $h$  keeps increasing and reaches its maxima at  $t_1$ ,  $z$  decreases from its positive maxima to 0; in the interval  $[t_1, t_2]$ , both  $h$  and  $z$  decrease, and  $z$  reaches its negative minima at  $t_2$ ; in the interval  $[t_2, \infty)$ ,  $h$  decreases and  $z$  increases, and  $(h, z) \rightarrow (0, 0)$  as  $t \rightarrow \infty$ . This can be summarized as follows:

$$\begin{aligned} z(t) \uparrow_0, \quad h(t) \uparrow_0, \quad t \in [0, t_0] \\ z(t) \downarrow_0, \quad h(t) \uparrow, \quad t \in [t_0, t_1] \\ z(t) \downarrow^0, \quad h(t) \downarrow, \quad t \in [t_1, t_2] \\ z(t) \uparrow^0, \quad h(t) \downarrow_0, \quad t \in [t_2, \infty). \end{aligned}$$

It gives from the above analysis that there exists a finite constant  $C = C(\gamma, M)$  such that

$$0 \leq h(t) \leq C \quad \text{for } t \geq 0. \quad (5.2)$$

In view of (2.9) and (2.11), we then see that

$$(1+t)^{1/(3\gamma-1)} \leq \tilde{\eta}_r(t) \leq K(1+t)^{1/(3\gamma-1)}.$$

On the other hand, equation (2.10) can be rewritten as

$$\begin{aligned} \tilde{\eta}_{rtt} + \tilde{\eta}_{rt} - \tilde{\eta}_r^{2-3\gamma}/(3\gamma - 1) &= 0, \quad t > 0, \\ \tilde{\eta}_r(t = 0) &= 1, \quad \tilde{\eta}_{rt}(t = 0) = 1/(3\gamma - 1). \end{aligned} \quad (5.3)$$

Then, we have by solving (5.3) that

$$\tilde{\eta}_{rt}(t) = \frac{1}{3\gamma - 1}e^{-t} + \frac{1}{3\gamma - 1} \int_0^t e^{-(t-s)} \tilde{\eta}_r^{2-3\gamma}(s) ds \geq 0. \quad \square \quad (5.4)$$

**Proof of (2.13)<sub>2</sub>.** We use the mathematical induction to prove (2.13)<sub>2</sub>. First, it follows from (5.4) that

$$\begin{aligned} (3\gamma - 1)\tilde{\eta}_{rt}(t) &= e^{-t} + \int_0^{t/2} e^{-(t-s)} \tilde{\eta}_r^{2-3\gamma}(s) ds + \int_{t/2}^t e^{-(t-s)} \tilde{\eta}_r^{2-3\gamma}(s) ds \\ &\leq e^{-t} + e^{-t/2} \int_0^{t/2} (1+s)^{\frac{2-3\gamma}{3\gamma-1}} ds + \left(1 + \frac{t}{2}\right)^{\frac{2-3\gamma}{3\gamma-1}} \int_{t/2}^t e^{-(t-s)} ds \\ &\leq e^{-t} + C e^{-t/2} \left(1 + \frac{t}{2}\right)^{\frac{1}{3\gamma-1}} + \left(1 + \frac{t}{2}\right)^{\frac{2-3\gamma}{3\gamma-1}} \\ &\leq C(1+t)^{\frac{2-3\gamma}{3\gamma-1}}, \quad t \geq 0, \end{aligned} \quad (5.5)$$

for some constant  $C$  independent of  $t$ . This proves (2.13)<sub>2</sub> when  $k = 1$ . Suppose that (2.13)<sub>2</sub> holds for all  $k = 1, 2, \dots, m-1$ , that is,

$$\left| \frac{d^k \tilde{\eta}_r(t)}{dt^k} \right| \leq C(m) (1+t)^{\frac{1}{3\gamma-1}-k}, \quad k = 1, 2, \dots, m-1. \quad (5.6)$$

It suffices to prove (2.13)<sub>2</sub> holds for  $k = m$ . We derive from (5.3) that for  $m = 1, \dots, k$ ,

$$\frac{d^{m+1}}{dt^{m+1}} \tilde{\eta}_r(t) + \frac{d^m}{dt^m} \tilde{\eta}_r(t) - \frac{1}{3\gamma - 1} \frac{d^{m-1}}{dt^{m-1}} \tilde{\eta}_r^{2-3\gamma}(t) = 0, \quad t \geq 0,$$

so that

$$\frac{d^m}{dt^m} \tilde{\eta}_r(t) = e^{-t} \frac{d^m}{dt^m} \tilde{\eta}_r(0) + \frac{1}{3\gamma - 1} \int_0^t e^{-(t-s)} \frac{d^{m-1} \tilde{\eta}_r^{2-3\gamma}}{ds^{m-1}}(s) ds, \quad t \geq 0, \quad (5.7)$$

where  $(d^m/dt^m)\tilde{\eta}_r(0)$  is finite, which can be determined by the equation inductively. In view of (2.13)<sub>1</sub> and (5.6), we see that

$$\begin{aligned} \left| \frac{d}{dt} \tilde{\eta}_r^{2-3\gamma}(t) \right| &\lesssim \left| \tilde{\eta}_r^{1-3\gamma}(t) \frac{d}{dt} \tilde{\eta}_r(t) \right| \lesssim (1+t)^{\frac{1}{3\gamma-1}-2}, \\ \left| \frac{d^2}{dt^2} \tilde{\eta}_r^{2-3\gamma}(t) \right| &\lesssim \left| \tilde{\eta}_r^{-3\gamma}(t) \left( \frac{d}{dt} \tilde{\eta}_r(t) \right)^2 \right| + \left| \tilde{\eta}_r^{1-3\gamma}(t) \frac{d^2}{dt^2} \tilde{\eta}_r(t) \right| \lesssim (1+t)^{\frac{1}{3\gamma-1}-3} \end{aligned}$$



$$\left| \frac{d^{m-1}}{dt^{m-1}} \tilde{\eta}_r^{2-3\gamma}(t) \right| \leq C(\gamma, m)(1+t)^{\frac{1}{3\gamma-1}-m}. \quad (5.8)$$

Similar to deriving (5.5), we can obtain, noting (5.7) and (5.8), that

$$\left| \frac{d^m \tilde{\eta}_r(t)}{dt^m} \right| \leq C(\gamma, m)(1+t)^{\frac{1}{3\gamma-1}-m}.$$

This finishes the proof of (2.13)<sub>2</sub>.  $\square$

**Proof of (2.14).** We may write the equation for  $h$ , (2.10), as

$$h_t + \frac{1}{3\gamma-1}(1+t)^{-\frac{3\gamma-2}{3\gamma-1}} \left[ 1 - \left( 1 + (1+t)^{-\frac{1}{3\gamma-1}} h \right)^{2-3\gamma} \right] = -\tilde{\eta}_{rtt}, \quad t > 0 \quad (5.9)$$

Notice that

$$\left( 1 + (1+t)^{-\frac{1}{3\gamma-1}} h \right)^{2-3\gamma} \leq 1 + (2-3\gamma)(1+t)^{-\frac{1}{3\gamma-1}} h + \frac{(2-3\gamma)(1-3\gamma)}{2}(1+t)^{-\frac{2}{3\gamma-1}} h^2,$$

due to  $h \geq 0$ . We then obtain, in view of (2.13)<sub>2</sub>, that

$$h_t + \frac{3\gamma-2}{3\gamma-1}(1+t)^{-1} h \leq \frac{3\gamma-2}{2}(1+t)^{-\frac{3\gamma}{3\gamma-1}} h^2 + C(1+t)^{\frac{1}{3\gamma-1}-2}.$$

Thus,

$$h(t) \leq C(1+t)^{-\frac{3\gamma-2}{3\gamma-1}} \int_0^t \left( (1+s)^{-\frac{2}{3\gamma-1}} h^2(s) + (1+s)^{-1} \right) ds. \quad (5.10)$$

We use an iteration to prove (2.14). First, since  $h$  is bounded due to (5.2), we have

$$h(t) \leq C(1+t)^{-\frac{3\gamma-2}{3\gamma-1}} \int_0^t (1+s)^{-\frac{2}{3\gamma-1}} ds \leq C(1+t)^{-\frac{1}{3\gamma-1}}. \quad (5.11)$$

Substituting this into (5.10), we obtain

$$h(t) \leq C(1+t)^{-\frac{3\gamma-2}{3\gamma-1}} \int_0^t \left( (1+s)^{-\frac{4}{3\gamma-1}} + (1+s)^{-1} \right) ds;$$

which implies

$$h(t) \leq \begin{cases} C(1+t)^{-\frac{3\gamma-2}{3\gamma-1}} \ln(1+t) & \text{if } \gamma \leq 5/3, \\ C(1+t)^{-\frac{3}{3\gamma-1}} & \text{if } \gamma > 5/3. \end{cases}$$

If  $\gamma \leq 5/3$ , then the first part of (2.14) has been proved. If  $\gamma > 5/3$ , we repeat this procedure and obtain

$$h(t) \leq C(1+t)^{-\frac{3\gamma-2}{3\gamma-1}} \int_0^t \left( (1+s)^{-\frac{8}{3\gamma-1}} + (1+s)^{-1} \right) ds;$$

which implies

$$h(t) \leq \begin{cases} C(1+t)^{-\frac{3\gamma-2}{3\gamma-1}} \ln(1+t) & \text{if } \gamma \leq 3, \\ C(1+t)^{-\frac{7}{3\gamma-1}} & \text{if } \gamma > 3. \end{cases}$$

For general  $\gamma$ , we repeat this procedure  $k$  times to obtain

$$h(t) \leq C(1+t)^{-\frac{3\gamma-2}{3\gamma-1}} \ln(1+t).$$

This, together with (5.2), proves the first part of (2.14), which in turn implies the second part of (2.14), by virtue of (5.9) and (2.13)<sub>2</sub>.  $\square$

**Proof of (3.27).** Recall that  $j \geq 0$ ,  $i \geq 1$  and  $i + j \leq l$ . Let  $n \in [0, j]$ ,  $m \in [0, i - 1]$  and  $q \in [0, n]$  be integers. Denote

$$\mathcal{H} := \left\| r\sigma^{\frac{\alpha+i-1}{2}} (|r\partial_t^{n-q}\partial_r^{m+1}\zeta| + |\partial_t^{n-q}\partial_r^m\zeta|) \left( |\sigma r\partial_t^{j-n}\partial_r^{i-m+1}\zeta| + \sum_{\iota=0}^{i-m} |\partial_t^{j-n}\partial_r^\iota\zeta| \right) \right\|^2.$$

**Case 1.** Assume  $2n + 4m \geq 2i + j + q$ . We first note that

$$\alpha + (2m + n) - (i + j) + 2 \geq \alpha - \frac{j}{2} + \frac{q}{2} + 2 \geq \alpha - \frac{l-1}{2} + 2 \geq 0, \quad (5.12)$$

$$i + j - (n + m) \leq l - 2. \quad (5.13)$$

(Indeed, if  $i + j - (n + m) = l$ , then  $i + j = l$  and  $n + m = 0$ , so that it is a contradiction due to

$$0 = 4(n + m) \geq 2n + 4m \geq 2i + j + q \geq i + j = l;$$

if  $i + j - (n + m) = l - 1$ , then  $i + j = l - 1$  and  $n + m = 0$  or  $i + j = l$  and  $n + m = 1$ , so that it is also a contradiction because of

$$0 = 4(n + m) \geq 2n + 4m \geq 2i + j + q \geq i + j = l - 1 > 0$$

or

$$4 = 4(n + m) \geq 2n + 4m \geq 2i + j + q \geq i + i + j \geq 1 + l = 5 + [\alpha] \geq 5.$$

So, (5.13) holds.)

When  $2i + j \leq 2m + n + 3$ , it follows from (3.1) and (5.13) that

$$\begin{aligned} \mathcal{H} &\lesssim \epsilon_0^2 (1+t)^{2n-2j} \left\| r\sigma^{\frac{\alpha+i-1}{2}} (|r\partial_t^{n-q}\partial_r^{m+1}\zeta| + |\partial_t^{n-q}\partial_r^m\zeta|) \right\|^2 \\ &\lesssim \epsilon_0^2 (1+t)^{2q-2j} (\mathcal{E}_{n-q,m+1} + \mathcal{E}_{n-q,m}). \end{aligned} \quad (5.14)$$

When  $2i + j \geq 2m + n + 4$ , it follows from (3.1) and (5.13) that

$$\begin{aligned} \mathcal{H} &\lesssim \epsilon_0^2 (1+t)^{2n-2j} \left\| r\sigma^{\frac{\alpha+i-1}{2} - \frac{j+2i-(n+2m)-3}{2}} (|r\partial_t^{n-q}\partial_r^{m+1}\zeta| + |\partial_t^{n-q}\partial_r^m\zeta|) \right\|^2 \\ &= \epsilon_0^2 (1+t)^{2n-2j} \left\| r\sigma^{\frac{\alpha+m+(n+m-(i+j)+2)}{2}} (|r\partial_t^{n-q}\partial_r^{m+1}\zeta| + |\partial_t^{n-q}\partial_r^m\zeta|) \right\|^2; \end{aligned}$$

which implies for  $n + m - (i + j) + 2 \geq 0$ ,

$$\begin{aligned} \mathcal{H} &\lesssim \epsilon_0^2 (1+t)^{2n-2j} \left\| r \sigma^{\frac{\alpha+m}{2}} (|\partial_t^{n-q} \partial_r^{m+1} \zeta| + |\partial_t^{n-q} \partial_r^m \zeta|) \right\|^2 \\ &\lesssim \epsilon_0^2 (1+t)^{2q-2j} (\mathcal{E}_{n-q,m+1} + \mathcal{E}_{n-q,m}) \end{aligned} \quad (5.15)$$

and for  $n + m - (i + j) + 2 \leq -1$ ,

$$\begin{aligned} \mathcal{H} &\lesssim \epsilon_0^2 (1+t)^{2n-2j} \left( \left\| r^2 \partial_t^{n-q} \partial_r^{m+1} \zeta \right\|_{L^2(\mathcal{I}_o)}^2 + \left\| r \partial_t^{n-q} \partial_r^m \zeta \right\|_{L^2(\mathcal{I}_o)}^2 \right. \\ &\quad \left. + \left\| \sigma^{\frac{\alpha+m+(n+m-(i+j)+2)}{2}} (|\partial_t^{n-q} \partial_r^{m+1} \zeta| + |\partial_t^{n-q} \partial_r^m \zeta|) \right\|_{L^2(\mathcal{I}_b)}^2 \right) \\ &\lesssim \epsilon_0^2 (1+t)^{2n-2j} \left( \left\| r^2 \sigma^{\frac{\alpha+m+1}{2}} \partial_t^{n-q} \partial_r^{m+1} \zeta \right\|_{L^2(\mathcal{I}_o)}^2 + \left\| r \sigma^{\frac{\alpha+m-1}{2}} \partial_t^{n-q} \partial_r^m \zeta \right\|_{L^2(\mathcal{I}_o)}^2 \right) \\ &\quad + \epsilon_0^2 (1+t)^{2q-2j} \sum_{h=m}^{i+j-n+q-1} \mathcal{E}_{n-q,h} \lesssim \epsilon_0^2 (1+t)^{2q-2j} \sum_{h=m}^{i+j-n+q-1} \mathcal{E}_{n-q,h}. \end{aligned} \quad (5.16)$$

Here we have used (5.12) and the Hardy inequality (3.6) to derive

$$\begin{aligned} &\left\| \sigma^{\frac{\alpha+m+(n+m-(i+j)+2)}{2}} (|\partial_t^{n-q} \partial_r^{m+1} \zeta| + |\partial_t^{n-q} \partial_r^m \zeta|) \right\|_{L^2(\mathcal{I}_b)}^2 \\ &\lesssim \sum_{h=0}^1 \left\| \sigma^{\frac{\alpha+m+(n+m-(i+j)+2)}{2}} \partial_t^{n-q} \partial_r^{m+h} \zeta \right\|_{L^2(\mathcal{I}_b)}^2 \lesssim \sum_{h=0}^2 \left\| \sigma^{\frac{\alpha+m+(n+m-(i+j)+2)+2}{2}} \partial_t^{n-q} \partial_r^{m+h} \zeta \right\|_{L^2(\mathcal{I}_b)}^2 \\ &\lesssim \dots \lesssim \sum_{h=0}^{i+j-(n+m)+q} \left\| \sigma^{\frac{\alpha+m+(n+m-(i+j)+2)+2(i+j-(n+m)+q-1)}{2}} \partial_t^{n-q} \partial_r^{m+h} \zeta \right\|_{L^2(\mathcal{I}_b)}^2 \\ &= \sum_{h=0}^{i+j-(n+m)+q} \left\| \sigma^{\frac{\alpha+i+j-n+2q}{2}} \partial_t^{n-q} \partial_r^{m+h} \zeta \right\|_{L^2(\mathcal{I}_b)}^2 \\ &\lesssim \sum_{h=0}^{i+j-(n+m)+q} \left\| r^2 \sigma^{\frac{\alpha+i+j-n+q}{2}} \partial_t^{n-q} \partial_r^{m+h} \zeta \right\|_{L^2(\mathcal{I}_b)}^2 \lesssim (1+t)^{2q-2n} \sum_{h=m}^{i+j-n+q-1} \mathcal{E}_{n-q,h}, \end{aligned}$$

which implies (5.16). Therefore, it gives from (5.14), (5.15) and (5.16) that

$$\mathcal{H} \lesssim \epsilon_0^2 (1+t)^{2q-2j} \left( \mathcal{E}_{n-q,m} + \mathcal{E}_{n-q,m+1} + \sum_{h=m}^{i+j-n+q-1} \mathcal{E}_{n-q,h} \right). \quad (5.17)$$

**Case 2.** Assume  $2n + 4m < 2i + j + q$ . In this case, we can use the similar way to dealing with Case 1 to obtain

$$\mathcal{H} \lesssim \epsilon_0^2 (1+t)^{2q-2j} \sum_{h=0}^{i+n-j} \mathcal{E}_{j-n,h}(t). \quad (5.18)$$

In view of (5.17) and (5.18), we prove (3.27).  $\square$

**Acknowledgements** The author is grateful to Professor Tao Luo for many helpful discussions. This work was supported in part by NSFC under grants 11301293 and 11271219, and the Center of Mathematical Sciences and Applications, Harvard University.

## References

- [1] Adams, R.: *Sobolev Spaces*, New York: Academic Press 1975.
- [2] Alazard, T., Burq, N., Zuily, C.: On the Cauchy problem for gravity water waves, *Invent. Math.* **198**, 71–163 (2014).
- [3] Ambrose, D., Masmoudi, N.: The zero surface tension limit of three-dimensional water waves, *Indiana Univ. Math. J.* **58**, 479–521 (2009).
- [4] Barenblatt, G.: On one class of solutions of the one-dimensional problem of non-stationary filtration of a gas in a porous medium, *Prikl. Mat. i. Mekh.* **17**, 739–742 (1953).
- [5] Chandrasekhar, S.: *Introduction to the Stellar Structure*, University of Chicago Press (1939).
- [6] Christodoulou, D., Lindblad, H.: On the motion of the free surface of a liquid, *Comm. Pure Appl. Math.* **53**, 1536–1602 (2000).
- [7] Coutand, D., Lindblad, H., Shkoller, S.: A priori estimates for the free-boundary 3-D compressible Euler equations in physical vacuum, *Comm. Math. Phys.* **296**, 559–587 (2010).
- [8] Coutand, D., Shkoller, S.: Well-posedness of the free-surface incompressible Euler equations with or without surface tension, *J. Am. Math. Soc.* **20**, 829–930 (2007).
- [9] Coutand, D., Shkoller, S.: Well-posedness in smooth function spaces for the moving-boundary 1-D compressible Euler equations in physical vacuum, *Comm. Pure Appl. Math.* **64**, 328–366 (2011).
- [10] Coutand, D., Shkoller, S.: Well-Posedness in Smooth Function Spaces for the Moving-Boundary Three-Dimensional Compressible Euler Equations in Physical Vacuum, *Arch. Ration. Mech. Anal.* **206**, 515–616 (2012).
- [11] Cox, J., Giuli, R.: *Principles of stellar structure, I.,II.*, New York: Gordon and Breach, 1968.
- [12] Gu, X., Lei, Z.: Well-posedness of 1-D compressible Euler-Poisson equations with physical vacuum, *J. Diff. Equ.* **252**, 2160–2188 (2012).
- [13] Gu, X., Lei, Z.: Local Well-posedness of the three dimensional compressible Euler–Poisson equations with physical vacuum, *arXiv:1405.4612*.
- [14] Hanouzet B., Natalini R.: Global existence of smooth solutions for partially dissipative hyperbolic systems with a convex entropy, *Archive for Rational Mechanics and Analysis* **169** (2003) 89–117.
- [15] Hsiao, L., Liu, T.: Convergence to nonlinear diffusion waves for solutions of a system of hyperbolic conservation laws with damping, *Comm. Math. Phys.* **143**, 599–605 (1992).

- [16] Huang, F., Marcati, P., Pan, R.: Convergence to the Barenblatt solution for the compressible Euler equations with damping and vacuum, *Arch. Ration. Mech. Anal.* **176**, 1–24 (2005).
- [17] Huang, H., Pan, R., Wang, Z.:  $L^1$  convergence to the Barenblatt solution for compressible Euler equations with damping, *Arch. Ration. Mech. Anal.* **200**, 665–689 (2011).
- [18] Germain, P., Masmoudi, N., Shatah, J.: Global solutions for the gravity water waves equation in dimension 3, *Ann. of Math.* **175**, 691–754 (2012).
- [19] Ionescu, A, Pusateri, F.: Global solutions for the gravity water waves system in 2d, *Invent. Math.* (forthcoming), DOI 10.1007/s00222-014-0521-4.
- [20] Jang, J., Masmoudi, N.: Well-posedness for compressible Euler with physical vacuum singularity, *Comm. Pure Appl. Math.* **62**, 1327–1385 (2009).
- [21] Jang, J., Masmoudi, N.: Well-posedness of compressible Euler equations in a physical vacuum, *Comm. Pure Appl. Math.* 68 (2015), no. 1, 61C111.
- [22] Jang, J., Masmoudi, N.: Well and ill-posedness for compressible Euler equations with vacuum, *J. Math. Phys.* **53**, 115625, 11 pp (2012).
- [23] Jang, J.: Nonlinear Instability Theory of Lane-Emden stars, *Comm. Pure Appl. Math.* **67**, 1418–1465 (2014).
- [24] Kreiss, H.: Initial boundary value problems for hyperbolic systems, *Comm. Pure Appl. Math.* **23**, 277–296 (1970).
- [25] Kufner, A, Maligranda, L., Persson, L. E.: *The Hardy inequality*, Vydavatelsky Servis, Plzen, 2007. About its history and some related results.
- [26] Lannes, D.: Well-posedness of the water-waves equations, *J. Am. Math. Soc.* **18**, 605–654 (2005).
- [27] Li T., *Global Classical Solutions for Quasilinear Hyperbolic Systems*, Masson/John Wiley, 1994.
- [28] Lindblad, H.: Well-posedness for the motion of an incompressible liquid with free surface boundary, *Ann. Math.* **162**, 109–194 (2005).
- [29] Lindblad, H.: Well posedness for the motion of a compressible liquid with free surface boundary, *Comm. Math. Phys.* **260**, 319–392 (2005).
- [30] Liu, C.; Qu, P.: Global classical solution to partially dissipative quasilinear hyperbolic systems, *J. Math. Pures Appl.* **97** (2012), no. 3, 262–281.
- [31] Liu, T.: Compressible flow with damping and vacuum, *Jpn. J. Appl. Math.* **13**, 25–32 (1996).

- [32] Liu, T., Yang, T.: Compressible Euler equations with vacuum, *J. Differ. Equ.* **140**, 223–237 (1997).
- [33] Liu, T., Yang, T.: Compressible flow with vacuum and physical singularity. *Methods Appl. Anal.* **7**, 495–310 (2000).
- [34] Luo, T., Zeng, H.: Global Existence of Smooth Solutions and Convergence to Barenblatt Solutions for the Physical Vacuum Free Boundary Problem of Compressible Euler Equations with Damping, *Comm. Pure & Appl. Math.*, *forthcoming*, *Article first published online: 2 FEB 2015* — DOI: 10.1002/cpa.21562.
- [35] Luo, T., Xin, Z., Zeng, H.: Well-Posedness for the Motion of Physical Vacuum of the Three-dimensional Compressible Euler Equations with or without Self-Gravitation, *Arch. Rational Mech. Anal* **213**, 763–831 (2014).
- [36] Shatah, J., Zeng, C.: Geometry and a priori estimates for free boundary problems of the Euler equation, *Comm. Pure Appl. Math.* **61**, 698–744 (2008).
- [37] Trakhinin, Y.: Local existence for the free boundary problem for the non-relativistic and relativistic compressible Euler equations with a vacuum boundary condition, *Comm. Pure Appl. Math.* **62**, 1551–1594 (2009) .
- [38] Wu, S.: Well-posedness in Sobolev spaces of the full waterwave problem in 2-D. *Invent. Math.* **130**, 39–72 (1997).
- [39] Wu, S.: Well-posedness in Sobolev spaces of the full water wave problem in 3-D. *J. Am. Math. Soc.* **12**, 445–495 (1999).
- [40] Wu, S.: Almost global wellposedness of the 2-D full water wave problem. *Invent. Math.* **177**, 45–135 (2009).
- [41] Wu, S.: Global wellposedness of the 3-D full water wave problem. *Invent. Math.* **184**, 125–220 (2011).
- [42] Xu, C., Yang, T.: Local existence with physical vacuum boundary condition to Euler equations with damping, *J. Differ. Equ.* **210**, 217–231 (2005).
- [43] Yang, T.: Singular behavior of vacuum states for compressible fluids, *J. Comput. Appl. Math.* **190**, 211–231 (2006).
- [44] Yong W., Entropy and global existence for hyperbolic balance laws, *Archive for Rational Mechanics and Analysis* **172** (2004) 247–266.
- [45] Zeng Y.: Gas dynamics in thermal nonequilibrium and general hyperbolic systems with relaxation, *Archive for Rational Mechanics and Analysis* **150** (1999) 225C279.
- [46] Zeng Y.: Gas flows with several thermal nonequilibrium modes, *Archive for Rational Mechanics and Analysis* **196** (2010) 191C225.
- [47] Zhang, P., Zhang, Z.: On the free boundary problem of three-dimensional incompressible Euler equations, *Comm. Pure Appl. Math.* **61**, 877–940 (2008).