

FLOW ON SWEEPING NETWORKS*

PIERRE DEGOND[†], MICHAEL HERTY[‡], AND JIAN-GUO LIU[§]

Abstract. We introduce a cellular automaton model coupled with a transport equation for flows on graphs. The direction of the flow is described by a switching process where the switching probability dynamically changes according to the value of the transported quantity in the neighboring cells. A motivation is pedestrian dynamics during panic situations in a small corridor where the propagation of people in a part of the corridor can be either left- or right-going. Under the assumptions of propagation of chaos and mean-field limit, we derive a master equation and the corresponding mean-field kinetic and macroscopic models. Steady-states are computed and analyzed and exhibit the possibility of multiple metastable states and hysteresis.

Key words. cellular automata, pedestrian dynamics, networks, master equation, kinetic model, hydrodynamical model, multiple metastable states, hysteresis

AMS subject classifications. 35Q20, 82C40, 82C31, 90B15, 90B18, 90B20, 94C10, 60J20, 82C20

DOI. 10.1137/130927061

1. Introduction. We are interested in the prediction of qualitative properties and large time behavior of cellular automata (CA) as appearing, for example, in research on traffic and pedestrian flow [41, 42]. A typical CA is described by a finite set of states lying on a regular lattice and some rules on how to change those within a given time step. In contrast to existing approaches [1, 20, 41, 42], we investigate more general graph geometries, and we couple this dynamic to a deterministic flow equation for an additional quantity, for example, a density. The flow rates in the additional equation depend on the states of the CA and vice versa. Specifically, we assume that the density sweeps from one cell to one of the neighboring cells according to the state of the CA, hence the term “sweeping network.” On the other hand, the cell-states can switch from one state to another, according to a probability which depends on an average of the sweeping quantity over the neighboring cells.

Several examples of applications of such sweeping networks can be envisioned. Our first motivation is the modeling of pedestrian flows in corridors in a situation of escape panic. There, the sweeping quantity is the density of pedestrians in a cell, whereas the CA is the ensemble of the cell-states describing in which direction (left or right) pedestrians can move. In a situation of escape panic, pedestrians may not know which is the correct way to go (right or left) to escape the danger. Then, the

*Received by the editors July 1, 2013; accepted for publication (in revised form) February 10, 2014; published electronically April 17, 2014. This work was supported by KI-Net NSF RNMS grant 1107444, grants HE5386/7-1 and DAAD 54365630, and the French “Agence Nationale pour la Recherche (ANR)” in the frame of the contract “MOTIMO” (ANR-11-MONU-009-01). The second and third authors are grateful for the opportunity to stay and work at Université Toulouse Paul Sabatier in fall 2012, under sponsorship of Centre National de la Recherche Scientifique and Université Toulouse Paul Sabatier.

<http://www.siam.org/journals/mms/12-2/92706.html>

[†]CNRS, Institut de Mathématiques de Toulouse, F-31062 Toulouse, France. Current address: Department of Mathematics, Imperial College London, London SW7 2AZ, UK (pdegond@imperial.ac.uk).

[‡]Department of Mathematics, RWTH Aachen University, D-52062 Aachen, Germany (herty@igpm.rwth-aachen.de).

[§]Department of Physics and Department of Mathematics, Duke University, Durham, NC 27708 (jliu@phy.duke.edu).

choice of the direction of motion may be constrained by the flow direction of the majority. But the direction of the majority flow may change from one location to the next. Here, we will assume that the local direction of the flow (i.e., the cell-state) has a higher probability of changing if, in the neighboring location, the direction of the flow is opposite. We will see that this rule may lead to a fast synchronization of the flow direction in the entire corridor.

Another example consists of traffic or information networks whose characteristics change with load or occupation. In this case, the sweeping quantity is the load or occupation of the network: it obeys a flow equation whose flow direction is given by the state of the CA at each node. Here, the state of the CA at one node is the index of the neighboring nodes towards which the outgoing flow from the considered node is directed. Hence, the state of the CA does not belong to the set $\{-1, 1\}$, but it is still a finite set (which may differ from one node to the next). Therefore, the corresponding dynamical system is not a CA in the restricted sense but shares similar features with CA, such as the discreteness of the cell-states. For simplicity, we will still refer to it as a CA.

One more example is that of supply networks. Such networks are used to describe the flow of parts along a production line. In some instances, there may exist several suppliers or several clients, and the procedure by which the supplier or the client is chosen corresponds to the state of the network. This state may be influenced by variables attached to the nodes of the network (such as loads, delays, and financial reliability). The choice of the flow direction may in turn determine the flow of products or money along the network.

Further applications include molecular transport in cell biology or bacterial motion [21].

In this paper we present a unified approach to such coupled problems. We consider a CA coupled to a transport equation for a density attached to each cell of the CA, where the flux function in the transport is dictated by the CA cell-states locally. Further, the CA cell-states may switch randomly from one value to another according to a switching rate which depends on an average of the density over the neighboring cells. We first investigate a simple one-dimensional system, where the cells are arranged along a line and have a periodic structure. Then, the cell-states are just the variable $z_j \in \{-1, 1\}$, where j is the cell index and $z_j = +1$ (resp., $z_j = -1$) corresponds to sweeping the density towards the neighboring node to the right (resp., to the left), while the j th cell density is denoted by $\rho_j \in \mathbb{R}_+$.

From the discrete dynamics, we derive a master equation using a presentation similar to that in [13]. The master equation provides the deterministic time evolution of the joint N -cell probability distribution function (pdf) $\mathcal{F}(z_1, \dots, z_N, \rho_1, \dots, \rho_N, t)$, where N is the total number of cells of the CA. One distinctive feature of the dynamical systems investigated here lies in the coupling of a stochastic system (the dynamics of the states z_j of the CA) with that of a deterministic system (the transport equation for the cell-densities ρ_j). However, the stochasticity of the CA makes the dynamics of the cell-densities random as well. This is why the resulting master equation is posed on the high dimensional space $(z_1, \dots, z_N, \rho_1, \dots, \rho_N) \in \{-1, 1\}^N \times \mathbb{R}_+^N$ which encompasses both the cell-state random variables z_j and the cell-densities ρ_j . This master equation takes the form of a transport equation in the continuous density variables (ρ_1, \dots, ρ_N) and rate equations for the discrete CA cell-state variables (z_1, \dots, z_N) . To our knowledge, this form of a master equation has not been found elsewhere.

The master equation is posed on a huge dimensional space when N is large and leads to overwhelming numerical complexity for practical use. Additionally, it is

difficult to retrieve direct qualitative information, such as analytical solutions and asymptotic behavior of the system, from this complex equation. This is the reason why lower dimensional reductions of this equation are desirable. The goal of this paper is to derive a hierarchy of lower dimensional descriptions of the system. This requires some simplifying assumptions, which, in model cases, can be rigorously proved, but which, for the present complex problem, can only be assumed at this stage.

The first model reduction consists of averaging the N -cell pdf over $N - 1$ variables, keeping only information on the state of a single cell j by means of the 1-cell pdf $\mathbf{f}_j(z_j, \rho_j, t)$. We do not assume cell indistinguishability so that the 1-cell pdf of different cells may be different. An equation for \mathbf{f}_j is easily deduced from the master equation by integrating it over all cell variables (z_k, ρ_k) for $k = 1, \dots, N$ except $k = j$. However, this operation does not lead to a closed equation for \mathbf{f}_j unless a suitable ansatz is made for the N -cell pdf. This ansatz is the so-called propagation of chaos which assumes that the cell-states have probability distributions independent from each other, i.e.,

$$\mathcal{F}(z_1, \dots, z_N, \rho_1, \dots, \rho_N, t) \approx \prod_{j=1}^N \mathbf{f}_j(z_j, \rho_j, t).$$

The resulting equation for \mathbf{f}_j has a form similar to that of the master equation: it comprises a transport equation in ρ_j and a rate equation for the z_j -dependence. But, in contrast to the master equation, it is posed on the low dimensional space $(z_j, \rho_j) \in \{-1, 1\} \times \mathbb{R}_+$. Propagation of chaos can be proved in model cases, such as the Boltzmann equation [23, 29, 33, 34], its caricature proposed by Kač [28], and models of swarming behavior [13, 12] (see also [44]).

The second model reduction is to take the limit of an infinite number of cells, i.e., taking the cell-spacing h to zero, while looking at large time scales, of order h^{-1} . This has several consequences. The first is to legitimize the use of a mean-field formula for the switching probabilities for the cell-states. Indeed, as the cell-spacing goes to zero, more and more neighboring cells are included in the computation of the switching probability, leading, through a law of large numbers, to a mean-field evaluation. The second consequence, related to the change of time scale, is to make the dynamics in ρ -space instantaneously convergent to a deterministic dynamics; i.e., the pdf \mathbf{f} becomes a Dirac delta in ρ at its mean value $\bar{\rho}(x, t)$ which evolves at the macroscopic time scale according to a classical continuity equation. The flux in this density equation can be expressed in terms of a mean velocity, whose evolution is dictated by an ordinary differential equation derived from the mean-field equation for the switching probabilities. To our knowledge, this is the first demonstration of the fact that a sweeping dynamics on a network converges to a monokinetic distribution. This directly provides a hydrodynamic closure.

The resulting model is a deterministic system of partial differential equations from which all the stochasticity of the original model has disappeared. It bears similarities to the Euler equations of compressible fluid dynamics in that it comprises a continuity equation for the cell-density and an evolution equation for the mean velocity. However, there is an important difference in that the velocity equation is a pure ordinary differential equation expressing a relaxation towards a local velocity obtained through some nonlocal density average. The fact that there is no transport in the velocity equation originates from the fact that the direction of the flux in the sweeping process is determined purely from local quantities at the considered time. Again, we have not found a similar model elsewhere. It is likely, though, that adding

a time delay in the evaluation of the switching probabilities would restore the spatial transport in the velocity equation. This point will be investigated in future work.

These general results are then applied to a model of a pedestrian flow under panic. The steady-states of the corresponding fluid model are analyzed. According to the strength of the coupling between the density and the cell-states, we may get multiple steady-states and various kinds of phase transitions (either continuous or discontinuous) between them leading to a hysteresis phenomenon. Metastable states and hysteresis are well-documented phenomena in car traffic [6] and in pedestrian traffic [26]. This allows one to establish some qualitative properties analytically. In particular, the occurrence of phase transitions is reminiscent of similar phenomena arising in consensus formation models [18]. The model also bears analogies to the locust model of [20], but the consideration of cell-states in the present paper is original.

Finally, the presented technique is further refined to treat the case of connected nodes and flows on graphs. Under the propagation of chaos assumption, a similar equation for the 1-cell pdf is obtained. However, the large N limit is not considered because this would necessitate the passage from a discrete network to a continuous space. This limit is outside the scope of the present paper. Still, the equation for the discrete 1-cell pdf is interesting, as it couples the pdf of the neighboring nodes within the flux of the transport term in density space, a feature which we have not observed before.

CA are widely used models in car traffic [14, 36, 40, 41, 42, 43] and pedestrian traffic [9, 10, 11, 37]. Standard supply chain models are discrete event simulators [5], which bear strong analogies to CA.

Among individual-based models, i.e., models which follow each agent in the course of time, alternatives to CA are particle models such as follow-the-leader models in car traffic [22] and pedestrian traffic [30], or more complex models based on behavioral heuristics [35]. CA provide an Eulerian description of the system where volume exclusion can be easily enforced, while particle models provide a Lagrangian description. CA have been widely used in traffic because it is easier to monitor a stretch of the road or corridor than to follow the agents themselves. However, within the Mobile Millenium project,¹ data from individual car drivers (agents) also have been collected and used to model traffic flow [7]. Further, flocking field studies tend to follow the motion of the agents, using GPS devices, for instance.

Kinetic models provide a statistical (and consequently coarser) description of the ensemble of agents. They have been proposed for car traffic in [39] and for pedestrian traffic in, e.g., [27]. Finally, fluid models provide the coarsest—and consequently least computationally intensive—description of traffic systems and have been developed for car traffic in [4, 32, 38]. They have been recently adapted to pedestrian traffic in [1]. We refer the reader to [25] and [15] for reviews on traffic and pedestrian dynamics and on supply chain modeling, respectively.

The question of proving a rigorous connection between individual-based, kinetic, and fluid models has been treated in, e.g., [3, 25] for car traffic, [17, 24] for pedestrian traffic, and [2, 19] for supply chain modeling. In connection with CA of traffic, it has been investigated in particular in [14, 20]. But, to the best of our knowledge, the present paper provides the first derivation of a fluid model for a CA coupled with the deterministic evolution of a sweeping variable.

The paper is organized as follows. In section 2, we present our sweeping model in one dimension and derive its master equation. In section 3, we use the propagation

¹See <http://traffic.berkeley.edu>.

of chaos and mean-field assumptions to derive a single-particle closure of the kinetic equation and the hydrodynamic model in the limit of a large number of particles and cells. Section 4 is devoted to an application to pedestrian traffic where metastable multiple equilibria and phase transitions are exemplified. Section 5 is concerned with the extension of the model to a general graph topology. Finally, section 6 provides a conclusion and some perspectives.

2. A one-dimensional sweeping model and its master equation.

2.1. The one-dimensional sweeping model. We are interested in a one-dimensional network consisting of connected cells $j = 1, \dots, N$. Each cell contains a certain density $\rho_j \geq 0$ of a given number of people, animals, data, goods, particles, etc., which are able to move or sweep from one cell to the next. For the simplicity of the presentation, we assume a periodic domain of size equal to 1, each cell being of size $1/N$. Each cell has a state $z_j \in \{-1, 1\}$ describing the possible direction of the flow (from left to right ($z_j = 1$) or from right to left ($z_j = -1$)). For simplicity we assume that all particles in cell j move according to the state of the cell j at discrete times $t^n = n\Delta t$, with a time step Δt and for $n \in \mathbb{N}$. Hence, the flow of particles $\Psi_{j+\frac{1}{2}}$ across the cell boundary with the $(j+1)$ th cell is given by

$$(2.1) \quad \Psi_{j+\frac{1}{2}}^n = \rho_j^n \max\{z_j^n, 0\} + \rho_{j+1}^n \min\{z_{j+1}^n, 0\},$$

where the superscript n indicates that the associated quantities are evaluated at time t^n . In order to simplify the following discussion, we consider a periodic setting $\rho_{j+N}^n = \rho_j^n$.

We assume the cell j changes state according to a Poisson process with rate γ_j^n where γ_j^n depends on all the cell-states $(z_i^n)_{i=1, \dots, N}$ and cell-densities $(\rho_i^n)_{i=1, \dots, N}$. To be more precise, within a given time interval Δt the probability of changing the state of cell j is $1 - \exp(-\gamma_j^n \Delta t)$, i.e.,

$$z_j^{n+1} = z_j^n \zeta_j^n,$$

where ζ_j^n is a random variable taking values in $\{-1, 1\}$ with probability

$$P(\zeta_j^n = 1) = e^{-\gamma_j^n \Delta t} \quad \text{and} \quad P(\zeta_j^n = -1) = 1 - e^{-\gamma_j^n \Delta t}.$$

Given some initial data z_j^0 and ρ_j^0 for $j = 1, \dots, N$, the microscopic model for $n \in \mathbb{N}$ is given by

$$(2.2) \quad \rho_j^{n+1} = \rho_j^n + N\Delta t(\Psi_{j-\frac{1}{2}}^n - \Psi_{j+\frac{1}{2}}^n), \quad \rho_{j+N}^n = \rho_j^n.$$

The factor N in front highlights the fact that the densities change over one time step by an $\mathcal{O}(N\Delta t)$ quantity. This choice is consistent with the choice of the kinetic time scale for the evolution of the cell-states z^n which will be made below. We note the analogy of (2.2) with standard upwind schemes for conservation laws [31].

We note that the total number of particles is conserved:

$$\sum_{j=1}^N \rho_j^n = \sum_{j=1}^N \rho_j^0.$$

The particle density ρ_j^n is nonnegative as soon as the initial density ρ_j^0 is so, provided that the time step satisfies the Courant–Friedrichs–Lewy (CFL) stability condition

$N\Delta t \leq 1/2$. We refer the reader to [31] for stability conditions of upwind schemes for conservation laws.

Remark 2.1. Many practical networks have finite capacity. This means that the magnitude of the flux is bounded by a maximal value $\Psi^* > 0$ and that the expression (2.1) must be cut off by this maximal value when it exceeds this value. The modifications of the present theory induced by such a cut-off will be discussed in future work.

We now derive a master equation for this process using the weak formulation as in [13]. Here, the number N of cells will be kept fixed. Later on, we will make $N \rightarrow \infty$ in the resulting master equation. In a first section, we derive the master equation for the cell-states, ignoring the dependencies of the rates upon the cell-densities.

2.2. A simple CA for the cell-states and corresponding master equation. In this section, we first restrict ourselves to the case where the rates γ_j^n are independent of the cell-densities $(\rho_i^n)_{i=1,\dots,N}$. In this case, the dynamics of the cell-states is independent of the cell-densities, and the latter can be ignored in the derivation of the master equation for the former. Therefore, the random variables are only the states of the cells z_j^n at time t^n , and the framework is that of a CA. The discrete state-space at any time for N cells is therefore Σ^N with $\Sigma := \{-1, 1\}$. We denote by $\vec{z} = (z_i)_{i=1}^N$ an element of Σ^N . A measure ϕ on Σ^N is defined by the discrete duality with a test function φ as

$$\langle \phi, \varphi \rangle_{\Sigma^N} := \sum_{i=1}^N \sum_{z_i \in \{-1, 1\}} \phi(\vec{z}) \varphi(\vec{z}).$$

The model is a Markov process. We adapt the classical Markov transition operator formalism to derive the master equation (see, e.g., [13]). The pdf of \vec{z} at time t^n is denoted by $\mathcal{F}^n(\vec{z})$. Let φ be any smooth test function on Σ^N with values in \mathbb{R} , and let \mathbb{E} be the expected value of a random variable. By definition the expectation of the random variable $\varphi(\vec{z}^n)$ for all realizations of \vec{z}^n with distribution \mathcal{F}^n is therefore

$$(2.3) \quad \mathbb{E} \{ \varphi(\vec{z}^n) \} = \langle \mathcal{F}^n, \varphi \rangle_{\Sigma^N}.$$

We denote the Markov transition operator from state \vec{z}^n to \vec{z}^{n+1} by Q . By definition, it is a bounded nonnegativity preserving linear operator on $\ell^2(\Sigma^N)$ with the function $\varphi(\vec{z}) \equiv 1$ as a fixed point. The operator Q applied on the test function φ from t^n to t^{n+1} and evaluated at state \vec{z} is given by

$$(2.4) \quad Q\varphi(\vec{z}) = \mathbb{E} \{ \varphi(\vec{z}^{n+1}) \mid \vec{z}^n = \vec{z} \},$$

where the expectation is to be taken over all random processes leading from the known state \vec{z}^n to \vec{z}^{n+1} . Hence, $Q\varphi(\vec{z}^n)$ is a random variable for all realizations of \vec{z}^n with distribution \mathcal{F}^n . Therefore, its expectation is

$$\mathbb{E} \{ Q\varphi(\vec{z}^n) \} = \langle \mathcal{F}^n, Q\varphi \rangle_{\Sigma^N} = \langle Q^* \mathcal{F}^n, \varphi \rangle_{\Sigma^N},$$

where Q^* is the ℓ^2 adjoint operator to Q . Due to the property of the conditional expectation, we also have

$$\mathbb{E} \{ Q\varphi(\vec{z}^n) \} = \mathbb{E} \{ \mathbb{E} \{ \varphi(\vec{z}^{n+1}) \mid \vec{z}^n \} \} = \mathbb{E} \{ \varphi(\vec{z}^{n+1}) \} = \langle \mathcal{F}^{n+1}, \varphi \rangle_{\Sigma^N}.$$

Noting that the previous equations hold for all functions φ , we have

$$(2.5) \quad \mathcal{F}^{n+1}(\vec{z}) = Q^* \mathcal{F}^n(\vec{z}).$$

We will show that $(Q^* - \text{Id})\mathcal{F} = O(N\Delta t)$. Therefore, the rate of change of the pdf over one time step is

$$\frac{\mathcal{F}^{n+1} - \mathcal{F}^n}{N\Delta t}(\vec{z}) = \frac{1}{N\Delta t}(Q^* - \text{Id})\mathcal{F}^n(\vec{z}) = O(1).$$

In the limit $N\Delta t \rightarrow 0$, with $nN\Delta t \rightarrow t$, we have $\mathcal{F}^n(\vec{z}) \rightarrow \mathcal{F}(\vec{z}, t)$ with

$$(2.6) \quad \frac{\partial \mathcal{F}}{\partial t}(\vec{z}, t) = \lim_{N\Delta t \rightarrow 0} \frac{1}{N\Delta t}(Q^* - \text{Id})\mathcal{F}(\vec{z}, t) = L^*\mathcal{F}(\vec{z}, t).$$

This is the so-called time-continuous master equation of the process, and the operator L (the adjoint operator to L^*) is called the Markov generator. This choice of time scale is called the kinetic time scale. It corresponds to each particle colliding on average once during one time step Δt .

PROPOSITION 2.1 (formal). *The master equation for the time-continuous version of the CA described in section 2.1 when the rates γ_j are independent of the cell-densities $(\rho_i)_{i=1, \dots, N}$ is given, at the kinetic time scale, by*

$$(2.7) \quad \frac{\partial}{\partial t}\mathcal{F}(\vec{z}, t) = \frac{1}{N} \sum_{j=1}^N (\gamma_j(-z_j, \hat{z}_j)\mathcal{F}(-z_j, \hat{z}_j, t) - \gamma_j(z_j, \hat{z}_j)\mathcal{F}(z_j, \hat{z}_j, t)),$$

where $\mathcal{F}(\vec{z}, t)$ is the time-continuous joint pdf of the cell-states and where we denote $\hat{z}_j = (z_1, \dots, z_{j-1}, z_{j+1}, \dots, z_N)$ and, for a function $\phi(\vec{z})$, $\phi(z_j, \hat{z}_j) = \phi(\vec{z})$ and $\phi(-z_j, \hat{z}_j) = \phi(z_1, \dots, z_{j-1}, -z_j, z_{j+1}, \dots, z_N)$.

The operator on the right-hand side of (2.7) contains two terms. The first term is positive and describes the increase of the pdf due to cells which reach the state \vec{z} after switching from a different state (namely from the j th cell-state $-z_j$). The increase occurs at rate $\gamma_j(-z_j, \hat{z}_j)$. The second term is negative and describes the decrease of the pdf due to cells which leave the state \vec{z} for a different one (namely the j th cell-state z_j). The decrease occurs with rate $\gamma_j(z_j, \hat{z}_j)$. The resulting expression has to be summed up over all possible cells $j \in [1, N]$. The weighting factor $\frac{1}{N}$ is there to ensure that the proper time scale has been chosen to ensure the finiteness of the right-hand side in the limit $N \rightarrow \infty$. This is the so-called kinetic time scale, where, on average, a given cell changes state only a finite number of times over a finite time interval.

Proof of Proposition 2.1. Let φ be a smooth test function. We have

$$\begin{aligned} (2.8) \quad & \langle (Q^* - \text{Id})\mathcal{F}^n, \varphi \rangle_{\Sigma^N} = \mathbb{E} \left\{ \mathbb{E} \left\{ \varphi(\vec{z}^{n+1}) - \varphi(\vec{z}^n) \mid \vec{z}^n \right\} \vec{z}^n \right\} \\ & = \left\langle \mathcal{F}^n(\vec{z}), \sum_{j=1}^N (\varphi(-z_j, \hat{z}_j) - \varphi(z_j, \hat{z}_j))(1 - e^{-\gamma_j(\vec{z})\Delta t}) \prod_{i \neq j} e^{-\gamma_i(\vec{z})\Delta t} \right\rangle_{\Sigma^N} \\ & \quad + O((N\Delta t)^2) \\ & = \Delta t \sum_{j=1}^N \langle \mathcal{F}^n(z_j, \hat{z}_j), (\varphi(-z_j, \hat{z}_j) - \varphi(z_j, \hat{z}_j)) \gamma_j(z_j, \hat{z}_j) \rangle_{\Sigma^N} + O((N\Delta t)^2) \\ & = \Delta t \left\langle \sum_{j=1}^N \{ \gamma_j(-z_j, \hat{z}_j)\mathcal{F}^n(-z_j, \hat{z}_j) - \gamma_j(z_j, \hat{z}_j)\mathcal{F}^n(z_j, \hat{z}_j) \}, \varphi(\vec{z}) \right\rangle_{\Sigma^N} + O((N\Delta t)^2). \end{aligned}$$

To derive (2.8), we note that the probability that a given k -tuple of cells switches states is $O(\Delta t^k)$, but there are $O(N^k)$ possible k -tuples of cells. Hence, the total probability

that k cells change is $O((N\Delta t)^k)$. Therefore, the probability that there are strictly more than one change is $O((N\Delta t)^2)$, while that of only one change is $O(N\Delta t)$. We note that the probability of no change is dropped out by the subtraction. Then, we have

$$\frac{\mathcal{F}^{n+1}(\vec{z}) - \mathcal{F}^n(\vec{z})}{N\Delta t} = \frac{1}{N} \sum_{j=1}^N (\gamma_j(-z_j, \hat{z}_j) \mathcal{F}^n(-z_j, \hat{z}_j) - \gamma_j(z_j, \hat{z}_j) \mathcal{F}^n(z_j, \hat{z}_j)) + O(N\Delta t),$$

and, in the limit $N\Delta t \rightarrow 0$, we get (2.7). \square

In the next section, we consider the full process where the rates γ_j^n depend on the cell-densities $(\rho_i^n)_{i=1, \dots, N}$.

2.3. The master equation for the sweeping process. We now consider the full sweeping process as described in section 2. The random variables are now the states of the cells $z_i \in \{-1, 1\}$ and the number of particles within each cell $\rho_i \in \mathbb{R}_+$. The discrete state-space for N cells is therefore \mathbb{A}^N with $\mathbb{A} := \{-1, 1\} \times \mathbb{R}_+$. We still denote $\vec{z} = (z_i)_{i=1}^N$ and similarly $\vec{\rho} = (\rho_i)_{i=1}^N$. A measure ϕ on \mathbb{A}^N is defined by its action on a continuous function φ on \mathbb{A}^N by

$$\langle \phi, \varphi \rangle_{\mathbb{A}^N} = \sum_{\vec{z} \in \{-1, 1\}^N} \int_{\vec{\rho} \in \mathbb{R}_+^N} \phi(\vec{z}, \vec{\rho}) \varphi(\vec{z}, \vec{\rho}) d\rho_1 \dots d\rho_N.$$

We also denote $\gamma_i = \gamma_i(z_i, \hat{z}_i, \vec{\rho})$, $\vec{\Psi}_+ = (\Psi_{1+\frac{1}{2}}, \dots, \Psi_{i+\frac{1}{2}}, \dots, \Psi_{N+\frac{1}{2}})$, $\vec{\Psi}_- = (\Psi_{1-\frac{1}{2}}, \dots, \Psi_{i-\frac{1}{2}}, \dots, \Psi_{N-\frac{1}{2}})$. Then, the vector version of the density update is

$$(2.9) \quad \vec{\rho}^{n+1} - \vec{\rho}^n + N\Delta t (\vec{\Psi}_+^n - \vec{\Psi}_-^n) = 0.$$

PROPOSITION 2.2 (formal). *The master equation for the time-continuous version of the sweeping process described in section 2.1 when the rates γ_j depend on both the cell-states $(z_i)_{i=1, \dots, N}$ and the cell-densities $(\rho_i)_{i=1, \dots, N}$ is given, at the kinetic time-scale, by*

$$(2.10) \quad \left(\frac{\partial}{\partial t} \mathcal{F} - \nabla_{\vec{\rho}} \cdot ((\vec{\Psi}_+ - \vec{\Psi}_-) \mathcal{F}) \right) (\vec{z}, \vec{\rho}, t) = \frac{1}{N} \sum_{j=1}^N (\gamma_j(-z_j, \hat{z}_j, \vec{\rho}, t) \mathcal{F}(-z_j, \hat{z}_j, \vec{\rho}, t) - \gamma_j(z_j, \hat{z}_j, \vec{\rho}, t) \mathcal{F}(z_j, \hat{z}_j, \vec{\rho}, t))$$

in strong form or

$$(2.11) \quad \left\langle \frac{\partial \mathcal{F}}{\partial t}, \varphi \right\rangle_{\mathbb{A}^N} = - \langle \mathcal{F}, \nabla_{\vec{\rho}} \varphi \cdot (\vec{\Psi}_+ - \vec{\Psi}_-) \rangle_{\mathbb{A}^N} + \frac{1}{N} \sum_{j=1}^N \langle \mathcal{F}(z_j, \hat{z}_j, \vec{\rho}), \gamma_j(z_j, \hat{z}_j, \vec{\rho}) \{ \varphi(-z_j, \hat{z}_j, \vec{\rho}) - \varphi(z_j, \hat{z}_j, \vec{\rho}) \} \rangle_{\mathbb{A}^N}$$

for any smooth test function φ on \mathbb{A}^N with values in \mathbb{R} in weak form. We have noted $\nabla_{\vec{\rho}} \varphi \cdot \vec{g} = \sum_{j=1}^N g_j \partial_{\rho_j} \varphi$ and $\nabla_{\vec{\rho}} \cdot \vec{g} \varphi = \varphi \sum_{j=1}^N \partial_{\rho_j} g_j$ for any functions $\varphi(\vec{\rho})$ and $\vec{g}(\vec{\rho}) = (g_j(\vec{\rho}))_{j=1, \dots, N}$.

The right-hand side of (2.10) has the same structure as that of (2.7). We refer the reader to the paragraph following Proposition 2.1 for its interpretation. The time-derivative on the left-hand side is now supplemented with a first order differential term in $\vec{\rho}$ space (the second term). Due to (2.1), the coefficient $(\vec{\Psi}_+ - \vec{\Psi}_-)$ inside this derivative couples the neighboring nodes of each cell j . It expresses how the density evolves as a consequence of the density in cell j sweeping to one of its neighboring cells and the density in the neighboring cells sweeping into the j th cell. Because the stochasticity of the dynamics of the cell-states z_j is propagated to the densities ρ_j , the description of the densities is through the pdf \mathcal{F} . Therefore, the density evolution translates into a transport equation in density space for the pdf.

Proof. Let φ be any smooth test function on \mathbb{A}^N with values in \mathbb{R} . We write

$$\begin{aligned} (2.12) \quad \langle \mathcal{F}^{n+1} - \mathcal{F}^n, \varphi \rangle_{\mathbb{A}^N} &= \mathbb{E} \left\{ \mathbb{E} \left\{ \varphi(\vec{z}^{n+1}, \vec{\rho}^{n+1}) - \varphi(\vec{z}^n, \vec{\rho}^n) \mid (\vec{z}^n, \vec{\rho}^n) \right\} (\vec{z}^n, \vec{\rho}^n) \right\} \\ &= \mathbb{E} \left\{ \mathbb{E} \left\{ \varphi(\vec{z}^{n+1}, \vec{\rho}^{n+1}) - \varphi(\vec{z}^{n+1}, \vec{\rho}^n) \mid (\vec{z}^n, \vec{\rho}^n) \right\} (\vec{z}^n, \vec{\rho}^n) \right\} \\ &\quad + \mathbb{E} \left\{ \mathbb{E} \left\{ \varphi(\vec{z}^{n+1}, \vec{\rho}^n) - \varphi(\vec{z}^n, \vec{\rho}^n) \mid (\vec{z}^n, \vec{\rho}^n) \right\} (\vec{z}^n, \vec{\rho}^n) \right\} \\ &= I + II, \end{aligned}$$

together with (2.9). Then, we have

$$\begin{aligned} (2.13) \quad I &= \mathbb{E} \left\{ \mathbb{E} \left\{ \nabla_{\vec{\rho}} \varphi(\vec{z}^{n+1}, \vec{\rho}^n) (\vec{\rho}^{n+1} - \vec{\rho}^n) \mid (\vec{z}^n, \vec{\rho}^n) \right\} (\vec{z}^n, \vec{\rho}^n) \right\} + \mathcal{O}((N\Delta t)^2) \\ &= -N\Delta t \mathbb{E} \left\{ \mathbb{E} \left\{ \nabla_{\vec{\rho}} \varphi(\vec{z}^{n+1}, \vec{\rho}^n) (\vec{\Psi}_+^n - \vec{\Psi}_-^n) \mid (\vec{z}^n, \vec{\rho}^n) \right\} (\vec{z}^n, \vec{\rho}^n) \right\} + \mathcal{O}((N\Delta t)^2) \\ &= -N\Delta t \mathbb{E} \left\{ \mathbb{E} \left\{ \nabla_{\vec{\rho}} \varphi(\vec{z}^n, \vec{\rho}^n) (\vec{\Psi}_+^n - \vec{\Psi}_-^n) \mid (\vec{z}^n, \vec{\rho}^n) \right\} (\vec{z}^n, \vec{\rho}^n) \right\} + \mathcal{O}((N\Delta t)^2), \end{aligned}$$

where for the second equality, we have used (2.9). For the third, we note that the probability of no state change is given by $\prod_{j=1}^N \exp(-\gamma_j^n \Delta t) = 1 - \mathcal{O}(N\Delta t)$, and therefore, the probability of at least one change is $\mathcal{O}(N\Delta t)$. Then we can remove the inner expectation in (2.13) because there is no change involved. Using the definition of the outer expectation, we can recast (2.13) as follows:

$$(2.14) \quad I = -N\Delta t \langle \mathcal{F}^n, \nabla_{\vec{\rho}} \varphi \cdot (\vec{\Psi}_+ - \vec{\Psi}_-) \rangle_{\mathbb{A}^N} + \mathcal{O}((N\Delta t)^2)$$

$$(2.15) \quad = N\Delta t \langle \nabla_{\vec{\rho}} \cdot ((\vec{\Psi}_+ - \vec{\Psi}_-) \mathcal{F}^n), \varphi \rangle_{\mathbb{A}^N} + \mathcal{O}((N\Delta t)^2).$$

For the second term, the algebra proceeds exactly as in section 2.2. Details are omitted. As an outcome we get

$$(2.16)$$

$$\begin{aligned} II &= \Delta t \sum_{j=1}^N \left\langle \mathcal{F}^n(z_j, \hat{z}_j, \vec{\rho}), \gamma_j(z_j, \hat{z}_j, \vec{\rho}) \left\{ \varphi(-z_j, \hat{z}_j, \vec{\rho}) - \varphi(\vec{z}, \vec{\rho}) \right\} \right\rangle_{\mathbb{A}^N} \\ &\quad + \mathcal{O}((N\Delta t)^2) \end{aligned}$$

$$(2.17)$$

$$\begin{aligned} &= \Delta t \left\langle \sum_{j=1}^N \left\{ \gamma_j(-z_j, \hat{z}_j, \vec{\rho}) \mathcal{F}^n(-z_j, \hat{z}_j, \vec{\rho}) - \gamma_j(z_j, \hat{z}_j, \vec{\rho}) \mathcal{F}^n(z_j, \hat{z}_j, \vec{\rho}) \right\}, \varphi(\vec{z}, \vec{\rho}) \right\rangle_{\mathbb{A}^N} \\ &\quad + \mathcal{O}((N\Delta t)^2). \end{aligned}$$

Inserting (2.15) and (2.17) into (2.12) leads to

$$(2.18) \quad \left\langle \frac{\mathcal{F}^{n+1} - \mathcal{F}^n}{N\Delta t}, \varphi \right\rangle_{\mathbb{A}^N} = -\langle \mathcal{F}^n, \nabla_{\vec{\rho}} \varphi \cdot (\vec{\Psi}_+ - \vec{\Psi}_-) \rangle_{\mathbb{A}^N} + \frac{1}{N} \sum_{j=1}^N \langle \mathcal{F}^n(z_j, \hat{z}_j, \vec{\rho}), \gamma_j(z_j, \hat{z}_j, \vec{\rho}) \{ \varphi(-z_j, \hat{z}_j, \vec{\rho}) - \varphi(\vec{z}, \vec{\rho}) \} \rangle_{\mathbb{A}^N} + \mathcal{O}(N\Delta t).$$

Now, letting $N\Delta t \rightarrow 0$ in (2.18) and $nN\Delta t \rightarrow t$, we find the weak form (2.11) of the master equation. Then, since the test function φ is arbitrary, using (2.14) and (2.16), we get the strong form (2.10) of the master equation. \square

3. Single-particle closure and macroscopic model.

3.1. Goal. The description of the system by means of the N -cell pdf is too complicated and cannot be practically used, neither numerically nor analytically. The goal of this section is to propose a reduction of the system to a 1-cell pdf (i.e., the 1-cell marginal of the pdf $\mathcal{F}(\vec{z}, \vec{\rho}, t)$) and to compute its time evolution. A straightforward integration of the master equation does not lead to a closed equation for the 1-cell pdf. The goal of this section is to propose a closure of this equation by assuming that propagation of chaos holds. Then we investigate the limit of $N \rightarrow \infty$ and postulate that the rates can be approximated by mean-field approximation. In this limit, we find a system of hydrodynamic equations.

We first define the marginals of the pdf as follows.

DEFINITION 3.1. For any $j \in \{1, 2, \dots, N\}$, we define the marginal density \mathbf{f}_j on \mathbb{A} by

$$(3.1) \quad \mathbf{f}_j(z_j, \rho_j, t) = \langle \mathcal{F}(z_j, \hat{z}_j, \rho_j, \hat{\rho}_j, t), 1 \rangle_{\hat{\mathbb{A}}_j},$$

where $\langle \cdot, \cdot \rangle_{\hat{\mathbb{A}}_j}$ denotes the duality between measures and functions of the variables $(\hat{z}_j, \hat{\rho}_j)$ in \mathbb{A}^{N-1} (and \mathbb{A}^{N-1} is denoted by $\hat{\mathbb{A}}_j$ when such a duality is considered).

We note that (3.1) is equivalent to saying that for any smooth function $\varphi_j(z_j, \rho_j)$ of the single variables $(z_j, \rho_j) \in \mathbb{A}$, we have

$$\langle \mathcal{F}(z_j, \hat{z}_j, \rho_j, \hat{\rho}_j, t), \varphi_j(z_j, \rho_j) \rangle_{\mathbb{A}^N} = \langle \mathbf{f}_j(z_j, \rho_j, t), \varphi_j(z_j, \rho_j) \rangle_{\mathbb{A}}.$$

To get an equation for \mathbf{f}_j at the kinetic time scale, we use the master equation in weak form (2.11) with a test function $\varphi_j(z_j, \rho_j)$ of the single variables $(z_j, \rho_j) \in \mathbb{A}$. The resulting equation is given in section 3.2. It is not a closed equation because its coefficients depend on the full joint pdf \mathcal{F} .

In order to obtain a closed system of equations, we make the assumption of propagation of chaos. Here, in the perspective of letting $N \rightarrow \infty$, we introduce a spatial variable $x_j = j/N$ and the cell-size $h = \frac{1}{N}$. We write $\mathbf{f}_j(z, \rho, t) = \mathbf{f}^h(x_j, z, \rho, t)$, where $(z, \rho) \in \mathbb{A}$. With this notation, the assumption of propagation of chaos reads as follows.

Assumption 3.1. We assume that the joint pdf $\mathcal{F}(\vec{z}, \vec{\rho}, t)$ is written as

$$\mathcal{F}(\vec{z}, \vec{\rho}, t) = \prod_{j=1}^N \mathbf{f}^h(x_j, z_j, \rho_j, t).$$

This assumption states that the cell-states and cell-densities at different points are statistically independent. As a result, we obtain a closed kinetic equation for

the one-particle marginal $\mathbf{f}^h(x_j, z_j, \rho_j, t)$ for a fixed number of cells N in section 3.3. The next step is to make the number of cells $N \rightarrow \infty$ or, equivalently, the cell-spacing $h = \frac{1}{N} \rightarrow 0$. For this purpose, we make the following mean-field assumption for the rates.

Assumption 3.2. We assume that as $h \rightarrow 0$ (or $N \rightarrow \infty$), and for any fixed x and any subsequence $x_j = \frac{j}{N}$ such that $x_j \rightarrow x$, the limit $\bar{\gamma}^h(x_j, z, \rho, t) \rightarrow \bar{\gamma}(x, z, \rho, t)$ exists, where

$$(3.2) \quad \bar{\gamma}^h(x_j, z_j, \rho_j, t) := \left\langle \prod_{i \in \{1, \dots, N\}, i \neq j} \mathbf{f}^h(x_i, z_i, \rho_i, t), \gamma_j \right\rangle_{\hat{\mathbb{A}}_j}.$$

With these assumptions, we can first derive equations for the moments of the one-particle marginal in section 3.4 and then prove the convergence of the one-particle marginal distribution to a Dirac delta modeling a monokinetic distribution function in section 3.5. The final result is stated below.

THEOREM 3.2 (formal). *We consider the one-particle marginal distribution*

$$\tilde{\mathbf{f}}^h(x, z, \rho, t) = \mathbf{f}^h\left(x, z, \rho, \frac{t}{h}\right)$$

and let $h \rightarrow 0$. We assume that $\tilde{\mathbf{f}}^h \rightarrow \tilde{\mathbf{f}}$, where $\tilde{\mathbf{f}}$ is a measure of (x, z, ρ, t) , and that the convergence is as smooth as needed. We also assume the propagation of chaos assumption (Assumption 3.1) and the mean-field limit assumption for the rates (Assumption 3.2). Then, formally, we have

$$\tilde{\mathbf{f}}(x, \pm 1, \rho, t) = p_{\pm}(x, t) \delta(\rho - \bar{\rho}(x, t)),$$

where $\bar{\rho}(x, t)$ and $p_{\pm}(x, t)$ satisfy the following system:

$$(3.3) \quad \partial_t \bar{\rho} + \partial_x(\bar{\rho} u) = 0,$$

$$(3.4) \quad \partial_t u = \gamma_t(u_{coll} - u),$$

with

$$(3.5) \quad \gamma_t = \tilde{\gamma}_- + \tilde{\gamma}_+, \quad u_{coll} = \frac{\tilde{\gamma}_- - \tilde{\gamma}_+}{\tilde{\gamma}_- + \tilde{\gamma}_+},$$

$$\tilde{\gamma}_{\pm}(x, t) = \bar{\gamma}(x, \pm 1, \bar{\rho}(x, t), t),$$

and with $\bar{\gamma}$ given by Assumption 3.2, i.e.,

$$\bar{\gamma}^h(x_j, z, \rho, t) \rightarrow \bar{\gamma}(x, z, \rho, t) \quad \text{as } h \rightarrow 0,$$

$$\bar{\gamma}^h(x_j, z_j, \rho_j, t) := \left\langle \prod_{i \in \{1, \dots, N\}, i \neq j} p_{z_i}(x_i, t) \delta(\rho_i - \bar{\rho}(x_i, t)), \gamma_j \right\rangle_{\hat{\mathbb{A}}_j}.$$

Additionally, we have

$$(3.6) \quad p_+ = \frac{1+u}{2}, \quad p_- = \frac{1-u}{2}.$$

The time rescaling (i.e., t replaced by t/h in the 1-cell pdf) is needed to find the correct time scale over which the pdf relaxes to an equilibrium. This time scale

is called the hydrodynamic time scale, because it gives rise to the hydrodynamic model (3.3), (3.4) (see comment below). It is a longer time scale than the kinetic time scale considered so far. This is because this relaxation is very slow and requires much longer time units to be observable. This hydrodynamic rescaling is classical in kinetic theory (see, e.g., the review [16]).

Theorem 3.2 states that in the limit $h \rightarrow 0$, the 1-cell pdf $\tilde{\mathbf{f}}^h(x, z, \rho, t)$ observed at the hydrodynamic time scale converges to a deterministic pdf in the density variable ρ , i.e., a Dirac delta located at the mean density $\bar{\rho}$. Both values of the pdf for the cell-states $+1$ and -1 are proportional to the same Dirac delta, with proportionality coefficients p_{\pm} meaning that among the $\bar{\rho}(x, t) dx$ particles located in the neighborhood dx of position x at time t , a proportion $p_+(x, t)$ (resp., $p_-(x, t)$) corresponds to right-going (resp., left-going) pedestrians (with $p_+(x, t) + p_-(x, t) = 1$). Both the mean density $\bar{\rho}$ and the proportions p_{\pm} depend on (x, t) . Their evolution is described by system (3.3), (3.4). The mean velocity u is given by (3.6), which shows that it is proportional to the imbalance between the right- and left-going pedestrians $u = p_+ - p_-$.

Equation (3.3) is a classical continuity equation. It expresses that the total mass $M_{[a,b]}(t)$ contained in the interval $[a, b]$ at time t and given by $M_{[a,b]}(t) = \int_a^b \rho(x, t) dx$ evolves due to particles leaving or entering $[a, b]$ through its boundaries. Indeed, integrating (3.3) with respect to $x \in [a, b]$, we get that

$$\frac{d}{dt} M_{[a,b]}(t) = (\rho u)(a, t) - (\rho u)(b, t).$$

The quantities $(\rho u)(a, t)$ and $(\rho u)(b, t)$ are the particle fluxes through a and b , respectively. These particle fluxes (counted as positive if they are directed in the positive x direction) contribute to an increase of the mass at a and a decrease of the mass at b . Therefore, (3.3) describes a simple particle budget.

By contrast, (3.4) is a simple ordinary differential equation describing the relaxation of the local velocity $u(x, t)$ to a velocity $u_{\text{coll}}(x, t)$ expressing a collective consensus. We will refer to this velocity as the collective consensus velocity. It depends on the state of the CA in a possibly large neighborhood of x at time t . It is computed through (3.5) in terms of the switching rates of the cell corresponding to point x . More precisely, $u_{\text{coll}}(x, t)$ depends on the normalized difference between the switching rates for switching from state -1 to state $+1$ and for switching from state $+1$ to state -1 . Indeed, this difference is the phenomenon producing a nonzero collective consensus velocity. There might be multiple solutions of the equation $u = u_{\text{coll}}$. These multiple solutions are associated to collective decision making about the direction of the motion which can be, according to the state of the CA, either left-going or right-going. In general, the actual velocity u is different from the collective consensus velocity u_{coll} and (3.4) states that u relaxes to u_{coll} at rate γ_t equal to the sum of the switching rates. We will provide examples of these features in the next section. The fact that there is no spatial transport in (3.4) results from the instantaneous evaluation of the switching rates within the original CA. More sophisticated CA may result in the restoration of spatial transport in (3.4). Such dynamics will be studied in future work.

Sections 3.2–3.5 are devoted to the proof of Theorem 3.2.

3.2. Equation for the single-particle marginal distribution. We remind the reader that, in order to get an equation for \mathbf{f}_j at the kinetic time scale, we use the master equation in weak form (2.11) with a test function $\varphi_j(z_j, \rho_j)$ of the single

variables $(z_j, \rho_j) \in \mathbb{A}$. We have the following proposition, the proof of which is immediate and left to the reader.

PROPOSITION 3.3. *Define*

$$(3.7) \quad \begin{aligned} \overline{\Delta\Psi_j}(t) \mathbf{f}_j(t) &:= \langle \mathcal{F}(t), \Psi_{j+\frac{1}{2}} - \Psi_{j-\frac{1}{2}} \rangle_{\hat{\mathbb{A}}_j}, \\ \bar{\gamma}_j^h(t) \mathbf{f}_j(t) &:= \langle \mathcal{F}(t), \gamma_j \rangle_{\hat{\mathbb{A}}_j}. \end{aligned}$$

The functions $\overline{\Delta\Psi_j}(t)$ and $\bar{\gamma}_j^h(t)$ are functions of (z_j, ρ_j) only. Then the equation for the marginal \mathbf{f}_j is written in weak form as

$$\left\langle \frac{\partial \mathbf{f}_j}{\partial t}, \varphi_j \right\rangle_{\mathbb{A}} = - \langle \mathbf{f}_j, \overline{\Delta\Psi_j} \partial_{\rho_j} \varphi_j \rangle_{\mathbb{A}} + \frac{1}{N} \langle \mathbf{f}_j, \bar{\gamma}_j^h \{ \varphi_j(-z_j, \rho_j) - \varphi_j(z_j, \rho_j) \} \rangle_{\mathbb{A}}$$

and in strong form as

$$(3.8) \quad \begin{aligned} &\left(\frac{\partial}{\partial t} \mathbf{f}_j - \partial_{\rho_j} (\overline{\Delta\Psi_j} \mathbf{f}_j) \right) (x_j, z_j, \rho_j, t) \\ &= \frac{1}{N} (\bar{\gamma}_j^h(x_j, -z_j, \rho_j, t) \mathbf{f}_j(x_j, -z_j, \rho_j, t) - \bar{\gamma}_j^h(x_j, z_j, \rho_j, t) \mathbf{f}_j(x_j, z_j, \rho_j, t)). \end{aligned}$$

We introduce the following definition of moments and velocity.

DEFINITION 3.4. *The probability of having right-going (resp., left-going) particles at (x, t) is denoted by $p_+(x, t)$ (resp., $p_-(x, t)$). The average right-going (resp., left-going) particle density at (x, t) is denoted by $\bar{\rho}_+(x, t)$ (resp., $\bar{\rho}_-(x, t)$). They are defined by*

$$(3.9) \quad p_{\pm}(x, t) = \int_0^{\infty} \mathbf{f}(x, \pm 1, \rho, t) d\rho, \quad \bar{\rho}_{\pm}(x, t) = \int_0^{\infty} \mathbf{f}(x, \pm 1, \rho, t) \rho d\rho.$$

The average velocity of the particles at (x, t) is defined by

$$(3.10) \quad u(x, t) = (p_+ - p_-)(x, t).$$

We note that p_{\pm} and $\bar{\rho}_{\pm}$ are nonnegative quantities and that $p_+ + p_- = 1$. We define $\bar{\rho} = \bar{\rho}_+ + \bar{\rho}_-$, the total particle density at (x, t) .

In the following section, we use the propagation of chaos assumption to close the kinetic equation (3.8) for the one-particle marginal distribution.

3.3. Propagation of chaos assumption and closed kinetic equation for the one-particle marginal distribution. We now make the propagation of chaos assumption (Assumption 3.1). With this assumption we can simplify the expressions of the flux (3.7). We have the following.

LEMMA 3.5. *Under the chaos assumption (Assumption 3.1), the flux (3.7) is given by*

$$(3.11) \quad \overline{\Delta\Psi_j}(t) = \overline{\Delta\Psi_j}(\rho_j, t) = \rho_j - \bar{\rho}_-(x_j + h, t) - \bar{\rho}_+(x_j - h, t).$$

Proof. By direct computation from (2.1), we have

$$\Psi_{j+\frac{1}{2}} - \Psi_{j-\frac{1}{2}} = \rho_j + \rho_{j+1} \min\{z_{j+1}, 0\} - \rho_{j-1} \max\{z_{j-1}, 0\}.$$

So, now,

$$(3.12) \quad \begin{aligned} &\langle \mathcal{F}(t), \Psi_{j+\frac{1}{2}} - \Psi_{j-\frac{1}{2}} \rangle_{\hat{\mathbb{A}}_j} \\ &= \mathbf{f}(x_j, z_j, \rho_j, t) \langle \mathbf{f}(x_{j-1}, z_{j-1}, \rho_{j-1}, t) \mathbf{f}(x_{j+1}, z_{j+1}, \rho_{j+1}, t), \\ &\quad \rho_j + \rho_{j+1} \min\{z_{j+1}, 0\} - \rho_{j-1} \max\{z_{j-1}, 0\} \rangle_{\mathbb{A}_{j-1} \otimes \mathbb{A}_{j+1}}, \end{aligned}$$

where $\langle \cdot, \cdot \rangle_{\mathbb{A}_{j-1} \otimes \mathbb{A}_{j+1}}$ denotes the duality between measures and functions on \mathbb{A}^2 with respect to the variables $(z_{j-1}, \rho_{j-1}, z_{j+1}, \rho_{j+1})$. Then, using the definitions of the moments (3.9), the evaluation of the right-hand side of (3.12) leads to (3.11). \square

As in the previous section we index the one-particle marginal distribution by $h = \frac{1}{N}$ and denote it by \mathbf{f}^h , and, similarly, we denote $\overline{\Delta\Psi}^h(x_j, z, \rho, t) = \overline{\Delta\Psi}_j(\rho, t)$. With Lemma 3.5, we can get a closed equation for \mathbf{f}^h . More precisely, we have the following.

PROPOSITION 3.6. *Under the propagation of chaos assumption (Assumption 3.1), the single-particle marginal distribution function \mathbf{f}^h satisfies the closed kinetic equation*

$$(3.13) \quad \left(\frac{\partial}{\partial t} \mathbf{f}^h - \partial_\rho (\overline{\Delta\Psi}^h \mathbf{f}^h) \right) (x_j, z, \rho, t) = h (\bar{\gamma}^h(x_j, -z, \rho, t) \mathbf{f}^h(x_j, -z, \rho, t) - \bar{\gamma}^h(x_j, z, \rho, t) \mathbf{f}^h(x_j, z, \rho, t)),$$

with rates given by (3.2).

Now we make a change of time scale to the macroscopic time scale. We let $t' = ht$. The rationale for this change is that both $\overline{\Delta\Psi}^h$ and the right-hand side of (3.13) formally tend to zero as $h \rightarrow 0$. In order to recover a meaningful dynamics for the one-particle marginal, we have to observe it on a time interval of length $1/h$. Performing this change of variables in (3.13) and dropping the primes for simplicity, we are led to the following problem:

$$\left(\frac{\partial}{\partial t} \mathbf{f}^h - \partial_\rho \left(\frac{1}{h} \overline{\Delta\Psi}^h \mathbf{f}^h \right) \right) (x_j, z, \rho, t) = \bar{\gamma}^h(x_j, -z, \rho, t) \mathbf{f}^h(x_j, -z, \rho, t) - \bar{\gamma}^h(x_j, z, \rho, t) \mathbf{f}^h(x_j, z, \rho, t).$$

In the next section, we investigate the $h \rightarrow 0$ limit. A key assumption will be that the rates converge to their mean-field limit, as stated in Assumption 3.2.

3.4. Large cell-number mean-field limit and macroscopic moments.

In this section, we make the formal limit of a large number of cells $N \rightarrow \infty$ or $h \rightarrow 0$. We assume that $\mathbf{f}^h \rightarrow \mathbf{f}$, where \mathbf{f} is a measure of (x, z, ρ, t) , and that the convergence is as smooth as needed. The goal of this section is to compute the dynamics of \mathbf{f} . For this purpose, we need Assumption 3.2, which assumes that the rates converge to their mean-field limit. This assumption will be shown for some example in section 4. We first consider the equations for the total density $\bar{\rho}$ given by (3.9) and the mean velocity $u(x, t)$ given by (3.10). We have the following lemma.

LEMMA 3.7 (formal). *When $h \rightarrow 0$, we formally have $\bar{\rho}^h \rightarrow \bar{\rho}$ and $u^h \rightarrow u$, where $\bar{\rho}$ and u satisfy*

$$(3.14) \quad \partial_t \bar{\rho} + \partial_x (\bar{\rho} u) = 0,$$

$$(3.15) \quad \partial_t u = \gamma_t (u_{coll} - u),$$

with

$$\gamma_t = \gamma_-^{(0)} + \gamma_+^{(0)}, \quad u_{coll} = \frac{\gamma_-^{(0)} - \gamma_+^{(0)}}{\gamma_-^{(0)} + \gamma_+^{(0)}}$$

defining

$$(3.16) \quad \gamma_\pm^{(k)}(x, t) = \frac{\int_0^\infty \bar{\gamma}(x, \pm 1, \rho, t) \mathbf{f}(x, \pm 1, \rho, t) \rho^k d\rho}{\int_0^\infty \mathbf{f}(x, \pm 1, \rho, t) \rho^k d\rho}.$$

We note that the denominator of the expression (3.16) with $k = 0$ of $\gamma_{\pm}^{(0)}$ is p_{\pm} and that

$$(3.17) \quad p_+ = \frac{1+u}{2}, \quad p_- = \frac{1-u}{2}.$$

Remark 3.1. The model (3.14), (3.16) possesses analogies to the locust model (4.27)–(4.29) of [20], but with some differences. The major one is that our equation (3.15) does not involve any spatial transport, while the corresponding equation (4.28) of [20] involves such a transport. The reason for this difference is that in the model of [20], the state of motion is carried by the agents themselves and is transported along with their motion, while in the present model, the states of motion are carried by the underlying network and are immobile.

Proof. By Taylor expansion and since $\bar{\rho} = \bar{\rho}_+ + \bar{\rho}_-$, we have

$$\frac{1}{h} \overline{\Delta \Psi_j^h}(\rho_j, t) = \frac{1}{h} (\rho_j - \bar{\rho}(x_j, t)) + \partial_x (\bar{\rho}_+ - \bar{\rho}_-)(x_j, t) + o(h).$$

Inserting this expansion into (3.8) and using the mean-field assumption for rates (Assumption 3.2), we have

$$(3.18) \quad \left(\frac{\partial}{\partial t} \mathbf{f}^h - \partial_{\rho} (\partial_x (\bar{\rho}_+ - \bar{\rho}_-) \mathbf{f}^h) \right) (x, z, \rho, t) = \frac{1}{h} \partial_{\rho} \left((\rho - \bar{\rho}(x, t)) \mathbf{f}^h \right) (x, z, \rho, t) \\ + \bar{\gamma}(x, -z, \rho, t) \mathbf{f}^h(x, -z, \rho, t) - \bar{\gamma}(x, z, \rho, t) \mathbf{f}^h(x, z, \rho, t) + o(h).$$

This equation shares some features of standard kinetic equation such as the appearance of some “collision operator” (the term in factor of h) which will determine the leading order behavior of the solution as $h \rightarrow 0$. However, there is a significant difference, namely the fact that the transport operator on the left-hand side describes a transport in ρ -space, the same ρ -space on which the collision operator acts. In standard kinetic models, the collision operator and the transport operator generally operate on different variables, such as space and velocity.

Now, multiplying (3.18) by ρ and integrating with respect to $\rho \in \mathbb{R}_+$ fixing z to the values $z = +1$ and $z = -1$ successively, we get

$$\left(\frac{\partial}{\partial t} \bar{\rho}_+^h + p_+^h (\partial_x (\bar{\rho}_+ - \bar{\rho}_-)) \right) (x, t) = -\frac{1}{h} (\bar{\rho}_+^h - p_+^h \bar{\rho}) (x, t) \\ + (\gamma_-^{(1)} \bar{\rho}_-^h - \gamma_+^{(1)} \bar{\rho}_+^h) (x, t) + o(h), \\ \left(\frac{\partial}{\partial t} \bar{\rho}_-^h + p_-^h (\partial_x (\bar{\rho}_+ - \bar{\rho}_-)) \right) (x, t) = -\frac{1}{h} (\bar{\rho}_-^h - p_-^h \bar{\rho}) (x, t) \\ + (\gamma_+^{(1)} \bar{\rho}_+^h - \gamma_-^{(1)} \bar{\rho}_-^h) (x, t) + o(h).$$

Adding and subtracting these two equations, we get

$$(3.19) \quad \left(\frac{\partial}{\partial t} \bar{\rho}^h + \partial_x (\bar{\rho}_+ - \bar{\rho}_-) \right) (x, t) = o(h),$$

$$(3.20) \quad \left(\frac{\partial}{\partial t} (\bar{\rho}_+^h - \bar{\rho}_-^h) + (p_+^h - p_-^h) (\partial_x (\bar{\rho}_+ - \bar{\rho}_-)) \right) (x, t) \\ = -\frac{1}{h} (\bar{\rho}_+^h - \bar{\rho}_-^h - (p_+^h - p_-^h) \bar{\rho}) (x, t) + 2(\gamma_-^{(1)} \bar{\rho}_-^h - \bar{\gamma}_+^{(1)} \bar{\rho}_+^h) (x, t) + o(h).$$

Now, letting $h \rightarrow 0$ in (3.20) leads to

$$\bar{\rho}_+ - \bar{\rho}_- = (p_+ - p_-)\bar{\rho} = u\bar{\rho},$$

and inserting it into (3.19) leads to the conservation equation (3.14).

Now, multiplying (3.18) by 1, integrating with respect to $\rho \in \mathbb{R}_+$, fixing z to the value $z = 1$, and letting $h \rightarrow 0$, we get

$$\frac{\partial}{\partial t} p_+(x, t) = (\gamma_-^{(0)} p_- - \gamma_+^{(0)} p_+)(x, t) + o(h).$$

Now, using that $p_- = 1 - p_+$, simple algebraic manipulations lead to (3.15). Finally, equations (3.17) are obvious from what precedes. This ends the proof. \square

So far, system (3.14), (3.15) is not closed because we are lacking a simple expression of $\gamma_{\pm}^{(0)}$ in terms of $\bar{\rho}$ and u . In the next subsection, we provide such a closure relation by taking the limit $h \rightarrow 0$ in the kinetic equation (3.18).

3.5. Local equilibrium closure and macroscopic model. We now consider (3.18) and let $h \rightarrow 0$ in it. We have the following.

LEMMA 3.8 (formal). *Let $\mathbf{f} = \lim_{h \rightarrow 0} \mathbf{f}^h$. Then, \mathbf{f} is written as*

$$(3.21) \quad \mathbf{f}(x, 1, \rho, t) = p_+(x, t) \delta(\rho - \bar{\rho}(x, t)), \quad \mathbf{f}(x, -1, \rho, t) = p_-(x, t) \delta(\rho - \bar{\rho}(x, t)),$$

where p_{\pm} and $\bar{\rho}$ are the moments defined in Definition 3.4. This leads to the following expression of $\gamma_{\pm}^{(0)}$:

$$(3.22) \quad \gamma_{\pm}^{(0)}(x, t) = \frac{1}{p_{\pm}(x, t)} \bar{\gamma}(x, \pm 1, \bar{\rho}(x, t), t).$$

Proof. Taking $h \rightarrow 0$ in (3.18), we are led to the fact that \mathbf{f} satisfies

$$\partial_{\rho} \left((\rho - \bar{\rho}) \mathbf{f} \right) = 0,$$

which implies, since \mathbf{f} must be a positive measure, that

$$(3.23) \quad \mathbf{f}(x, z, \rho, t) = p(x, z, t) \delta(\rho - \bar{\rho}(x, t)),$$

with a convenient $p(x, z, t)$. Additionally, if we focus on the leading order term, we can consider the simplified problem

$$\frac{\partial}{\partial t} \mathbf{f}^h - \frac{1}{h} \partial_{\rho} \left((\rho - \bar{\rho}) \mathbf{f}^h \right) = 0.$$

This is a first order partial differential equation which can be solved by characteristics. We denote by $\rho(t)$ an arbitrary characteristic. It is obtained by solving the equation

$$\dot{\rho}(t) = -\frac{1}{h} (\rho - \bar{\rho}).$$

Its solution converges in exponential time with time scale $O(h)$ towards the fixed point $\bar{\rho}$. Therefore, \mathbf{f}^h itself converges in exponential time towards a distribution of the form (3.23). Now, by taking the moments of (3.23), we realize that the local equilibrium necessarily has the form (3.21). Inserting this expression into (3.16) (with $k = 0$) leads to (3.22). \square

This lemma completes the proof of Theorem 3.2.

4. Example: A model for pedestrian escape in corridors. Here, we are interested in pedestrian dynamics within a corridor in panic escape. We assume that the corridor is artificially decomposed into small stretches (the cells) and that within a given cell, the flow of pedestrians is either left- or right-going. Indeed, in a panic situation, pedestrians may push in both directions, but only one of the directions wins. The pedestrians moving in the “successful” direction drag the opposing pedestrians with them. The orientation of the flow in this cell is described by the variable z_j^n ($z_j^n = +1$ if the flow is right-going, and $z_j^n = -1$ if the flow is left-going). It is controlled by the flow orientation in the neighboring cells. If, in these neighboring cells, the right-going flux is larger than the left-going one, then the probability that the state of the considered cell will be given by $z = +1$ increases; i.e., if the state is already $z = +1$, it will have a larger probability of staying at this value, while if the state is originally $z = -1$, the probability of a state change to the value $z = +1$ increases. This corresponds to a herding behavior where pedestrians in panic, ignoring which direction will allow them to escape danger, will decide to go in the direction where the largest number of people go. The main characteristic of this dynamics is that the result is unpredictable, due to the possible outcome of phase transitions and hysteresis, as shown in the present example.

To model this rule, we assume that the rate of change for cell j at time t_n can be given by

$$(4.1) \quad \gamma_j^n := \gamma_j(z_j^n, \hat{z}_j^n, \rho^n) = \gamma_0 + b|z_j^n - \langle z \rangle_j^n|^\alpha,$$

where

$$(4.2) \quad \langle z \rangle_j^n = \frac{\frac{1}{N} \sum_{i=1}^N z_i^n w\left(\frac{i-j}{N}\right) \pi(\rho_i^n)}{\frac{1}{N} \sum_{i=1}^N w\left(\frac{i-j}{N}\right) \pi(\rho_i^n)}.$$

The coefficients γ_0 and b are supposed to be nonnegative (and might as well depend on j and n). We assume that $\alpha \geq 0$ and that the density-sensing function π is supposed to be monotone increasing with $\pi(0) = 0$. The weight $w : [0, 1] \rightarrow \mathbb{R}_+$ is a smooth function. We note that, because $z_j^n = \pm 1$, $-1 \leq \langle z \rangle_j^n \leq 1$. For simplicity, we still consider a periodic domain and consider $\frac{i-j}{N}$ in (4.2) as a real number modulo 1.

The rationale for (4.1), (4.2) is as follows. The quantity $\langle z \rangle_j^n$ describes the state of the given cell and the neighboring ones, defined by those which are in the support of the function w . This average weights the cells with a large density more strongly than it weights those with a low density thanks to the density-sensing function π . Now, the probability for a cell-state change decreases if the actual state variable z_j^n is close to the average $\langle z \rangle_j^n$, while it increases if the distance to the average $\langle z \rangle_j^n$ increases. This increase is linear if $\alpha = 1$ and superlinear if $\alpha > 1$. A superlinear increase triggers self-organization, as we will see below, while a linear increase does not. In addition to cell-state changes due to pedestrian interaction as just described, we add a certain level of fluctuations described by a constant rate of cell-state changes equal to γ_0 . Many modeling choices for the kernel w can be envisioned. For instance, a symmetric weighting function w parameterized by a sensing radius $r > 0$ of the form

$$w(x) = w_r(x) = \frac{1}{\sqrt{\pi}r} \exp\left(-\frac{x^2}{r^2}\right)$$

can be chosen. Here, r is kept fixed and $O(1)$.

In this example, we verify the mean-field assumption for the rates (Assumption 3.2), as shown in the following.

LEMMA 4.1 (formal). *In the limit $h \rightarrow 0$, we formally have $\bar{\gamma}^h(x_j, z, \rho, t) \rightarrow \bar{\gamma}(x_j, z, t)$ with*

$$(4.3) \quad \bar{\gamma}(x, z, t) = \gamma_0 + b \left| z - \frac{\sum_{\zeta=\pm 1} \int_{(y,\xi) \in [0,1] \times \mathbb{R}_+} \zeta w(y-x) \pi(\xi) \mathbf{f}(y, \zeta, \xi, t) d\xi dy}{\sum_{\zeta=\pm 1} \int_{(y,\xi) \in [0,1] \times \mathbb{R}_+} w(y-x) \pi(\xi) \mathbf{f}(y, \zeta, \xi, t) d\xi dy} \right|^\alpha.$$

In particular, $\bar{\gamma}(x, z, t)$ does not depend on ρ .

Proof. Formula (3.2) can be written as

$$\begin{aligned} &\bar{\gamma}^h(x_j, z_j, \rho_j, t) \\ &= \sum_{\hat{z}_j \in \{-1, 1\}^{N-1}} \int_{(\hat{x}_j, \hat{\rho}_j) \in ([0,1] \times \mathbb{R}_+)^{N-1}} \left(\gamma_0 + b \left| z_j - \frac{\frac{1}{N} \sum_{i=1}^N z_i w(x_i - x_j) \pi(\rho_i)}{\frac{1}{N} \sum_{i=1}^N w(x_i - x_j) \pi(\rho_i)} \right|^\alpha \right) \\ &\quad \times \prod_{k=1, k \neq j}^N \mathbf{f}^h(x_k, z_k, \rho_k, t) d\hat{x}_j d\hat{\rho}_j. \end{aligned}$$

The numerator and denominator of the fraction inside the integral are mean values of the functions $(y, \zeta, \xi) \in [0, 1] \times \{-1, 1\} \times \mathbb{R}_+ \rightarrow \zeta w(y-x) \pi(\xi)$ and $(y, \zeta, \xi) \rightarrow w(y-x) \pi(\xi)$, respectively, over $N-1$ independent and identically distributed random variables (x_i, z_i, ρ_i) drawn according to the probability distribution $\mathbf{f} = \lim_{h \rightarrow 0} \mathbf{f}^h$. Therefore, for large N , they converge to the average value of these functions, respectively, which make the numerator and denominator of (4.3). Then by formal manipulation, we deduce (4.3). The proof of this result, which requires the central limit theorem, is beyond the scope of this paper. We refer the reader to the appendix of [20], where a similar result is proved in a slightly simpler situation. \square

Within this example, Theorem 3.2 holds true with $u = p_+ - p_-$ and

$$(4.4) \quad \begin{aligned} \tilde{\gamma}_+(x, t) &= \gamma_0 + b |1 - \langle u \rangle(x, t)|^\alpha, & \tilde{\gamma}_-(x, t) &= \gamma_0 + b |-1 - \langle u \rangle(x, t)|^\alpha, \\ \langle u \rangle(x, t) &= \frac{\int_{y \in [0,1]} w(y-x) \pi(\bar{\rho}(y, t)) u(y, t) dy}{\int_{y \in [0,1]} w(y-x) \pi(\bar{\rho}(y, t)) dy}. \end{aligned}$$

In the case $\alpha = 2$, we note that $\gamma_t = 2(\gamma_0 + b(1 + \langle u \rangle^2))$ and $u_{\text{coll}} = \frac{2b\langle u \rangle}{\gamma_0 + b(1 + \langle u \rangle^2)}$. If we restrict ourselves to spatially homogeneous solutions, then $\bar{\rho}$ is uniform and constant, and $\langle u \rangle = u$ depends only on time. Furthermore, $\langle u \rangle$ is independent of π . Then, inserting this into (4.4) leads to the first order differential equation with cubic nonlinearity:

$$(4.5) \quad \partial_t u = 2bu \left[1 - \frac{\gamma_0}{b} - u^2 \right].$$

The parameter γ_0/b , which describes the ratio of the noise to consensus force, can be seen as a bifurcation parameter for this ordinary differential equation, which has a pitchfork bifurcation with critical point $\gamma_0/b = 1$. Indeed, the equilibrium solution of this equation when $t \rightarrow \infty$ is $u_\infty = 0$ or $u_\infty^2 = \frac{b-\gamma_0}{b}$. Therefore, if $\gamma_0 > b$, $u_\infty = 0$ is the only stationary equilibrium, and it can be seen that it is a stable one (the right-hand side of (4.5) has opposite sign to u). By contrast, if $\gamma_0 < b$, two other




FIG. 1. Left: Pitchfork bifurcation diagram for the stationary equilibrium solution u_∞ for $\alpha = 2$. Right: Subcritical pitchfork bifurcation diagram for the stationary equilibrium solution u_∞ for $\alpha > 6$. The arrows highlight the hysteresis loop.

stationary equilibria exist: $u_\infty = \pm \frac{\sqrt{b-\gamma_0}}{\sqrt{b}}$. Then, it is readily seen by inspection of (4.5) that the equilibrium $u_\infty = 0$ is unstable, while the two equilibria $u_\infty = \pm \frac{\sqrt{b-\gamma_0}}{\sqrt{b}}$ are stable. In this case, the stable equilibrium describes the formation of a consensus about one direction of motion. This consensus is obeyed by more people if the random state change frequency γ_0 is close to 0. This analysis shows that there exists a phase transition from disordered to ordered motion when b (which describes the consensus force) crosses γ_0 . The bifurcation diagram is shown in Figure 1 (left). The upper half of the curve (which provides the order parameter $|u_\infty|$ versus the noise level γ_0/b) can be regarded as the standard phase transition diagram. In this case, this diagram indicates a second order (or continuous) phase transition with critical exponent $1/2$.

By contrast, in the case $\alpha = 1$, we find $\bar{\gamma}_t = 2(\gamma_0 + b)$ and $u_{\text{coll}} = \frac{b(u)}{\gamma_0 + b}$. If a spatially homogeneous solution is sought, it is given by

$$(4.6) \quad \partial_t u = -2\gamma_0 u.$$

Then, the stationary equilibrium solution $u_\infty = 0$ is the only solution. There is no possibility of ordered motion. The motion stays disordered whatever the value of the consensus force b is. Therefore, an exponent strictly larger than 1 is necessary for the appearance of consensus.

In the case $\alpha \geq 1$ and for spatially homogeneous problems, a general formulation of the equation for u is available as follows:

$$\partial_t u = 2bu \left[-\frac{\gamma_0}{b} + (1-u^2) \frac{(1+u)^{\alpha-1} - (1-u)^{\alpha-1}}{2u} \right].$$

We recover (4.6) and (4.5) in the cases $\alpha = 1$ and $\alpha = 2$, respectively. For integers $\alpha = 3$ to $\alpha = 5$, the behavior is the same as for $\alpha = 2$ with the critical point becoming $(\gamma_0/b)_c = \alpha - 1$. For integers $\alpha > 6$, there is another critical point $(\gamma_0/b)_s > (\gamma_0/b)_c = \alpha - 1$, and the bifurcation diagram shows a subcritical pitchfork bifurcation as depicted in Figure 1 (right). Arrows in Figure 1 (right) indicate the existence of a hysteresis loop. As before, the upper half of this diagram provides the phase transition diagram giving the order parameter $|u_\infty|$ as a function of the noise level γ_0/b . In this case,

this diagram indicates a first order (or discontinuous) phase transition as shown by the occurrence of a jump at the value $(\gamma_0/b)_s$.

Remark 4.1. In the context of a spatially homogeneous situation as analyzed here, the present model and that of [20] coincide. However, their analysis is restricted to the case $\alpha = 2$, and they only get a second phase transition (or, equivalently, a pitchfork bifurcation diagram). We show that for $\alpha > 6$, the phase transition becomes first order (or, equivalently, a subcritical pitchfork bifurcation arises) with a hysteresis loop.

5. Networks.

5.1. Graph framework. The goal of this section is to extend the previous CA and its associated mean-field and hydrodynamic limits to more general network topologies. We consider a network as a graph $(\mathcal{J}, \mathcal{A})$, where \mathcal{A} is the set of graph edges and $\mathcal{J} = \{1, \dots, J\}$ is the set of graph nodes. We denote $J = \text{Card } \mathcal{J}$. We define the adjacency matrix $(a_{jk})_{j,k \in \mathcal{J}}$, i.e., the matrix such that $a_{jk} = a_{kj} = 1$ when node j is connected to node k and 0 otherwise. We assume that each node j is not connected to itself: $a_{jj} = 0$. We consider only undirected graphs. This is because we want to allow the particles to flow equally well in both directions.

We denote by d_j the degree of node j , i.e., $d_j = \sum_{k \in \mathcal{J}} a_{jk}$. We assume that the graph is connected; i.e., for any pair of nodes (j, k) with $k \neq j$, there exists a path within the graph which connects j and k . For each node j , we define the set \mathcal{N}_j of nodes connected to it, i.e.,

$$\mathcal{N}_j = \{k \in \mathcal{J} \mid a_{jk} = 1\},$$

with $\text{Card } \mathcal{N}_j = d_j$.

Each node $j \in \mathcal{J}$ contains the density $\rho_j \geq 0$ of the sweeping quantity. Indeed, this quantity is able to sweep from node j to any other (directly) connected node k (i.e., such that $a_{jk} = 1$). We assume that the whole quantity ρ_j sweeps entirely to one of the neighboring nodes. Of course, a more complex model can be envisioned, but we wish to keep the setting as simple as possible for this presentation. We denote by Ψ_{jk}^n the outgoing flux from j to k at time t^n . Each node carries the index of the neighboring node to which it sweeps $z_j \in \mathcal{N}_j$. Then, the outgoing flow (counted algebraically) from j to k , denoted by Ψ_{jk}^n , is given by

$$(5.1) \quad \Psi_{jk}^n = \rho_j^n \delta_{z_j k} - \rho_k^n \delta_{z_k j},$$

where δ_{ij} is the Kronecker index: $\delta_{ij} = 1$ if $i = j$ and 0 otherwise. The convention is that the flux between j and k is positive when it is outgoing from j and negative when it is incoming. With this convention, we have $\Psi_{kj}^n = -\Psi_{jk}^n$.

Now, node j changes state according to a Poisson process with rate γ_j^n depending on the states and densities of the nodes in the vicinity of j . Within a given time interval Δt the probability of changing the state of node j is $1 - \exp(-\gamma_j^n \Delta t)$, and the change from state z_j to any other state $z'_j \in \mathcal{N}_j \setminus \{z_j\}$ occurs with uniform probability. This means that

$$z_j^{n+1} = \begin{cases} z_j^n & \text{with probability } e^{-\gamma_j^n \Delta t}, \\ z'_j \in \mathcal{N}_j \setminus \{z_j\} & \text{uniformly in } \mathcal{N}_j \setminus \{z_j\} \text{ each with probability } \frac{1 - e^{-\gamma_j^n \Delta t}}{d_j - 1}. \end{cases}$$

Given some initial data z_j^0 and ρ_j^0 for $j \in \mathcal{J}$, the discrete time update algorithm for ρ_j^n

is given at any discrete time index $n \in \mathbb{N}$ by

$$\rho_j^{n+1} = \rho_j^n - J\Delta t \sum_{k \in \mathcal{N}_j} \Psi_{jk}^n.$$

Remark 5.1. In the one-dimensional case of section 2, the vertices of the graph are the centers of the cells.

Now, we have the following proposition, which shows that the total number of particles is conserved.

PROPOSITION 5.1. (i) *The total number of particles is conserved, i.e.,*

$$\sum_{j \in \mathcal{J}} \rho_j^n = \sum_{j \in \mathcal{J}} \rho_j^0.$$

(ii) *(Positivity preservation.) Introduce $d = \max_{j \in \mathcal{J}} d_j$, the maximal degree of the nodes. Suppose that the CFL condition $J\Delta t \leq \frac{1}{d}$ is satisfied. Then, we have*

$$\rho_j^n \geq 0 \quad \forall j \in \mathcal{J} \quad \implies \quad \rho_j^{n+1} \geq 0 \quad \forall j \in \mathcal{J}.$$

Proof. The proof of (i) follows immediately from the antisymmetry of the flux Ψ_{jk}^n . To prove (ii), we notice that when $\Psi_{jk}^n > 0$, it takes the value ρ_j^n , and when $\Psi_{jk}^n < 0$, it takes the value $-\rho_k^n$. Then, we have

$$\begin{aligned} \rho_j^{n+1} &= \rho_j^n - J\Delta t \left(\sum_{k \in \mathcal{N}_j, \Psi_{jk}^n > 0} \rho_j^n - \sum_{k \in \mathcal{N}_j, \Psi_{jk}^n < 0} \rho_k^n \right) \\ &= \rho_j^n \left(1 - J\Delta t \sum_{k \in \mathcal{N}_j, \Psi_{jk}^n > 0} 1 \right) + J\Delta t \sum_{k \in \mathcal{N}_j, \Psi_{jk}^n < 0} \rho_k^n. \end{aligned}$$

Now, since $\sum_{k \in \mathcal{N}_j, \Psi_{jk}^n > 0} 1 \leq d_j$, the first term is nonnegative under the CFL condition. The second term is nonnegative by assumption. This ends the proof. \square

5.2. A simple CA on networks. Again, as in section 2.2, we first consider the case where the rates γ_j^n are independent of the node-densities $(\rho_j^n)_{j \in \mathcal{J}}$. Then the node-densities can be ignored. The random variables consist of the node-states z_j for $j \in \mathcal{J}$, and the discrete state-space is given by $\Sigma^J = \prod_{j \in \mathcal{J}} \mathcal{N}_j$. We denote by $\vec{z} = (z_j)_{j \in \mathcal{J}}$ an element of Σ^J . A measure ϕ on Σ^J is defined as in section 2.2 by

$$(5.2) \quad \langle \phi, \varphi \rangle_{\Sigma^J} := \sum_{\vec{z} \in \Sigma^J} \phi(\vec{z}) \varphi(\vec{z}).$$

The pdf of \vec{z} at time t_n is still denoted by $\mathcal{F}^n(\vec{z})$. Let φ be a smooth test function on Σ^J with values in \mathbb{R} . As before, the expectation of the random variable $\varphi(\vec{z}^n)$ for all realizations of \vec{z}^n with distribution \mathcal{F}^n is given by (2.3) (with N replaced by J). We define the Markov transition operator Q from state \vec{z}^n to \vec{z}^{n+1} by (2.4), and we get (2.5) (again with N replaced by J). In the limit $J\Delta t \rightarrow 0$, with $nJ\Delta t \rightarrow t$, we have $\mathcal{F}^n(\vec{z}) \rightarrow \mathcal{F}(\vec{z}, t)$, where $\mathcal{F}(\vec{z}, t)$ satisfies the time-continuous master equation (2.6) associated to the adjoint operator Q^* to Q . It is written as

$$\frac{\partial \mathcal{F}}{\partial t}(\vec{z}, t) = L^* \mathcal{F}(\vec{z}, t),$$

with

$$L^* = \lim_{J\Delta t \rightarrow 0} \frac{1}{J\Delta t} (Q^* - \text{Id}) \quad \text{and} \quad Q\varphi(\vec{z}) = \mathbb{E} \{ \varphi(\vec{z}^{n+1}) \mid \vec{z}^n = \vec{z} \}.$$

We write the master equation explicitly in the next proposition.

PROPOSITION 5.2 (formal). *The master equation for the time-continuous version of the CA described above when the rates γ_j are independent of the node-densities $(\rho_j)_{j \in \mathcal{J}}$ is given by*

$$(5.3) \quad \frac{\partial}{\partial t} \mathcal{F}(\vec{z}, t) = \frac{1}{J} \sum_{j \in \mathcal{J}} \frac{1}{d_j - 1} \sum_{z'_j \in \mathcal{N}_j \setminus \{z_j\}} (\gamma_j(z'_j, \hat{z}_j) \mathcal{F}(z'_j, \hat{z}_j, t) - \gamma_j(z_j, \hat{z}_j) \mathcal{F}(z_j, \hat{z}_j, t)),$$

where we denote by \hat{z}_j the vector of length $J - 1$ collecting all node-states except that corresponding to node j , and by (z'_j, \hat{z}_j) a state vector where the state of the j th node is $z'_j \in \mathcal{N}_j \setminus \{z_j\}$ and the states of the other nodes are given by \hat{z}_j .

This equation has a form and meaning similar to that of (2.7) (except that now more than two nodes may be connected to a given node), and we refer to the paragraph following Proposition 2.1 for its interpretation.

Proof. The proof follows the same strategy as that of Proposition 2.1. Let φ be a smooth test function. Again the probability that a given k -tuple of cells switches states is $\mathcal{O}((J\Delta t)^k)$. Therefore, the probability that there are strictly more than one change is $\mathcal{O}((J\Delta t)^2)$, while that of only one change is $\mathcal{O}(J\Delta t)$. This leads to

$$\begin{aligned} \langle (Q^* - \text{Id})\mathcal{F}^n, \varphi \rangle_{\Sigma^J} &= \mathbb{E} \{ \mathbb{E} \{ \varphi(\vec{z}^{n+1}) - \varphi(\vec{z}^n) \mid \vec{z}^n \} \vec{z}^n \} \\ &= \left\langle \mathcal{F}^n(\vec{z}), \sum_{j \in \mathcal{J}} \frac{1}{d_j - 1} \sum_{z'_j \in \mathcal{N}_j \setminus \{z_j\}} (\varphi(z'_j, \hat{z}_j) - \varphi(z_j, \hat{z}_j)) \right. \\ &\quad \left. \times (1 - e^{-\gamma_j(z_j, \hat{z}_j)\Delta t}) \prod_{i \neq j} e^{-\gamma_i(z_i, \hat{z}_i)\Delta t} \right\rangle_{\Sigma^J} + \mathcal{O}((J\Delta t)^2). \end{aligned}$$

Using (5.2), Taylor expansion when $J\Delta t \ll 1$, and the fact that the restriction to $\mathcal{N}_j \setminus \{z_j\}$ in the second sum can be removed since the added term is simply zero, we get

$$\begin{aligned} \langle (Q^* - \text{Id})\mathcal{F}^n, \varphi \rangle_{\Sigma^J} &= \Delta t \sum_{z \in \Sigma^J} \sum_{j \in \mathcal{J}} \frac{1}{d_j - 1} \sum_{z'_j \in \mathcal{N}_j} (\varphi(z'_j, \hat{z}_j) - \varphi(z_j, \hat{z}_j)) \\ &\quad \times \gamma_j(z_j, \hat{z}_j) \mathcal{F}^n(z_j, \hat{z}_j) + \mathcal{O}((J\Delta t)^2). \end{aligned}$$

Now, pulling the summation over j out and decomposing the summation over \vec{z} into a summation over \hat{z}_j and a summation over z_j , we get

$$\begin{aligned} \langle (Q^* - \text{Id})\mathcal{F}^n, \varphi \rangle_{\Sigma^J} &= \Delta t \sum_{j \in \mathcal{J}} \frac{1}{d_j - 1} \sum_{\hat{z}_j \in \Sigma^J \setminus \{j\}} \sum_{z_j \in \mathcal{N}_j} \sum_{z'_j \in \mathcal{N}_j} (\varphi(z'_j, \hat{z}_j) - \varphi(z_j, \hat{z}_j)) \\ &\quad \times \gamma_j(z_j, \hat{z}_j) \mathcal{F}^n(z_j, \hat{z}_j) + \mathcal{O}((J\Delta t)^2). \end{aligned}$$

We can now exchange z_j and z'_j in the first term and obtain

$$\begin{aligned} \langle (Q^* - \text{Id})\mathcal{F}^n, \varphi \rangle_{\Sigma^J} &= \Delta t \sum_{j \in \mathcal{J}} \frac{1}{d_j - 1} \sum_{\hat{z}_j \in \Sigma^J \setminus \{j\}} \sum_{z_j \in \mathcal{N}_j} \sum_{z'_j \in \mathcal{N}_j} \varphi(z_j, \hat{z}_j) \\ &\quad \times (\gamma_j(z'_j, \hat{z}_j) \mathcal{F}^n(z'_j, \hat{z}_j) - \gamma_j(z_j, \hat{z}_j) \mathcal{F}^n(z_j, \hat{z}_j)) + \mathcal{O}((J\Delta t^2)). \end{aligned}$$

Collecting the summation over \hat{z}_j and over z_j into a summation over \vec{z} , pulling this summation out, and using again (5.2), we finally find

$$\begin{aligned} \langle (Q^* - \text{Id})\mathcal{F}^n, \varphi \rangle_{\Sigma^J} &= \Delta t \left\langle \sum_{j \in \mathcal{J}} \frac{1}{d_j - 1} \sum_{z'_j \in \mathcal{N}_j} (\gamma_j(z'_j, \hat{z}_j) \mathcal{F}^n(z'_j, \hat{z}_j) - \gamma_j(z_j, \hat{z}_j) \mathcal{F}^n(z_j, \hat{z}_j)), \varphi(\vec{z}) \right\rangle_{\Sigma^J} \\ &\quad + \mathcal{O}((J\Delta t^2)). \end{aligned}$$

This ends the proof. \square

In the next section, we consider the case where the rates γ_j^n depend on the node-densities $(\rho_j^n)_{j \in \mathcal{J}}$.

5.3. The master equation for the sweeping process on networks. We now consider the full sweeping process on the network as described above. The random variables are now the node-states $\vec{z} \in \Sigma^J$ and the node-densities $\vec{\rho} = (\rho_j)_{j \in \mathcal{J}}$ with $\rho_j \in \mathbb{R}_+$. The discrete state-space for J nodes is therefore $\mathbb{A}^J = \Sigma^J \times \mathbb{R}_+^J$. A measure ϕ on \mathbb{A}^J is defined by its action on a continuous function φ on \mathbb{A}^J by

$$\langle \phi, \varphi \rangle_{\mathbb{A}^J} = \sum_{\vec{z} \in \Sigma^J} \int_{\mathbb{R}_+^J} \phi(\vec{z}, \vec{\rho}) \varphi(\vec{z}, \vec{\rho}) d\vec{\rho}.$$

We also denote $\gamma_j = \gamma_j(\vec{z}, \vec{\rho})$ and $\vec{\Xi}^n = (\xi_j^n)_{j \in \mathcal{J}}$, with

$$\xi_j^n = \sum_{k \in \mathcal{N}_j} \Psi_{jk}^n.$$

Then, the vector version of the density update is

$$\vec{\rho}^{n+1} - \vec{\rho}^n + J\Delta t \vec{\Xi}^n = 0.$$

In the limit $J\Delta t \rightarrow 0$, we have the following proposition, whose proof is identical to that of Proposition 2.2 and is left to the reader.

PROPOSITION 5.3 (formal). *The master equation for the time-continuous version of the sweeping process described above when the rates γ_j depend on both the node-states \vec{z} and node-densities $\vec{\rho}$ is given by*

$$\begin{aligned} (5.4) \quad &\left(\frac{\partial}{\partial t} \mathcal{F} - \nabla_{\vec{\rho}} \cdot (\vec{\Xi} \mathcal{F}) \right) (\vec{z}, \vec{\rho}, t) \\ &= \frac{1}{J} \sum_{j \in \mathcal{J}} \frac{1}{d_j - 1} \sum_{z'_j \in \mathcal{N}_j \setminus \{z_j\}} (\gamma_j(z'_j, \hat{z}_j, \vec{\rho}) \mathcal{F}(z'_j, \hat{z}_j, \vec{\rho}, t) - \gamma_j(z_j, \hat{z}_j, \vec{\rho}) \mathcal{F}(z_j, \hat{z}_j, \vec{\rho}, t)) \end{aligned}$$

in strong form or

$$\begin{aligned} (5.5) \quad &\left\langle \frac{\partial \mathcal{F}}{\partial t}, \varphi \right\rangle_{\mathbb{A}^J} = -\langle \mathcal{F}, \nabla_{\vec{\rho}} \varphi \cdot \vec{\Xi} \rangle_{\mathbb{A}^J} \\ &+ \left\langle \mathcal{F}(\vec{z}), \frac{1}{J} \sum_{j \in \mathcal{J}} \frac{1}{d_j - 1} \gamma_j(z_j, \hat{z}_j, \vec{\rho}) \sum_{z'_j \in \mathcal{N}_j} (\varphi(z'_j, \hat{z}_j, \vec{\rho}) - \varphi(z_j, \hat{z}_j, \vec{\rho})) \right\rangle_{\mathbb{A}^J} \end{aligned}$$

for any smooth test function φ on \mathbb{A}^J with values in \mathbb{R} in weak form, where we recall that

$$(5.6) \quad \vec{\Xi} = (\xi_j)_{j \in \mathcal{J}}, \quad \text{with} \quad \xi_j = \sum_{k \in \mathcal{N}_j} \Psi_{jk},$$

$$(5.7) \quad \Psi_{jk} = \rho_j \delta_{z_j k} - \rho_k \delta_{z_k j}.$$

We have noted that $\nabla_{\vec{\rho}} \varphi \cdot \vec{g} = \sum_{j \in \mathcal{J}} g_j \partial_{\rho_j} \varphi$ and $\nabla_{\vec{\rho}} \cdot \vec{g} \varphi = \varphi \sum_{j \in \mathcal{J}} \partial_{\rho_j} g_j$ for any functions $\varphi(\vec{\rho})$ and $\vec{g}(\vec{\rho}) = (g_j(\vec{\rho}))_{j \in \mathcal{J}}$.

Again, the form and meaning of (5.4) are the same as those of (2.10), and we refer to the paragraph following Proposition 2.2 for its interpretation. The only remark worth being made is that now, the total flux ξ_j at node j does not take the form of a simple difference of neighboring fluxes, as in (2.10), but has the more complex expression (5.6). However, it is readily seen that this expression reduces to $\vec{\Psi}_+ - \vec{\Psi}_-$ in the one-dimensional case.

5.4. Single-particle closure for networks. The goal of this section is again to compute a closed system of equations for the one-particle marginals of $\mathcal{F}(z, \vec{\rho}, t)$. We define the marginals according to the following definition.

DEFINITION 5.4. For any $j \in \mathcal{J}$, we define the marginal density \mathbf{f}_j on \mathbb{A} by duality by

$$\langle \mathbf{f}_j(z_j, \rho_j, t), \varphi_j(z_j, \rho_j) \rangle_{\mathbb{A}^J} = \langle \mathcal{F}(z_j, \hat{z}_j, \rho_j, \hat{\rho}_j, t), \varphi_j(z_j, \rho_j) \rangle_{\mathbb{A}^J},$$

where $\hat{\rho}_j$ is a $(J - 1)$ -dimensional vector collecting all ρ_m for $m \in \mathcal{J}$, with $m \neq j$, and with any test function $\varphi_j(z_j, \rho_j)$ of the single variables (z_j, ρ_j) . Equivalently, we have

$$\mathbf{f}_j(z_j, \rho_j, t) = \langle \mathcal{F}(z_j, \hat{z}_j, \rho_j, \hat{\rho}_j, t), 1 \rangle_{\hat{\mathbb{A}}_j},$$

where $\langle \cdot, \cdot \rangle_{\hat{\mathbb{A}}_j}$ denotes the duality between measures and functions of the variables $(\hat{z}_j, \hat{\rho}_j)$ in \mathbb{A}^{J-1} (and \mathbb{A}^{J-1} is denoted by $\hat{\mathbb{A}}_j$ when such a duality is considered).

To get an equation for \mathbf{f}_j , we use the master equation in weak form (5.5) with a test function $\varphi_j(z_j, \rho_j)$ of the single variables (z_j, ρ_j) . The resulting equation is given by the following proposition.

PROPOSITION 5.5. Define

$$(5.8) \quad \begin{aligned} \bar{\xi}_j(t) \mathbf{f}_j(t) &:= \langle \mathcal{F}(t), \xi_j \rangle_{\hat{\mathbb{A}}_j}, \\ \bar{\gamma}_j(t) \mathbf{f}_j(t) &:= \langle \mathcal{F}(t), \gamma_j \rangle_{\hat{\mathbb{A}}_j}. \end{aligned}$$

The functions $\bar{\xi}_j(t)$ and $\bar{\gamma}_j(t)$ are functions of (z_j, ρ_j) only. Then, the equation for the marginal \mathbf{f}_j is written in weak form as

$$\begin{aligned} \left\langle \frac{\partial \mathbf{f}_j}{\partial t}, \varphi_j \right\rangle_{\mathbb{A}_j} &= - \langle \mathbf{f}_j, \bar{\xi}_j(t) \partial_{\rho_j} \varphi_j \rangle_{\mathbb{A}_j} \\ &\quad + \frac{1}{J} \frac{1}{d_j - 1} \left\langle \mathbf{f}_j, \bar{\gamma}_j(z_j, \rho_j, t) \sum_{z'_j \in \mathcal{N}_j} (\varphi_j(z'_j, \rho_j) - \varphi_j(z_j, \rho_j)) \right\rangle_{\mathbb{A}_j}, \end{aligned}$$

where $\mathbb{A}_j = \mathcal{N}_j \times \mathbb{R}_+$, and in strong form as

$$\begin{aligned} & \left(\frac{\partial}{\partial t} \mathbf{f}_j - \partial_{\rho_j} (\overline{\xi}_j(t) \mathbf{f}_j) \right) (z_j, \rho_j, t) \\ &= \frac{1}{J} \frac{1}{d_j - 1} \sum_{z'_j \in \mathcal{N}_j} (\bar{\gamma}_j(z'_j, \rho_j, t) \mathbf{f}_j(z'_j, \rho_j, t) - \bar{\gamma}_j(z_j, \rho_j, t) \mathbf{f}_j(z_j, \rho_j, t)). \end{aligned}$$

We now make the propagation of chaos assumption, which in the network framework reads as follows.

Assumption 5.1. We assume that the joint pdf $\mathcal{F}(\vec{z}, \vec{\rho}, t)$ is written as

$$(5.9) \quad \mathcal{F}(\vec{z}, \vec{\rho}, t) = \prod_{j \in \mathcal{J}} \mathbf{f}_j(z_j, \rho_j, t).$$

With this assumption, we can simplify the expressions of the flux (5.8). We have the following.

LEMMA 5.6. *Under the chaos assumption (Assumption 5.1), the flux (5.8) is given by*

$$(5.10) \quad \overline{\xi}_j(z_j, \rho_j, t) = \rho_j - \sum_{k \in \mathcal{N}_j} \int_0^\infty \rho_k \mathbf{f}_k(j, \rho_k, t) d\rho_k.$$

Proof. From (5.8), we have

$$\overline{\xi}_j(z_j, \rho_j, t) \mathbf{f}_j(z_j, \rho_j, t) = \left\langle \prod_{\ell \in \mathcal{J}} \mathbf{f}_\ell(z_\ell, \rho_\ell, t), \xi_j \right\rangle_{\hat{\mathbb{A}}_j}.$$

Inserting (5.6), (5.7) into this equation leads to

$$\overline{\xi}_j(z_j, \rho_j, t) = \left\langle \prod_{\ell \in \mathcal{J}, \ell \neq j} \mathbf{f}_\ell(z_\ell, \rho_\ell, t), \sum_{k \in \mathcal{N}_j} (\rho_j \delta_{z_j k} - \rho_k \delta_{z_k j}) \right\rangle_{\hat{\mathbb{A}}_j}.$$

Interchanging the summation over k and over $\hat{\mathbb{A}}_j$, we get

$$(5.11) \quad \overline{\xi}_j(z_j, \rho_j, t) = \sum_{k \in \mathcal{N}_j} \left\langle \prod_{\ell \in \mathcal{J}, \ell \neq j} \mathbf{f}_\ell(z_\ell, \rho_\ell, t), (\rho_j \delta_{z_j k} - \rho_k \delta_{z_k j}) \right\rangle_{\hat{\mathbb{A}}_j}.$$

The term $\rho_j \delta_{z_j k}$ depends only on the state of the j th node. Therefore, it can be taken out of the bracket over $\hat{\mathbb{A}}_j$. There remains $\langle \prod_{\ell \in \mathcal{J}, \ell \neq j} \mathbf{f}_\ell(z_\ell, \rho_\ell, t), 1 \rangle_{\hat{\mathbb{A}}_j}$ which is equal to 1 because each \mathbf{f}_ℓ is a probability density. Therefore, the positive term on the right-hand side of (5.11) reduces to $\rho_j \sum_{k \in \mathcal{N}_j} \delta_{z_j k}$. Since there is only one node $k \in \mathcal{N}_j$ such that the state z_j of node j is equal to k , we have $\sum_{k \in \mathcal{N}_j} \delta_{z_j k} = 1$. Finally, the production term reduces to ρ_j .

The expression of the negative term on the right-hand side of (5.11) follows from the fact that

$$\begin{aligned} & \sum_{k \in \mathcal{N}_j} \left\langle \prod_{\ell \in \mathcal{J}, \ell \neq j} \mathbf{f}_\ell(z_\ell, \rho_\ell, t), \rho_k \delta_{z_k j} \right\rangle_{\hat{\mathbb{A}}_j} \\ &= \sum_{k \in \mathcal{N}_j} \sum_{\hat{z}_j \in \Sigma^{J-1}} \int_{\hat{\rho}_j \in \mathbb{R}_+^{J-1}} \rho_k \delta_{z_k j} \prod_{\ell \neq j, \ell \in \mathcal{J}} \mathbf{f}_\ell(z_\ell, \rho_\ell, t) d\hat{\rho}_j. \end{aligned}$$

In the previous formula, only the sum over z_k and the integral over ρ_k are different from 1 because again, each \mathbf{f}_ℓ is a probability on the state-space (z_ℓ, ρ_ℓ) . Now, because of the multiplication by $\delta_{z_k, j}$, the sum over z_k has only one nonzero contribution, that corresponding to $z_k = j$. The resulting value of the sink term is therefore equal to

$$\sum_{k \in \mathcal{N}_j} \left\langle \prod_{\ell \in \mathcal{J}, \ell \neq j} \mathbf{f}_\ell(z_\ell, \rho_\ell, t), \rho_k \delta_{z_k, j} \right\rangle_{\hat{\mathbb{A}}_j} = \sum_{k \in \mathcal{N}_j} \int_{\rho_k \in \mathbb{R}_+} \rho_k \mathbf{f}_k(j, \rho_k, t) d\rho_k.$$

Collecting the calculations of the production and sink terms, we are led to (5.10), which ends the proof. \square

Now collecting the results of Proposition 5.5 and Lemma 5.6, we can state the following theorem.

THEOREM 5.7 (formal). *Under the closure assumption (5.9), the equation for the marginal \mathbf{f}_j is written in weak form as*

$$\begin{aligned} \left\langle \frac{\partial \mathbf{f}_j}{\partial t}, \varphi_j \right\rangle_{\mathbb{A}_j} &= - \left\langle \mathbf{f}_j, \left(\rho_j - \sum_{k \in \mathcal{N}_j} \int_0^\infty \rho_k \mathbf{f}_k(j, \rho_k, t) d\rho_k \right) \partial_{\rho_j} \varphi_j \right\rangle_{\mathbb{A}_j} \\ &\quad + \frac{1}{J} \frac{1}{d_j - 1} \left\langle \mathbf{f}_j, \bar{\gamma}_j(z_j, \rho_j, t) \sum_{z'_j \in \mathcal{N}_j} (\varphi_j(z'_j, \rho_j) - \varphi_j(z_j, \rho_j)) \right\rangle_{\mathbb{A}_j}, \end{aligned}$$

where $\mathbb{A}_j = \mathcal{N}_j \times \mathbb{R}_+$, and in strong form as

$$\begin{aligned} (5.12) \quad &\left(\frac{\partial}{\partial t} \mathbf{f}_j - \partial_{\rho_j} \left(\left(\rho_j - \sum_{k \in \mathcal{N}_j} \int_0^\infty \rho_k \mathbf{f}_k(j, \rho_k, t) d\rho_k \right) \mathbf{f}_j \right) \right) (z_j, \rho_j, t) \\ &= \frac{1}{J} \frac{1}{d_j - 1} \sum_{z'_j \in \mathcal{N}_j} (\bar{\gamma}_j(z'_j, \rho_j, t) \mathbf{f}_j(z'_j, \rho_j, t) - \bar{\gamma}_j(z_j, \rho_j, t) \mathbf{f}_j(z_j, \rho_j, t)). \end{aligned}$$

Here, $\bar{\gamma}_j(z_j, \rho_j, t)$ is given by

$$(5.13) \quad \bar{\gamma}_j(z_j, \rho_j, t) = \left\langle \prod_{\ell \in \mathcal{J}, \ell \neq j} \mathbf{f}_\ell(z_\ell, \rho_\ell, t), \gamma_j \right\rangle_{\hat{\mathbb{A}}_j}.$$

Equation (5.12) provides the evolution of the 1-node pdf in the phase-space consisting of the j th cell-state space \mathcal{N}_j for z_j and the density space \mathbb{R}_+ for ρ_j . It takes the form of a transport equation in the ρ_j variable (the left-hand side), with a collision term describing the rate of change of the j th cell-states z_j (the right-hand side). The collision operator has a form and meaning similar to that of the right-hand side of (2.7) or (5.3) (but for the passage from the J -node pdf to the 1-node pdf), and we refer to the paragraph following Proposition 2.1 for its interpretation. The interesting feature in (5.12) is the transport operator. Indeed, the flux term (inside the ∂_{ρ_j} derivative) in the j th cell pdf is given in terms of the average density in neighboring cells. This average density is obtained through integrating the neighboring cell pdf f_k over the density variable ρ_k . Therefore, the various pdf's are coupled together by this flux term in an integral fashion. To some extent, this coupling resembles a mean-field coupling as in Vlasov-type models. Another source of coupling of the various 1-cell pdf's is through the evaluation of the switching rates $\bar{\gamma}_j$, which depend on the pdf's of some of the neighboring cells through the mean-field evaluation (5.13).

6. Summary and perspectives. We present a derivation of macroscopic equations for the large time behavior of microscopic sweeping processes coupled to density evolutions. Within the derivation a general master equation is considered, and under a mean-field assumption kinetic equations are derived. We applied the general calculus to an example of pedestrian flow in small corridors. An extension of the ideas to flows on networks has also been presented.

In future work we discuss equations arising from a Chapman–Enskog-like expansion for the cell-width going to zero. Further, it would be interesting to analyze a Taylor expansion of the rate equation (4.3) for strongly confined kernels w . Another open problem is the combination of the mean-field assumption and kernel localized within a finite number of cells (such as, e.g., a nearest neighbor interaction) leading to possibly correlated particle distributions. Physically more sophisticated CA may be envisioned. For instance, we could introduce different particle densities for left- and right-going particles and, according to the state of the cell, move one of the populations while the other population stays immobile. Other improvements would consist of taking into account finite network capacity or, more generally, more complex rules for the computation of the switching rates. For instance, time delays could be introduced to model the finiteness of the information propagation speed. Finally, the hypotheses made here, i.e., propagation of chaos and mean-field limit, need to be validated by intensive numerical simulations.

REFERENCES

- [1] C. APPERT-ROLLAND, P. DEGOND, AND S. MOTSCH, *Two-way multi-lane traffic model for pedestrians in corridors*, Netw. Heterog. Media, 6 (2011), pp. 351–381.
- [2] D. ARMBRUSTER, P. DEGOND, AND C. RINGHOFER, *Kinetic and fluid models for supply chains supporting policy attributes*, Bull. Inst. Math. Acad. Sin. (N.S.), 2 (2007), pp. 433–460.
- [3] A. AW, A. KLAR, A. MATERNE, AND M. RASCLE, *Derivation of continuum traffic flow models from microscopic follow-the-leader models*, SIAM J. Appl. Math., 63 (2002), pp. 259–278.
- [4] A. AW AND M. RASCLE, *Resurrection of “second order” models of traffic flow*, SIAM J. Appl. Math., 60 (2000), pp. 916–938.
- [5] J. BANKS, J. CARSON II, AND B. NELSON, *Discrete Event System Simulation*, Prentice–Hall, Englewood Cliffs, NJ, 1999.
- [6] R. BARLOVIC, L. SANTEN, A. SCHADSCHNEIDER, AND M. SCHRECKENBERG, *Metastable states in cellular automata for traffic flow*, Eur. Phys. J. B, 5 (1998), pp. 793–800.
- [7] C. G. CLAUDEL AND A. M. BAYEN, *Convex formulations of data assimilation problems for a class of Hamilton–Jacobi equations*, SIAM J. Control Optim., 49 (2011), pp. 383–402.
- [8] F. BERTHELIN, P. DEGOND, M. DELITALA, AND M. RASCLE, *A model for the formation and evolution of traffic jams*, Arch. Ration. Mech. Anal., 187 (2008), pp. 185–220.
- [9] V. J. BLUE AND J. L. ADLER, *Cellular automata microsimulation for modeling bi-directional pedestrian walkways*, Transportation Res. Part B, 35 (2001), pp. 293–312.
- [10] M. BURGER, P. A. MARKOWICH, AND J.-F. PIETSCHMANN, *Continuous limit of a crowd motion and herding model: Analysis and numerical simulations*, Kinet. Relat. Models, 4 (2011), pp. 1025–1047.
- [11] C. BURSTEDDE, K. KLAUCK, A. SCHADSCHNEIDER, AND J. ZITTARTZ, *Simulation of pedestrian dynamics using a two-dimensional cellular automaton*, Phys. A, 295 (2001), pp. 507–525.
- [12] E. CARLEN, R. CHATELIN, P. DEGOND, AND B. WENNBERG, *Kinetic hierarchy and propagation of chaos in biological swarm models*, Phys. D, 260 (2013), pp. 90–111.
- [13] E. CARLEN, P. DEGOND, AND B. WENNBERG, *Kinetic limits for pair-interaction driven master equations and biological swarm models*, Math. Models Methods Appl. Sci., 23 (2013), pp. 1339–1376.
- [14] A. CHERTOCK, A. KURGANOV, A. POLIZZI, AND I. TIMOFEYEV, *Pedestrian flow models with slowdown interactions*, Math. Models Methods Appl. Sci., 24 (2014), pp. 249–275.
- [15] C. DAGANZO, *A Theory of Supply Chains*, Lecture Notes in Econom. and Math. Systems 526, Springer-Verlag, Berlin, 2003.
- [16] P. DEGOND, *Macroscopic limits of the Boltzmann equation: A review*, in Modeling and Computational Methods for Kinetic Equations, Model. Simul. Sci. Eng. Technol., P. Degond,

- L. Pareschi, and G. Russo, eds., Birkhäuser Boston, Boston, MA, 2004, pp. 3–57.
- [17] P. DEGOND, C. APPERT-ROLLAND, M. MOUSSAÏD, J. PETTRE, AND G. THERAULAZ, *A hierarchy of heuristic-based models of crowd dynamics*, J. Stat. Phys., 152 (2013), pp. 1033–1068.
- [18] P. DEGOND, A. FROUVELLE, AND J.-G. LIU, *Macroscopic limits and phase transition in a system of self-propelled particles*, J. Nonlinear Sci., 23 (2013), pp. 427–456.
- [19] P. DEGOND AND C. RINGHOFER, *Stochastic dynamics of long supply chains with random breakdowns*, SIAM J. Appl. Math., 68 (2007), pp. 59–79.
- [20] R. ERBAN AND J. HAŠKOVEC, *From individual to collective behaviour of coupled velocity jump processes: A locust example*, Kinet. Relat. Models, 5 (2012), pp. 817–842.
- [21] R. ERBAN AND H. G. OTHMER, *From individual to collective behavior in bacterial chemotaxis*, SIAM J. Appl. Math., 65 (2004), pp. 361–391.
- [22] D. C. GAZIS, R. HERMAN, AND R. W. ROTHERY, *Nonlinear follow-the-leader models of traffic flow*, Oper. Res., 9 (1961), pp. 545–567.
- [23] J. HAŠKOVEC AND C. SCHMEISER, *Convergence of a stochastic particle approximation for measure solutions of the 2D Keller-Segel system*, Comm. Partial Differential Equations, 36 (2011), pp. 940–960.
- [24] D. HELBING, *A fluid-dynamic model for the movement of pedestrians*, Complex Systems, 6 (1992), pp. 391–415.
- [25] D. HELBING, *Traffic and related self-driven many-particle systems*, Rev. Modern Phys., 73 (2001), pp. 1067–1141.
- [26] D. HELBING, P. MOLNÁR, I. J. FARKAS, AND K. BOLAY, *Self-organizing pedestrian movement*, Environ. Plann. B: Plann. Design, 28 (2001), pp. 361–383.
- [27] S. HOOGENDOORN AND P. H. L. BOVY, *Gas-kinetic modeling and simulation of pedestrian flows*, Transp. Res. Rec., 1710 (2007), pp. 28–36.
- [28] M. KAČ, *Foundations of kinetic theory*, in Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability, 1954–1955, Vol. III, University of California Press, Berkeley, Los Angeles, 1956, pp. 171–197.
- [29] O. E. LANFORD III, *On a derivation of the Boltzmann equation*, in International Conference on Dynamical Systems in Mathematical Physics (Rennes, 1975), Astérisque 40, Soc. Math. France, Paris, 1976, pp. 117–137.
- [30] S. LEMERCIER, A. JELIC, R. KULPA, J. HUA, J. FEHRENBACH, P. DEGOND, C. APPERT-ROLLAND, S. DONIKIAN, AND J. PETTRÉ, *Realistic following behaviors for crowd simulation*, Comput. Graph. Forum, 31 (2012), pp. 489–498.
- [31] R. J. LEVEQUE, *Numerical Methods for Conservation Laws*, Lectures Math. ETH Zürich, Birkhäuser Verlag, Basel, 1992.
- [32] M. J. LIGHTHILL AND G. B. WHITHAM, *On kinematic waves. II. A theory of traffic flow on long crowded roads*, Proc. Roy. Soc. Lond. Ser. A, 229 (1955), pp. 317–345.
- [33] S. MISCHLER AND C. MOUHOT, *Quantitative Uniform in Time Chaos Propagation for Boltzmann Collision Processes*, preprint, arXiv:1001.2994 [math.AP], 2010.
- [34] S. MISCHLER, C. MOUHOT, AND B. WENNBURG, *A new approach to quantitative propagation of chaos for drift, diffusion and jump processes*, Probab. Theory Related Fields, to appear.
- [35] M. MOUSSAÏD, D. HELBING, AND G. THERAULAZ, *How simple rules determine pedestrian behavior and crowd disasters*, Proc. Natl. Acad. Sci. USA, 108 (2011), pp. 6884–6888.
- [36] K. NAGEL AND M. SCHRECKENBERG, *A cellular automaton model for freeway traffic*, J. Phys. I France, 2 (1992), pp. 2221–2229.
- [37] K. NISHINARI, A. KIRCHNER, A. NAMAZI, AND A. SCHADSCHNEIDER, *Extended floor field CA model for evacuation dynamics*, IEICE Trans. Inform. Syst., E87-D (2004), pp. 726–732.
- [38] H. J. PAYNE, *Models of freeway traffic and control*, in Mathematical Models of Public Systems, Simulation Councils Proc. Ser. 1, Society for Computer Simulation, La Jolla, CA, 1971, pp. 51–61.
- [39] I. PRIGOGINE AND R. HERMAN, *Kinetic Theory of Vehicular Traffic*, Elsevier, New York, 1971.
- [40] M. RICKERT, K. NAGEL, M. SCHRECKENBERG, AND A. LATOUR, *Two lane traffic simulations using cellular automata*, Phys. A, 231 (1996), pp. 534–550.
- [41] A. SCHADSCHNEIDER AND A. SEYFRIED, *Empirical results for pedestrian dynamics and their implications for modeling*, Netw. Heterog. Media, 6 (2011), pp. 545–560.
- [42] A. SOPASAKIS, *Formal asymptotic models of vehicular traffic. Model closures*, SIAM J. Appl. Math., 63 (2003), pp. 1561–1584.
- [43] A. SOPASAKIS AND M. A. KATSOUKAKIS, *Stochastic modeling and simulation of traffic flow: Asymmetric single exclusion process with Arrhenius look-ahead dynamics*, SIAM J. Appl. Math., 66 (2006), pp. 921–944.
- [44] A.-S. SZNITMAN, *Topics in propagation of chaos*, in Ecole d’Été de Probabilités de Saint-Flour XIX, Lecture Notes in Math. 1464, Springer, Berlin, 1991, pp. 165–251.