

## ON THE CONTACT CLASS IN HEEGAARD FLOER HOMOLOGY

KO HONDA, WILLIAM H. KAZEZ & GORDANA MATIĆ

### Abstract

We present an alternate description of the Ozsváth-Szabó contact class in Heegaard Floer homology. Using our contact class, we prove that if a contact structure  $(M, \xi)$  has an adapted open book decomposition whose page  $S$  is a once-punctured torus, then the monodromy is right-veering if and only if the contact structure is tight.

### 1. Introduction

In the paper [OS5], Ozsváth and Szabó defined an invariant of a contact 3-manifold  $(M, \xi)$  which lives in the Heegaard Floer homology  $\widehat{HF}(-M)$  of the manifold  $M$  with reversed orientation. It is defined via the work of Giroux [Gi2], who showed that there is a 1-1 correspondence between isomorphism classes of open book decompositions modulo positive stabilization and isomorphism classes of contact structures on closed 3-manifolds. Ozsváth and Szabó associated an element in Heegaard Floer homology to an open book decomposition and showed that its homology class is independent of the choice of the open book compatible with the given contact structure. They also showed that this invariant  $c(\xi)$  is zero if the contact structure is overtwisted, and that it is nonzero if the contact structure is symplectically fillable. The *contact class*  $c(\xi)$  has proven to be extremely powerful at (i) proving the tightness of various contact structures and (ii) distinguishing tight contact structures, especially in the hands of Lisca-Stipsicz [LS1, LS2] and Ghiggini [Gh].

The goal of this paper is to introduce an alternate, more hands-on, description of the contact class in Heegaard Floer homology and to use it in the context of our program of relating right-veering diffeomorphisms to tight contact structures.

---

KH supported by an Alfred P. Sloan Fellowship and an NSF CAREER Award (DMS-0237386); GM supported by NSF grant DMS-0410066; WHK supported by NSF grant DMS-0406158.

Received 10/16/2006.

We briefly recall the notion of a *right-veering* diffeomorphism which was introduced in [HKM2]. Let  $S$  be a compact oriented surface with nonempty boundary (sometimes called a “bordered surface”) and  $h : S \xrightarrow{\sim} S$  be a diffeomorphism such that  $h|_{\partial S} = id$ . Given two properly embedded oriented arcs  $\alpha$  and  $\beta$  with the same initial point  $x \in \partial S$ , we say  $\alpha$  is *to the left* of  $\beta$  if the following holds: Isotop  $\alpha$  and  $\beta$ , while fixing their endpoints, so that they intersect transversely (this include the endpoints) and with the fewest possible number of intersections. We then say  $\alpha$  is *to the left* of  $\beta$  if either  $\alpha = \beta$  or the tangent vectors  $(\dot{\beta}(0), \dot{\alpha}(0))$  define the orientation on  $S$  at  $x$ . Then  $h$  is *right-veering* if for every choice of basepoint  $x \in \partial S$  and every choice of properly embedded oriented arc  $\alpha$  based at  $x$ ,  $h(\alpha)$  is to the right of  $\alpha$ .

In [HKM2] we proved that if  $(S, h)$  is an open book decomposition compatible with a tight contact structure, then  $h$  is right-veering. In [HKM3] we continued the study of the monoid  $Veer(S, \partial S)$  of right-veering diffeomorphisms and investigated its relationship with symplectic fillability in the pseudo-Anosov case. In particular we proved the following:

**Theorem 1.1.** *Let  $S$  be a bordered surface with connected boundary and  $h$  be pseudo-Anosov with fractional Dehn twist coefficient  $c$ . If  $c \geq 1$ , then the contact structure  $\xi_{(S, h)}$  supported by  $(S, h)$  is isotopic to a perturbation of a taut foliation. Hence  $(S, h)$  is (weakly) symplectically fillable and universally tight if  $c \geq 1$ .*

Hence, when a contact structure is supported by an open book with “sufficiently” right-veering monodromy, it is symplectically fillable and therefore tight as a consequence of a theorem of Eliashberg and Gromov [El]. Unfortunately, a right-veering diffeomorphism with a small amount of rotation does not always correspond to a tight contact structure. In fact, any open book can be stabilized to a right-veering one (see Goodman [Go], as well as [HKM2]). However, we might optimistically conjecture that a minimal (i.e., not destabilizable) right-veering open book defines a tight contact structure. If we specialize to the case of a once-punctured torus, then we can use our description of the contact class to prove this conjecture.

**Theorem 1.2.** *Let  $(M, \xi)$  be a contact 3-manifold which is supported by an open book decomposition  $(S, h)$ , where  $S$  is a once-punctured torus. Then  $\xi$  is tight if and only if  $h$  is right-veering.*

Very recently John Baldwin [Ba] also obtained results similar to Theorem 1.2.

The paper is organized as follows. In Section 2, we review the standard definition of  $c(\xi)$ . Then, in Section 3, we describe the class  $EH(\xi) \in \widehat{HF}(-M)$ , which arose in discussions between John Etnyre

and the first author. We also prove that the class  $EH(\xi)$  equals the Ozsváth-Szabó contact class  $c(\xi)$ , and hence  $EH(\xi)$  is a contact invariant. In Section 4, the class  $EH(\xi)$  is applied to contact structures with compatible genus one open book decompositions with connected boundary to prove Theorem 1.2.

**Acknowledgements.** The authors are grateful to John Etnyre for discussions which led to the alternate description of the contact class. We also thank the referee for helpful comments and suggestions.

### 2. Open books and Ozsváth-Szabó contact invariants

In [OS1, OS2], Ozsváth and Szabó defined invariants of closed oriented 3-manifolds  $M$  which they called *Heegaard Floer homology*. Among the several versions of Heegaard Floer homology defined by Ozsváth and Szabó, we concentrate on the simplest one, namely  $\widehat{HF}(M)$ . It is defined as the homology associated to a chain complex determined by a Heegaard decomposition of  $M$ . Consider a Heegaard decomposition  $(\Sigma, \alpha = \{\alpha_1, \dots, \alpha_g\}, \beta = \{\beta_1, \dots, \beta_g\})$  of genus  $g$ . Here  $\Sigma$  is the Heegaard surface, i.e., a closed oriented surface of genus  $g$  which splits  $M$  into two handlebodies  $H_1$  and  $H_2$ ,  $\Sigma = \partial H_1 = -\partial H_2$ ,  $\alpha_i$  are the boundaries of the compressing disks of  $H_1$ , and  $\beta_i$  are the boundaries of the compressing disks of  $H_2$ . Then consider two tori  $\mathbb{T}_\alpha = \alpha_1 \times \dots \times \alpha_g$  and  $\mathbb{T}_\beta = \beta_1 \times \dots \times \beta_g$  in  $Sym^g(\Sigma)$ . Also pick a basepoint  $z \in \Sigma - \cup_{i=1}^g \alpha_i - \cup_{i=1}^g \beta_i$ . The complex  $\widehat{CF}(M)$  is defined to be the free  $\mathbb{Z}$ -module generated by the points  $\mathbf{x} = \{x_1, \dots, x_g\}$  of  $\mathbb{T}_\alpha \cap \mathbb{T}_\beta$ . The boundary map is defined by counting points in certain 0-dimensional moduli spaces of holomorphic maps of the unit disk into  $Sym^g(\Sigma)$ . It is, very roughly, defined as follows. Denote by  $\mathcal{M}_{\mathbf{x}, \mathbf{y}}$  the 0-dimensional (after quotienting by the natural  $\mathbb{R}$ -action) moduli space of holomorphic maps  $u$  from the unit disk  $D^2 \subset \mathbb{C}$  (with complex coordinate  $\zeta$ ) to  $Sym^g(\Sigma)$  that (i) send  $1 \mapsto \mathbf{x}$ ,  $-1 \mapsto \mathbf{y}$ ,  $S^1 \cap \{\text{Im } \zeta \geq 0\}$  to  $\mathbb{T}_\alpha$  and  $S^1 \cap \{\text{Im } \zeta \leq 0\}$  to  $\mathbb{T}_\beta$ , and (ii) avoid  $\{z\} \times Sym^{g-1}(\Sigma) \subset Sym^g(\Sigma)$ . Then define

$$\partial \mathbf{x} = \sum_{\mu(\mathbf{x}, \mathbf{y})=1} \#(\mathcal{M}_{\mathbf{x}, \mathbf{y}}) \mathbf{y},$$

where  $\mu(\mathbf{x}, \mathbf{y})$  is the relative Maslov index of the pair and  $\#(\mathcal{M}_{\mathbf{x}, \mathbf{y}})$  is a signed count of points in  $\mathcal{M}_{\mathbf{x}, \mathbf{y}}$ . The homology of this complex  $\widehat{HF}(M)$  is shown to be independent of the various choices made in the definition. In particular, it is independent of the choice of a “weakly admissible” Heegaard decomposition.


Each intersection point  $\mathbf{x}$  in  $\mathbb{T}_\alpha \cap \mathbb{T}_\beta$  defines a  $\text{Spin}^c$  structure  $\mathbf{s}_\mathbf{x}$  on  $M$ . If there is a topological disk from  $\mathbf{x}$  to  $\mathbf{y}$  which satisfies (i) and (ii) in the previous paragraph, then the two  $\text{Spin}^c$  structures agree. Hence,

the complex (as well as the homology of the complex) splits according to  $\text{Spin}^c$  structures. The Heegaard Floer homology decomposes as a direct sum

$$\widehat{HF}(M) = \bigoplus_{\mathfrak{s}} \widehat{HF}(M, \mathfrak{s}).$$

Given a contact structure  $\xi$  on  $M$ , let  $(S, h)$  be a compatible open book decomposition, where  $S$  is a surface of genus  $g$  (here genus means the genus of the surface capped off with disks) and  $h$  is a diffeomorphism of  $S$  which is the identity on  $\partial S$ . The manifold  $M$  is homeomorphic to  $(S \times [0, 1]) / \sim$  and the binding  $K$  is given by  $(\partial S \times [0, 1]) / \sim$ . Here, the equivalence relation  $\sim$  is generated by  $(x, 1) \sim (h(x), 0)$  for  $x \in S$  and  $(y, t) \sim (y, t')$  for  $y \in \partial S$ ,  $t, t' \in [0, 1]$ . From the above description of  $M$  we immediately see a natural Heegaard splitting of  $M$  by setting  $H_1 = (S \times [0, \frac{1}{2}]) / \sim$  and  $H_2 = (S \times [\frac{1}{2}, 1]) / \sim$ . This gives a Heegaard decomposition of genus  $2g$  with the splitting surface  $\Sigma = S_{1/2} \cup -S_0$ , where  $S_t = S \times \{t\}$ . A set of  $2g$  properly embedded disjoint arcs  $a_1, \dots, a_{2g}$  which cut  $S$  into a disk defines a set of compressing disks  $a_i \times [0, \frac{1}{2}]$ ,  $i = 1, \dots, 2g$ , in  $H_1$  and a set of compressing disks  $a_i \times [\frac{1}{2}, 1]$ ,  $i = 1, \dots, 2g$ , in  $H_2$ . We then set  $\alpha_i = \partial(a_i \times [0, \frac{1}{2}])$  and  $\beta_i = \partial(a_i \times [\frac{1}{2}, 1])$ , for  $i = 1, \dots, 2g$ . See Figure 1.

Given the contact manifold  $(M, \xi)$ , we denote the associated  $\text{Spin}^c$  structure by  $\mathfrak{s}_\xi$ . Ozsváth and Szabó define in [OS5] an element  $c(\xi) \in \widehat{HF}(-M, \mathfrak{s}_\xi) / (\pm 1)$  by using a Heegaard splitting associated to an open book decomposition  $(S, h)$  compatible with  $\xi$ . (At the time of the writing of the paper, the  $\pm 1$  ambiguity still exists. It is possible, however, that a careful study of orientations would remove this ambiguity. The  $\pm 1$  issue does not arise in Seiberg-Witten Floer homology. To avoid writing  $\pm 1$  everywhere, we either work with  $\mathbb{Z}/2\mathbb{Z}$ -coefficients or tacitly assume that  $c(\xi)$  is well-defined up to a sign when  $\mathbb{Z}$ -coefficients are used.) The Heegaard splitting given in the previous paragraph is not quite the Heegaard splitting that Ozsváth and Szabó consider when defining  $c(\xi)$ . Instead, they use a Heegaard surface that can be viewed simultaneously as a Heegaard surface for  $M$  and for  $M_0(K)$ , the zero surgery along the binding  $K$ . (They also assume that  $\partial S$  is connected and hence  $K$  is a knot.) The contact element in  $\widehat{HF}(-M)$  can be seen on this Heegaard surface as the image of a class in  $\widehat{HF}(-M_0(K))$  (provided  $g(K) > 1$ ), or, equivalently, as the image of a class in  $\widehat{HF}K(-M, K, F, -g)$ . To construct such a splitting, take a disk  $D \subset \text{int}(S)$  which is contained in a small neighborhood of  $\partial S$ , dig  $D \times [0, \frac{1}{2}]$  out of  $H_1$ , and then attach it to  $H_2$ . This produces two new handlebodies  $H'_1$  and  $H'_2$ . On  $H'_2$  we keep the same set of  $\beta$ -curves  $\beta_1, \dots, \beta_{2g}$  as  $H_2$  and add  $\beta_0 = \partial D \times \{\frac{1}{4}\}$ . Next, let  $d$  be a short arc connecting between the two boundary components of  $S - D$ , and let  $\{b_1, \dots, b_{2g}\}$  be a set of arcs with endpoints on  $\partial D$  which are



**Figure 1.** The left-hand portion of the figure shows the decomposition into the two handlebodies  $H_1$  and  $H_2$  and a compressing disk on each corresponding to  $a_1$ . The upper right portion shows  $-\Sigma = -S_{1/2} \cup S_0$  and the boundaries of two compressing disks. We draw just the lower right portion to indicate the Heegaard decomposition and the effect of the monodromy on arcs.

“dual” to  $\{a_1, \dots, a_{2g}\}$ . (By this we mean  $a_{2i+1} \cap b_j = \emptyset$  if  $j \neq 2i$  and  $a_{2i+1} \cap b_{2i} = \{x_{2i+1}\}$ ; also  $a_{2i} \cap b_j = \emptyset$  if  $j \neq 2i+1$  and  $a_{2i} \cap b_{2i+1} = \{x_{2i}\}$ .) Then on  $H'_1$ , we let  $\alpha_0 = \partial(d \times [0, \frac{1}{2}])$  and  $\alpha_i = \partial(b_i \times [0, \frac{1}{2}])$ . Also let  $\alpha_0 \cap \beta_0 = \{x_0\}$ .

The above choices determine a special point  $\mathbf{x} = \{x_0, x_1, \dots, x_{2g}\}$  in  $\mathbb{T}_\alpha \cap \mathbb{T}_\beta \subset \text{Sym}^{2g+1}(\Sigma)$ . (Here,  $x_i$  means  $(x_i, \frac{1}{2})$ , for  $i > 0$ .) This point (after modifying the Heegaard diagram by winding in a region that does not affect  $\mathbf{x}$  to adjust for admissibility) defines the special cycle in Heegaard Floer homology. The homology class of  $\mathbf{x}$  is defined as the contact class  $c(\xi)$  by Ozsváth-Szabó. They show that  $\widehat{HF\bar{K}}(-M, K, F, -g)$ , the knot Floer homology for  $(-M, K)$  at the lowest possible filtration level


$-g$ , is isomorphic to  $\mathbb{Z}$  and is generated by  $\mathbf{x}$ . Then  $c(\xi)$  is defined to be the image of this generator in  $\widehat{HF}(-M)$ . For details, including the figures describing this decomposition and the corresponding generator of  $c(\xi)$ , see [OS5].

### 3. An alternate description of the contact element

**3.1. Definition and main theorem.** Let  $S$  be a bordered surface whose boundary is not necessarily connected. Let  $\{a_1, \dots, a_r\}$  be a collection of disjoint, properly embedded arcs of  $S$  so that  $S - \bigcup_{i=1}^r a_i$  is a single polygon. We will call such a collection a *basis for  $S$* . Observe that every arc  $a_i$  of a basis is a nonseparating arc of  $S$ . Next let  $b_i$  be an arc which is isotopic to  $a_i$  by a small isotopy so that the following hold:

- 1) The endpoints of  $a_i$  are isotoped along  $\partial S$ , in the direction given by the boundary orientation of  $S$ .
- 2)  $a_i$  and  $b_i$  intersect transversely in one point in the interior of  $S$ .
- 3) If we orient  $a_i$ , and  $b_i$  is given the induced orientation from the isotopy, then the sign of the intersection  $a_i \cap b_i$  is  $+1$ .

See Figure 2.



**Figure 2.** The arcs  $a_i$  and  $b_i$  for a once-punctured torus  $S$ .

Let  $M = M_{(S,h)}$  be the 3-manifold with open book decomposition  $(S, h)$ . Recall the Heegaard decomposition for  $M$  described in the previous section, where  $\Sigma = S_{1/2} \cup -S_0$ . We choose the compressing disks to be  $\alpha_i = \partial(a_i \times [0, \frac{1}{2}])$  and  $\beta_i = \partial(b_i \times [\frac{1}{2}, 1])$ . We will sometimes write  $\alpha_i = (a_i, a_i)$  and  $\beta_i = (b_i, h(b_i))$ , where the first entry is the arc on  $S_{1/2}$  and the second entry is the arc on  $S_0$ . Let  $x_i$  be the intersection point  $(a_i \cap b_i) \times \{\frac{1}{2}\}$  lying in  $S_{1/2} \subset \Sigma$ , and let  $z$  be the basepoint which sits on  $S_{1/2}$  and lies *outside* the thin strips of isotopy between the  $a_i$ 's and the  $b_i$ 's. Then  $(\Sigma, \beta, \alpha, z)$  gives a weakly admissible Heegaard diagram, namely every periodic domain has positive and negative

components. This is due to the fact that every periodic domain which involves  $\alpha_i$  crosses  $x_i$ , at which point the sign of the connected component of  $\Sigma - \bigcup_{i=1}^r \alpha_i - \bigcup_{i=1}^r \beta_i$  changes.

Let  $\mathcal{D}_0, \dots, \mathcal{D}_m$  be the closures of the components of  $\Sigma - \bigcup_{i=1}^r \alpha_i - \bigcup_{i=1}^r \beta_i$ , and let  $z_i, i = 0, \dots, m$ , be a point in the interior of  $\mathcal{D}_i$ . Assume additionally that  $z_0$  is the basepoint  $z$ . Given a Whitney disk  $u : D \rightarrow \text{Sym}^r(\Sigma)$ , the *domain associated to  $u$*  is the formal linear combination

$$\mathcal{D}(u) = \sum_{i=0}^m n_{z_i}(u) \mathcal{D}_i,$$

where  $n_{z_i}(u)$  is the algebraic intersection number of  $u$  with  $\{z_i\} \times \text{Sym}^{r-1}(\Sigma)$ . To a holomorphic map  $u : D \rightarrow \text{Sym}^r(\Sigma)$  there is the associated map  $\tilde{u} : \tilde{D} \rightarrow \Sigma$  of a branched cover of  $D$  to  $\Sigma$ . The domain  $\mathcal{D}(u)$  can also be interpreted as the (homotopy class of the) image of  $\tilde{u}$ . In the definition of the boundary map in the  $\widehat{HF}$  theory, we only count holomorphic disks  $u : D \rightarrow \text{Sym}^r(\Sigma)$  that miss  $\{z\} \times \text{Sym}^{r-1}(\Sigma)$ , i.e., where the coefficient of  $\mathcal{D}_0$  is zero. For such disks the intersection of the support of  $\mathcal{D}(u)$  with  $S_{1/2}$  is thus constrained to lie in the thin strips of isotopy of the  $a_i$  to  $b_i$ .

We claim that  $\mathbf{x} = \{x_1, \dots, x_r\} \in \widehat{CF}(\Sigma, \beta, \alpha, z)$  is a cycle, thanks to the fortuitous placement of the basepoint  $z$ . (We write  $\widehat{CF}(\Sigma, \beta, \alpha, z)$  instead of  $\widehat{CF}(\Sigma, \alpha, \beta, z)$  to indicate homology on  $-M$ .) Suppose  $u : D \rightarrow \text{Sym}^r(\Sigma)$  contributes nontrivially to  $\partial \mathbf{x}$ ; in particular it is a holomorphic disk from  $\mathbf{x}$  to  $\mathbf{y}$  with  $\mathbf{y} \neq \mathbf{x}$ . Since  $\mathbf{y} = (y_1, \dots, y_r) \neq \mathbf{x} = (x_1, \dots, x_r)$ , at least one  $y_i$  lies on  $S \times \{0\}$ . There is an arc connecting  $y_i$  to some component  $x_i$  of  $\mathbf{x}$  that is in the image of the boundary of the corresponding holomorphic map  $\tilde{u}$ . Because of the orientation of the holomorphic disk that starts at  $\mathbf{x}$  (recall that  $S^1 \cap \{\text{Im } \zeta \geq 0\}$  maps to  $\mathbb{T}_\beta$  and  $S^1 \cap \{\text{Im } \zeta \leq 0\}$  maps to  $\mathbb{T}_\alpha$ ), the image of  $\tilde{u}$  near  $x_i$  has to come out of  $x_i$  in the region containing  $z$  and hence the coefficient of  $\mathcal{D}_0$  is nonzero, which contradicts the assumption that it contributes to the boundary operator.

Define  $EH(S, h, \{a_1, \dots, a_r\})$  to be the homology class of the generator  $\mathbf{x}$ . The following is the main theorem of this section:

**Theorem 3.1.**  *$EH(S, h, \{a_1, \dots, a_r\})$  is an invariant of the contact structure and equals  $c(\xi_{(S,h)})$ .*


In particular,  $EH(S, h, \{a_1, \dots, a_r\})$  is independent of the choice of basis, and it will often be denoted by  $EH(S, h)$ .

In Theorem 3.1 we are not assuming that  $\partial S$  is connected.

**Examples:** To give some intuition for the class  $EH(S, h)$ , we give three examples when  $S$  is an annulus. Refer to Figure 3. The leftmost

diagram gives  $a$  and  $b$  on  $S_{1/2}$ . The subsequent diagrams give  $S_0$  for (1), (2), and (3) below (from left to right).

- (1) If  $h$  is the identity, then  $(M, \xi)$  is the standard tight contact structure on  $S^1 \times S^2$ . Since there are two holomorphic disks from  $y$  to  $x$ , it follows that  $EH(S, h) \neq 0$ . One of the holomorphic disks from  $y$  to  $x$  has been shaded in Figure 3.
- (2) If  $h$  is a positive Dehn twist about the core curve, then  $(M, \xi)$  is the standard tight contact structure on  $S^3$ . Since  $x$  is the unique intersection point on  $\Sigma = T^2$ ,  $EH(S, h) \neq 0$ .
- (3) If  $h$  is a negative Dehn twist about the core curve, then  $(M, \xi)$  is an overtwisted contact structure on  $S^3$ . We have  $\partial y_1 = \partial y_2 = x$ ; hence  $EH(S, h) = 0$ .



**Figure 3.** Examples when  $S$  is an annulus.

The following lemma echoes our result in [HKM2], which states that  $\xi_{(S,h)}$  is overtwisted if  $h$  is not right-veering.

**Lemma 3.2.** *If  $h$  is not right-veering, then  $EH(S, h) = 0$ .*

*Proof.* If  $h$  is not right-veering, then there exists an arc  $a_1$  on  $S$  so that  $h(a_1)$  is to the left of  $a_1$ . If  $a_1$  is nonseparating, then it can be completed to a basis  $\{a_1, \dots, a_r\}$ . There exists an intersection point  $y_1 \in \alpha_1 \cap \beta_1$  and a unique (up to translation) holomorphic map from the unit disk  $D \subset \mathbb{C}$  (with complex coordinate  $\zeta$ ) to  $\Sigma$ , where  $1 \mapsto y_1$ ,  $-1 \mapsto x_1$ ,  $\partial D \cap \{\text{Im } \zeta \geq 0\}$  maps to  $\beta_i$  and  $\partial D \cap \{\text{Im } \zeta \leq 0\}$  maps to  $\alpha_i$ . Since  $z$  forces any holomorphic disk  $u : D \rightarrow \text{Sym}^r(\Sigma)$  which contributes to  $\partial\{y_1, x_2, \dots, x_n\}$  to be constant near  $x_i$ ,  $i = 2, \dots, r$ , all the  $\alpha_i$  and  $\beta_i$ ,  $i = 2, \dots, r$ , are “used up”, and the only holomorphic disk that remains is the unique one from  $y_1$  to  $x_1$ . Hence  $\partial\{y_1, x_2, \dots, x_n\} = \{x_1, x_2, \dots, x_n\}$ .

If the arc  $a_1$  is separating, then let us call its initial point  $p$ . Suppose the arcs  $h(a_1)$  and  $a_1$  intersect efficiently. Then  $h(a_1)$  and  $a_1$  must intersect at some point in the interior of  $h(a_1)$ ; otherwise  $h(a_1)$  will cut

off a strictly smaller subsurface of  $S$  inside a subsurface of  $S$  cut off by  $a_1$ . Let  $q$  be the first point of intersection between  $h(a_1)$  and  $a_1$  in  $\text{int}(h(a_1))$ ,  $c$  be the subarc of  $a_1$  from  $p$  to  $q$ , and  $c'$  be the subarc of  $h(a_1)$  from  $p$  to  $q$ . Then either  $c(c')^{-1}$  is separating or it is not. If  $c(c')^{-1}$  separates a region  $S'$  to the left of  $a_1$ , then there is a nonseparating arc  $b \subset S'$  which begins and ends at  $p$ . On the other hand, if  $c(c')^{-1}$  is nonseparating, then we let  $b = c(c')^{-1}$ . In either case, since  $b$  is strictly to the left of  $a_1$  and strictly to the right of  $h(a_1)$ , it follows that  $h(b)$  is strictly to the left of  $b$ . By separating the endpoints of  $b$  a little, we obtain a nonseparating, properly embedded, oriented arc which is moved to the left under  $h$ . q.e.d.

In view of Theorem 3.1 and the fact that every overtwisted contact structure admits an open book that is not right-veering, Lemma 3.2 immediately implies that  $c(\xi) = 0$  for an overtwisted contact structure.

*Proof of Theorem 3.1.* Let us denote a positive Dehn twist about a closed curve  $\gamma$  by  $\phi_\gamma$ . Assume  $\partial S$  is connected. We first prove the theorem for a special case, namely when  $h = \phi_{\partial S}^n$  with  $n > 0$ , in Section 3.2. Next, in Section 3.3 we prove that  $EH(S, h, \{a_1, \dots, a_r\})$  only depends on the isotopy class of  $h$  (relative to the boundary), and in Section 3.4 we show that  $EH(S, h, \{a_1, \dots, a_r\})$  is independent of the choice of basis by using handleslides. Then in Section 3.5 we prove that the class  $EH(S, h)$  is mapped to the class  $EH(S, \phi_\gamma^{-1} \circ h)$  under the natural map

$$\widehat{HF}(-M_{(S,h)}) \rightarrow \widehat{HF}(-M_{(S,\phi_\gamma^{-1} \circ h)}),$$


which corresponds to a Legendrian (+1)-surgery. We then start with  $\phi_{\partial S}^n$  with  $n \gg 0$  and apply a sequence of negative Dehn twists until we reach the desired monodromy map  $h$ . In Section 3.6 we reduce the case of multiple boundary components to the case when  $\partial S$  is connected. q.e.d.

**3.2. Primordial Example.** Let  $S$  be a once-punctured torus and  $h = \phi_{\partial S}$ , i.e., a positive Dehn twist about  $\partial S$ . The same argument works if  $S$  is a genus  $g$  surface with one boundary component and  $h = \phi_{\partial S}^n$ ,  $n > 0$ .

The subarcs of  $\alpha_i$  and  $\beta_i$  that live in  $S_0$  are given in Figure 4. We change notation and the constituent points of  $\mathbf{x}$  representing  $EH(S, h)$  will be denoted  $x_0 = x'_0$  and  $y_0 = y'_0$  as in Figure 4. Although  $x_0 = x'_0$  and  $y_0 = y'_0$ , strictly speaking, live on  $S_{1/2}$ , we view them as sitting on  $\partial S_0$ . (Also, the points  $x_0$  and  $x'_0$ , as well as  $y_0$  and  $y'_0$ , are drawn as distinct points on  $\partial S_0$ , but we hope this will not cause any confusion for the reader.)


We then place the basepoint  $w$  on  $S_0$  as indicated in Figure 4. Observe that  $z$  and  $w$  together represent the binding  $K$ . The binding  $K$  is

isotopic to the dotted curve  $\gamma_0$  which consists of two subarcs  $c_1$  and  $c_2$  between  $z$  and  $w$ , where  $c_1$  intersects only the  $\alpha$ -curves and  $c_2$  intersects only the  $\beta$ -curves. Then  $(\Sigma, \beta, \alpha, z, w)$  is a doubly-pointed Heegaard diagram for the knot Floer homology of  $K$ .



**Figure 4.**  $S_0$  when  $h = \phi_{\partial S}$ , and a zoomed-in ammonite-like region.

If we stabilize this Heegaard splitting by digging a handle in  $S \times [0, \frac{1}{2}]$  which is parallel to the arc  $c_2$ , then we obtain a Heegaard surface  $\Sigma'$  on which we can see both  $-M$  and  $-M_0(K)$ . See Figure 5. Here  $-M$  is given by  $\{\beta_0\} \cup \beta$  and  $\{\alpha_0\} \cup \alpha$ , whereas  $-M_0(K)$  is given by  $\gamma = \{\gamma_0\} \cup \beta$  and  $\{\alpha_0\} \cup \alpha$ . (Here  $\gamma_0$  is viewed as a curve that passes through the handle once.) The stabilization sends  $\mathbf{x} = \{x_0, y_0\}$  to  $\mathbf{x}' = \{z_0, x_0, y_0\}$ , where  $z_0$  is the intersection of the two new compressing curves  $\alpha_0$  and  $\beta_0$ .



**Figure 5.** Part of the stabilized Heegaard surface  $-\Sigma'$ . The domain  $F$  has been shaded.

As a first step in exploiting the Ozsváth-Szabó characterization of  $c(\xi)$ , we show that the lowest filtration level is generated by  $\mathbf{x}' = \{z_0, x_0, y_0\}$  as well as the other intersection points  $\mathbf{y} = \{z_0, x, y\}$ , where  $x$  and  $y$  live inside the dotted lines of Figure 4. The filtration level is

computed by first letting  $F \subset \Sigma'$  be the domain bounded by  $\alpha_0$  and  $\gamma_0$  which does not intersect  $S_{1/2}$  (and hence lives mostly on  $S_0$ ). We additionally assume that  $F$  is oriented so that the surface  $\hat{F}$ , obtained from  $F$  by capping off  $\partial F$ , is an oriented fiber of the fibration of  $M_0(K)$ . In order to find generators  $\mathbf{y}$  which are at the lowest filtration level, we minimize  $\langle c_1(\mathbf{s}_{\mathbf{y}'}, [\hat{F}]) \rangle$ . Here  $\mathbf{y}' = \{z'_0, x, y\}$  and  $z'_0$  is the intersection point on  $\alpha_0 \cap \gamma_0$  which forms a small triangle with  $\Theta \in \beta_0 \cap \gamma_0$  and  $z_0$  as in Figure 5. (Keep in mind that since we are dealing with  $\widehat{HF}$  of  $-M$  and  $-M_0(K)$ , the Heegaard surface is  $-\Sigma'$ ; otherwise our calculations will be off by a negative sign.)

To this end, we recall the formula for the first Chern class (Section 7.1 of [OS2]; for some details, see Rasmussen [Ra]):

$$\langle c_1(\mathbf{s}_{\mathbf{y}'}, [A]) \rangle = \chi(\mathcal{P}) - 2\bar{n}_z(\mathcal{P}) + 2 \sum_{p \in \mathbf{y}'} \bar{n}_p(\mathcal{P}).$$

Here  $[A] \in H_2(M_0(K), \mathbb{Z})$ ,  $\mathbf{s}_{\mathbf{y}'}$  is a  $\text{Spin}^c$  structure corresponding to  $\mathbf{y}'$ ,  $\mathcal{P}$  is the periodic domain for  $[A]$  (where we do not require that  $\mathcal{P}$  avoid  $z$ ) and  $\chi$  is the Euler measure. Let  $\mathcal{D}$  be a component of  $(-\Sigma') - \bigcup_i \alpha_i - \bigcup_i \gamma_i$ . Then  $\bar{n}_p(\mathcal{D})$  equals (i) 1 if  $p$  is in the interior of  $\mathcal{D}$ , (ii) 0 if  $p$  does not intersect  $\mathcal{D}$ , (iii)  $\frac{1}{2}$  if  $p$  is on an edge of  $\mathcal{D}$  (but not a corner), and (iv)  $\frac{1}{4}$  if  $p$  is on a corner of  $\mathcal{D}$ . We then extend  $\bar{n}_p$  linearly to  $\mathcal{P}$ .


In the case at hand, the possible  $x$ 's and  $y$ 's are either in the interior of  $F$  or not in  $F$ , and therefore they either contribute 1 or 0. On the other hand,  $\bar{n}_z(\mathcal{P}) = -2$ ,  $\chi(\mathcal{P}) = -2g(S)$ , and  $\bar{n}_{z'_0}(\mathcal{P}) = -1$  are constant, and it follows that  $\langle c_1(\mathbf{s}_{\mathbf{y}'}, [\hat{F}]) \rangle = 2 - 2g(S)$  is the minimal value and it is attained when both  $x$  and  $y$  are not in  $F$ . (In fact,  $(\{\beta_0\} \cup \beta, \{\alpha_0\} \cup \alpha, z, w)$  is a “sutured Heegaard diagram” in the sense of [Ni].)

The graded complex for calculating  $\widehat{HFK}(-M, K, -1)$  is generated by:

$$\begin{aligned} &\{z_0, x_0, y_0\}, \{z_0, x_0, y_2\}, \{z_0, x'_1, y_1\}, \{z_0, x_1, y_1\}, \\ &\{z_0, x_2, y_0\}, \{z_0, x_2, y_2\}, \{z_0, x_3, y_1\}. \end{aligned}$$

Our task is to identify  $\mathbf{x}' = \{z_0, x_0, y_0\}$  as a generator of  $\widehat{HFK}(-M, K, -1) \simeq \mathbb{Z}$ . We will show that all the generators besides  $\mathbf{x}'$  correspond to  $\text{Spin}^c$  structures which are different from that of the contact structure  $\xi$ . An easy computation shows that  $H_2(M; \mathbb{Z}) \simeq \mathbb{Z}^2$  and is generated by tori  $T_\delta$  of the form  $(\delta \times [0, 1]) / \sim$ , where  $\delta$  is any nonseparating curve on  $S$  and  $(x, 1) \sim (h(x), 0)$  as before. Since  $\xi$  is close to the foliation  $S \times \{t\}$  on  $(S \times [0, 1]) / \sim$ , it follows that  $\langle c(\xi), [T_\delta] \rangle = 0$ . Now, let  $\delta_1$  be a  $(0, 1)$ -curve on  $S$  and  $\delta_2$  be a  $(1, 0)$ -curve. Then  $[T_{\delta_1}]$  is given by the periodic domain  $\mathcal{P}_{\delta_1}$ , which consists of two rectangles  $y_0 y_2 y'_4 y'_2$  and  $y'_0 y'_2 y_4 y_2$  with opposite signs, shown in Figure 6. Similarly,  $[T_{\delta_2}]$  is

represented by  $\mathcal{P}_{\delta_2}$ , consisting of  $x_0x_2x'_4x'_2$  and  $x'_0x'_2x_4x_2$  with opposite signs.



**Figure 6.** Cover of a neighborhood of  $\partial S$ .

Now refer to Figure 6, which is a cover of an annular neighborhood of  $\partial S \subset S$ . The dotted curve is (a lift of)  $c_1c_2$ . Points below the dotted curve are not in  $F$ , so only they have the proper filtration level to represent generators.

We will show  $\mathbf{s}_{\{z_0,x,y\}} \neq \mathbf{s}_{\{z_0,x_0,y_0\}}$  if  $\{x,y\} \neq \{x_0,y_0\}$ , by showing that  $\langle c(\mathbf{s}_{\{z_0,x,y\}}), [T_{\delta_i}] \rangle \neq 0$  for  $i = 1$  and  $2$  if  $\{x,y\} \neq \{x_0,y_0\}$ .

First consider the intersection points on the vertical lines starting at  $x_0$  and at  $x'_0$ . Suppose that  $\langle c(\mathbf{s}_{\{z_0,x,y\}}), [T_{\delta_2}] \rangle = 0$ . The rectangle  $x'_0x'_2x_4x_2$  of the periodic domain  $\mathcal{P}_{\delta_2}$  contributes  $\frac{1}{2}$  if  $x = x_3$  or  $x = x'_1$ . Since there is no value of  $y$  below the dotted curve with a contribution of  $-\frac{1}{2}$  from the rectangle  $x_0x_2x'_4x'_2$  to cancel the  $\frac{1}{2}$ , the possibilities  $x = x_3, x'_1$  are eliminated. Since  $x_0x_2x'_4x'_2$  gives a contribution of  $-\frac{1}{2}$  to  $x_1$ , and  $x'_0x'_2x_4x_2$  contributes  $0$  to  $y_0$ ,  $\frac{1}{2}$  to  $y_1$  and  $1$  to  $y_1$ , the only generator containing  $x_1$  that is allowed is  $\{z_0, x_1, y_1\}$ . Any generator containing  $y_2$  is also disallowed since  $x'_0x'_2x_4x_2$  contributes  $1$  to  $y_2$ , and there is no  $x$  value that will offset it from the  $x_0x_2x'_4x'_2$  rectangle. The only generator allowed to contain  $y_1$  is again  $\{z_0, x_1, y_1\}$ . The same rectangle gives  $x'_1$  a contribution of  $-\frac{1}{2}$  that cannot be offset.

It therefore remains to consider the generator  $\{z_0, x_1, y_1\}$ , as well as pairs with  $x = x_0$  or  $x_2$ . Moreover, the only possible  $y$ -coordinates are  $y_0$  and  $y_1$ , and  $\{z_0, x_1, y_1\}$  is the only option allowed for  $y = y_1$ . Now use the periodic domain  $\mathcal{P}_{\delta_1}$ . The rectangle  $y_0y_2y'_4y'_2$  contributes  $-1$  to  $\{z_0, x_1, y_1\}$ , thus eliminating it as a possibility. The only other option

different from  $\{z_0, x_0, y_0\}$  is  $\{z_0, x_2, y_0\}$  (since  $y_2$  was banned) which gets a nonzero contribution from  $y_0 y_2 y'_4 y'_2$ .

To show how this argument generalizes to higher genus surfaces, let us examine the genus two case. The generators will have the form  $\{z_0, x, y, u, v\}$ , and there will be 8 intersection points on each vertical segment in a picture analogous to Figure 6. Denote the points on the boundary by  $u_0, v_0, u'_0, v'_0, x_0, y_0, x'_0, y'_0$  going from right to left. Start by considering the rectangles  $u_0 u_2 u'_8 u'_6$  and  $u'_0 u'_6 u_8 u_2$ . We eliminate  $u_3, \dots, u_7$  and all the  $u'$  values besides  $u'_0$ , by noticing that there is no allowable  $v$  value to offset the  $\frac{1}{2}$  contribution from  $u'_0 u'_6 u_8 u_2$ . The contribution of 1 from the same rectangle eliminates all values of  $v$  other than  $v_0$  and  $v_1$  (though no  $v'_i$  are yet disallowed). If  $v = v_1$ , only generators of the form  $\{z_0, x, y, u_1, v_1\}$  are allowed.

Now use the periodic domain represented by the rectangles  $v_0 v_2 v'_8 v'_6$  and  $v'_0 v'_6 v_8 v_2$ . The generators of the form  $\{z_0, x, y, u_1, v_1\}$  get a contribution of  $-1$  from  $v_0 v_2 v'_8 v'_6$  and there is no positive contribution from the allowable  $x, y$  coordinates that can be gained from  $v'_0 v'_6 v_8 v_2$ ; therefore all the  $\{z_0, x, y, u_1, v_1\}$  are eliminated. Next,  $u_2$  gets a contribution of  $-1$  from  $v_0 v_2 v'_8 v'_6$  that cannot be canceled since there is no  $v$  value that gets a contribution of 1 needed from  $v'_0 v'_6 v_8 v_2$ . It follows that  $u_0$  is the only allowable  $u$ -coordinate. Generators  $\{z_0, x, y, u_0, v'_i\}$ ,  $i \neq 0$ , are eliminated since  $v'_i$  gets a contribution of  $\frac{1}{2}$  from  $v'_0 v'_6 v_8 v_2$  that cannot be canceled. Therefore we are left with  $\{z_0, x, y, u_0, v_0\}$ . The argument is now reduced to eliminating the possible  $x, y$  coordinates, and this follows just as in the genus one argument given above.

This shows how the proof works for arbitrary genus. The inductive step is done in the same way by eliminating all extra options in the two new coordinates, thus reducing to the case of lower genus.

Since the contact invariant is the image of a generator of  $\widehat{HFK}(-M, K, -g)$  in  $\widehat{HF}(-M)$ , it follows that  $c(\xi_{(S,h)}) = EH(S, h)$ . It is not hard to see how the above argument generalizes to the  $h = \phi^n_{\partial S}$ ,  $n > 0$  case.

**3.3. Isotopy.** In this subsection we prove the following:

**Lemma 3.3.** *If  $h_t : S \xrightarrow{\sim} S$ ,  $t \in [0, 1]$ , is a 1-parameter family of diffeomorphisms which restrict to the identity on  $\partial S$ , then  $EH(S, h_0, \{a_1, \dots, a_r\}) = EH(S, h_1, \{a_1, \dots, a_r\})$ .*

*Proof.* Let  $\alpha_i = (a_i, a_i)$  and  $\beta_i^t = (b_i, h_t(b_i))$ . In other words, we fix the  $\alpha_i$  and isotop the  $\beta_i$ . Observe that the  $\beta_i^t$  remain constant on  $S \times \{1\}$ . According to Theorem 7.3 of [OS1], we can reduce to the case where  $h_t$  is a Hamiltonian isotopy. Let  $\Psi_t : \Sigma \xrightarrow{\sim} \Sigma$  be the Hamiltonian isotopy which restricts to the identity on  $S \times \{1\}$  and restricts to  $h_t$  on  $S \times \{0\}$ . We use the same notation for the induced isotopy on  $Sym^r(\Sigma)$ . Then the chain map  $\Phi : \widehat{CF}(\beta^0, \alpha) \rightarrow \widehat{CF}(\beta^1, \alpha)$  is obtained by counting holomorphic disks  $u : [0, 1] \times \mathbb{R} \rightarrow Sym^r(\Sigma)$  which satisfy  $\lim_{t \rightarrow +\infty} u(s +$

$it) = \mathbf{x}$ ,  $\lim_{t \rightarrow -\infty} u(s + it) = \mathbf{x}'$ ,  $u(0 + it) \in \Psi_t(\mathbb{T}_\beta)$ , and  $u(1 + it) \in \mathbb{T}_\alpha$ , and avoid  $\{z\} \times \text{Sym}^{r-1}(\Sigma)$ . Here  $\mathbf{x} \in \widehat{CF}(\beta^0, \alpha)$  and  $\mathbf{x}' \in \widehat{CF}(\beta^1, \alpha)$ . Now, if  $\mathbf{x}$  is the unique  $r$ -tuple of points on  $S \times \{1\}$  representing the generator of  $EH(S, h_0, \{a_1, \dots, a_r\})$ , then the only holomorphic disks of the above type are constant holomorphic disks, due to the placement of the basepoint  $z$ . This implies that  $EH(S, h_0, \{a_1, \dots, a_r\})$  is mapped to  $EH(S, h_1, \{a_1, \dots, a_r\})$  under the isomorphism  $\Phi : \widehat{HF}(\beta^0, \alpha) \xrightarrow{\sim} \widehat{HF}(\beta^1, \alpha)$ . q.e.d.

**3.4. Change of basis.** In this subsection we prove the following proposition:

**Proposition 3.4.**  *$EH(S, h, \{a_1, \dots, a_r\})$  is independent of the choice of basis  $\{a_1, \dots, a_r\}$ .*

Let  $\{a_1, a_2, \dots, a_r\}$  be a basis for  $S$ . After possibly reordering the  $a_i$ 's, suppose  $a_1$  and  $a_2$  are adjacent arcs on  $\partial S$ , i.e., there is an arc  $\tau \subset \partial S$  with endpoints on  $a_1$  and  $a_2$  such that  $\tau$  does not intersect any  $a_i$  in  $\text{int}(\tau)$ . Define  $a_1 + a_2$  to be the isotopy class of  $a_1 \cup \tau \cup a_2$ , relative to the endpoints. Then the modification  $\{a_1, a_2, \dots, a_r\} \mapsto \{a_1 + a_2, a_2, \dots, a_r\}$  is called an *arc slide*.

Proposition 3.4 is immediate from the following two lemmas.

**Lemma 3.5.**  *$EH(S, h)$  is invariant under an arc slide  $\{a_1, a_2, \dots, a_r\} \mapsto \{a_1 + a_2, a_2, \dots, a_r\}$ .*

*Proof.* Without loss of generality, consider the case where  $S$  is a once-punctured torus. We show that the chain map which corresponds to an arc slide takes the representative of  $EH(S, h, \{a_1, a_2\})$  determined by  $\mathbf{x} = \{x_1, x_2\}$  to the representative of  $EH(S, h, \{a_1 + a_2, a_2\})$  determined by the intersection point  $\mathbf{w} = \{w_1, w_2\}$ . Observe that an arc slide corresponds to a sequence of two handleslides for the corresponding Heegaard splitting.

Let  $(\Sigma, \beta, \alpha, z)$  be the pointed Heegaard diagram corresponding to  $a_i, b_i$  as described above, with  $z$  a point in  $S_{1/2}$  lying outside the thin strips of isotopy between the  $a_i$ 's and the  $b_i$ 's. If we slide  $\alpha_2$  over  $\alpha_1$  along a path parallel to  $\partial S$ , then we obtain a new pair  $\gamma = \{\gamma_1, \gamma_2\}$ , where  $\gamma_1 = (a_1 + a_2, a_1 + a_2)$  and  $\gamma_2$  is a suitable pushoff of  $(a_2, a_2)$  as in the proof of the invariance of Heegaard Floer homology under handleslides in [OS1]. Figure 7 depicts the case where  $a_1$  is to the right of  $a_2$  with respect to  $\tau$ ; the case where  $a_2$  is to the right of  $a_1$  is treated similarly.

We claim that  $(\Sigma, \gamma, \beta, \alpha, z)$  is a weakly admissible Heegaard triple-diagram. Recall that a triple-diagram is *weakly admissible* if each non-trivial triply-periodic domain which can be written as a sum of doubly-periodic domains has both positive and negative coefficients. First let us restrict to a neighborhood  $\mathcal{R}$  of the labeled regions of  $\Sigma - \cup_i \alpha_i - \cup_i \beta_i - \cup_i \gamma_i$  on the right-hand side of Figure 7. Due to the placement

of  $z$ , the only potential doubly-periodic region involving  $\beta, \alpha$  on  $\mathcal{R}$  is  $D_2 + D_3 - D_5 - D_6$ . (Here  $D_i$  is the domain labeled  $i$ .) Similarly, for  $\gamma, \beta$  we have  $D_1 + D_2 - D_4 - D_5$  and for  $\alpha, \gamma$  we have  $D_1 + D_6 - D_3 - D_4$ . Taking linear combinations, we have


$$\begin{aligned} & a(D_2 + D_3 - D_5 - D_6) + b(D_1 + D_2 - D_4 - D_5) \\ & \quad + c(D_1 + D_6 - D_3 - D_4) \\ = & (b + c)D_1 + (a + b)D_2 + (a - c)D_3 \\ & \quad - (b + c)D_4 - (a + b)D_5 + (-a + c)D_6. \end{aligned}$$

Since the coefficients come in pairs, e.g.,  $a + b$  and  $-(a + b)$ , if any of  $a + b, b + c, a - c$  does not vanish, then the triply-periodic domain has both positive and negative coefficients. Hence, if any of  $\alpha_1, \beta_1$  and  $\gamma_1$  is used, then we are done. Otherwise, we may assume that none of  $\alpha_1, \beta_1$  and  $\gamma_1$  is used in the periodic domain. This allows us to erase all three, and apply the above considerations to  $\alpha_2, \beta_2$ , and  $\gamma_2$ . The verifications of weak admissibility of all other triple-diagrams in this paper are identical, and are omitted.

Let  $\Theta = \{\Theta_1, \Theta_2\}$  be the top generator of  $\widehat{HF}(\#(S^1 \times S^2)) = \widehat{HF}(\alpha, \gamma)$ . Define the map

$$\psi : \widehat{HF}(\beta, \alpha) \otimes \widehat{HF}(\alpha, \gamma) \rightarrow \widehat{HF}(\beta, \gamma),$$

where  $\psi(\mathbf{y} \otimes \mathbf{y}')$  counts holomorphic triangles, two of whose vertices are  $\mathbf{y} \in \widehat{CF}(\beta, \alpha)$  and  $\mathbf{y}' \in \widehat{CF}(\alpha, \gamma)$ . Then the isomorphism  $g : \widehat{HF}(\beta, \alpha) \xrightarrow{\sim} \widehat{HF}(\beta, \gamma)$  is given by  $g(\mathbf{y}) = \psi(\mathbf{y} \otimes \Theta)$ .



**Figure 7.** The first handleslide.


We claim that the representative  $\mathbf{x} = \{x_1, x_2\}$  of  $EH(S, h, \{a_1, a_2\})$  gets mapped to  $\mathbf{y} = \{y_1, y_2\} \in \widehat{CF}(\beta, \gamma)$  given in Figure 7. By the

placement of  $z$ , we see that the unique domain which has  $x_1$  and some  $\Theta_i$  as corners and avoids  $z$  must be a triangle with vertices  $x_1, \Theta_1, y_1$ . Now that  $\alpha_1, \beta_1$ , and  $\gamma_1$  are used up, it easily follows that the unique domain which involves  $x_2$  and  $\Theta_2$  and avoids  $z$  is a triangle with vertices  $x_2, \Theta_2, y_2$ . This proves the claim.

Let us now consider the effect of the second handleslide, depicted in Figure 8. Let  $\delta = \{\delta_1, \delta_2\}$ , where  $\delta_1$  and  $\delta_2$  are suitable pushoffs of  $(a_1 + a_2, h(a_1 + a_2))$  and  $(a_2, h(a_2))$ , respectively. A similar argument as above shows that, under the map

$$\widehat{HF}(\delta, \beta) \otimes \widehat{HF}(\beta, \gamma) \rightarrow \widehat{HF}(\delta, \gamma),$$

$\Theta \otimes \mathbf{y}$  gets mapped to  $\mathbf{w}$ . This shows that  $\mathbf{x}$  and  $\mathbf{w}$  determine the same element in Heegaard Floer homology, and consequently  $EH(S, h, \{a_1, a_2\}) = EH(S, h, \{a_1 + a_2, a_2\})$ . q.e.d.



**Figure 8.** The second handleslide.


**Lemma 3.6.** *Let  $\{a_1, \dots, a_r\}$  and  $\{b_1, \dots, b_r\}$  be two bases for  $S$ . Then there is a sequence of arc slides that takes  $\{a_1, \dots, a_r\}$  to  $\{b_1, \dots, b_r\}$ .*

We do not need to assume that  $\partial S$  is connected.

*Proof.* We argue that we can reduce the total number of intersections of  $\bigcup_i a_i$  and  $\bigcup_i b_i$  by replacing  $\{a_1, \dots, a_r\}$  with  $\{a'_1, \dots, a'_r\}$ , which is obtained from  $\{a_1, \dots, a_r\}$  by a sequence of arc slides. By inducting on the number of intersection points, this shows that we can perform a sequence of arc slides until  $\bigcup_i a_i$  and  $\bigcup_i b_i$  become disjoint. We then show that two disjoint bases can be brought one into another by a sequence of arc slides.

Let  $P = S - \bigcup_i a_i$ . Then  $P$  is a polygon whose boundary  $\partial P$  consists of  $4r$  arcs,  $2r$  of which are  $a_i$  or  $a_i^{-1}$  and  $2r$  of which are arcs  $\tau_1, \dots, \tau_{2r}$  of  $\partial S$ .

Suppose  $(\bigcup_i a_i) \cap (\bigcup_i b_i) \neq \emptyset$ , where we are assuming efficient intersections. After possibly reordering the arcs, there is a subarc  $b_1^0 \subset b_1$  which starts on  $\tau_1 \subset \partial S$  and ends on  $a_1$ , and whose interior  $\text{int}(b_1^0)$  does not intersect  $\bigcup_i a_i$ . (In other words,  $b_1^0$  is a properly embedded arc of  $P$ .) We may assume that  $a_1$  is not adjacent to  $\tau_1$ ; otherwise, isotop the relevant endpoint of  $b_1$  along  $\tau_1$ . The subarc  $b_1^0$  separates the polygon  $P$  into two regions  $P_1$  and  $P_2$ , only one of which contains a boundary arc that is labeled  $a_1^{-1}$  (say  $P_2$ ). We can then slide  $a_1$  over all the arcs of type  $a_i$  or  $a_i^{-1}$  in the other region  $P_1$ , and obtain the new curve  $a'_1$  as in Figure 9 so that the new basis  $\{a'_1, a_2, \dots, a_r\}$  has fewer intersections with  $\bigcup_i b_i$ . (Note that trying to slide over  $a_1^{-1}$  presents a problem, so we



**Figure 9.** Simplifying the intersections of  $\bigcup a_i$  and  $\bigcup b_i$ .

must go the other way around.) There is one situation when the above strategy needs a little more thought, namely when  $\partial P_2$  only intersects  $a_1$  and  $a_1^{-1}$  (among all the  $a_i$  and  $a_i^{-1}$ ). In this case,  $b_1$  exits the polygon  $P$  along  $a_1$  and reenters through  $a_1^{-1}$ . Eventually we find a subarc of  $b_1$  which starts on some  $\tau_2$  and ends on an adjacent  $a_1^{-1}$ , a contradiction. We now apply the same procedure to  $\{a'_1, a_2, \dots, a_r\}$  and  $\{b_1, \dots, b_r\}$  until they become disjoint.

Now suppose that the two bases  $\{a_1, \dots, a_r\}$  and  $\{b_1, \dots, b_r\}$  are disjoint. We consider the polygon  $P = S - \bigcup_i a_i$ . Some of the  $b_i$  arcs may be parallel to  $a_j$  or  $a_j^{-1}$ . An arc  $b_1$  that is not parallel to any of the  $a_i$  will cut  $P$  into two components  $P_1$  and  $P_2$ , each containing more than

one of  $a_i, a_i^{-1}$ ,  $i = 1, \dots, r$ . Recall that  $b_1$  is nonseparating. One can easily verify that  $b_1$  being nonseparating is equivalent to the existence of some  $a_i$  such that  $a_i \in P_1$  and  $a_i^{-1} \in P_2$  (or vice versa). (If there is some  $a_i$ , then take an arc  $c$  in  $P$  from  $a_i \subset P_1$  to  $a_i^{-1} \subset P_2$ . The closed curve in  $S$  obtained by gluing up  $c$  is dual to  $b_1$ .) If each such  $a_i$  is parallel to some  $b_j$ , then  $S - \bigcup_i b_i$  would be disconnected. Hence we may additionally assume that there is some  $a_i$  which is not parallel to any  $b_j$ . Now we slide  $a_i$  across all the arcs of type  $a_j, a_j^{-1}$  in  $P_1$  until it becomes parallel to  $b_1$ . q.e.d.


**3.5. Legendrian surgery.** Let  $\delta$  be a nonseparating curve and  $\phi_\delta^{-1}$  be a negative Dehn twist about  $\delta$ . We now transfer  $EH$  from  $M = M_{(S,h)}$  to  $M' = M_{(S,\phi_\delta^{-1} \circ h)}$ . Recall that there is a natural map

$$f : \widehat{HF}(-M) \rightarrow \widehat{HF}(-M'),$$

which arises from tensoring with the top generator  $\Theta$  of  $\widehat{HF}(\#(S^1 \times S^2))$ .

**Proposition 3.7.**  $f(EH(S, h)) = EH(S, \phi_\delta^{-1} \circ h)$ .

*Proof.* By Proposition 3.4 we may take a basis  $\{a_1, \dots, a_r\}$  for  $S$  so that  $\delta$  is disjoint from  $h(b_2), \dots, h(b_r)$ , intersects  $h(b_1)$  exactly once, and is parallel to  $h(b_2)$ . Then the result of performing (+1)-surgery along  $\delta$  (or, equivalently, a negative Dehn twist along  $\delta$ ) is given by Figure 10.



**Figure 10.** Legendrian (+1)-surgery. The second figure shows curves and their *relative* position correctly, but they are positioned on  $S_0$  as if  $h^{-1}$  had been applied to each of them.

The  $\alpha$ -curves and  $\beta$ -curves are as before, and we define the  $\gamma$ -curves as follows: Let  $\gamma_1 = (b_1, \phi_\delta^{-1} \circ h(b_1))$  and  $\gamma_i = (b_i, h(b_i))$  for  $i > 1$ . Let  $\Theta \in \widehat{HF}(\gamma, \beta)$  be the top generator of  $\#(S^1 \times S^2)$ , given in Figure 10. Define the map

$$\phi : \widehat{HF}(\gamma, \beta) \otimes \widehat{HF}(\beta, \alpha) \rightarrow \widehat{HF}(\gamma, \alpha),$$

where  $\phi(\mathbf{y} \otimes \mathbf{y}')$  counts holomorphic triangles, two of whose vertices are  $\mathbf{y}$  and  $\mathbf{y}'$ . Then the map  $f : \widehat{HF}(\beta, \alpha) \rightarrow \widehat{HF}(\gamma, \alpha)$  is given by  $f(\mathbf{y}) = \phi(\Theta \otimes \mathbf{y})$ . By the convenient placement of  $z$ , it follows that we only have small triangles in the Heegaard diagram. Hence if  $[\mathbf{x}] = EH(S, h, \{a_1, a_2\})$ , then  $\phi([\Theta \otimes \mathbf{x}]) = EH(S, \phi_\delta^{-1} \circ h, \{a_1, a_2\})$ . q.e.d.

**3.6. Multiple boundary components.** Consider  $(S, h)$  where  $S$  has disconnected boundary. For simplicity, assume  $S$  has two boundary components. Pick a basis  $\{a_1, \dots, a_r\}$  for  $S$ . Next consider  $(S', h\#id)$ , where  $S'$  is obtained from  $S$  by attaching a 1-handle between the two boundary components and we are extending  $h$  by the identity. If  $a_0$  is the cocore of the 1-handle, then  $\{a_0, \dots, a_r\}$  is a basis for  $S'$ . Our argument is similar to that of Lemma 4.4 of [OS3]. The natural map

$$F_U : \widehat{HF}((-M_{(S,h)}) \# (S^1 \times S^2)) \rightarrow \widehat{HF}(-M_{(S,h)}),$$

which corresponds to the cobordism  $U$  attaching a 3-handle as in Section 4.3 of [OS4], sends

$$EH(S', h\#id, \{a_0, a_1, \dots, a_r\}) \mapsto EH(S, h, \{a_1, \dots, a_r\}).$$

Since  $S'$  has only one boundary component, we already know that

$$c(S', h\#id) = EH(S', h\#id, \{a_0, a_1, \dots, a_r\}).$$

Moreover, if  $\delta$  is a closed curve on  $S'$  which is “dual” to  $a_0$ , then there is a natural map

$$F_W : \widehat{HF}(-M_{(S,h)}) \rightarrow \widehat{HF}((-M_{(S,h)}) \# (S^1 \times S^2))$$

which maps  $c(S, h)$  to  $c(S', h\#id)$ . Here  $(S, h)$  and  $(S', \phi_\delta \circ (h\#id))$  represent the same 3-manifold, and  $W$  is the cobordism from  $-M_{(S', \phi_\delta \circ (h\#id))}$  to  $-M_{(S', h\#id)}$ , obtained by attaching a 2-handle along the curve  $\delta$ . Finally,  $U \circ W \simeq [0, 1] \times M_{(S,h)}$ , so

$$c(S, h) = F_U \circ F_W(c(S, h)) = F_U(c(S', h\#id)) = EH(S, h, \{a_1, \dots, a_r\}).$$

#### 4. Right-veering and holomorphic disks

In this section we prove Theorem 1.2.

*Proof of Theorem 1.2.* Let  $S$  be a once-punctured torus.

Suppose first that  $h$  has pseudo-Anosov monodromy. If the fractional Dehn twist coefficient  $c \geq 1$ , then the contact structure is already symplectically fillable and universally tight. It also follows that  $c(\xi_{(S,h)}) \neq 0$ . If  $c = \frac{1}{2}$ , then  $c(\xi_{(S,h)}) \neq 0$  follows from Theorem 4.1 below. If  $c \leq 0$ , then  $\xi$  is overtwisted since  $S$  is not right-veering. (See [HKM2].)

If  $h$  is periodic, then  $\xi$  is right-veering if and only if  $h$  is a product of positive Dehn twists by [HKM3].

If  $h$  is reducible, then  $c(\xi_{(S,h)}) \neq 0$  follows from Theorem 4.3 below.

q.e.d.

**Theorem 4.1.** *Let  $(S, h)$  be an open book decomposition for  $M$ , where  $S$  is a once-punctured torus and  $h$  is pseudo-Anosov with fractional Dehn twist coefficient  $c = \frac{1}{2}$ . Then  $c(\xi_{(S,h)}) = EH(S, h) \neq 0$ , and hence the contact structure  $\xi_{(S,h)}$  is tight.*

*Proof.* We show that  $EH(S, h) \neq 0$  by choosing a basis for  $S$  for which there are no holomorphic disks in the corresponding Heegaard diagram that map to the generator  $\mathbf{x} = \{x_0, y_0\}$  defining  $EH(S, h)$ .

The following lemma furnishes us with a convenient basis:

**Lemma 4.2.** *Let  $A \in SL(2, \mathbb{Z})$  be a matrix with  $\text{tr}(A) < -2$ . Then  $A$  is conjugate in  $SL(2, \mathbb{Z})$  to a matrix  $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ , where  $(a, c)$  and  $(b, d)$  are in the third quadrant.*

*Proof.* Let  $\Lambda^s$  and  $\Lambda^u$  be the stable and unstable laminations for  $A$ . The slopes of  $\Lambda^s$  and  $\Lambda^u$  will be written  $\text{slope}(\Lambda^s)$  and  $\text{slope}(\Lambda^u)$ . (Recall that these slopes are irrational.) Let us consider the Farey tessellation on the hyperbolic unit disk  $D^2$ . Pick a vertex  $s_1$  on the clockwise edge along  $\partial D^2$  from  $\text{slope}(\Lambda^s)$  to  $\text{slope}(\Lambda^u)$ , and pick a vertex  $s_2$  on the counterclockwise edge from  $\text{slope}(\Lambda^s)$  to  $\text{slope}(\Lambda^u)$ , so that there is an edge of the Farey tessellation between  $s_1$  and  $s_2$ . (The existence of such a pair  $s_1, s_2$  is an exercise.) Then  $A(s_1)$  (resp.  $A(s_2)$ ) is closer to  $\text{slope}(\Lambda^s)$  than  $s_1$  (resp.  $s_2$ ) is. An oriented basis corresponding to  $(s_1, s_2)$  will have the desired property. q.e.d.

With the choice of basis as above, we can represent  $M = M_{(S,h)}$  by the Heegaard diagram below. We have drawn a picture of the diagram corresponding to  $A = \begin{pmatrix} -1 & -1 \\ -1 & -2 \end{pmatrix}$ , but the same argument works for any such  $A$  as described in the previous lemma. We prove that there is no holomorphic disk from any  $\mathbf{y}$  to  $\mathbf{x} = \{x_0, y_0\}$ . Suppose on the contrary that there is such a holomorphic disk  $u$ . Let  $\partial\mathcal{D}(u)$  be the boundary of the support of  $\mathcal{D}(u)$ . Assuming  $\partial\mathcal{D}(u)$  is connected, it is given by a subarc of  $a_1$  from some  $x_i \in a_1 \cap h(a_2)$  to  $x_0$ , followed by a subarc of  $h(a_1)$  from  $x_0$  to some  $y_j \in a_2 \cap h(a_1)$ , followed by a subarc of  $a_2$  from  $y_j$  to  $y_0$  (you either turn left or turn right at  $y_j$ ), and then by a subarc of  $h(a_2)$  from  $y_0$  to  $x_i$ . If we lift  $\partial\mathcal{D}(u)$  to the universal cover of the capped off surface  $T^2 = S \cup D^2$ , then in all cases we see that  $\partial\mathcal{D}(u)$  is not contractible. This implies that  $\partial\mathcal{D}(u)$  cannot bound a surface in  $S$ . We argue similarly when  $\partial\mathcal{D}(u)$  has two components. It follows that the class  $EH(S, h)$  of  $\mathbf{x} = \{x_0, y_0\}$  is nonzero. q.e.d.

**Theorem 4.3.**  *$EH(S, h) \neq 0$  if  $h$  is reducible and right-veering.*

*Proof.* Suppose  $h$  is reducible. Let  $g$  be an element of  $\text{Aut}(S, \partial S)$  which is the minimally right-veering representative for the matrix  $A = -id$ . (In terms of positive Dehn twists,  $g = (A_1 A_2 A_1)^2$ , where  $A_1 = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$  and  $A_2 = \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}$ .) After changing bases if necessary,  $h = g^n \phi_\gamma^m$ ,




Figure 11

where  $n$  is a positive integer,  $m$  is an integer, and  $\phi_\gamma$  is a positive Dehn twist about a  $(0, 1)$ -curve  $\gamma$ . If  $m$  is nonnegative, then  $h$  is a product of positive Dehn twists, and  $EH(S, h) \neq 0$ .

Suppose  $m < 0$ . It suffices to prove the theorem for  $n = 1$ , since the contact structures corresponding to larger  $n$  are obtained from the  $n = 1$  case by Legendrian surgery. Take a basis corresponding to slopes  $0, \infty$  and matrix  $A = \begin{pmatrix} -1 & 0 \\ -m & -1 \end{pmatrix}$ . Then  $EH(S, h)$  is nonzero by the same argument as in Theorem 4.1. q.e.d.

### References

- [Ba] J. Baldwin, *Tight contact structures and genus one fibered knots*, *Algebr. Geom. Topol.* **7** (2007), 701–735, MR 2308961, Zbl pre05220891.
- [El] Y. Eliashberg, *Filling by holomorphic discs and its applications*, *Geometry of low-dimensional manifolds, 2* (Durham, 1989), 45–67, *London Math. Soc. Lecture Note Ser.* **151**, Cambridge Univ. Press, Cambridge, 1990, MR 1171908, Zbl 0731.53036.
- [Gh] P. Ghiggini, *Strongly fillable contact 3-manifolds without Stein fillings*, *Geom. Topol.* **9** (2005), 1677–1687 (electronic), MR 2175155, Zbl 1091.57018.
- [Gi1] E. Giroux, *Convexit  en topologie de contact*, *Comment. Math. Helv.* **66** (1991), 637–677, MR 1129802, Zbl 0766.53028.
- [Gi2] E. Giroux, *G om trie de contact: de la dimension trois vers les dimensions sup rieures*, *Proceedings of the International Congress of Mathematicians, Vol. II* (Beijing, 2002), 405–414, Higher Ed. Press, Beijing, 2002, MR 1957051, Zbl 1015.53049.
- [Go] N. Goodman, *Overtwisted open books from sobering arcs*, *Algebr. Geom. Topol.* **5** (2005), 1173–1195 (electronic), MR 2171807, Zbl 1090.57020.

- [H1] K. Honda, *On the classification of tight contact structures I*, *Geom. Topol.* **4** (2000), 309–368 (electronic), MR 1786111, Zbl 0980.57010.
- [HKM1] K. Honda, W. Kazez & G. Matić, *Tight contact structures on fibered hyperbolic 3-manifolds*, *J. Differential Geom.* **64** (2003), 305–358, MR 2029907, Zbl 1083.53082.
- [HKM2] K. Honda, W. Kazez & G. Matić, *Right-veering diffeomorphisms of compact surfaces with boundary*, *Invent. Math.* **169** (2007), 427–449, MR 2318562, Zbl pre05199715.
- [HKM3] K. Honda, W. Kazez & G. Matić, *Right-veering diffeomorphisms of compact surfaces with boundary II*, *Geom. Topol.* **12** (2008), 2057–2094, MR 2431016, Zbl pre05344746.
- [LS1] P. Lisca & A. Stipsicz, *Ozsváth-Szabó invariants and tight contact three-manifolds. I*, *Geom. Topol.* **8** (2004), 925–945 (electronic), MR 2087073, Zbl 1059.57017.
- [LS2] P. Lisca & A. Stipsicz, *Ozsváth-Szabó invariants and tight contact three-manifolds. II*, *J. Differential Geom.* **75** (2007), 109–141, MR 2282726, Zbl 1112.57005.
- [Ni] Y. Ni, *Sutured Heegaard diagrams for knots*, *Algebr. Geom. Topol.* **6** (2006), 513–537 (electronic), MR 2220687, Zbl 1103.57021.
- [Oh] Y. Oh, *Fredholm theory of holomorphic discs under the perturbation of boundary conditions*, *Math. Z.* **222** (1996), 505–520, MR 1400206, Zbl 0863.53024.
- [OS1] P. Ozsváth & Z. Szabó, *Holomorphic disks and topological invariants for closed three-manifolds*, *Ann. of Math. (2)* **159** (2004), 1027–1158, MR 2113019, Zbl 1073.57009.
- [OS2] P. Ozsváth & Z. Szabó, *Holomorphic disks and three-manifold invariants: properties and applications*, *Ann. of Math. (2)* **159** (2004), 1159–1245, MR 2113020, Zbl 1081.57013.
- [OS3] P. Ozsváth & Z. Szabó, *Holomorphic disks and knot invariants*, *Adv. Math.* **186** (2004), 58–116, MR 2065507, Zbl 1062.57019.
- [OS4] P. Ozsváth & Z. Szabó, *Holomorphic triangles and invariants for smooth four-manifolds*, *Adv. Math.* **202** (2006), 326–400, MR 2222356, Zbl 1099.53058.
- [OS5] P. Ozsváth & Z. Szabó, *Heegaard Floer homology and contact structures*, *Duke Math. J.* **129** (2005), 39–61, MR 2153455, Zbl 1083.57042.
- [Ra] J. Rasmussen, *Floer homology and knot complements*, Ph.D. Thesis 2003. ArXiv:math.GT/0306378.
- [TW] W. Thurston & H. Winkelnkemper, *On the existence of contact forms*, *Proc. Amer. Math. Soc.* **52** (1975), 345–347, MR 0375366, Zbl 0312.53028.

UNIVERSITY OF SOUTHERN CALIFORNIA  
LOS ANGELES, CA 90089

*E-mail address:* khonda@usc.edu

UNIVERSITY OF GEORGIA  
ATHENS, GA 30602

*E-mail address:* will@math.uga.edu

UNIVERSITY OF GEORGIA  
ATHENS, GA 30602  
*E-mail address:* [gordana@math.uga.edu](mailto:gordana@math.uga.edu)