

EXPLICIT BIRATIONAL GEOMETRY OF 3-FOLDS OF GENERAL TYPE, II

JUNGKAI A. CHEN & MENG CHEN

Abstract

Let V be a complex nonsingular projective 3-fold of general type. We shall give a detailed classification up to baskets of singularities on a minimal model of V . We show that the m -canonical map of V is birational for all $m \geq 73$ and that the canonical volume $\text{Vol}(V) \geq \frac{1}{2660}$. When $\chi(\mathcal{O}_V) \leq 1$, our result is $\text{Vol}(V) \geq \frac{1}{420}$, which is optimal. Other effective results are also included in the paper.

1. Introduction

Let Y be a nonsingular projective variety of dimension n . It is said to be of general type if the pluricanonical map φ_m corresponding to the linear system $|mK_Y|$ is birational into a projective space for $m \gg 0$. Thus it is natural and important to find a constant $c(n)$, depending only on dimension, so that φ_m is birational onto its image for all $m \geq c(n)$ and for all Y with $\dim Y = n$.

It was classically known that, when $\dim Y = 1$, $|mK_Y|$ gives an embedding of Y into a projective space for $m \geq 3$. When $\dim Y = 2$, Kodaira-Bombieri's theorem [2] implies that $|mK_Y|$ gives a birational map onto the image for $m \geq 5$. A recent result of Hacon and McKernan [10], Takayama [23], and Tsuji [25] shows the existence of $c(n)$, which is however non-explicit.

This is the continuation of our previous paper [4]. The aim of this paper is to prove a practical constant $c(3)$, which is not too far from being sharp. Other effective results are included in this paper as well.

Recall that we have proved the following result in [4].

Theorem 1. ([4, Theorem 1.1]) *Let V be a nonsingular projective 3-fold of general type. Then:*

- (1) $P_{12} > 0$;
- (2) $P_{m_0} \geq 2$ for some positive integer $m_0 \leq 24$.

Our main theorems of this paper are as follows.

Received 1/8/2010.

Theorem 1.1. *Let V be a nonsingular projective 3-fold of general type. Then:*

- (1) $P_m > 0$ for all $m \geq 27$.
- (2) $P_{24} \geq 2$ and $P_{m_0} \geq 2$ for some positive integer $m_0 \leq 18$.
- (3) φ_m is birational for all $m \geq 73$, and in case $\chi(\mathcal{O}_X) \leq 1$, φ_m is birational for all $m \geq 40$.

Here is our result on the volume.

Theorem 1.2. *Let V be a non-singular projective 3-fold of general type. Then:*

- (1) $\text{Vol}(V) \geq \frac{1}{2660}$. Furthermore, $\text{Vol}(V) = \frac{1}{2660}$ if and only if $P_2 = 0$ and either $\chi(\mathcal{O}_V) = 3$, $\mathcal{B}(X) = \{9 \times \frac{1}{2}(1, -1, 1), 2 \times \frac{1}{7}(1, -1, 3), \frac{1}{19}(1, -1, 7), 3 \times \frac{1}{3}(1, -1, 1), \frac{1}{10}(1, -1, 3), \frac{1}{4}(1, -1, 1), \frac{1}{5}(1, -1, 1)\}$ or $\chi(\mathcal{O}_V) = 2$, $\mathcal{B}(X) = \{2 \times \frac{1}{2}(1, -1, 1), 2 \times \frac{1}{7}(1, -1, 3), 2 \times \frac{1}{5}(1, -1, 2), \frac{1}{19}(1, -1, 7), \frac{1}{4}(1, -1, 1)\}$ where $\mathcal{B}(X)$ is the basket of singularities on a minimal model X of V .
- (2) In case $\chi(\mathcal{O}_V) \leq 1$, $\text{Vol}(V) \geq \frac{1}{420}$, which is an optimal lower bound. Furthermore, $\text{Vol}(V) = \frac{1}{420}$ if and only if the basket of singularities on any minimal model X of V is $\{3 \times \frac{1}{2}(1, -1, 1), \frac{1}{7}(1, -1, 3), \frac{1}{5}(1, -1, 2), \frac{1}{4}(1, -1, 1), \frac{1}{6}(1, -1, 1)\}$.

Theorem 1.2 (2) is optimal due to the following example:

Example 1.3. ([12, page 151, no. 23]) The canonical hypersurface $X_{46} \subset \mathbb{P}(4, 5, 6, 7, 23)$ has 7 terminal quotient singularities and the canonical volume $K_{X_{46}}^3 = \frac{1}{420}$. One knows $\chi(\mathcal{O}_{X_{46}}) = 1$ since $p_g(X_{46}) = q(X_{46}) = h^2(\mathcal{O}_{X_{46}}) = 0$. Furthermore, it is known that φ_m is birational for all $m \geq 27$, but φ_{26} is not birational.

We now briefly sketch the main idea of this article. A general approach to study pluricanonical maps in higher dimensions is by utilizing vanishing theorems. The difficulty is usually reduced to bound from below the canonical volume

$$\text{Vol}(Y) := \limsup_{\{m \in \mathbb{Z}^+\}} \left\{ \frac{n!}{m^n} \dim_{\mathbb{C}} H^0(Y, \mathcal{O}_Y(mK_Y)) \right\}.$$

The volume is an integer when $\dim Y \leq 2$, and hence a naive lower bound 1 is obtained. However, it's a rational number in dimension three or higher. This is an essential difficulty of high-dimensional birational geometry.

Another technical approach is the induction approach initiated by Kollár [15], who proved that φ_{11m_0+5} is birational provided $P_{m_0} \geq 2$ for 3-folds of general type. Kollár's method has been generalized in several directions by Chen [7], Chen-Hacon [3], Chen-Zuo [8], Chen-Chen [5], and so on. Therefore, it remains to consider 3-folds with

small plurigenera. One notices that the plurigenus $P_m(Y)$ is nothing but the Euler characteristic $\chi(X, mK_X)$ of its minimal model X thanks to the vanishing theorem, and moreover if the minimal model is non-singular or Gorenstein, then $\chi(\mathcal{O}_Y) < 0$. One obtains $P_2 \geq 4$ easily by the Riemann-Roch formula.

Reid introduced the notion of *baskets of singularities* which are local deformation of singularities into cyclic quotients and derived a singular Riemann-Roch formula for threefolds with at worst canonical singularities. Roughly speaking, the “singular” Riemann-Roch formula computes the Euler characteristic $\chi(Y, mK_Y)$ by the usual Riemann-Roch terms and the contribution from singularities which is computed by baskets. The key new ingredient is our systematical study of baskets of singularities in [4]. Our method provides a concrete way to determine an approximation of a basket with given leading Euler characteristics $\chi(K_Y)$, $\chi(2K_Y)$, \dots , etc. As a consequence, we are able to prove the finiteness of baskets with small leading Euler characteristics. It is even possible to give explicit classification up to baskets, which is exactly what we have done in this paper.

The article is organized as follows. In Section 2, we summarize some results on the geometry of $|mK|$, which substantially extend the above mentioned technique. Combining with the technique on baskets of singularities developed in [4], we will give a successful classification in case $\chi(\mathcal{O}) = 1$ in Section 3. In Section 4, we classify baskets such that $P_m \leq 1$ for all $1 < m \leq 12$ and $\chi(\mathcal{O}) > 1$. We get 63 classes of baskets of 3-folds in Table C. All these classification allows us to find a practical number $n_1 > 0$ such that $P_{n_1} \geq 2$. Therefore, we are able to prove our main theorems.

Throughout, we will frequently use those definitions, equalities, and inequalities about formal baskets in our previous paper (see [4, Sections 3 and 4]). We prefer to use “ \equiv ” to denote numerical equivalence, while “ \sim ” represents linear equivalence. Roundup operator “ $\lceil * \rceil$ ” is defined to be “ $-\lfloor -* \rfloor$ ”, where rounddown “ $\lfloor * \rfloor$ ” means taking the integral part.

We are very grateful to an anonymous referee whose keen suggestion makes this paper be much better organized.

Acknowledgments. The first author was partially supported by TIMS, NCTS/TPE and National Science Council of Taiwan. The second author was supported by National Outstanding Young Scientist Foundation (#10625103) and NNSFC Key project (#10731030).

2. Technical preparation

In this section, we set up some notions and principles evolved in our detailed study. We shall prove some general results on pluricanonical birationality and the lower bound of canonical volume. Though the

method has already appeared in several previous works, the way of applying it is resultful to the effect that we are able to treat various situations while proving our main theorems.

2.1. Reduction to problems on minimal 3-folds. Let V be a non-singular projective 3-fold of general type. By the 3-dimensional Minimal Model Program (see, for instance, [14, 16, 20]), V has a minimal model X (with K_X nef and admitting \mathbb{Q} -factorial terminal singularities). Denote by K_X a canonical divisor of X . A basic fact is that $\text{Vol}(V) = K_X^3 > 0$. From the view point of birational geometry, it suffices to prove main theorem for minimal 3-folds X .

Definition 2.2. (1) The number $\rho_i = \rho_i(X)$ denotes the minimal positive integer such that $P_m(X) > i$ for all $m \geq \rho_i$, where $i = 0, 1$.

(2) The number $\mu_i = \mu_i(X)$ denotes the minimal positive integer with $P_{\mu_i} = P_{\mu_i}(X) > i$ where $i = 0, 1, 2$.

(3) Denote by $\mathcal{B}(X)$ the basket of singularities on X (according to Reid [21]), and by $r(X)$ the Cartier index of X .

By our definition, we see $\rho_0 \leq \rho_1$ and $\mu_0 \leq \mu_1 \leq \rho_1$. The existence of ρ_1 can be guaranteed by Theorem 1.

Now suppose we have $P_{m_0} \geq 2$ for certain positive integer m_0 . We may study the geometry of the rational map $\varphi_{m_0} := \Phi_{|m_0 K_X|}$.

2.3. Set up for φ_{m_0} . We study the m_0 -canonical map of X :

$$\varphi_{m_0} : X \dashrightarrow \mathbb{P}^{P_{m_0}-1},$$

which is a rational map. First of all we fix an effective Weil divisor $K_{m_0} \sim m_0 K_X$. By Hironaka's big theorem, we can take successive blow-ups $\pi : X' \rightarrow X$ such that:

- (i) X' is smooth;
- (ii) the movable part of $|m_0 K_{X'}|$ is base point free;
- (iii) the support of the union of $\pi^*(K_{m_0})$ and the exceptional divisors is of simple normal crossings.

Set $g_{m_0} := \varphi_{m_0} \circ \pi$. Then g_{m_0} is a morphism by assumption. Let $X' \xrightarrow{f} \Gamma \xrightarrow{s} W'$ be the Stein factorization of g_{m_0} with W' the image of X' through g_{m_0} . In summary, we have the following commutative diagram:

$$\begin{array}{ccc} X' & \xrightarrow{f} & \Gamma \\ \pi \downarrow & \searrow^{g_{m_0}} & \downarrow s \\ X & \xrightarrow{\varphi_{m_0}} & W' \end{array}$$

Recall that

$$\pi^*(K_X) := K_{X'} - \frac{1}{r(X)}E_\pi$$

with E_π effective since X is terminal. So we always have

$$\lceil m\pi^*(K_X) \rceil := \lceil mK_{X'} - \frac{m}{r(X)}E_\pi \rceil \leq mK_{X'}$$

for any integer $m > 0$. Denote by M_{m_0} the movable part of $|m_0K_{X'}|$. One has

$$m_0\pi^*(K_X) = M_{m_0} + E'_{m_0}$$

for an effective \mathbb{Q} -divisor E'_{m_0} . In total, since

$$h^0(X', \lceil m_0\pi^*(K_X) \rceil) = h^0(X', \lceil m_0\pi^*(K_X) \rceil) = P_{m_0}(X') = P_{m_0}(X),$$

one has

$$m_0K_{X'} = M_{m_0} + (E'_{m_0} + \frac{m_0}{r(X)}E_\pi)$$

where $E'_{m_0} + \frac{m_0}{r(X)}E_\pi$ is exactly the fixed part of $|m_0K_{X'}|$.

If $\dim(\Gamma) \geq 2$, a general member S of $|M_{m_0}|$ is a nonsingular projective surface of general type by Bertini's theorem and by the easy addition formula for Kodaira dimension.

If $\dim(\Gamma) = 1$, a general fiber S of f is an irreducible smooth projective surface of general type, still by the easy addition formula for Kodaira dimension. We may write

$$M_{m_0} = \sum_{i=1}^{a_{m_0}} S_i \equiv a_{m_0}S$$

where S_i are smooth fibers of f for all i and $a_{m_0} \geq \min\{2P_{m_0} - 2, P_{m_0} + g(\Gamma) - 1\}$, by considering the degree of the divisor $f_*(M_0)$ on Γ .

Definition 2.4. We call S (in 2.3) a *generic irreducible element* of the linear system $|M_{m_0}|$. Denote by $\sigma : S \rightarrow S_0$ the blow-down onto the smooth minimal model S_0 . By abuse of concepts, we define a *generic irreducible element* of an arbitrary movable linear system on any projective variety in a similar way.

Definition 2.5. (1) Define the positive integer $p = p(m_0)$ as follows:

$$p = \begin{cases} 1 & \text{if } \dim(\Gamma) \geq 2, \\ a_{m_0} & \text{if } \dim(\Gamma) = 1. \end{cases}$$

(2) To simplify our statements, we say that the fibration f is of type III (resp. II, I) if $\dim \Gamma = 3$ (resp. 2, 1). According to our needs, we

would like to divide type I into subclasses:

$$f \text{ is of type } \begin{cases} I_q & \text{if } g(\Gamma) > 0, \\ I_3 & \text{if } g(\Gamma) = 0, P_{m_0} \geq 3, \\ I_p & \text{if } g(\Gamma) = 0, p_g(S) > 0, \\ I_n & \text{if } g(\Gamma) = 0, p_g(S) = 0. \end{cases}$$

2.6. Invariants of the fibration. Let V be a smooth projective 3-fold and $f : V \rightarrow \Gamma$ a fibration onto a nonsingular curve Γ . Leray spectral sequence tells that

$$E_2^{p,q} := H^p(\Gamma, R^q f_* \omega_V) \implies E^n := H^n(V, \omega_V).$$

By Serre duality and [15, Corollary 3.2, Proposition 7.6], one has the torsion-freeness of the sheaves $R^i f_* \omega_V$ and the following formulae:

$$\begin{aligned} h^2(\mathcal{O}_V) &= h^1(\Gamma, f_* \omega_V) + h^0(\Gamma, R^1 f_* \omega_V), \\ q(V) &:= h^1(\mathcal{O}_V) = g(\Gamma) + h^1(\Gamma, R^1 f_* \omega_V). \end{aligned}$$

2.7. Birationality principles. Let Y be a nonsingular projective variety on which there are two divisors D and M . Assume that $|M|$ is base point free. Take the Stein factorization of $\Phi_{|M|} : Y \xrightarrow{f} W \rightarrow \mathbb{P}^{h^0(Y,M)-1}$ where f is a fibration onto a normal variety W . Then the rational map $\Phi_{|D+M|}$ is birational onto its image if one of the following conditions is satisfied:

- (i) ([24, Lemma 2]) $\dim \Phi_{|M|}(Y) \geq 2$, $|D| \neq \emptyset$ and $\Phi_{|D+M|}|_S$ is birational for a general member S of $|M|$.
- (ii) ([6, §2.1]) $\dim \Phi_{|M|}(Y) = 1$, $\Phi_{|D+M|}$ can separate different general fibers of f and $\Phi_{|D+M|}|_F$ is birational for a general fiber F of f .

Remark 2.8. For the condition 2.7(ii), one knows that $\Phi_{|D+M|}$ can separate different general fibers of f whenever $\dim \Phi_{|M|}(Y) = 1$, W is a rational curve and D is an effective divisor. (In fact, since $|M|$ can separate different fibers of f , so can $|D + M|$.)

2.9. Assumptions. Let m be a positive integer. Let $|G|$ be a base point free linear system on S . Denote by C a generic irreducible element of $|G|$. Assume:

- (1) The linear system $|mK_{X'}|$ distinguishes different generic irreducible elements of $|M_{m_0}|$ (namely, $\Phi_{|mK_{X'}|}(S') \neq \Phi_{|mK_{X'}|}(S'')$ for two different generic irreducible elements S', S'' of $|M_{m_0}|$).
- (2) The linear system $|mK_{X'}|_S$ on S (as a sub-linear system of $|mK_{X'}|$) distinguishes different generic irreducible elements of $|G|$. (Or sufficiently, the complete linear system

$$|K_S + [(m - 1)\pi^*(K_X) - S - \frac{1}{p}E'_{m_0}]_S|$$

distinguishes different generic irreducible elements of $|G|$.)

2.10. A lower bound of K^3 . We keep the same notation as above. Since π^*K_X is nef and big, there is a rational number $\beta > 0$ such that $\pi^*(K_X)|_S - \beta C$ is numerically equivalent to an effective \mathbb{Q} -divisor on S .

We further define the following quantities:

$$\begin{aligned} \xi &:= (\pi^*(K_X) \cdot C)_{X'}; \\ \alpha &:= (m - 1 - \frac{m_0}{p} - \frac{1}{\beta})\xi; \\ \alpha_0 &:= \lceil \alpha \rceil. \end{aligned}$$

One has

$$K^3 \geq \frac{p}{m_0} \pi^*(K_X)^2 \cdot S \geq \frac{p\beta}{m_0} (\pi^*(K_X) \cdot C) = \frac{p\beta}{m_0} \xi. \tag{2.1}$$

So it is essential to estimate the rational number $\xi := (\pi^*(K_X) \cdot C)_{X'}$ in order to obtain the lower bound of K^3 . We recall the following:

Theorem 2.11. ([8, Theorem 3.1]) *Keep the notation as above. The inequality*

$$\xi \geq \frac{\deg(K_C) + \alpha_0}{m}$$

holds if one of the following conditions is satisfied:

- (i) $\alpha > 1$;
- (ii) $\alpha > 0$ and C is an even divisor, i.e., $C \sim 2H$ for a divisor H on S .

Furthermore, under Assumptions 2.9(1) and 2.9(2), the map $\varphi_m := \Phi_{|mK_{X'}|}$ is birational onto its image if one of the following conditions is satisfied:

- (i) $\alpha > 2$;
- (ii) $\alpha \geq 2$ and C is not a hyper-elliptic curve on S .

Remark 2.12. In particular, the inequality $\xi \geq \frac{\deg(K_C) + \alpha_0}{m}$ in Theorem 2.11 implies

$$\xi \geq \frac{\deg(K_C)}{1 + \frac{m_0}{p} + \frac{1}{\beta}} \tag{2.2}$$

since, whenever m is big enough so that $\alpha > 1$,

$$m\xi \geq \deg(K_C) + \alpha_0 \geq \deg(K_C) + (m - 1 - \frac{m_0}{p} - \frac{1}{\beta})\xi.$$

As long as we have fixed a linear system $|G|$ on S , we are able to prove the effective non-vanishing of plurigenera as follows.

Proposition 2.13. *Assume $P_{m_0} \geq 2$ for some positive integer m_0 . Then $P_m(X) > 1$ for all integers $m > 1 + \frac{m_0}{p} + \frac{1}{\beta}$. In particular, $\rho_0 \leq \rho_1 \leq \lfloor 2 + \frac{m_0}{p} + \frac{1}{\beta} \rfloor$.*

Proof. Assume $m > 1 + \frac{m_0}{p} + \frac{1}{\beta}$. Keep the same notation as in 2.3. Put

$$\mathcal{L}_m := (m-1)\pi^*(K_X) - \frac{1}{p}E'_{m_0}.$$

Then we have $|K_{X'} + \lceil \mathcal{L}_m \rceil| \subset |mK_{X'}|$. Noting that

$$\mathcal{L}_m - S \equiv (m-1 - \frac{m_0}{p})\pi^*(K_X)|_S$$

is nef and big, the Kawamata-Viehweg vanishing theorem ([13, 26]) yields the surjective map

$$H^0(X', K_{X'} + \lceil \mathcal{L}_m \rceil) \rightarrow H^0(S, (K_{X'} + \lceil \mathcal{L}_m \rceil)|_S). \quad (2.3)$$

Since S is a generic irreducible element of a free linear system, one has $\lceil * \rceil|_S \geq \lceil *_S \rceil$ for any divisor $*$ on X' . It follows that

$$(K_{X'} + \lceil \mathcal{L}_m \rceil)|_S \geq K_{X'}|_S + \lceil \mathcal{L}_m|_S \sim K_S + \lceil (\mathcal{L}_m - S)|_S \rceil. \quad (2.4)$$

Note that there is an effective \mathbb{Q} -divisor \hat{H} on S such that $\frac{1}{\beta}\pi^*(K_X)|_S \equiv C + \hat{H}$. We consider

$$\mathcal{D}_m := (\mathcal{L}_m - S)|_S - \hat{H}$$

on S . Then, by assumption, the divisor $\mathcal{D}_m - C \equiv (m-1 - \frac{m_0}{p} - \frac{1}{\beta})\pi^*(K_X)|_S$ is nef and big. Thus the Kawamata-Viehweg vanishing theorem again gives the surjective map

$$H^0(S, K_S + \lceil \mathcal{D}_m \rceil) \longrightarrow H^0(C, K_C + D), \quad (2.5)$$

where $D := \lceil \mathcal{D}_m - C \rceil|_C$ is a divisor on C . Because C is a generic irreducible element of a free linear system, we have $D \geq \lceil (\mathcal{D}_m - C)|_C \rceil$. A simple calculation gives

$$\deg(D) \geq (\mathcal{D}_m - C) \cdot C = (m-1 - \frac{m_0}{p} - \frac{1}{\beta})\xi = \alpha > 0.$$

Noting that $g(C) \geq 2$ since S is of general type, Riemann-Roch formula on C gives $h^0(C, K_C + D) \geq 2$. Finally, surjective maps (2.3), (2.5) and inequality (2.4) imply the statement. q.e.d.

We need the following lemma while studying type I_p , I_n , and I_3 cases.

Lemma 2.14. *Let S be a non-singular projective surface of general type. Denote by $\sigma : S \rightarrow S_0$ the blow-down onto its minimal model S_0 . Let Q be a \mathbb{Q} -divisor on S . Then $h^0(S, K_S + \lceil Q \rceil) \geq 2$ under one of the following conditions:*

- (i) $p_g(S) > 0$, $Q \equiv \sigma^*(K_{S_0}) + Q_1$ for some nef and big \mathbb{Q} -divisor Q_1 on S ;
- (ii) $p_g(S) = 0$, $Q \equiv 2\sigma^*(K_{S_0}) + Q_2$ for some nef and big \mathbb{Q} -divisor Q_2 on S .

Proof. First of all, $h^0(S, 2K_S) = h^0(S, 2K_{S_0}) > 0$ by the Riemann-Roch theorem on S , which is a surface of general type. Fix an effective divisor $R_0 \sim l\sigma^*(K_{S_0})$, where $l = 1, 2$ in cases (i) and (ii), respectively. Then R_0 is nef and big and R_0 is 1-connected by [17, Lemma 2.6]. The Kawamata-Viehweg vanishing theorem says $H^1(S, K_S + [Q] - R_0) = 0$, which gives the surjective map

$$H^0(S, K_S + [Q]) \longrightarrow H^0(R_0, K_{R_0} + G_{R_0})$$

where $G_{R_0} := ([Q] - R_0)|_{R_0}$ with $\deg(G_{R_0}) \geq (Q - R_0)R_0 = Q_l \cdot R_0 > 0$. The 1-connectedness of R_0 allows us to utilize the Riemann-Roch (see [1], Chapter II) as in the usual way. Note that S is of general type. So $K_{S_0}^2 > 0$ and $\deg(K_{R_0}) = 2p_a(R_0) - 2 = (K_S + R_0)R_0 \geq 2$. By the Riemann-Roch theorem on the 1-connected curve R_0 , we have

$$h^0(R_0, K_{R_0} + G_{R_0}) \geq \deg(K_{R_0} + G_{R_0}) + 1 - p_a(R_0) \geq p_a(R_0) \geq 2.$$

Hence $h^0(S, K_S + [Q]) \geq 2$. q.e.d.

Proposition 2.15. *Assume $P_{m_0} \geq 2$ for some positive integer m_0 . Then $P_m \geq 2$ for $m \geq h(m_0)$ under one of the following situations:*

- (i) $h(m_0) = 2m_0 + 3$ when f is of type I_p ;
- (ii) $h(m_0) = 3m_0 + 4$ when f is of type I_n ;
- (iii) $h(m_0) = \lfloor \frac{3m_0}{2} \rfloor + 4$ when f is of type I_3 .

In particular, $\rho_0 \leq \rho_1 \leq 2m_0 + 3, 3m_0 + 4, \lfloor \frac{3m_0}{2} \rfloor + 4$, respectively.

Proof. Keep the same notation as in 2.3. When f is of type I, we have $p = a_{m_0}$. By [8, Lemma 3.3], there is a sequence of rational numbers $\{\hat{\beta}_n\}$ with $\hat{\beta}_n \mapsto \frac{p}{m_0+p} \geq \frac{1}{m_0+1}$ such that

$$\pi^*(K_X)|_S - \hat{\beta}_n\sigma^*(K_{S_0}) \equiv H_n$$

for an effective \mathbb{Q} -divisors H_n .

We consider

$$\mathcal{D}'_m := (\mathcal{L}_m - S)|_S - (m - 1 - \frac{m_0}{p})H_n \equiv (m - 1 - \frac{m_0}{p})\hat{\beta}_n\sigma^*(K_{S_0}).$$

If, for $m > 0$, $h^0(S, K_S + [\mathcal{D}'_m]) \geq 2$, then $h^0(S, K_S + [(\mathcal{L}_m - S)|_S]) \geq 2$. It follows then that $P_m \geq 2$ by surjective map (2.3) and inequality (2.4). We can choose $h(m_0)$ according to the type of f .

When f is of type I_p , we can pick a big number n so that $\hat{\beta}_n \geq \frac{1}{m_0+1} - \delta$ for some $0 < \delta \ll 1$. For $m \geq 2m_0 + 3$, we see $(m - 1 - \frac{m_0}{p})\hat{\beta}_n > 1$. By Lemma 2.14 and since $p_g(S) > 0$, we know $h^0(S, K_S + [\mathcal{D}'_m]) \geq 2$. Thus we may take $h(m_0) = 2m_0 + 3$.

When f is of type I_n , we still take a big number n so that $\hat{\beta}_n \geq \frac{1}{m_0+1} - \delta$ for some $0 < \delta \ll 1$. But, for $m \geq 3m_0 + 4$, we have $(m - 1 - \frac{m_0}{p})\hat{\beta}_n > 2$. By Lemma 2.14 again, we see $h^0(S, K_S + [\mathcal{D}'_m]) \geq 2$. Thus we may take $h(m_0) = 3m_0 + 4$.

Finally, when f is of type I_3 , we have $p \geq 2$. One may take a big number n so that $\hat{\beta}_n \geq \frac{2}{m_0+2} - \delta$ for some $0 < \delta \ll 1$. For $m \geq \lfloor \frac{3m_0}{2} \rfloor + 4$, we have $(m-1 - \frac{m_0}{p})\hat{\beta}_n > 2$. Lemma 2.14 implies $h^0(S, K_S + \lceil \mathcal{D}'_m \rceil) \geq 2$. Thus we may take $h(m_0) = \lfloor \frac{3m_0}{2} \rfloor + 4$. This completes the proof. q.e.d.

Lemma 2.16. *Assume $P_{m_0}(X) \geq 2$ for some positive integer m_0 . Keep the same notation as in 2.3. Then, for $m \geq \rho_0 + m_0$, Assumptions 2.9 (1) is satisfied if f is of type III, II, I_3 , I_p , or I_n .*

Proof. Let $t > 0$ be an integer. We consider the linear system $|K_{X'} + [t\pi^*(K_X)] + M_{m_0}| \subset |(m_0 + t + 1)K_{X'}|$. Since $K_{X'} + [t\pi^*(K_X)] \geq (t + 1)\pi^*(K_X)$, we see that $K_{X'} + [t\pi^*(K_X)]$ is effective whenever $t + 1 \geq \rho_0$.

When f is of type I_3 , I_p or I_n , we necessarily have $g(\Gamma) = 0$. Thus, by [24, Lemma 2] and Remark 2.8, the linear system $|K_{X'} + [t\pi^*(K_X)] + M_{m_0}|$ can separate different generic irreducible elements S of $|M_{m_0}|$. q.e.d.

Lemma 2.17. *Let T be a non-singular projective surface of general type on which there is a base point free linear system $|G|$. Let Q be an arbitrary \mathbb{Q} -divisor on T . Then the linear system $|K_T + \lceil Q \rceil + G|$ can distinguish different generic irreducible elements of $|G|$ under one of the following conditions:*

- (i) $K_T + \lceil Q \rceil$ is effective and $|G|$ is not composed with an irrational pencil of curves;
- (ii) Q is nef and big and $|G|$ is composed with an irreducible pencil of curves.

Proof. Statement (i) follows from [24, Lemma 2] and Remark 2.8.

For statement (ii), we pick up a generic irreducible element C of $|G|$. Then $G \equiv sC$ where $s \geq 2$ and $C^2 = 0$. Let C' be another generic irreducible element. The Kawamata-Viehweg vanishing theorem gives the surjective map

$$H^0(T, K_T + \lceil Q \rceil + G) \longrightarrow H^0(C, K_C + D) \oplus H^0(C', K_{C'} + D')$$

where $D := (\lceil Q \rceil + G - C)|_C$ and $D' := (\lceil Q \rceil + G - C')|_{C'}$ with $\deg(D) > 0$, $\deg(D') > 0$. Since T is of general type, both C and C' are curves of genus ≥ 2 . Thus $h^0(C, K_C + D) = h^0(C', K_{C'} + D') > 1$. Thus $|K_T + \lceil Q \rceil + G|$ can distinguish C and C' . q.e.d.

Lemma 2.18. *Assume $P_{m_0}(X) \geq 2$ for some positive integer m_0 . Keep the same notation as in 2.3. Take $G := S|_S$ for a generic irreducible element S of $|M_{m_0}|$. Then Assumptions 2.9 (2) is satisfied under one of the following situations:*

- (i) f is of type III and $m \geq \rho_0 + m_0$.
- (ii) f is of type II and $m \geq \max\{\rho_0 + m_0, 2m_0 + 2\}$.

Proof. Since

$$\begin{aligned} & K_S + \lceil (m-1)\pi^*(K_X) - S - \frac{1}{p}E'_{m_0} \rceil_S \\ \geq & K_S + (m-1)\pi^*(K_X)|_S - (S + E'_{m_0})|_S \\ = & K_S + (m - m_0 - 1)\pi^*(K_X)|_S \\ \geq & (m - m_0)\pi^*(K_X)|_S + G \end{aligned}$$

and

$$\begin{aligned} & K_S + (m - m_0 - 1)\pi^*(K_X)|_S \\ \geq & K_S + (m - 2m_0 - 1)\pi^*(K_X)|_S + G, \end{aligned}$$

Lemma 2.17 implies that $|K_S + \lceil (m-1)\pi^*(K_X) - S - \frac{1}{p}E'_{m_0} \rceil_S|$ can distinguish different generic irreducible elements of $|G|$ respectively. Note that, if f is of type *III*, $|G|$ is not composed with a pencil of curves. We are done. q.e.d.

Under the condition $P_{m_0} \geq 2$, we study the pluricanonical map φ_m according to the type of f .

2.19. Type *III*.

When f is of type *III*, we have $p = 1$ by definition. In this case, $S \sim M_{m_0}$ and $|S|$ gives a generically finite morphism. We take $G := S|_S$. Then $|G|$ is base point free and $\varphi_{|G|}$ gives a generically finite map. So a generic irreducible element $C \sim G$ is a smooth curve.

If $\varphi_{|G|}$ gives a birational map, then $\dim \varphi_{|G|}(C) = 1$ for a general member C . The Riemann-Roch and Clifford's theorem on C says $C^2 = G \cdot C \geq 2$. If $\varphi_{|G|}$ gives a generically finite map of degree ≥ 2 , since $h^0(S, G) \geq h^0(X', S) - 1 \geq 3$, one gets $C^2 \geq 2(h^0(S, G) - 2) \geq 2$. Either way, we have $C^2 \geq 2$. So $\deg(K_C) = (K_S + C) \cdot C > 2C^2 \geq 4$. We see $\deg(K_C) \geq 6$ since it is an even number.

One may take $\beta = \frac{1}{m_0}$ since $m_0\pi^*(K_X)|_S \geq C$.

Now inequality (2.2) gives $\xi \geq \frac{6}{2m_0+1}$. Take $m = 3m_0 + 2$. Then $\alpha = (m - 2m_0 - 1)\xi > 3$. So, by Theorem 2.11, $\xi \geq \frac{10}{3m_0+2}$. It follows from inequality (2.1) that $K^3 \geq \frac{10}{(3m_0+2)m_0^2}$.

We now consider the non-vanishing of plurigenera. By Proposition 2.13, we have $P_m \geq 2$ for all $m > 2m_0 + 1$. Now, if $m = 2m_0 + 1$, the surjective map (2.3) and inequality (2.4) lead us to compute $h^0(S, K_S + \lceil m_0\pi^*K_X|_S \rceil)$. Let L be a generic irreducible element in $|S|_S$. Then L is effective and nef. Since $h^2(K_S + L) = 0$, one has $h^0(S, K_S + L) \geq \chi(S, K_S + L) = \frac{1}{2}(K_S \cdot L + L^2) + \chi(\mathcal{O}_S) \geq 2$ by Riemann-Roch theorem. Hence $P_{2m_0+1} \geq 2$. Also, $P_{2m_0} \geq P_{m_0} \geq 2$. Therefore, we have $P_m > 1$ for all $m \geq 2m_0$. In particular, $\rho_0 \leq \rho_1 \leq 2m_0$.

By Lemmas 2.16 and 2.18, Assumptions 2.9(1) and 2.9(2) are satisfied if $m \geq 3m_0$. Now $\alpha = (m - 2m_0 - 1)\xi \geq (m - 2m_0 - 1)\frac{10}{3m_0+2}$. One sees

that $\alpha > 2$ if $m > \frac{13m_0+7}{5}$. Hence φ_m is birational if

$$m > \max\{3m_0 - 1, \frac{13m_0 + 7}{5}\}.$$

We conclude the following:

Theorem 2.20. *Assume $P_{m_0}(X) \geq 2$ for some positive integer m_0 . If the induced map f is of type III. Then:*

- 1) $\rho_0 \leq \rho_1 \leq 2m_0$.
- 2) $K^3 \geq \frac{10}{(3m_0+2)m_0^2}$.
- 3) φ_m is birational if $m > \max\{3m_0 - 1, \frac{13m_0+7}{5}\}$.

2.21. Type II.

When f is of type II, we see that $S \sim M_{m_0}$. Take $|G| := |S|_S$, which is, clearly, composed with a pencil of curves.

Since a generic irreducible element C of $|G|$ is a smooth curve of genus ≥ 2 , we have $\deg(K_C) \geq 2$. Furthermore, we have $h^0(S, G) \geq h^0(X', S) - 1 \geq 2$. So $G \equiv \tilde{a}C$ where $\tilde{a} \geq h^0(S, G) - 1 \geq 1$. This means that $m_0\pi^*(K_X)|_S \geq S|_S \geq_{\text{num}} C$. So we may take $\beta = \frac{1}{m_0}$.

Now inequality (2.2) gives $\xi \geq \frac{2}{2m_0+1}$. Take $m = 3m_0 + 2$. Then $\alpha > 1$. One gets $\xi \geq \frac{4}{3m_0+2}$ by Theorem 2.11. So inequality (2.1) implies $K^3 \geq \frac{4}{(3m_0+2)m_0^2}$.

Exactly the same proof as in Type III shows that $\rho_0 \leq \rho_1 \leq 2m_0$.

By Lemmas 2.16 and 2.18, Assumptions 2.9(1) and 2.9(2) are satisfied if $m \geq 3m_0$. Now $\alpha = (m - 2m_0 - 1)\xi \geq (m - 2m_0 - 1)\frac{4}{3m_0+2}$. One sees that $\alpha > 2$ if $m > \frac{7m_0+4}{2}$. Since $\frac{7m_0+4}{2} > 3m_0$, φ_m is birational if $m > \frac{7m_0+4}{2}$.

We conclude the following:

Theorem 2.22. *Assume $P_{m_0}(X) \geq 2$ for some positive integer m_0 . If the induced map f is of type II, then:*

- 1) $\rho_0 \leq \rho_1 \leq 2m_0$.
- 2) $K^3 \geq \frac{4}{(3m_0+2)m_0^2}$.
- 3) φ_m is birational if $m > \frac{7m_0+4}{2}$.

2.23. Type I_q .

Since $g(\Gamma) > 0$, one sees $q(X) > 0$ and hence X is irregular. This case is particularly well-behaved. It's known that φ_m is birational for all $m \geq 7$ (see [3]). Also $K_X^3 \geq \frac{1}{22}$ (see [5]).

2.24. Type I_p .

We have an induced fibration $f : X' \rightarrow \Gamma$ with $g(\Gamma) = 0$. By definition, $p = a_{m_0} \geq 1$. By assumption, $p_g(S) > 0$ for a general fiber S of f . We take $G := 2\sigma^*(K_{S_0})$. Then one knows that $|G|$ is base point free (see [9, Theorem 3.1]). Thus $|G|$ is not composed with a pencil and

a generic irreducible element C is smooth. By [8, Lemma 3.3], we can find a sequence of rational numbers $\{\beta_n\}$ with $\beta_n \mapsto \frac{p}{m_0+p}$ such that $\pi^*(K_X)|_S - \frac{\beta_n}{2}C \equiv H_n$ for effective \mathbb{Q} -divisors H_n . We may assume that $\beta \geq \frac{1}{2(m_0+1)} - \delta$ for some $0 < \delta \ll 1$.

Since $C \sim 2\sigma^*(K_{S_0})$,

$$\deg(K_C) = (K_S + C) \cdot C \geq (\pi^*(K_X)|_S + C) \cdot C > C^2 \geq 4.$$

Since $\deg(K_C)$ is even, we see $\deg(K_C) \geq 6$.

Now inequality (2.2) gives $\xi \geq \frac{6}{3m_0+3}$. Take $m = 4m_0 + 5$. Then $\alpha = (m - 1 - m_0 - \frac{1}{\beta})\xi > 2$ and Theorem 2.11 gives $\xi \geq \frac{9}{4m_0+5}$. So, by inequality (2.1), one gets $K^3 \geq \frac{9}{2m_0(m_0+1)(4m_0+5)}$.

Note that

$$\begin{aligned} & K_S + [(m - 1)\pi^*(K_X) - S - \frac{1}{p}E'_{m_0}]|_S \\ & \geq K_S + [(m - m_0 - 1)\pi^*(K_X)|_S] \\ & \geq K_S + [(m - m_0 - 1)\pi^*(K_X)|_S - \frac{3}{\beta_n}H_n] \\ & = K_S + [Q_1] + \sigma^*(K_{S_0}) + C \end{aligned} \tag{2.6}$$

where $Q_1 := (m - m_0 - 1)\pi^*(K_X)|_S - C - \sigma^*(K_{S_0}) - \frac{3}{\beta_n}H_n \equiv (m - m_0 - 1 - \frac{3}{\beta_n})\pi^*(K_X)|_S$ is nef and big whenever $m \geq 4m_0 + 5$. By Lemma 2.14(i), $K_S + [Q_1] + \sigma^*(K_{S_0})$ is effective. Thus, according to [24, Lemma 2], Assumption 2.9 (2) is satisfied for $m \geq 4m_0 + 5$. Since Proposition 2.15 (ii) implies $\rho_0 \leq 2m_0 + 3$, Lemma 2.16 (ii) tells that Assumption 2.9(1) is satisfied as long as $m \geq 3m_0 + 3$. Take $m \geq 4m_0 + 5$. Then $\alpha \geq (m - 3m_0 - 3)\xi \geq \frac{2m_0+4}{m_0+1} > 2$. So Theorem 2.11 implies that φ_m is birational for all $m \geq 4m_0 + 5$.

We thus summarize:

Theorem 2.25. *Assume $P_{m_0}(X) \geq 2$ for some positive integer m_0 . If the induced map f is of type I_p , then:*

- 1) $\rho_0 \leq \rho_1 \leq 2m_0 + 3$.
- 2) $K^3 \geq \frac{9}{2m_0(m_0+1)(4m_0+5)}$.
- 3) φ_m is birational if $m \geq 4m_0 + 5$.

2.26. Type I_n .

Similar to the type I_p case, we have $p \geq 1$. We take $|G| := |4\sigma^*(K_{S_0})|$ which is base point free by a well-known result in [2]. Thus $|G|$ is not composed with a pencil and a generic irreducible element C is smooth. Similarly, we can find a sequence of rational numbers $\{\beta_n\}$ with $\beta_n \mapsto \frac{p}{m_0+p}$ such that $\pi^*(K_X)|_S - \frac{\beta_n}{4}C \equiv H_n$ for effective \mathbb{Q} -divisors H_n . We may assume that $\beta \geq \frac{1}{4(m_0+1)} - \delta$ for some $0 < \delta \ll 1$.

Since $\deg(K_C) > 16\sigma^*(K_{S_0})^2 \geq 16$ and $\deg(K_C)$ is even, inequality (2.2) gives $\xi \geq \frac{18}{5m_0+5}$. Take $m = 6m_0 + 6$. Then $\alpha = (m - 1 - m_0 - \frac{1}{\beta})\xi =$

$\frac{18}{5} > 3$ and Theorem 2.11 gives $\xi \geq \frac{11}{3m_0+3}$. So, by inequality (2.1), one gets $K^3 \geq \frac{11}{12m_0(m_0+1)^2}$.

By Proposition 2.15, we have $P_m \geq 2$ for all $m \geq 3m_0 + 4$. Thus we have the following:

Theorem 2.27. *Assume $P_{m_0}(X) \geq 2$ for some positive integer m_0 . If the induced map f is of type I_n , then:*

- 1) $\rho_0 \leq \rho_1 \leq 3m_0 + 4$.
- 2) $K^3 \geq \frac{11}{12m_0(m_0+1)^2}$.
- 3) φ_m is birational if $m \geq 5m_0 + 6$ (cf. [7, Theorem 0.1]).

2.28. Type I_3 .

We take $G_1 = 4\sigma^*(K_{S_0})$ so as to estimate K_X^3 . Then, as seen in 2.26, $\deg(K_C) \geq 18$. Being in a better situation with $p = a_{m_0} - 1 \geq 2$, a better number β can be found. In fact, by [8, Lemma 3.3], one may take a number sequence $\{\beta_n\}$ with $\beta_n \mapsto \frac{p}{4(m_0+p)} \geq \frac{1}{2(m_0+2)}$ such that $\pi^*(K_X)|_S - \beta_n C$ is numerically equivalent to an effective \mathbb{Q} -divisor. Namely, one may take a number $\beta \geq \frac{1}{2(m_0+2)} - \delta$ for some $0 < \delta \ll 1$. Now inequality (2.2) gives $\xi \geq \frac{18}{1 + \frac{m_0+1}{2} + \beta}$, i.e., $\xi \geq \frac{36}{5(m_0+2)}$ by taking the limit. Hence inequality (2.1) implies $K^3 \geq \frac{36}{5m_0(m_0+2)^2}$.

We take a different $|G|$ on S to study the birationality. In fact, we will take $|G|$ to be the movable part of $|2\sigma^*(K_{S_0})|$. A different point from previous ones is that $|G|$ is not always base point free. But since we have the induced fibration $f : X' \rightarrow \Gamma$, we can consider the relative bi-canonical map of f , namely, the rational map $\Psi : X' \dashrightarrow \mathbf{P}$ over Γ . First we can blow up the indeterminacy of Ψ on X' . Then we can assume, in the birational equivalence sense, that Ψ is a morphism over B . By further modifying π , we can even finally assume that π dominates Ψ . With this assumption (or by taking a sufficiently good π), we see that $|G|$ is base point free since $|G|$ gives the bicanonical morphism for each general fiber S of f .

By Proposition 2.15 and Lemma 2.16, Assumption 2.9(1) is satisfied for $m \geq \lfloor \frac{5m_0}{2} \rfloor + 4$. Recall that we have $p = a_{m_0} \geq 2$.

Claim A. Assumption 2.9(2) is satisfied for $m \geq \min\{3m_0 + 6, \rho_0 + 2m_0 + 2\}$.

In fact, the argument of 2.24 works here. A different place is that we have a better bound for β_n since $p \geq 2$, but we only have $\deg(K_C) \geq 2$. By [8, Lemma 3.3], we can find a sequence of rational numbers $\{\beta_n\}$ with $\beta_n \mapsto \frac{p}{2(m_0+p)}$ such that $\pi^*(K_X)|_S - \beta_n(2\sigma^*(K_{S_0})) \equiv H_n$ for effective \mathbb{Q} -divisors H_n . We may assume that $\beta \geq \frac{1}{m_0+2} - \delta$ for some $0 < \delta \ll 1$.

Now the last three terms of inequality (2.6) can be replaced by

$$\begin{aligned} & K_S + \lceil (m - m_0 - 1)\pi^*(K_X)|_S \rceil \\ \geq & K_S + \lceil (m - m_0 - 1)\pi^*(K_X)|_S - \frac{2}{\beta_n}H_n \rceil \\ = & K_S + \lceil Q_2 \rceil + 4\sigma^*(K_{S_0}) \end{aligned}$$

where $Q_2 := (m - m_0 - 1)\pi^*(K_X)|_S - 4\sigma^*(K_{S_0}) - \frac{2}{\beta_n}H_n \equiv (m - m_0 - 1 - \frac{2}{\beta_n})\pi^*(K_X)|_S$ is nef and big whenever $m \geq 3m_0 + 6$. According to a theorem of Xiao [28], $|G|$ is either not composed with a pencil or composed with a rational pencil. Thus, according to [24, Lemma 2] and Remark 2.8, Assumption 2.9(2) is satisfied for $m \geq 3m_0 + 6$. On the other hand, we have an inclusion, $\mathcal{O}_\Gamma(2) \hookrightarrow f_*\omega_{X'}^{m_0}$, which naturally gives rise to the inclusion $f_*\omega_{X'/\Gamma}^2 \hookrightarrow f_*\omega_{X'}^{2m_0+2}$. Now Viehweg's semi-positivity theorem [27] implies that $f_*\omega_{X'/\Gamma}^2$ is generated by global sections. Thus $\lceil (2m_0 + 2)K_{X'}|_S \rceil$ can distinguish different generic irreducible elements of $|G|$. So Assumption 2.9(2) is naturally satisfied for all $m \geq \rho_0 + 2m_0 + 2$. We have proved Claim A.

Finally, we consider the value of α . Recall that we may take $\beta \mapsto \frac{p}{2m_0+2p} \geq \frac{1}{m_0+2}$. Inequality (2.2) gives $\xi \geq \frac{2}{1+\frac{m_0}{2}+m_0+2} = \frac{4}{3(m_0+2)}$. If we take $m = 3m_0 + 4$. Then $\alpha > 1$. Theorem 2.11 says $\xi \geq \frac{4}{3m_0+4}$. Eventually, take $m \geq 3m_0 + 6$. Then $\alpha > 2$. Theorem 2.11 implies that φ_m is birational for all $m \geq 3m_0 + 6$.

We thus conclude the following:

Theorem 2.29. *Assume $P_{m_0}(X) \geq 3$ for some positive integer m_0 . If the induced map f is of type I_3 , then:*

- 1) $\rho_0 \leq \rho_1 \leq \lfloor \frac{3m_0}{2} \rfloor + 4$.
- 2) $K^3 \geq \frac{36}{5m_0(m_0+2)^2}$.
- 3) φ_m is birational if $m \geq 3m_0 + 6$.

By collecting all above results, we have the following:

Corollary 2.30. *Assume $P_{m_0}(X) \geq 2$ for some positive integer m_0 . Then $K^3 \geq \frac{11}{12m_0(m_0+1)^2}$.*

2.31. Volume optimization.

Indeed, when m_0 is small, the estimation of K_X^3 could be optimized by recursively applying Theorem 2.11 with a suitable m .

For example, suppose $m_0 = 11$ and f is of type III . Then inequality (2.2) gives $\xi \geq \frac{6}{23}$. Take $m = 27$. By Theorem 2.11, we get $\xi \geq \frac{8}{27}$. So inequality (2.1) gives $K^3 \geq \frac{8}{3267} > \frac{10}{m_0^2(3m_0+2)}$.

Let's consider another example with $m_0 = 8$ and f being of type II . Then we may take $\beta = \frac{1}{8}$. Inequality (2.2) gives $\xi \geq \frac{2}{17}$. Take $m = 26$. Then $\alpha \geq \frac{18}{17} > 1$. Theorem 2.11 gives $\xi \geq \frac{2}{13}$. Take $m = 24$. Then

$\alpha > 1$. Again, one gets $\xi \geq \frac{1}{6}$. So inequality (2.1) implies $K^3 \geq \frac{1}{384} > \frac{4}{m_0^2(3m_0+2)}$.

With the idea mentioned above, a patient reader should have no difficulty to check the following table on the lower bound of K^3 for small m_0 .

Table A

m_0	2	3	4	5	6	7
<i>III</i>	1/3	8/81	1/22	8/325	1/72	4/441
<i>II</i>	1/8	2/45	1/52	1/100	1/162	4/1029
$P_{m_0} \geq 3$	1/8	2/45	1/52	1/100	1/162	4/1029
$P_{m_0} \geq 2$	5/96	5/264	1/108	1/192	5/1554	5/2408
m_0	8	9	10	11	12	
<i>III</i>	1/160	4/891	2/625	8/3267	1/522	
<i>II</i>	1/384	2/1053	1/725	1/968	1/1224	
$P_{m_0} \geq 3$	1/384	2/1053	1/725	1/968	1/1224	
$P_{m_0} \geq 2$	5/3456	1/954	1/1276	5/8448	5/10764	

Lemma 2.32. *If f is of type I_n and $q(X) = 0$, then $\chi(\mathcal{O}_X) \leq 1$.*

Proof. We have an induced fibration $f : X' \rightarrow \Gamma$ onto the rational curve Γ . A general fiber S of f is a non-singular projective surface of general type with $p_g(S) = 0$. Because $\chi(\mathcal{O}_S) > 0$, we see $q(S) = 0$. This means $f_*\omega_{X'} = 0$ and $R^1f_*\omega_{X'} = 0$ since they are both torsion free by [15]. Thus we get by 2.6 the following formulae:

$$h^2(\mathcal{O}_X) = h^2(\mathcal{O}_{X'}) = h^1(f_*\omega_{X'}) + h^0(R^1f_*\omega_{X'}) = 0;$$

$$q(X) = q(X') = g(\Gamma) + h^1(R^1f_*\omega_{X'}) = 0.$$

So we see $\chi(\mathcal{O}_X) = 1 - q(X) + h^2(\mathcal{O}_X) - p_g(X) \leq 1$. q.e.d.

2.33. Miyaoka-Reid inequality on $\mathcal{B}(X)$. We refer to [4, Section 2] for the definition of baskets. Assume that Reid’s basket of singularities on X is $B_X := \mathcal{B}(X) = \{(b_i, r_i)\}$. According to [21, 10.3], one has

$$\frac{1}{12}K_X \cdot c_2(X) = -2\chi(\mathcal{O}_X) + \sum_i \frac{r_i^2 - 1}{12r_i}$$

where $c_2(X)$ is defined via the intersection theory by taking a resolution of singularities of X . On the other hand, [18, Corollary 6.7] says $K_X \cdot c_2(X) \geq 0$. Thus one has the following inequality:

$$\sum_i r_i - 24\chi(\mathcal{O}_X) \geq \sum_i \frac{1}{r_i}. \tag{2.7}$$

A direct application of inequality (2.7) is the following:

Corollary 2.34. *Suppose that we have a packing between formal baskets $\mathbf{B} := (B, \chi(\mathcal{O}_X), P_2) \succcurlyeq \mathbf{B}' := (B', \chi(\mathcal{O}_X), \tilde{P}_2)$ and that inequality (2.7) fails for \mathbf{B}' . Then (2.7) fails for \mathbf{B} .*

3. General type 3-folds with $\chi = 1$

In this section, we always assume $\chi(\mathcal{O}_X) = 1$. If there is a small number m_0 such that $P_{m_0} > 1$, then one can detect the birational geometry of X by studying φ_{m_0} . Thus a natural question is what practical number m_0 can be found such that $P_{m_0} > 1$. This is exactly the motivation of this section. Equivalently, we shall give a complete classification of baskets to those X with $P_m \leq 1$ for $m \leq 6$.

3.1. Assumption: $P_m(X) \leq 1$ for $1 \leq m \leq 6$.

In fact, P_m satisfies the following geometric condition.

Lemma 3.2. *Assume $\chi(\mathcal{O}_X) = 1$. Then $P_{m+2} \geq P_m + P_2$ for all $m \geq 2$.*

Proof. By Reid’s formula ([21]), we have

$$P_{m+2} - P_m - P_2 = (m^2 + m)K_X^3 - \chi(\mathcal{O}_X) + (l(m + 2) - l(m) - l(2)).$$

By [11, Lemma 3.1], one sees $l(m + 2) - l(m) - l(2) \geq 0$. Since $K_X^3 > 0$ and $\chi(\mathcal{O}_X) = 1$, we have $P_{m+2} - P_m - P_2 > -1$. q.e.d.

We consider the formal basket

$$\mathbf{B} := (B, \chi(\mathcal{O}_X), P_2(X))$$

where $B = \mathcal{B}(X)$. As we have seen in [4, Section 3],

- (i) $K^3(\mathbf{B}) = K^3(B) = K_X^3 > 0$;
- (ii) $P_m(\mathbf{B}) = P_m(X)$ for all $m \geq 2$.

By Lemma 3.2, we see $P_4 \geq 2$ if $P_2 > 0$. Thus under Assumption 3.1, we have $P_2 = 0$. We can also get $P_{m+2} > 0$ whenever $P_m > 0$. Thus, in practice, we only need to study the following types: $P_2 = 0$ and

$$\begin{aligned} (P_3, P_4, P_5, P_6) = & (0, 0, 0, 0), (0, 0, 0, 1), (0, 0, 1, 0), (0, 0, 1, 1), \\ & (0, 1, 0, 1), (0, 1, 1, 1), (1, 0, 1, 1), (1, 1, 1, 1). \end{aligned} \tag{3.1}$$

Now we consider the formal basket $\mathbf{B} := (B, 1, 0)$. We might abuse the notation of baskets and formal baskets in this section for we always have $\chi = 1, P_2 = 0$ in this section. We keep the notation as in [4].

With explicit values of (P_3, P_4, P_5, P_6) , we are able to determine $\mathcal{B}^{(5)}(B)$ (cf. [4, Sections 3,4]). Our main task is to search all possible minimal (with regard to \succ) positive baskets B_{min} dominated by $\mathcal{B}^{(5)}(B)$. Take $\mathbf{B}^5 := (\mathcal{B}^{(5)}(B), 1, 0)$ and $\mathbf{B}_{min} := (B_{min}, 1, 0)$. Then we see $\mathbf{B}^5 \succ \mathbf{B} \succ \mathbf{B}_{min}$.

Now we classify all minimal positive geometric baskets B_{min} .

3.3. Case I: $P_3 = P_4 = P_5 = P_6 = 0$ (impossible)

We have $\sigma = 10, \tau = 4, \Delta^3 = 5, \Delta^4 = 14, \epsilon = 0, \sigma_5 = 0$, and $\epsilon_5 = 2$. The only possible initial basket is $\{5 \times (1, 2), 4 \times (1, 3), (1, 4)\}$. And

$B^{(5)} = \{3 \times (1, 2), 2 \times (2, 5), 2 \times (1, 3), (1, 4)\}$ with $K^3 = \frac{1}{60}$. We shall calculate B_{min} of $B^{(5)}$.

If we pack $\{(1, 2), (2, 5)\}$ into $\{(3, 7)\}$. Then we get:

I-1. $B_{1,1} = \{2 \times (1, 2), (3, 7), (2, 5), 2 \times (1, 3), (1, 4)\}$, $K^3 = \frac{1}{420}$, which admits no further prime packing into positive baskets. Hence $B_{1,1}$ is minimal positive.

We consider those baskets with $(1, 2)$ unpacked because otherwise it's dominated by $B_{1,1}$. So we consider the packing

$$\{3 \times (1, 2), (2, 5), (3, 8), (1, 3), (1, 4)\}$$

with $K^3 = \frac{1}{120}$. This basket allows two further packings to minimal positive ones:

I-2. $B_{1,2} = \{3 \times (1, 2), (2, 5), (4, 11), (1, 4)\}$, $K^3 = \frac{1}{220}$.

I-3. $B_{1,3} = \{3 \times (1, 2), (5, 13), (1, 3), (1, 4)\}$, $K^3 = \frac{1}{156}$.

Finally we consider the case that both $(1, 2)$ and $(2, 5)$ remain unpacked. We get one more basket which is indeed minimal positive

I-4. $B_{1,4} = \{3 \times (1, 2), 2 \times (2, 5), (1, 3), (2, 7)\}$, $K^3 = \frac{1}{210}$.

A direct calculation shows that none of $B_{1,1}$, $B_{1,2}$, $B_{1,3}$, and $B_{1,4}$ satisfy inequality (2.7). Hence B does not satisfy (2.7), a contradiction. This proves that Case I is impossible.

3.4. Case II: $P_3 = P_4 = P_5 = 0, P_6 = 1 (\Rightarrow B_{2,1}, B_{2,2})$

Now we have $\sigma = 10, \tau = 4, \Delta^3 = 5, \Delta^4 = 14, \epsilon \leq 1$. If $\epsilon = 0$, then $\epsilon_5 = 1$; if $\epsilon = 1$, then $\epsilon_5 = 0$. Thus all possible initial baskets and $B^{(5)}$ are as follows:

II-i. $B^{(0)} = \{5 \times (1, 2), 4 \times (1, 3), (1, 4)\} \succ B^{(5)} = \{4 \times (1, 2), (2, 5), 3 \times (1, 3), (1, 4)\}$, with $K^3(B^{(5)}) = \frac{1}{20}$.

II-ii. $B^{(0)} = \{5 \times (1, 2), 4 \times (1, 3), (1, 5)\} \succ B^{(5)} = \{5 \times (1, 2), 4 \times (1, 3), (1, 5)\}$, with $K^3(B^{(5)}) = \frac{1}{30}$.

In Case II-i, we first consider the situation that all single baskets $(1, 2)$ are packed into $\{(6, 13), 3 \times (1, 3), (1, 4)\}$, which gives a unique minimal positive basket:

II-1. $B_{2,1} = \{(6, 13), (1, 3), (3, 10)\}$, $K^3 = \frac{1}{390}, P_9 = 2, P_{13} = 3$.

We then consider the situation that at least one basket $(1, 2)$ remains unpacked. Then we get the following minimal positive basket:

II-2. $B_{2,2} = \{(1, 2), (5, 11), (4, 13)\}$, $K^3 = \frac{1}{286}, P_9 = 2, P_{13} = 3$.

Notice, however, that if $\{3 \times (1, 2), (3, 7), 3 \times (1, 3), (1, 4)\} \succ B$, then B dominates $B_{2,2}$. Thus it remains to consider the situation that all single baskets $(1, 2)$ are unpacked, but $(2, 5)$ must be packed with some $(1, 3)$. So we get the following minimal positive baskets:

II-3. $B_{2,3} = \{(4, 8), (3, 8), (3, 10)\}$, $K^3 = \frac{1}{40}$.

II-4. $B_{2,4} = \{(4, 8), (4, 11), (2, 7)\}$, $K^3 = \frac{2}{77}$.

II-5. $B_{2,5} = \{(4, 8), (5, 14), (1, 4)\}$, $K^3 = \frac{1}{28}$.

In Case II-ii, $B^{(5)}$ admits no further prime packing. Thus we get:

II-6. $B_{2,6} = \{(5, 10), (4, 12), (1, 5)\}$, $K^3 = \frac{1}{30}$.

One may check that $B_{2,3}, B_{2,4}, B_{2,5}, B_{2,6}$ do not satisfy inequality (2.7). Thus only **II-1** and **II-2** can happen.

3.5. Case III: $P_3 = P_4 = 0, P_5 = 1, P_6 = 0 (\Rightarrow B_{3,1} \sim B_{3,5})$

Now we have $\sigma = 10, \tau = 4, \Delta^3 = 5, \Delta^4 = 15$. Moreover, $P_7 \geq 1$, and hence $\epsilon = 0, \sigma_5 = 0$, and $\epsilon_5 = 4$. Thus the only possible initial basket and $B^{(5)}$ are

$$B^{(0)} = \{5 \times (1, 2), 5 \times (1, 3)\} \succ B^{(5)} = \{(1, 2), 4 \times (2, 5), (1, 3)\}.$$

So we get the following minimal positive baskets:

III-1. $B_{3,1} = \{(9, 22), (1, 3)\}$, $K^3 = \frac{1}{66}, P_9 = 2, P_{10} = 3$.

III-2. $B_{3,2} = \{(7, 17), (3, 8)\}$, $K^3 = \frac{1}{136}, P_{10} = 2, P_{12} = 3$.

III-3. $B_{3,3} = \{(5, 12), (5, 13)\}$, $K^3 = \frac{1}{156}, P_{10} = 2, P_{12} = 3$.

III-4. $B_{3,4} = \{(3, 7), (7, 18)\}$, $K^3 = \frac{1}{126}, P_{10} = 2, P_{12} = 3$.

III-5. $B_{3,5} = \{(1, 2), (9, 23)\}$, $K^3 = \frac{1}{46}, P_8 = 2, P_{10} = 4$.

3.6. Case IV: $P_3 = P_4 = 0, P_5 = 1, P_6 = 1 (\Rightarrow B_{3,1}, B_{3,2}, B_{3,4}, B_{3,5})$

Now we have $\sigma = 10, \tau = 4, \Delta^3 = 5, \Delta^4 = 15$. Moreover, the initial basket must have $n_{1,2}^0 = n_{1,3}^0 = 5$, and hence $n_{1,r}^0 = 0$ for all $r \geq 4$. It follows that $\epsilon = 0, \sigma_5 = 0$, and $\epsilon_5 = 3$. Thus the only possible initial basket and $B^{(5)}$ are

$$B^{(0)} = \{5 \times (1, 2), 5 \times (1, 3)\} \succ B^{(5)} = \{2 \times (1, 2), 3 \times (2, 5), 2 \times (1, 3)\}.$$

So we get the following minimal positive baskets:

IV-1. $\{(8, 19), (2, 6)\} \succ B_{3,1}$.

IV-2. $\{(6, 14), (4, 11)\} \succ B_{3,4}$.

IV-3. $\{(4, 9), (6, 16)\} \succ B_{3,2}$.

IV-4. $\{(2, 4), (8, 21)\} \succ B_{3,5}$.

3.7. Case V: $P_3 = 0, P_4 = 1, P_5 = 0, P_6 = 1. (\Rightarrow B_{5,1} \sim B_{5,3})$

We have $\sigma = 10, \tau = 4, \Delta^3 = 6, \Delta^4 = 13$, and $\sigma_5 \leq \epsilon \leq 2$. The initial baskets have 4 types:

V-i. $\{6 \times (1, 2), (1, 3), 3 \times (1, 4)\}$;

V-ii. $\{6 \times (1, 2), (1, 3), 2 \times (1, 4), (1, 5)\}$;

V-iii. $\{6 \times (1, 2), (1, 3), (1, 4), 2 \times (1, 5)\}$;

V-iv. $\{6 \times (1, 2), (1, 3), 2 \times (1, 4), (1, r)\}$ with $r \geq 6$.

Cases V-iii and V-iv are impossible since $K^3 \leq 0$. For Case V-i, we have $\epsilon_5 = 1$, and for V-ii, we have $\epsilon_5 = 0$. Hence $B^{(5)}$ has two possibilities, correspondingly:

- V-i.** $\{(5, 10), (2, 5), (3, 12)\}$;
V-ii. $\{(6, 12), (1, 3), (2, 8), (1, 5)\}$.

By computation, we get minimal positive baskets as follows:

- V-1.** $B_{5,1} = \{(7, 15), (3, 12)\}$, $K^3 = \frac{1}{60}$, $P_7 = 2$, $P_8 = 3$.
V-2. $B_{5,2} = \{(6, 12), (1, 3), (3, 13)\}$, $K^3 = \frac{1}{39}$, $P_8 = 3$.
V-3. $B_{5,3} = \{(6, 12), (3, 11), (1, 5)\}$, $K^3 = \frac{1}{55}$, $P_8 = 2$, $P_{10} = 4$.

3.8. Case VI: $P_3 = 0, P_4 = P_5 = P_6 = 1$ ($\Rightarrow B_{6,1} \sim B_{6,6}$)

We have $\sigma = 10$, $\tau = 4$, $\Delta^3 = 6$, $\Delta^4 = 14$. Also, $P_7 \geq 1$ and hence $\sigma_5 \leq \epsilon \leq 2$. The initial baskets have four types:

- VI-i.** $\{6 \times (1, 2), 2 \times (1, 3), 2 \times (1, 4)\}$;
VI-ii. $\{6 \times (1, 2), 2 \times (1, 3), (1, 4), (1, 5)\}$;
VI-iii. $\{6 \times (1, 2), 2 \times (1, 3), 2 \times (1, 5)\}$;
VI-iv. $\{6 \times (1, 2), 2 \times (1, 3), (1, 4), (1, r)\}$ with $r \geq 6$.

Since there are only 2 baskets of $(1, 3)$, we have $\epsilon_5 = 3 - \sigma_5 \leq 2$. Hence $\sigma_5 > 0$ and $\epsilon > 0$. Therefore, Case VI-i is impossible.

For Case VI-ii and $\epsilon_5 = 2$, we get:

- VI-ii.** $B^{(5)} = \{4 \times (1, 2), 2 \times (2, 5), (1, 4), (1, 5)\}$.
Hence we get minimal positive baskets as follows:
VI-1. $B_{6,1} = \{(1, 2), (7, 16), (2, 9)\}$, $K^3 = \frac{1}{144}$, $P_7 = 2$, $P_9 = 3$.
VI-2. $B_{6,2} = \{(6, 13), (2, 5), (2, 9)\}$, $K^3 = \frac{8}{585}$, $P_7 = 2$, $P_8 = 3$.
VI-3. $B_{6,3} = \{(8, 18), (1, 4), (1, 5)\}$, $K^3 = \frac{1}{180}$, $P_7 = 2$, $P_9 = 3$.

For Case VI-iii and $\epsilon_5 = 1$, we get:

- VI-ii.** $B^{(5)} = \{5 \times (1, 2), (2, 5), (1, 3), 2 \times (1, 5)\}$.
Hence we get minimal positive baskets as follows:
VI-4. $B_{6,4} = \{(1, 2), (6, 13), (1, 3), (2, 10)\}$, $K^3 = \frac{1}{390}$, $P_8 = 2$, $P_9 = 3$.
VI-5. $B_{6,5} = \{(5, 10), (3, 8), (2, 10)\}$, $K^3 = \frac{1}{40}$, $P_8 = 3$.

For Case VI-iv and $\epsilon_5 = 2$, we get:

- VI-iv.** $B^{(5)} = \{4 \times (1, 2), 2 \times (2, 5), (1, 4), (1, r)\}$ with $r \geq 6$.
Since $K^3(B^{(5)}) > 0$, we must have $r = 6$. Then we get the following minimal positive basket:
VI-6. $B_{6,6} = \{(3, 6), (3, 7), (2, 5), (1, 4), (1, 6)\}$, $K^3 = \frac{1}{420}$, $P_{10} = 2$, $P_{12} = 3$.

3.9. Case VII: $P_3 = 1, P_4 = 0, P_5 = P_6 = 1$ (impossible)

We have $\sigma = 9$, $\tau = 3$, $\Delta^3 = 1$, $\Delta^4 = 9$. Moreover, $P_7 \geq 1$ and hence $\epsilon = 0$. It follows that $\sigma_5 = 0$ and $\epsilon_5 = 2$. The initial basket is $B^{(0)} = \{(1, 2), 7 \times (1, 3), (1, 4)\}$.

Note that there is only one basket of type $(1, 2)$. However, since $\epsilon_5 = 2$, one has $1 \geq n_{2,5}^5 = 2$, a contradiction. Thus Case VII does not happen.

3.10. Case VIII: $P_3 = P_4 = P_5 = P_6 = 1$ ($\Rightarrow B_{8,1} \sim B_{8,3}$)

We have $\sigma = 9$, $\tau = 3$, $\Delta^3 = 2$, $\Delta^4 = 8$. Moreover, $P_7 \geq 1$ and then $\epsilon \leq 1$. If $\epsilon = 1$, then $\sigma_5 = 1$ and $\epsilon_5 = 1$. If $\epsilon = 0$, then $\sigma_5 = 0$ and $\epsilon_5 = 2$. The initial baskets and $B^{(5)}$ have 2 types:

VIII-i. $B^{(0)} = \{2 \times (1, 2), 4 \times (1, 3), 3 \times (1, 4)\} \succ B^{(5)} = \{2 \times (2, 5), 2 \times (1, 3), 3 \times (1, 4)\}$ with $K^3(B^{(5)}) = \frac{1}{60}$.

VIII-ii. $B^{(0)} = \{2 \times (1, 2), 4 \times (1, 3), 2 \times (1, 4), (1, 5)\} \succ B^{(5)} = \{(1, 2), (2, 5), 3 \times (1, 3), 2 \times (1, 4), (1, 5)\}$ with $K^3(B^{(5)}) = 0$.

Clearly, Case VIII-ii is impossible since K^3 is not positive.

For Case VIII-i, we first consider the situation that one single basket $(2, 5)$ is packed, so that we get the basket $\{(2, 5), (3, 8), (1, 3), (4, 12)\}$. We can get two minimal positive baskets as follows:

VIII-1. $B_{8,1} = \{(5, 13), (1, 3), (3, 12)\}$, $K^3 = \frac{1}{156}$, $P_7 = 2$, $P_8 = 3$.

VIII-2. $B_{8,2} = \{(2, 5), (4, 11), (3, 12)\}$, $K^3 = \frac{1}{220}$, $P_7 = 2$, $P_8 = 3$.

It remains to consider the situation that each single basket $(2, 5)$ remains unpacked. We then obtain the basket

$$B_{210} := \{(4, 10), (1, 3), (2, 7), (2, 8)\}$$

with $K^3 = \frac{1}{210}$, $P_7 = 2$, $P_{10} = 3$. After a one-step prime packing, we get the minimal positive basket:

VIII-3. $B_{8,3} = \{(4, 10), (1, 3), (3, 11), (1, 4)\}$, $K^3 = \frac{1}{660}$, $P_7 = 2$.

The detailed classification (3.3~ 3.10) makes it possible for us to study the birational geometry of X , of which the first application is the following theorem.

Theorem 3.11. *Assume $\chi(\mathcal{O}_X) = 1$. Then $K_X^3 \geq \frac{1}{420}$. Furthermore, $K_X^3 = \frac{1}{420}$ if, and only if, $\mathcal{B} = B_{6,6}$.*

Proof. If $\mu_1 \leq 6$, then Proposition 2.30 implies $K_X^3 \geq \frac{1}{294} \cdot \frac{11}{12} > \frac{1}{420}$.

We may assume that $P_m \leq 1$ for $m \leq 6$. We have seen $P_2 = 0$. Since $\mathbf{B}^5 \succ \mathbf{B} \succ \mathbf{B}_{\min}$ and by [4, Lemma 3.6], we have

$$K_X^3 = K^3(B) \geq K^3(B_{\min})$$

where B_{\min} is in the set $\{B_{2,1}, B_{2,2}, B_{3,1} \sim B_{3,5}, B_{5,1} \sim B_{5,3}, B_{6,1} \sim B_{6,6}, B_{8,1} \sim B_{8,3}\}$.

If $B_{\min} \neq B_{6,6}$, or $B_{8,3}$, then we have seen $K^3(B_{\min}) > \frac{1}{420}$.

If $B_{\min} = B_{8,3}$, we show $B \neq B_{8,3}$. In fact, if $B = B_{8,3}$, then $P_7(B) = 2$ as we have seen in 3.10. By Table A in Section 2, we have $K_X^3 = K^3(\mathbf{B}) \geq \frac{5}{2408} > \frac{1}{660}$, a contradiction. Hence $B \succ B_{8,3}$. Notice that $B_{8,3}$ is obtained, exactly, by one-step packing from

$$B_{210} := \{(4, 10), (1, 3), (2, 7), (2, 8)\}$$

and no other ways. This says $B \succ B_{210}$ and so $K_X^3 \geq K^3(B_{210}) = \frac{1}{210}$.

We have seen $K^3(B_{6,6}) = \frac{1}{420}$. We are done. q.e.d.

With a different approach, L. Zhu [29] also proved $K^3 \geq \frac{1}{420}$. The proof of the last theorem gives the following:

Corollary 3.12. *Assume $\chi(\mathcal{O}_X) = 1$ and $P_m \leq 1$ for all $m \leq 6$. Then $\mathcal{B}(X)$ either dominates a minimal basket in the set*

$$\{B_{2,1}, B_{2,2}, B_{3,1} \sim B_{3,5}, B_{5,1} \sim B_{5,3}, B_{6,1} \sim B_{6,6}, B_{8,1}, B_{8,2}\}$$

or dominates the basket B_{210} .

Corollary 3.13. *Assume $\chi(\mathcal{O}_X) = 1$. Then $P_{10}(X) \geq 2$ and, in particular, $\mu_1 \leq 10$.*

Proof. If $P_{m_0} \geq 2$ for some $m_0 \leq 6$, then, by Lemma 3.2, one can see $P_{10} \geq 2$. Otherwise, Corollary 3.12 and [4, Lemma 3.6] imply that $P_{10} = P_{10}(\mathcal{B}(X)) \geq P_{10}(B_*)$ where B_* denotes a minimal positive basket mentioned in Corollary 3.12. By a direct computation, we get $P_{10}(B_*) \geq 2$. q.e.d.

Example 1.3 shows that the statement in Corollary 3.13 is optimal since $P_9(X_{46}) = 1$.

Theorem 3.14. *Assume $\chi(\mathcal{O}_X) = 1$. Then:*

- (1) $\rho_0 \leq 7$.
- (2) *Either $P_5 > 0$ or $P_6 > 0$.*

Proof. (1) Recall that $\mu_0 := \min\{m | P_m > 0\}$. By 3.3, we see $\mu_0 \leq 6$.

When $\mu_0 \leq 3$, it is easy to deduce the statement by Lemma 3.2.

When $\mu_0 = 4$, Lemma 3.2 implies $P_{2k} > 0$ for all $k \geq 3$. If $P_7 > 0$, Lemma 3.2 implies $P_{2k+1} > 0$ for all $k \geq 3$ and the statement (1) is true. Assume $P_7 = 0$. Then $P_5 = 0$. Now $\epsilon_5 = 2 - P_6 - \sigma_5 \geq 0$ implies $\sigma_5 \leq 2 - P_6 \leq 1$. On the other hand, $\epsilon_6 = P_4 + P_6 - \epsilon = 0$ implies $\epsilon \geq 2$. This means $\sigma_5 = P_6 = P_4 = 1$ and the situation corresponds to 3.7. Thus $B \succ B_{\min}$ where $B_{\min} = B_{5,2}, B_{5,3}$. But the computation tells $P_7(B_{\min}) > 0$, a contradiction.

When $\mu_0 = 5$, we study P_8 . If $P_8 > 0$, then (1) is true by Lemma 3.2. Assume $P_8 = 0$. Then $P_6 = 0$. Now $\epsilon_6 = P_5 - P_7 - \epsilon = 0$ gives $\epsilon = 0$ and $P_5 = P_7$ since $P_7 \geq P_5$. Since $n_{1,4}^0 = 1 - P_5 \geq 0$, we see $P_5 = 1$. So the situation corresponds to 3.5. Since the computation shows $P_8 \geq P_8(B_{3,*}) > 0$, a contradiction.

Finally, when $\mu_0 = 6$, we study P_7 . If $P_7 > 0$, then Lemma 3.2 implies (1). Otherwise, $P_7 = 0$. Now $\epsilon_6 = P_6 - \epsilon = 0$ implies $\epsilon = P_6 > 0$. Besides, $\epsilon_5 = 2 - P_6 - \sigma_5 \geq 0$ says $P_6 \leq 1$ since $\sigma_5 > 0$. Hence $\epsilon = P_6 = 1$. The situation corresponds to 3.4. But the computation shows $P_7 \geq P_7(B_{2,1}) > 0$ or $P_7 \geq P_7(B_{2,2}) > 0$, a contradiction.

(2) Assume $P_5 = P_6 = 0$. Then Lemma 3.2 implies $P_3 = P_4 = 0$. The situation corresponds to 3.3, which is impossible as already seen there.

q.e.d.

D. Shin [22] proved the first statement in a different way.

4. General type 3-folds with $\chi > 1$

In this section, we assume $\chi(\mathcal{O}_X) > 1$. Again, we will frequently apply our formulae and inequalities in [4, Sections 3 and 4].

When $P_{m_0} \geq 2$ for some positive integer $m_0 \leq 12$, known theorems will give an effective lower bound of K_X^3 and a practical pluricanonical birationality. Therefore, similar to Section 3, we need to classify X up to baskets when preceding plurigenera are smaller. For this reason, we make the following:

4.1. Assumption: $P_m \leq 1$ for all $m \leq 12$.

According to [4, Lemma 4.8], we have seen that $P_2 = 0$ under Assumption 4.1. Note that inequality [4, (3.14)], for general-type 3-folds, is as follows:

$$2P_5 + 3P_6 + P_8 + P_{10} + P_{12} \geq \chi + 10P_2 + 4P_3 + P_7 + P_{11} + P_{13} + R \tag{4.1}$$

where

$$\begin{aligned} R &:= 14\sigma_5 - 12n_{1,5}^0 - 9n_{1,6}^0 - 8n_{1,7}^0 - 6n_{1,8}^0 - 4n_{1,9}^0 - 2n_{1,10}^0 - n_{1,11}^0 \\ &= 2n_{1,5}^0 + 5n_{1,6}^0 + 6n_{1,7}^0 + 8n_{1,8}^0 + 10n_{1,9}^0 + 12n_{1,10}^0 + 13n_{1,11}^0 \\ &\quad + 14 \sum_{r \geq 12} n_{1,r}^0 \end{aligned}$$

and $\sigma_5 = \sum_{r \geq 5} n_{1,r}^0$.

Inequality (4.1) and Assumption 4.1 imply that both χ and P_{13} are bounded from above. Thus our formulae in [4, Section 4] allows us to explicitly compute $B^{(12)}$. To be more solid, we prove the following:

Proposition 4.2. *Assume $\chi(\mathcal{O}_X) > 1$ and $P_m \leq 1$ for all $m \leq 12$. Then the formal basket $\mathbf{B} = \mathbf{B}(X) := (\mathcal{B}(X), \chi(\mathcal{O}_X), 0)$ has a finite number of possibilities.*

Proof. We study $n_{1,r}^0$ for $r \geq 6$. If there exists a number $r \geq 6$ such that $n_{1,r}^0 \neq 0$, then $R \geq 5$ by the definition of R in inequality (4.1). Hence, by (4.1), one has

$$8 \geq 2P_5 + 3P_6 + P_8 + P_{10} + P_{12} \geq \chi + 5 \geq 7.$$

This implies that $P_5 = P_6 = 1$. Hence $P_{11} = 1$. Now (4.1) again reads $5 + P_8 + P_{10} + P_{12} \geq 8 + P_7 + P_{13}$. It follows that $P_8 = P_{10} = P_{12} = 1$ and $P_7 = P_{13} = 0$. This gives a contradiction since $P_{13} \geq P_5 P_8 = 1$. So we conclude $n_{1,r}^0 = 0$ for all $r \geq 6$. In other words, [4, Assumption 3.8] is satisfied.

This essentially allows us to utilize those formulae in the last part of [4, Section 3]. In particular, one sees that each quantity there is bounded and hence $B^{(12)}$ has a finite number of possibilities. Dominated by $B^{(12)}$ (i.e., $B^{(12)} \succcurlyeq B$), $B = \mathcal{B}(X)$ also has a finite number of possibilities. We are done. q.e.d.

4.3. Complete classification of \mathbf{B} satisfying Assumption 4.1.

Note that, for all $0 < m, n \leq 12$, and $m + n \leq 13$,

$$P_{m+n} \geq P_m P_n \tag{4.2}$$

naturally holds since $P_m, P_n \leq 1$.

Suppose we have known $B^{(12)}$. Then we can determine all possible minimal positive baskets B_{\min} dominated by $B^{(12)}$, where $B_{\min} \in T$ (a finite set). Now the formal basket \mathbf{B} satisfies the following relation:

$$(B^{(12)}, \chi, 0) \succcurlyeq \mathbf{B} \succcurlyeq (B_{\min}, \chi, 0)$$

for some $B_{\min} \in T$. Therefore, by [4, Lemma 3.6], we have $K_X^3 = K^3(B) \geq K^3(B_{\min}) > 0$ and $P_m = P_m(B) \geq P_m(B_{\min})$. This is the whole strategy.

The calculation can be done by a simple computer program, or even by hand. Our main result is Table C, which is a complete list of all possibilities of $B^{(12)}$ and its minimal positive elements.

In fact, first we preset $P_m = 0, 1$ for $m = 3, \dots, 11$. Then $\epsilon_6 = 0$ gives the value of ϵ . So we know the value of $n_{1,5}^0$. By inequality (4.1) we get the upper bound of χ since $P_{13} \geq 0$. Since $n_{1,4}^7 \geq 0$, we get the upper bound of η . Similarly $n_{2,9}^9 \geq 0$ gives the upper bound of ζ . Also $n_{4,9}^{11} \geq 0$ yields $\alpha \leq \zeta$. Finally, $n_{3,8}^{11} \geq 0$ gives the upper bound of β . Now we set $P_{12} = 0, 1$. Then inequality (4.1) again gives the upper bound of P_{13} , noting that $\chi \geq 2$. Clearly there are, at most, finitely many solutions. With inequality (4.2) imposed, we can get about 80 cases. An important property to mention is the inequality $K^3(B^{(12)}) \geq K^3(B) = K_X^3 > 0$. With $K^3 > 0$ imposed on, we have 63 outputs, which is exactly Table C. Simultaneously, we have been able to calculate all those minimal positive baskets dominated by $B^{(12)}$, since $B^{(12)}$ is “nearly” minimal in most cases.

If one would like to take a direct calculation by hand, it is of course possible. Consider the no. 2 case in Table C as an example. Since $P_2 = 0, P_3 = \dots = P_7 = 0, P_8 = 1$, and $P_9 = P_{10} = P_{11} = 0$, [4, (3.10)] tells that $\epsilon = 0$ and thus $\sigma_5 = 0$, which means $R = 0$. Now inequality (4.1) gives $P_{12} + 1 \geq \chi + P_{13} \geq 2$. So $P_{12} = 1, \chi = 2$, and $P_{13} = 0$. Now the formula for ϵ_{10} gives $\epsilon_{10} = -\eta \geq 0$, which means $\eta = 0$. Similarly, $n_{1,5}^9 = \zeta - 1 \geq 0$. On the other hand, $n_{3,7}^9 = 1 - \zeta \geq 0$. Thus $\zeta = 1$. Now $n_{4,9}^{11} = \zeta - \alpha \geq 0$ gives $\alpha \leq 1$. $n_{3,11}^{11} = 1 - \zeta - \alpha - \beta \geq 0$ gives $\alpha = \beta = 0$. Finally, we get

$$\{n_{1,2}, n_{5,12}, \dots, n_{1,5}\} = \{4, 0, 1, 0, 0, 2, 1, 0, 3, 0, 0, 0, 2, 0, 0\}.$$

That is, $B^{(12)} = \{4 \times (1, 2), (4, 9), 2 \times (2, 5), (3, 8), 3 \times (1, 3), 2 \times (1, 4)\}$.

We see that $B^{(12)}$ admits only one prime packing of type

$$\{(2, 5), (3, 8)\} \succ \{(5, 13)\}$$

over the minimal positive basket $\{4 \times (1, 2), (4, 9), (2, 5), (5, 13), 3 \times (1, 3), 2 \times (1, 4)\}$. We simply write this as $\{(5, 13), *\}$ in Table C. It is now easy to calculate K^3 for both $B^{(12)}$ and the minimal positive basket $\{(5, 13), *\}$. Finally, we can directly calculate P_m . At the same time, μ_1 is given in the table. For our needs in this context, we also display the value of $P_{18} = P_{18}(B^{(12)})$ or $P_{18}(B_{\min})$ and $P_{24} = P_{24}(B^{(12)})$ or $P_{24}(B_{\min})$ in Table C, though the symbols P_{18} or P_{24} are misused here.

So theoretically we can finish our classification by detailed computations. We omit the details because all calculations are similar.

4.4. Notation. By abuse of notation, we denote by B_* the final basket corresponding to No.* in Table C. For example, $B_2 = \{4 \times (1, 2), (4, 9), 2 \times (2, 5), (3, 8), 3 \times (1, 3), 2 \times (1, 4)\}$ while $B_{2a} = \{4 \times (1, 2), (4, 9), (2, 5), (5, 13), 3 \times (1, 3), 2 \times (1, 4)\}$ is minimal positive. The relation is as follows:

$$B_2 \succ B \succ B_{2a}.$$

Clearly, for this case, we have $\frac{1}{360} = K^3(B_2) \geq K_X^3 \geq K^3(B_{2a}) = \frac{1}{1170}$.

Another typical example is No.63, where we have

$$B_{63} = \{5 \times (1, 2), (4, 9), 2 \times (3, 7), (2, 5), (3, 8), (4, 11), 3 \times (1, 3), (2, 7), (1, 5)\},$$

which is already minimal positive. So we have the relation

$$B^{(12)} = B_{63} = B = B_{\min}$$

and thus $K_X^3 = \frac{1}{5544}$. Of course, we will see that No.63 does not happen on any X .

Now we begin to analyze Table C and pick out “impossible” cases.

Proposition 4.5. *In Table C, $B \neq B_*$ for any B_* in the set*

$$\{B_{4a}, B_9, B_{16a}, B_{16c}, B_{18a}, B_{20a}, B_{21a}, B_{22}, B_{24}, B_{27a}, B_{29a}, B_{33a}, B_{44b}, B_{46a}, B_{47}, B_{52a}, B_{55}, B_{60a}, B_{61}, B_{63}\}.$$

In particular, cases No. 9, No. 22, No. 24, No. 47, No. 55, No. 61, and No. 63 do not happen at all.

Table C

<i>No.</i>	(P_3, \dots, P_{11})	P_{18}	P_{24}	μ_1	χ	$B^{(12)} = (n_{1,2}, n_{5,11}, \dots, n_{1,5})$ or B_{min}	K^3
1	(0, 0, 0, 0, 0, 0, 0, 1, 0)	4	8	14	2	(5, 0, 0, 1, 0, 3, 0, 0, 3, 0, 0, 1, 0, 0, 0)	$\frac{3}{770}$
2	(0, 0, 0, 0, 0, 1, 0, 0, 0)	3	7	15	2	(4, 0, 1, 0, 0, 2, 1, 0, 3, 0, 0, 0, 2, 0, 0)	$\frac{1}{360}$
2a		2	3	18		{(2, 5), (3, 8), *} > {(5, 13), *}	$\frac{1170}{23}$
3	(0, 0, 0, 0, 0, 1, 0, 1, 0)	3	7	15	3	(6, 1, 0, 0, 0, 4, 1, 0, 4, 0, 1, 0, 2, 0, 0)	$\frac{9240}{17}$
3a		2	3	18		{(2, 5), (3, 8), *} > {(5, 13), *}	$\frac{3080}{17}$
4	(0, 0, 0, 0, 0, 1, 0, 1, 0)	4	9	14	3	(7, 0, 1, 0, 0, 4, 0, 1, 3, 0, 1, 0, 2, 0, 0)	$\frac{3465}{17}$
4a		1	2	14		{(4, 11), (2, 6), *} > {(6, 17), *}	$\frac{5355}{17}$
5	(0, 0, 0, 0, 0, 1, 0, 1, 0)	5	10	14	3	(7, 0, 1, 0, 0, 4, 1, 0, 4, 0, 0, 1, 1, 0, 0)	$\frac{3960}{17}$
5a		4	3	15		{(8, 20), (3, 8), *} > {(11, 28), *}	$\frac{1386}{17}$
5b		3	3	15		{(5, 13), (4, 15), *}	$\frac{1170}{17}$
6	(0, 0, 0, 1, 0, 0, 0, 1, 0)	3	6	14	3	(9, 0, 0, 2, 0, 1, 0, 1, 4, 0, 2, 0, 0, 0, 1)	$\frac{462}{17}$
7	(0, 0, 0, 1, 0, 0, 1, 0, 0)	3	5	14	2	(5, 0, 1, 1, 0, 0, 0, 0, 5, 0, 1, 0, 0, 0, 1)	$\frac{630}{17}$
7a		2	3	14		{(4, 9), (3, 7), *} > {(7, 16), *}	$\frac{1680}{17}$
8	(0, 0, 0, 1, 0, 0, 1, 1, 0)	3	5	14	3	(7, 1, 0, 1, 0, 2, 0, 0, 6, 0, 2, 0, 0, 0, 1)	$\frac{770}{17}$
9	(0, 0, 0, 1, 0, 1, 0, 0, 0)	2	2	14	3	(9, 0, 0, 2, 0, 0, 1, 1, 4, 0, 1, 0, 0, 1, 0)	$\frac{5544}{17}$
10	(0, 0, 0, 1, 0, 1, 0, 0, 0)	3	6	14	3	(8, 0, 1, 1, 0, 0, 2, 0, 5, 0, 1, 0, 1, 0, 1)	$\frac{630}{17}$
10a		2	4	14		{(4, 9), (3, 7), *} > {(7, 16), *}	$\frac{1680}{17}$
11	(0, 0, 0, 1, 0, 1, 0, 1, 0)	2	4	14	3	(9, 0, 0, 2, 0, 0, 1, 1, 3, 1, 0, 0, 1, 0, 1)	$\frac{3080}{17}$
11a		2	3	14		{(3, 8), (4, 11), *} > {(7, 19), *}	$\frac{2660}{17}$
12	(0, 0, 0, 1, 0, 1, 0, 1, 0)	5	11	14	3	(9, 0, 1, 0, 0, 1, 2, 0, 4, 0, 2, 0, 0, 0, 1)	$\frac{252}{17}$
12a		4	6	14		{(2, 5), (6, 16), *} > {(8, 21), *}	$\frac{630}{17}$
13	(0, 0, 0, 1, 0, 1, 0, 1, 0)	3	4	14	4	(12, 0, 0, 2, 0, 2, 0, 2, 4, 0, 2, 0, 0, 1, 0)	$\frac{3465}{17}$
14	(0, 0, 0, 1, 0, 1, 0, 1, 0)	3	6	14	4	(10, 1, 0, 1, 0, 2, 2, 0, 6, 0, 2, 0, 1, 0, 1)	$\frac{770}{17}$
15	(0, 0, 0, 1, 0, 1, 0, 1, 0)	4	8	14	4	(11, 0, 1, 1, 0, 2, 1, 1, 5, 0, 2, 0, 1, 0, 1)	$\frac{71}{17}$
15a		2	4	14		{(4, 11), (1, 3), *} > {(5, 14), *}	$\frac{27720}{17}$
15b		3	4	14		{(2, 5), (3, 8), *} > {(5, 13), *}	$\frac{2520}{17}$
15c		3	5	14		{(7, 16), (7, 19), *}	$\frac{36036}{17}$
16	(0, 0, 0, 1, 0, 1, 0, 1, 0)	5	9	14	4	(11, 0, 1, 1, 0, 2, 2, 0, 6, 0, 1, 1, 0, 0, 1)	$\frac{31020}{17}$
16a		4	3	14		{(4, 10), (3, 8), *} > {(7, 18), *}	$\frac{43}{17}$
16b		4	4	14		{(2, 5), (6, 16), *} > {(8, 21), *}	$\frac{3080}{17}$
16c		3	3	14		{(7, 16), (5, 13), *}	$\frac{1386}{17}$
17	(0, 0, 0, 1, 0, 1, 0, 1, 1)	3	6	14	3	(9, 0, 0, 2, 0, 0, 0, 2, 3, 0, 1, 0, 1, 0, 1)	$\frac{16016}{17}$
18	(0, 0, 0, 1, 0, 1, 0, 1, 1)	4	7	14	3	(9, 0, 0, 2, 0, 0, 1, 1, 4, 0, 0, 1, 0, 0, 1)	$\frac{3}{1540}$
18a		2	3	14		{(4, 11), (1, 3), *} > {(5, 14), *}	$\frac{9240}{17}$
18b		4	6	14		{(3, 8), (4, 11), *} > {(7, 19), *}	$\frac{3080}{88}$
19	(0, 0, 0, 1, 0, 1, 1, 0, 0)	3	3	14	3	(8, 0, 1, 1, 0, 1, 0, 1, 5, 0, 1, 0, 0, 1, 0)	$\frac{43890}{17}$
20	(0, 0, 0, 1, 0, 1, 1, 0, 0)	4	7	14	3	(7, 0, 2, 0, 0, 1, 1, 0, 6, 0, 1, 0, 1, 0, 1)	$\frac{3465}{17}$
20a		3	3	18		{(2, 5), (3, 8), *} > {(5, 13), *}	$\frac{504}{17}$
21	(0, 0, 0, 1, 0, 1, 1, 1, 0)	4	8	14	2	(6, 0, 1, 0, 0, 0, 1, 0, 3, 1, 0, 0, 0, 0, 1)	$\frac{16380}{17}$
21a		2	3	16		{(1, 3), (3, 10), *} > {(4, 13), *}	$\frac{4680}{17}$
22	(0, 0, 0, 1, 0, 1, 1, 1, 0)	2	3	18	3	(7, 1, 0, 1, 0, 1, 1, 0, 5, 1, 0, 0, 1, 0, 1)	$\frac{9240}{17}$
23	(0, 0, 0, 1, 0, 1, 1, 1, 0)	3	5	14	3	(8, 0, 1, 1, 0, 1, 0, 1, 4, 1, 0, 0, 1, 0, 1)	$\frac{13860}{17}$
23a		2	3	14		{(4, 9), (3, 7), *} > {(7, 16), *}	$\frac{2640}{17}$
24	(0, 0, 0, 1, 0, 1, 1, 1, 0)	3	3	14	4	(10, 1, 0, 1, 0, 3, 0, 1, 6, 0, 2, 0, 0, 1, 0)	$\frac{3465}{17}$
25	(0, 0, 0, 1, 0, 1, 1, 1, 0)	4	7	14	4	(9, 1, 1, 0, 0, 3, 1, 0, 7, 0, 2, 0, 1, 0, 1)	$\frac{47}{27720}$
25a		4	6	14		{(5, 11), (4, 9), *} > {(9, 20), *}	$\frac{840}{17}$
26	(0, 0, 0, 1, 0, 1, 1, 1, 0)	5	9	14	4	(10, 0, 2, 0, 0, 3, 0, 1, 6, 0, 2, 0, 1, 0, 1)	$\frac{13860}{17}$
26a		3	5	14		{(4, 11), (1, 3), *} > {(5, 14), *}	$\frac{1260}{17}$
27	(0, 0, 0, 1, 0, 1, 1, 1, 0)	6	10	14	4	(10, 0, 2, 0, 0, 3, 1, 0, 7, 0, 1, 1, 0, 0, 1)	$\frac{27720}{17}$
27a		5	3	14		{(6, 15), (3, 8), *} > {(9, 23), *}	$\frac{19}{79695}$
27b		5	5	14		{(5, 13), (5, 18), *}	$\frac{1170}{17}$
28	(0, 0, 0, 1, 0, 1, 1, 1, 1)	4	8	14	2	(5, 1, 0, 0, 0, 0, 1, 0, 4, 0, 1, 0, 0, 0, 1)	$\frac{23}{9240}$
29	(0, 0, 0, 1, 0, 1, 1, 1, 1)	5	10	14	2	(6, 0, 1, 0, 0, 0, 0, 1, 3, 0, 1, 0, 0, 0, 1)	$\frac{13}{3465}$
29a		2	3	14		{(4, 11), (2, 6), *} > {(6, 17), *}	$\frac{5355}{17}$
30	(0, 0, 0, 1, 0, 1, 1, 1, 1)	3	5	14	3	(7, 1, 0, 1, 0, 1, 0, 1, 5, 0, 1, 0, 1, 0, 1)	$\frac{924}{17}$
31	(0, 0, 0, 1, 0, 1, 1, 1, 1)	4	6	14	3	(7, 1, 0, 1, 0, 1, 1, 0, 6, 0, 0, 1, 0, 0, 1)	$\frac{616}{17}$
32	(0, 0, 0, 1, 0, 1, 1, 1, 1)	5	8	14	3	(8, 0, 1, 1, 0, 1, 0, 1, 5, 0, 0, 1, 0, 0, 1)	$\frac{693}{17}$
32a		4	6	14		{(4, 9), (3, 7), *} > {(7, 16), *}	$\frac{528}{17}$
32b		2	2	14		{(4, 11), (1, 3), *} > {(5, 14), *}	$\frac{1386}{17}$
33	(0, 0, 0, 1, 1, 0, 0, 1, 0)	2	4	14	2	(5, 0, 0, 2, 0, 0, 1, 0, 1, 1, 1, 0, 0, 0, 0)	$\frac{840}{17}$
33a		1	3	14		{(3, 10), (2, 7), *} > {(5, 17), *}	$\frac{2856}{17}$

No.	(P_3, \dots, P_{11})	P_{18}	P_{24}	μ_1	χ	$(n_{1,2}, n_{4,9}, \dots, n_{1,5})$ or B_{min}	K^3
34	(0, 0, 0, 1, 1, 0, 0, 1, 0)	4	8	14	3	(7, 0, 1, 1, 0, 2, 1, 0, 3, 0, 3, 0, 0, 0, 0)	$\frac{1}{360}$
34a		3	6	14		{(4, 9), (3, 7), *} > {(7, 16), *}	$\frac{1}{560}$
34b		3	4	14		{(2, 5), (3, 8), *} > {(5, 13), *}	$\frac{1}{1170}$
35	(0, 0, 0, 1, 1, 0, 0, 1, 1)	3	6	14	2	(5, 0, 0, 2, 0, 0, 0, 1, 1, 0, 2, 0, 0, 0, 0)	$\frac{1}{462}$
36	(0, 0, 0, 1, 1, 0, 1, 1, 0)	3	5	14	2	(4, 0, 1, 1, 0, 1, 0, 0, 2, 1, 1, 0, 0, 0, 0)	$\frac{1}{630}$
36a		2	3	14		{(4, 9), (3, 7), *} > {(7, 16), *}	$\frac{1}{1680}$
36b		2	4	14		{(3, 10), (2, 7), *} > {(5, 17), *}	$\frac{1}{5355}$
37	(0, 0, 0, 1, 1, 0, 1, 1, 0)	5	9	14	3	(6, 0, 2, 0, 0, 3, 0, 0, 4, 0, 3, 0, 0, 0, 0)	$\frac{1}{315}$
38	(0, 0, 0, 1, 1, 0, 1, 1, 1)	3	5	14	2	(3, 1, 0, 1, 0, 1, 0, 0, 3, 0, 2, 0, 0, 0, 0)	$\frac{1}{770}$
39	(0, 0, 0, 1, 1, 1, 0, 1, 0)	3	6	14	3	(7, 0, 1, 1, 0, 1, 2, 0, 2, 1, 1, 0, 1, 0, 0)	$\frac{1}{630}$
39a		2	4	14		{(4, 9), (3, 7), *} > {(7, 16), *}	$\frac{1}{1680}$
39b		2	5	14		{(3, 10), (2, 7), *} > {(5, 17), *}	$\frac{1}{5355}$
40	(0, 0, 0, 1, 1, 1, 0, 1, 0)	5	10	14	4	(9, 0, 2, 0, 0, 3, 2, 0, 4, 0, 3, 0, 1, 0, 0)	$\frac{1}{315}$
40a		4	4	14		{(4, 10), (3, 8), *} > {(7, 18), *}	$\frac{1}{2520}$
40b		4	5	14		{(2, 5), (6, 16), *} > {(8, 21), *}	$\frac{1}{1260}$
41	(0, 0, 0, 1, 1, 1, 0, 1, 1)	5	11	13	2	(5, 0, 1, 0, 0, 0, 2, 0, 1, 0, 2, 0, 0, 0, 0)	$\frac{1}{252}$
42	(0, 0, 0, 1, 1, 1, 0, 1, 1)	3	6	14	3	(6, 1, 0, 1, 0, 1, 2, 0, 3, 0, 2, 0, 1, 0, 0)	$\frac{1}{770}$
43	(0, 0, 0, 1, 1, 1, 0, 1, 1)	4	8	14	3	(7, 0, 1, 1, 0, 1, 1, 1, 2, 0, 2, 0, 1, 0, 0)	$\frac{1}{27720}$
43a		2	4	14		{(4, 11), (1, 3), *} > {(5, 14), *}	$\frac{1}{2520}$
43b		3	4	14		{(2, 5), (3, 8), *} > {(5, 13), *}	$\frac{1}{36036}$
43c		3	5	14		{(7, 16), (7, 19), *}	$\frac{1}{31920}$
44	(0, 0, 0, 1, 1, 1, 0, 1, 1)	5	9	14	3	(7, 0, 1, 1, 0, 1, 2, 0, 3, 0, 1, 1, 0, 0, 0)	$\frac{1}{13860}$
44a		4	4	14		{(2, 5), (6, 16), *} > {(8, 21), *}	$\frac{1}{1386}$
44b		3	3	14		{(7, 16), (5, 13), *}	$\frac{1}{16016}$
44c		4	6	14		{(7, 16), (5, 18), *}	$\frac{1}{720}$
44d		4	4	14		{(5, 13), (5, 18), *}	$\frac{1}{2184}$
45	(0, 0, 0, 1, 1, 1, 1, 0, 1)	4	7	14	2	(3, 0, 2, 0, 0, 0, 1, 0, 3, 0, 1, 0, 1, 0, 0)	$\frac{1}{504}$
46	(0, 0, 0, 1, 1, 1, 1, 1, 0)	4	7	14	3	(6, 0, 2, 0, 0, 2, 1, 0, 3, 1, 1, 0, 1, 0, 0)	$\frac{1}{504}$
46a		3	3	16		{(2, 5), (3, 8), *} > {(5, 13), *}	$\frac{1}{16380}$
46b		3	6	14		{(3, 10), (2, 7), *} > {(5, 17), *}	$\frac{1}{6120}$
47	0, 0, 0, 1, 1, 1, 1, 1, 1)	2	3	16	2	(3, 1, 0, 1, 0, 0, 1, 0, 2, 1, 0, 0, 1, 0, 0)	$\frac{1}{9240}$
48	0, 0, 0, 1, 1, 1, 1, 1, 1)	3	5	14	2	(4, 0, 1, 1, 0, 0, 0, 1, 1, 1, 0, 0, 1, 0, 0)	$\frac{1}{13860}$
48a		2	3	14		{(4, 9), (3, 7), *} > {(7, 16), *}	$\frac{1}{2640}$
49	(0, 0, 0, 1, 1, 1, 1, 1, 1)	4	7	14	3	(5, 1, 1, 0, 0, 2, 1, 0, 4, 0, 2, 0, 1, 0, 0)	$\frac{1}{27720}$
49a		4	6	14		{(5, 11), (4, 9), *} > {(9, 20), *}	$\frac{1}{840}$
50	(0, 0, 0, 1, 1, 1, 1, 1, 1)	5	9	14	3	(6, 0, 2, 0, 0, 2, 0, 1, 3, 0, 2, 0, 1, 0, 0)	$\frac{1}{13860}$
50a		3	5	14		{(4, 11), (1, 3), *} > {(5, 14), *}	$\frac{1}{1260}$
51	(0, 0, 0, 1, 1, 1, 1, 1, 1)	6	10	14	3	(6, 0, 2, 0, 0, 2, 1, 0, 4, 0, 1, 1, 0, 0, 0)	$\frac{1}{27720}$
51a		5	4	14		{(4, 10), (3, 8), *} > {(7, 18), *}	$\frac{1}{1386}$
51b		5	5	14		{(5, 13), (5, 18), *}	$\frac{1}{1170}$
52	(0, 0, 1, 0, 0, 1, 0, 1, 0)	3	7	14	2	(4, 0, 0, 1, 0, 2, 2, 0, 2, 0, 0, 0, 0, 0, 1)	$\frac{1}{420}$
52a		2	3	18		{(2, 5), (3, 8), *} > {(5, 13), *}	$\frac{1}{2184}$
53	(0, 0, 1, 0, 0, 1, 1, 1, 0)	4	8	14	2	(3, 0, 1, 0, 0, 3, 1, 0, 3, 0, 0, 0, 0, 0, 1)	$\frac{1}{360}$
53a		3	4	15		{(2, 5), (3, 8), *} > {(5, 13), *}	$\frac{1}{1170}$
54	(0, 0, 1, 0, 1, 0, 0, 1, 0)	2	4	14	2	(2, 0, 0, 2, 0, 3, 1, 0, 1, 0, 1, 0, 0, 0, 0)	$\frac{1}{840}$
55	(0, 0, 1, 0, 1, 0, 0, 1, 0)	2	2	14	3	(4, 0, 0, 3, 0, 4, 1, 0, 3, 0, 0, 1, 0, 0, 0)	$\frac{1}{3080}$
56	(0, 0, 1, 0, 1, 0, 1, 1, 0)	3	5	14	2	(1, 0, 1, 1, 0, 4, 0, 0, 2, 0, 1, 0, 0, 0, 0)	$\frac{1}{630}$
56a		2	3	14		{(4, 9), (3, 7), *} > {(7, 16), *}	$\frac{1}{1680}$
57	(0, 0, 1, 0, 1, 0, 1, 1, 0)	3	3	14	3	(3, 0, 1, 2, 0, 5, 0, 0, 4, 0, 0, 1, 0, 0, 0)	$\frac{1}{1386}$
58	(0, 0, 1, 0, 1, 1, 0, 1, 0)	3	6	14	3	(4, 0, 1, 1, 0, 4, 2, 0, 2, 0, 1, 0, 1, 0, 0)	$\frac{1}{630}$
58a		2	4	14		{(4, 9), (3, 7), *} > {(7, 16), *}	$\frac{1}{1680}$
59	(0, 0, 1, 0, 1, 1, 0, 1, 1)	2	4	14	2	(2, 0, 0, 2, 0, 2, 1, 1, 0, 0, 0, 0, 1, 0, 0)	$\frac{1}{3080}$
59a		2	3	14		{(3, 8), (4, 11), *} > {(7, 19), *}	$\frac{1}{2660}$
60	(0, 0, 1, 0, 1, 1, 1, 1, 0)	4	7	14	3	(3, 0, 2, 0, 0, 5, 1, 0, 3, 0, 1, 0, 1, 0, 0)	$\frac{1}{504}$
60a		3	3	15		{(2, 5), (3, 8), *} > {(5, 13), *}	$\frac{1}{16380}$
61	(0, 0, 1, 0, 1, 1, 1, 1, 1)	2	3	15	2	(0, 1, 0, 1, 0, 3, 1, 0, 2, 0, 0, 0, 1, 0, 0)	$\frac{1}{9240}$
62	(0, 0, 1, 0, 1, 1, 1, 1, 1)	3	5	14	2	(1, 0, 1, 1, 0, 3, 0, 1, 1, 0, 0, 0, 1, 0, 0)	$\frac{1}{13860}$
62a		2	3	14		{(4, 9), (3, 7), *} > {(7, 16), *}	$\frac{1}{2640}$
63	(0, 0, 1, 1, 1, 1, 1, 1, 1)	3	4	14	3	(5, 0, 1, 2, 0, 1, 1, 1, 3, 0, 1, 0, 0, 0, 1)	$\frac{1}{5544}$

Proof. Assume $B = B_*$. We hope to deduce a contradiction.

(1). If $P_{14} \geq 2$, then Proposition 2.30 implies $K^3 \geq \frac{11}{37800} > \frac{1}{3437}$. Thus $B \neq B_{4a}, B_9, B_{16c}, B_{24}, B_{27a}, B_{29a}, B_{44b}, B_{63}$.

(2). If $P_{15} \geq 2$, then Proposition 2.30 implies $K^3 \geq \frac{11}{46080} > \frac{1}{4190}$. Hence $B \neq B_{60a}, B_{61}$.

(3). If $P_{16} \geq 2$, then Proposition 2.30 implies $K^3 \geq \frac{11}{55488} > \frac{1}{5045}$. Hence $B \neq B_{46a}, B_{47}$.

(4). If $P_{18} \geq 2$, then Proposition 2.30 implies $K^3 \geq \frac{11}{77976} > \frac{1}{7089}$. Thus $B \neq B_{20a}, B_{22}$.

(5). Besides, we see $P_6(B_{33a}) = 1$, $P_{16}(B_{33a}) = 2$ but $P_{22}(B_{33a}) = 1$, a contradiction. So $B \neq B_{33a}$.

(6). For cases 16a, 18a, 21a, 52a, and case 55, one has $P_{17}(B_*) = 0$. But since $P_8(B_{21a}) = P_9(B_{21a}) = 1$, $B \neq (B_{21a})$. Also for case 52a and case 55, since $P_5(B_*) = P_{12}(B_*) = 1$, we see $B \neq B_{52a}, B_{55}$. For case 18a, since $P_6(B_{18a}) = P_{11}(B_{18a}) = 1$, we see $B \neq B_{18a}$. Finally, since $P_{19}(B_{16a}) = -1$, we see $B \neq B_{16a}$. q.e.d.

Theorem 4.6. *Assume $\chi(\mathcal{O}_X) > 1$. Then $K_X^3 \geq \frac{1}{2660}$. Furthermore, $K_X^3 = \frac{1}{2660}$ if, and only if, $P_2 = 0$ and either $\chi = 3$, $B = B_{11a}$, or $\chi = 2$, B_{59a} .*

Proof. If $P_{m_0} \geq 2$ for some positive integer $m_0 \leq 12$, then Proposition 2.30 implies $K_X^3 \geq \frac{11}{24336} > \frac{1}{2213} > \frac{1}{2660}$.

Assume $P_m \leq 1$ for $m \leq 12$. Then we have seen $B \geq B_*$ where B_* is one in Table C excluding those cases listed in Proposition 4.5.

We can see $K^3(B_{11a}) = K^3(B_{59a}) = \frac{1}{2660}$.

We pick out those cases with $K^3(B_*) < \frac{1}{2660}$. They are cases 4a, 16a, 16c, 18a, 20a, 21a, 27a, 29a, 33a, 44b, 46a, and case 60a. In all these cases, Corollary 4.5 says $B \neq B_*$. Thus $B \succ B_*$. In order to prove the theorem, we need to study the one-step unpacking of B_* case by case.

First we consider case 4a and case 29a. It's obtained by 2-steps of packing from B_4 :

$$B_4 = \{(2, 6), (4, 11), *\} \succ B_{4.5} := \{(1, 3), (5, 14), *\} \succ \{(6, 17), *\} = B_{4a}.$$

By [4, Lemma 3.6], we get $K_X^3 = K^3(B) \geq K^3(B_{4.5}) = \frac{1}{630} > \frac{1}{2660}$. Similarly, we also get $K_X^3 > \frac{1}{2660}$ for case 29a.

Next, we consider cases 18a, 20a, 21a, 46a, 52a, 60a. The common property is that they are obtained by a 1-step packing from $B^{(12)}$. So the only possibility is $B^{(12)} = B$. Thus $K_X^3 = K^3(B_{18})$ or $K^3(B_{20})$ or $K^3(B_{21})$ or $K^3(B_{46})$ or $K^3(B_{52})$ or $K^3(B_{60})$. In a word, $K_X^3 > \frac{1}{2660}$.

The remaining cases are 16a, 16c, 27a, and 44b. For case 44b, there are two intermediate baskets dominating B_{44c} or B_{44d} , respectively. Thus, in particular, $K_X^3 > \frac{1}{2184}$. For case 27a, it's obtained from B_{27} by

3-steps of packing, namely,

$$B_{27} = \{3 \times (2, 5), (5, 8), *\} \succ \{2 \times (2, 5), (5, 13), *\} \\ \succ B_{27.5} := \{(2, 5), (7, 18), *\} \succ \{(9, 23), *\} = B_{27a}.$$

Thus we see $B \succ B_{27.5}$ and $K_X^3 \geq K^3(B_{27.5}) = \frac{1}{1386} > \frac{1}{2660}$. Finally, we consider cases 16a and 16c. We know

$$B_{16} = \{(4, 9), (3, 7), (2, 5), (2, 5), (3, 8), (3, 8), *\}.$$

The 1-step packing of B_{16} yields

$$B_{16.5} := \{(4, 9), (3, 7), (2, 5), (5, 13), (3, 8), *\},$$

and the 1-step prime packing of $B_{16.5}$ is either B_{16a} or B_{16c} . Thus, if $B \succ B_{16.5}$, then $K_X^3 \geq K^3(B_{16.5}) = \frac{85}{72072} > \frac{1}{848}$. The other intermediate basket dominating B_{16a} and B_{16c} is

$$B_{16.6} := \{(7, 16), (2, 5), (2, 5), (3, 8), (3, 8), *\}$$

with $K^3(B_{16.6}) = \frac{13}{6160} > \frac{1}{474}$. There are no other ways to obtain either B_{16a} or B_{16c} beginning from $B^{(16)}$. The theorem is proved. q.e.d.

Corollary 4.7. *Assume $\chi(\mathcal{O}_X) > 1$. Then $P_{24} \geq 2$.*

Proof. By [4, Theorem 4.15], we know either $P_{10} \geq 2$ or $P_{24} \geq 2$. When $q(X) > 0$, the statement follows from [3]. So we may assume $q(X) = 0$.

If $P_{10} \geq 2$, we take $m_0 = 10$ and study φ_{10} . Keep the same notation as in 2.3. By Lemma 2.32, f must be of type *III*, *II*, I_p . Proposition 2.15 (i), Theorem 2.20 (1), and Theorem 2.22 (1) imply $P_{24} \geq 2$. q.e.d.

Theorem 4.8. *Assume $\chi(\mathcal{O}_X) > 1$. Then $P_{m_0} \geq 2$ for some positive integer $m_0 \leq 18$. In particular, $\mu_1 \leq 18$.*

Proof. Assume $P_m \leq 1$ for all $m \leq 12$. Then Table C tells us that

$$B^{(12)} \succ B \succ B_{\min}$$

where B_{\min} is of certain type in Table C. Since, in Table C, we have seen $\mu_1(B_{\min}) \leq 18$, thus [4, Lemma 3.6] implies $\mu_1(X) \leq \mu_1(B_{\min}) \leq 18$. q.e.d.

Theorem 4.9. *Assume $\chi(\mathcal{O}_X) > 1$. Then $\rho_0(X) \leq 27$.*

Proof. The statement follows from [3] when $q(X) > 0$. Assume $q(X) = 0$ from now on.

If $P_{m_0} \geq 2$ for some $m_0 \leq 12$, then the induced fibration f from φ_{m_0} is of type *III*, *II*, or I_p by Lemma 2.32. Thus Proposition 2.15(i), Theorem 2.20(1), and Theorem 2.22(1) imply that $P_m > 0$ for all $m \geq 27$.

If $P_m \leq 1$ for all $m \leq 12$, we have a complete classification (cf. Table C). For each B_{\min} in Table C, we observed that $P_m > 0$ for all $47 \geq m \geq 24$. This is enough to assert $P_m > 0$ for all $m \geq 24$. We are done. q.e.d.

5. Pluricanonical birationality

In this section, we mainly study the birationality of φ_m . Then we can conclude our main theorems. Let X be a projective minimal 3-fold of general type. First, we recall several known theorems.

Theorem 5.1 ([3]). *Assume $q(X) := h^1(\mathcal{O}_X) > 0$. Then φ_m is birational for all $m \geq 7$.*

Theorem 5.2. ([7], [Theorem 0.1]) *Assume $P_{m_0} \geq 2$ for some positive integer m_0 . Then φ_m is birational onto its image for all $m \geq 5m_0 + 6$.*

Theorem 5.3 ([8]). *Assume $\chi(\mathcal{O}_X) \leq 0$. Then φ_m is birational for all $m \geq 14$.*

We need the following lemma to prove our main theorems.

Lemma 5.4. *Assume $P_{m_0}(X) \geq 2$ for some positive integer m_0 . Keep the same notation as in 2.3 and assume f is of type I_p or I_n . Suppose $|G|$ is a base point free linear system on S . If there exists an integer $m_1 > 0$ with $m_1\pi^*(K_X)|_S \geq G$, then Assumption 2.9(2) is satisfied for all integers*

$$m \geq \max\{\rho_0 + m_0 + m_1, m_0 + m_1 + 2\}.$$

Proof. Since

$$\begin{aligned} & K_S + \lceil (m-1)\pi^*(K_X) - S - \frac{1}{p}E'_{m_0} \rceil_S \\ & \geq K_S + (m-1)\pi^*(K_X)|_S - (S + E'_{m_0})|_S \\ & \geq (m-m_0)\pi^*(K_X)|_S \geq (m-m_0-m_1)\pi^*(K_X)|_S + G \end{aligned}$$

and

$$\begin{aligned} & K_S + (m-m_0-1)\pi^*(K_X)|_S \\ & \geq K_S + (m-m_0-m_1-1)\pi^*(K_X)|_S + G, \end{aligned}$$

Lemma 2.17 implies that $|K_S + \lceil (m-1)\pi^*(K_X) - S - \frac{1}{p}E'_{m_0} \rceil_S|$ can distinguish different generic irreducible elements of $|G|$ when $m \geq \rho_0 + m_0 + m_1$ and $m \geq m_0 + m_1 + 2$. q.e.d.

Theorem 5.5. *Let X be a projective minimal 3-fold of general type with $\chi(\mathcal{O}_X) = 1$. Then φ_m is birational for all $m \geq 40$.*

Proof. If $P_{m_0} \geq 2$ for some $m_0 \leq 6$, then, by Theorem 5.2, φ_m is birational for $m \geq 36$.

Assume $P_m \leq 1$ for all $m \leq 6$. Then, by Corollary 3.12, $\mathcal{B}(X)$ either dominates a minimal basket in

$$\{B_{2,1}, B_{2,2}, B_{3,1} \sim B_{3,5}, B_{5,1} \sim B_{5,3}, B_{6,1} \sim B_{6,6}, B_{8,1}, B_{8,2}\}$$

or dominates the basket B_{210} . We have known $P_m(X) \geq P_m(B_{*,*})$. By analyzing all the above baskets, we see a common property that there

is a pair of positive integers (n_0, n_1) satisfying $P_{n_0} \geq 2$, $P_{n_1} \geq 3$, and one of the following conditions:

- (1) $n_0 \leq 10$, $n_1 \leq 12$ (see cases III-2, III-3, III-4, VI-6);
- (2) $n_0 \leq 9$, $n_1 \leq 13$ (for the remaining cases).

By Theorem 3.14 (1), we know $\rho_0 \leq 7$. We set $m_0 = n_1$. Keep the same notation as in 2.3. Our proof is organized according to the type of f . Note that $P_{m_0} \geq 3$ and $m_0 \leq 13$. By Theorem 5.1, we only need to care about the situation $q(X) = 0$.

Case 1. f is of type I_3 .

Take G to be the movable part of $|2\sigma^*(K_{S_0})|$. Claim A implies that Assumption 2.9(2) is satisfied whenever $m \geq 35 \geq \rho_0 + 2m_0 + 2$. Clearly, by Lemma 2.16, Assumption 2.9(1) is also satisfied. As seen in the later part of 2.28, we can take a rational number $\beta \mapsto \frac{p}{2m_0+2p} \geq \frac{1}{m_0+2}$. Now inequality (2.2) gives $\xi \geq \frac{4}{45}$. Take $m = 35$. Then $\alpha = (35 - 1 - \frac{m_0}{2} - \frac{1}{\beta})\xi \geq \frac{10}{9} > 1$. Theorem 2.11 gives $\xi \geq \frac{4}{35}$. Take $m = 32$. Then $\alpha > 1$. We will see $\xi \geq \frac{1}{8}$ similarly. Now, for $m \geq 39$, $\alpha \geq (39 - 1 - \frac{13}{2} - 15)\xi \geq \frac{33}{16} > 2$. Theorem 2.11 says that φ_m is birational for all $m \geq 39$.

Case 2. f is of type II or III .

We take $\tilde{m}_0 = n_0$ and $m_1 = n_1$. We still use the mechanics of 2.3 to study $\varphi_{\tilde{m}_0}$ instead of φ_{m_0} . But most notations will use the symbol $\tilde{\cdot}$. Note that $\tilde{m}_0 \leq 10$ and $P_{\tilde{m}_0} \geq 2$.

If \tilde{f} is of type II or III , Theorem 2.20(3) and Theorem 2.22(3) imply that φ_m is birational for $m \geq 38$.

If \tilde{f} is of type I_n or I_p , we take \tilde{G} to be the movable part of $|M_{m_1}|_{\tilde{S}}$, where \tilde{S} is a generic irreducible element of $|M_{\tilde{m}_0}|$. Clearly, $h^0(\tilde{S}, M_{m_1}|_{\tilde{S}}) \geq 2$ since $\dim \varphi_{m_1}(X) \geq 2$. Thus we are in the situation with $m_1\pi^*(K_X)|_{\tilde{S}} \geq \tilde{G}$. We may always take a sufficiently good $\tilde{\pi}$ instead of π . Now Lemma 2.16 and Lemma 5.4 imply that Assumptions 2.9(1) and 2.9(2) are simultaneously satisfied for $m \geq 30 \geq \rho_0 + \tilde{m}_0 + m_1$. Finally, we study the value of α . Clearly, one may take $\tilde{\beta} = \frac{1}{m_1}$. Thus inequality (2.2) says $\xi \geq \frac{2}{1+\tilde{m}_0+m_1}$. For situations (1) and (2), we have $\xi \geq \frac{2}{23}$. Take $m = 35$. Then $\alpha \geq \frac{24}{23} > 1$. Theorem 2.11 gives $\xi \geq \frac{4}{35}$. Take $m = 32$. Then, similarly, we get $\xi \geq \frac{1}{8}$. Take $m \geq 40$. Then $\alpha \geq \frac{17}{8} > 2$. Theorem 2.11 implies that φ_m is birational for all $m \geq 40$. We are done. q.e.d.

L. Zhu [30] showed φ_m is birational for $m \geq 46$.

Theorem 5.6. *Let X be a projective minimal 3-fold of general type with $\chi(\mathcal{O}_X) > 1$. Then φ_m is birational for all $m \geq 73$.*

Proof. By Theorem 5.1, we only need to consider the situation $q(X) = 0$. According to Lemma 2.32, the induced fibration f from φ_{m_0} is of type III , II , or I_p .

If $P_{m_0} \geq 2$ for some $m_0 \leq 16$, then, by Theorems 2.20, 2.22, and 2.25, φ_m is birational for all $m \geq 69$. Assume $P_m \leq 1$ for all $m \leq 16$. Then we have a complete classification for B_{\min} as in Table C. More precisely, we see $B \succcurlyeq B_{2a}$, $B \succcurlyeq B_{3a}$ and $B \succ B_{20a}$, $B \succ B_{52a}$, noting that case No.22 doesn't happen by Proposition 4.5. As we have observed in the proof of Theorem 4.6, for cases No. 20a and No. 52a, we actually have $B = B_{20}$ and $B = B_{52}$. Thus we see $P_{14}(X) \geq 2$ in both cases, a contradiction. We are left to study cases No. 2a and No. 3a, which correspond to two formal baskets, $(B_{2a}, 2, 0)$ and $(B_{3a}, 3, 0)$, where

$$B_{2a} = \{4 \times (1, 2), (4, 9), (2, 5), (5, 13), 3 \times (1, 3), 2 \times (1, 4)\},$$

$$B_{3a} = \{6 \times (1, 2), (5, 11), 3 \times (2, 5), (5, 13), 4 \times (1, 3), (2, 7), 2 \times (1, 4)\}.$$

The computation gives the following datum:

	ρ_0	μ_1	μ_2	μ_3
B_{2a}	20	18	24	30
B_{3a}	20	18	20	30

When $B \succcurlyeq B_{3a}$, we have $P_{20}(X) = P_{20}(B) \geq P_{20}(B_{3a}) \geq 3$. Theorems 2.20 and 2.29 imply that φ_m is birational for all $m \geq 66$ unless f is type *II*. Indeed, if f is of type *II* and $m_0 = 20$, at least we have $\xi \geq \frac{2}{31}$, following the argument in 2.21. Take $m = 57$; we have $\alpha > 1$ and hence $\xi \geq \frac{4}{57}$. Now take $m \geq 70$; we have $\alpha = (70 - 41)\frac{4}{57} > 2$. Thus φ_m is birational for all $m \geq 70$.

Now the theorem follows from the following claim.

Claim B. When $B \succcurlyeq B_{2a}$, φ_m is birational for all $m \geq 73$.

The proof is similar to that of Theorem 5.5, Case 1 and Case 2. We have known $\rho_0 \leq 20$. We can find two numbers $n_0 \leq 18$ and $n_1 \leq 24$ with $P_{n_0}(X) \geq 2$ and $P_{n_1}(X) \geq 3$. First, we set $m_0 = n_1$. Keep the same notation as in 2.3. Our proof is organized according to the type of f . Note that $P_{m_0} \geq 3$ and $m_0 \leq 24$.

Case i. f is of type I_3 .

By Lemma 2.32, f must be of type I_p . Take $G = 2\sigma^*(K_{S_0})$. Claim A implies that Assumption 2.9(2) is satisfied whenever $m \geq 70 \geq \rho_0 + 2m_0 + 2$. Clearly, by Lemma 2.16, Assumption 2.9(1) is also satisfied. As seen in the latter part of 2.28, we can take a rational number $\beta \mapsto \frac{p}{2m_0+2p} \geq \frac{1}{m_0+2}$. Note that $|G|$ is base point free, and we have $\deg(K_C) \geq 6$. Now inequality (2.2) gives $\xi \geq \frac{2}{13}$. For $m \geq 70$, $\alpha \geq (70 - 1 - 12 - 26)\xi > 2$. Theorem 2.11 says that φ_m is birational for all $m \geq 70$.

Case ii. f is of type *II* or *III*.

We take $\tilde{m}_0 = n_0$ and $m_1 = n_1$. We still use the mechanics of 2.3 to study $\varphi_{\tilde{m}_0}$ instead of φ_{m_0} . Noting that $\tilde{m}_0 \leq 18$, when \tilde{f} is of type *III*

or *II*, Theorems 2.20 and 2.22 imply that φ_m is birational for all $m \geq 66$. We are left to study the situation with \tilde{f} being of type *I*. We take \tilde{G} to be the movable part of $|M_{m_1}|_{\tilde{S}}$. Clearly, $h^0(\tilde{S}, M_{m_1}|_{\tilde{S}}) \geq 2$ since $\dim \varphi_{m_1}(X) \geq 2$. Thus we are in the situation with $m_1\pi^*(K_X)|_{\tilde{S}} \geq \tilde{G}$. Now Lemma 2.16 and Lemma 5.4 imply that Assumptions 2.9(1) and 2.9(2) are simultaneously satisfied for $m \geq 62 \geq \rho_0 + \tilde{m}_0 + m_1$. Clearly, one may take $\tilde{\beta} = \frac{1}{m_1}$. Thus inequality (2.2) says $\xi \geq \frac{2}{1+\tilde{m}_0+m_1} \geq \frac{2}{43}$.

Take $m = 65$. Then $\alpha \geq \frac{44}{43} > 1$. Theorem 2.11 gives $\xi \geq \frac{4}{65}$. Take $m = 60$. Then similarly we get $\xi \geq \frac{1}{15}$. Take $m = 59$. Then we shall get $\xi \geq \frac{4}{59}$. Take $m = 58$ and we obtain $\xi \geq \frac{2}{29}$. Eventually, for $m \geq 73$, we see $\alpha > 2$ and Theorem 2.11 implies that φ_m is birational for all $m \geq 73$. We are done. q.e.d.

We have proved all the main results. Indeed, Theorem 1.1 follows from Theorem 5.1, Theorem 5.3, Theorem 5.5, and Theorem 5.6. Theorem 1.2 follows from Theorem 3.11 and Theorem 4.6.

Finally, we would like to ask the following:

Question 5.7. Can one find an optimal lower bound for K^3 ?

The following problem is very interesting.

Open Problem 5.8. Can one find a minimal 3-fold X of general type with $q(X) = 0$ and $\chi(\mathcal{O}_X) > 1$?

References

- [1] W. Barth, C. Peters & A. Van de Ven, *Compact complex surfaces*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) 4, Springer-Verlag, Berlin, 1984. x+304 pp., MR 0749574, Zbl 1036.14016.
- [2] E. Bombieri, *Canonical models of surfaces of general type*, Inst. Hautes Etudes Sci. Publ. Math., **42** (1973), 171–219, MR 0318163, Zbl 0259.14005.
- [3] J. A. Chen & C. D. Hacon, *Pluricanonical systems on irregular 3-folds of general type*, Math. Z., **255** (2007), no. 2, 343–355, MR 2262735, Zbl 1094.14502.
- [4] J. A. Chen & M. Chen, *Explicit birational geometry of 3-folds of general type, I*, Ann. Sci. École Norm. Sup., **43** (2010), 365–394.
- [5] J. A. Chen & M. Chen, *The Canonical volume of 3-folds of general type with $\chi \leq 0$* , J. London Math. Soc., **78** (2008), 693–706, MR 2456899, Zbl 1156.14009.
- [6] M. Chen, *Canonical stability in terms of singularity index for algebraic threefolds*, Math. Proc. Camb. Phil. Soc., **131** (2001), 241–264, MR 1857118, Zbl 1068.14045.
- [7] M. Chen, *On the \mathbf{Q} -divisor method and its application*, J. Pure Appl. Algebra, **191** (2004), 143–156, MR 2048311, Zbl 1049.14034.
- [8] M. Chen & K. Zuo, *Complex projective threefolds with non-negative canonical Euler-Poincare characteristic*, Comm. Anal. Geom., **16** (2008), 159–182, MR 2411471, Zbl 1149.14034.
- [9] C. Ciliberto, *The bicanonical map for surfaces of general type*, Proc. Symposia in Pure Math., **62** (1997), 57–83, MR 1492518, Zbl 0929.14022.

- [10] C. D. Hacon & J. McKernan, *Boundedness of pluricanonical maps of varieties of general type*, Invent. Math., **166** (2006), 1–25, MR 2242631, Zbl 1121.14011.
- [11] A. R. Iano-Fletcher. *Contributions to Riemann-Roch on Projective 3-folds with Only Canonical Singularities and Applications*, Proceedings of Symposia in Pure Mathematics, **46** (1987), 221–231, MR 0927958, Zbl 0662.14026.
- [12] A. R. Iano-Fletcher, *Working with weighted complete intersections*, In *Explicit birational geometry of 3-folds*, London Mathematical Society, Lecture Note Series, 281, Cambridge University Press, Cambridge, 2000, MR 1798982.
- [13] Y. Kawamata, *A generalization of Kodaira-Ramanujam's vanishing theorem*, Math. Ann., **261** (1982), 43–46, MR 0675204, Zbl 0476.14007.
- [14] Y. Kawamata, K. Matsuda & K. Matsuki, *Introduction to the minimal model problem*, Adv. Stud. Pure Math., **10** (1987), 283–360, MR 0946243, Zbl 0672.14006.
- [15] J. Kollár, *Higher direct images of dualizing sheaves I*, Ann. Math., **123**(1986), 11–42, MR 0825838, Zbl 0598.14015; II, *ibid.* **124** (1986), 171–202, MR 0847955, Zbl 0605.14014.
- [16] J. Kollár & S. Mori, *Birational geometry of algebraic varieties*, 1998, Cambridge Univ. Press, MR 1658959, Zbl 0926.14003.
- [17] M. Mendes Lopes, *Adjoint systems on surfaces*, Boll. Unione Mat. Ital., (7)**10-A** (1996), 169–179, MR 1386254, Zbl 0861.14023.
- [18] Y. Miyaoka, *The Chern classes and Kodaira dimension of a minimal variety*, Algebraic Geometry, Sendai, 1985. Adv. Stud. Pure Math., **10** (1987), 449–476, MR 0946247, Zbl 0648.14006.
- [19] M. Reid, *Canonical 3-folds*, in A. Beauville, editor, *Géométrie Algébrique Angers*, pages 273–310, Sijthoff & Noordhoff, 1980, MR 0605348, Zbl 0451.14014.
- [20] M. Reid, *Minimal models of canonical 3-folds*, Adv. Stud. Pure Math., **1** (1983), 131–180, MR 0715649, Zbl 0558.14028.
- [21] M. Reid, *Young person's guide to canonical singularities*, Proc. Symposia in pure Math., **46** (1987), 345–414, MR 0927963, Zbl 0634.14003.
- [22] D.-K. Shin, *On a computation of plurigenera of a canonical threefold*, J. Alg., **309** (2007), 559–568, MR 2303193, Zbl 1116.14032.
- [23] S. Takayama, *Pluricanonical systems on algebraic varieties of general type*, Invent. Math., **165** (2006), 551–587, MR 2242627, Zbl 1108.14031.
- [24] S. G. Tankeev, *On n -dimensional canonically polarized varieties and varieties of fundamental type*, Izv. A. N. SSSR, Ser. Math., **35** (1971), 31–44, MR 0277528, Zbl 0248.14005.
- [25] H. Tsuji, *Pluricanonical systems of projective varieties of general type, I*, Osaka J. Math., **43** (2006), no. 4, 967–995, MR 2303558, Zbl 1142.14012.
- [26] E. Viehweg, *Vanishing theorems*, J. reine angew. Math., **335** (1982), 1–8, MR 0667459, Zbl 0485.32019.
- [27] E. Viehweg, *Weak positivity and the additivity of the Kodaira dimension for certain fibre spaces*, Proc. Algebraic Varieties and Analytic Varieties, Tokyo 1981. Adv. Studies in Math., **1**, Kinokunya-North-Holland Publ., 1983, 329–353, MR 0715656, Zbl 0513.14019.
- [28] G. Xiao, *Finitude de l'application bicanonique des surfaces de type général*, Bull. Soc. Math. France, **113** (1985), 23–51, MR 0807825, Zbl 0611.14031.

- [29] L. Zhu, *The sharp lower bound for the volume of threefolds of general type with $\chi(\mathcal{O}_X) = 1$* , Math. Zeit., **261** (2009), no. 1, 123–141, MR 2452640, Zbl 1154.14033.
- [30] L. Zhu, *On birational geometry of algebraic 3-folds of general type*, PhD thesis, Fudan University, 2008.

DEPARTMENT OF MATHEMATICS
NATIONAL TAIWAN UNIVERSITY
TAIPEI, 106
TAIWAN

E-mail address: jkchen@math.ntu.edu.tw

SCHOOL OF MATHEMATICAL SCIENCES & LMNS
FUDAN UNIVERSITY
SHANGHAI, 200433
PEOPLE'S REPUBLIC OF CHINA

E-mail address: mchen@fudan.edu.cn