

# One condition for solution uniqueness and robustness of both $\ell_1$ -synthesis and $\ell_1$ -analysis minimizations

Hui Zhang<sup>1</sup> · Ming Yan<sup>2</sup> · Wotao Yin<sup>3</sup>

Received: 30 October 2014 / Accepted: 23 June 2016 /  
Published online: 6 July 2016  
© Springer Science+Business Media New York 2016

**Abstract** The  $\ell_1$ -synthesis model and the  $\ell_1$ -analysis model recover structured signals from their undersampled measurements. The solution of the former is a sparse sum of dictionary atoms, and that of the latter makes sparse correlations with dictionary atoms. This paper addresses the question: when can we trust these models to recover specific signals? We answer the question with a condition that is both necessary and sufficient to guarantee the recovery to be unique and exact and, in the presence of measurement noise, to be robust. The condition is one-for-all in the sense that it applies to both the  $\ell_1$ -synthesis and  $\ell_1$ -analysis models, to both constrained and unconstrained formulations, and to both the exact recovery and robust recovery cases. Furthermore, a convex infinity-norm optimization problem is introduced for numerically verifying the condition. A comprehensive comparison with related existing conditions is included.

---

Communicated by: Yuesheng Xu.

---

✉ Ming Yan  
yanm@math.msu.edu

Hui Zhang  
h.zhang1984@163.com

Wotao Yin  
wotaoyin@math.ucla.edu

<sup>1</sup> Department of Mathematics and Systems Science, College of Science, National University of Defense Technology, Changsha, Hunan 410073, China

<sup>2</sup> Department of Computational Mathematics, Science and Engineering; Department of Mathematics, Michigan State University, East Lansing, MI 48824, USA

<sup>3</sup> Department of Mathematics, University of California, Los Angeles, CA 90095, USA

**Keywords** Exact recovery · Robust recovery ·  $\ell_1$ -analysis ·  $\ell_1$ -synthesis · Sparse optimization · Compressive sensing

**Mathematics Subject Classification (2010)** 65K05 · 90C25

## 1 Introduction

With both synthesis and analysis models of sparse signal recovery [1], one is interested in when the recovery is successful, namely, whether the solution is unique and whether the solution error is proportional to the amount of noise in the measurements.

Various *sufficient* conditions have been proposed to guarantee successful recovery. Let us first discuss two types of recovery conditions. The *uniform recovery* conditions guarantee not only the successful recovery of one sparse signal but all signals that are sufficiently sparse, irrelevant of the locations of their nonzero entries. The *non-uniform recovery conditions*, however, focus on the recovery of a restricted set of sufficiently sparse signals, for example, the sparse signals with a specific support. Since non-uniform recovery conditions cover fewer signals, they are in general weaker and thus easier to hold than uniform recovery conditions. In addition, some non-uniform recovery conditions, especially those for fixed signal supports, are easier to verify numerically, whereas all the existing uniform recovery conditions are numerically intractable to verify on given sensing matrices. On the other hand, several kinds of random matrices (such as those with i.i.d. subgaussian entries) satisfy the uniform recovery conditions with high probability. The uniform conditions are useful in designing randomized linear measurements, whereas non-uniform conditions are useful for deterministic linear measurements on restricted sets of signals.

Non-uniform conditions for  $\ell_1$ -synthesis minimization include the non-uniform dual certificate condition [2] and the “RIPless” property [3]. Well-known examples of uniform conditions include the restricted isometry principle [4], the null space condition [5], the spherical section property [6], and others.

Because  $\ell_1$ -analysis minimization takes a more general form than  $\ell_1$ -synthesis minimization, some of the above non-uniform recovery conditions have been extended to the analysis case. Recent works [7–13] have made significant contributions.

This paper studies the so-called *dual certificate* condition, a type of non-uniform condition. We show that if a signal is a solution to any model among (1a)–(1c) described below, there exists a necessary and sufficient condition, same for any of the three models, that guarantees that the signal will be uniquely recovered, i.e., each of models (1a)–(1c) has the unique solution equal to the signal. While this result has been partially known in previous work for  $\ell_1$  minimization as a sufficient condition, we establish three new results:

- the condition previously known to be sufficient is, in fact, necessary;
- the condition guarantees robustness to noise. That is, under this condition, if the observed data is contaminated by arbitrary noise, the solution to either Eqs. 1b or 1c is robust to the noise in the sense that the solution error is proportional to the Euclidean norm of the noise;

- nearly the same condition but imposed on the support of the largest  $|I|$  entries of an approximately sparse signal guarantees the robust recovery by model (1c).

These results complete the theory of non-uniform recovery of the  $\ell_1$ -synthesis and  $\ell_1$ -analysis models. Very recently, the authors in [14] studied generic regularized optimization models that include the  $\ell_1$ -synthesis and  $\ell_1$ -analysis models as special cases. They stated that a non-degenerate dual certificate, which generalizes the dual certificate condition in this paper, is sufficient for solution uniqueness. However, it is not necessary in general cases [15]. Thus, a natural question is in which cases this condition is also necessary for solution uniqueness. Our first result provides a partial answer to this question.

The proposed condition is compared to existing non-uniform conditions in the literature, most of which are stronger than ours and are thus sufficient yet not necessary. Note that some of those stronger ones also give additional properties that ours does not. Technically, a part of our analysis is inspired by existing results in [7, 16–20], which bear certain similarity among themselves and will be mentioned in later sections.

The rest of the paper is organized as follows. Sections 2 and 3 formulate the problem and state the main results. Section 4 reviews several related results. Section 5 discusses condition verification. Proofs for the main results are given in Sections 6, 7, and 8.

## 2 Problem formulation and contributions

### 2.1 Notation

We equip  $\mathbb{R}^n$  with the canonical scalar product  $\langle \cdot, \cdot \rangle$  and the Euclidean norm  $\| \cdot \|_2$ . We let  $|\cdot|$  return the cardinality if the input is a set or the absolute value if the input is a number. For any  $x \in \mathbb{R}^n$ ,  $\text{supp}(x) = \{k : 1 \leq k \leq n, x_k \neq 0\}$  is the index set of the non-zero entries of  $x$ .  $\text{sign}(x)$  is the vector whose  $i$ th entry is the sign of  $x_i$ , taking a value among  $+1, -1$ , and  $0$ . For any  $p \geq 1$ , the  $\ell_p$ -norm of  $x \in \mathbb{R}^n$  is

$$\|x\|_p = \left( \sum_{i=1}^n |x_i|^p \right)^{1/p},$$

its  $\ell_0$ -“norm” is  $\|x\|_0 = |\text{supp}(x)|$ , and its  $\ell_\infty$ -norm is  $\|x\|_\infty = \max\{|x_i| : i = 1, \dots, n\}$ . For  $x \in \mathbb{R}^n$  and  $I \subset \{1, 2, \dots, n\}$ ,  $x_I$  denotes the vector formed by the entries  $x_i$  of  $x$  for  $i \in I$ , and  $I^c$  is the complement of  $I$ . Similarly,  $A_I$  is the submatrix formed by the columns of  $A$  indexed by  $I$ .  $A^T$  is the transpose of  $A$ . We use  $A_I^T$  for the transpose of submatrix  $A_I$ , not a submatrix of  $A^T$ . For a square matrix  $A$ ,  $\lambda_{\max}(A)$  and  $\lambda_{\min}(A)$  denote its largest and smallest eigenvalues, respectively,  $\text{Cond}(A)$  denotes its condition number, and  $\|A\|$  denotes its spectral norm. The null and column spaces of  $A$  are denoted by  $\text{Ker}(A)$  and  $\text{Im}(A)$ , respectively.

## 2.2 Problem formulations

Let  $x^* \in \mathbb{R}^n$  be a signal of interest. This paper studies when  $\ell_1$  minimization can uniquely and robustly recover  $x^*$  from its linear measurements

$$b = \Phi x^* + w,$$

where  $\Phi \in \mathbb{R}^{m \times n}$  is a certain matrix and  $w \in \mathbb{R}^m$  is noise. We focus on the setting  $m \leq n$ .

The results of this paper cover the following  $\ell_1$  minimization formulations:

$$\underset{x}{\text{minimize}} \|\Psi^\top x\|_1, \quad \text{subject to } \Phi x = b, \quad (1a)$$

$$\underset{x}{\text{minimize}} \|\Phi x - b\|_2^2 + \lambda \|\Psi^\top x\|_1, \quad (1b)$$

$$\underset{x}{\text{minimize}} \|\Psi^\top x\|_1, \quad \text{subject to } \|\Phi x - b\|_2 \leq \delta, \quad (1c)$$

where  $\Psi \in \mathbb{R}^{n \times l}$  and  $\delta$  and  $\lambda$  are positive parameters.

These three models are applied at different cases. Model (1a) is used if no noise is present, i.e.,  $w = 0$ . When there is noise in the measurements, models (1b) and (1c) are used, and models (1b) and (1c) are equivalent in the sense that given  $\lambda$  in model (1b), we can find  $\delta$  for model (1c) such that they have the same optimal solution set, and vice versa. Depending on the prior information we have, one model is chosen over the other model. Model (1c) is applied when the  $\ell_2$  norm of the noise is given or can be accurately estimated.

If  $\Psi = Id$ , the identify matrix, models in Eq. 1 are referred to as the  $\ell_1$  (or more generally,  $\ell_1$ -synthesis) models. If  $\Psi \neq Id$ , they are referred to as the  $\ell_1$ -analysis models, which are recently reviewed in [1]. The  $\ell_1$ -analysis models, including the cospase analysis model [21] and the total variation model [22] as widely known examples, have recently attracted a lot of attention. The underlying signal is expected to make sparse correlations with the columns (atoms) in an possibly overcomplete dictionary  $\Psi$ , i.e.,  $\Psi^\top x^*$  is sparse; see [21, 23, 24].

## 2.3 Geometry

It is worth noting that the underlying geometry of solution uniqueness is rather clear. Indeed, it was characterized in terms of polytope faces by Donoho in [25], and also of null spaces and tangent cones by Chandrasekaran et al. in [26]. However, there lacks a concrete *if-and-only-if* condition for one to check and the robustness results are difficult to obtain from geometry.

Let us describe the geometry of unique recovery under the simple setting:  $\Psi = Id$ . Then,  $x^*$  is the unique solution to Eq. 1a if the affine set  $\mathcal{F} = \{x : \Phi x = b\}$  touches the  $\ell_1$  polytope  $\{x : \|x\|_1 = \gamma\}$ , for some  $\gamma > 0$ , at a *unique* point on a face of the polytope. The uniqueness for Eqs. 1b and 1c is similar except  $\mathcal{F}$  is replaced by a “tube”  $\mathcal{T} = \{x : \|\Phi x - b\|_2 \leq \sigma\}$ . In both cases, the uniqueness is certified by (i) a kernel condition that ensures touching the face at just one point, and (ii) the existence of a hyperplane (dual certificate) that separates  $\mathcal{F}$  or  $\mathcal{T}$  from the polytope and creates

a positive gap between them outside the face. The same geometry applies to  $\ell_1$ -analysis minimization but involves the more complicated polytope  $\{x : \|\Psi^\top x\|_1 = \gamma\}$ ; see [27, 28].

This work not only formalizes these geometrical explanations with a concrete *if-and-only-if* condition, but also establishes robustness bounds under the same condition, which cannot be explained by the above geometry. Suppose that noise is present in the measurements. Then, the point where  $\mathcal{F}$  or  $\mathcal{T}$  touches the polytope will move, and the point may or may not stay in the same face as in the noise-free case. The above geometry cannot clearly explain how far the point will move due to noise.

### 3 Main condition and results

#### 3.1 Main condition

**Condition 1** Given  $\bar{x} \in \mathbb{R}^n$ , index sets  $I = \text{supp}(\Psi^\top \bar{x}) \subset \{1, \dots, l\}$  and  $J = I^c$  satisfy

- (1)  $\text{Ker}(\Psi_J^\top) \cap \text{Ker}(\Phi) = \{0\}$ ;
- (2) There exists  $y \in \mathbb{R}^l$  such that  $\Psi y \in \text{Im}(\Phi^\top)$ ,  $y_I = \text{sign}(\Psi_I^\top \bar{x})$ , and  $\|y_J\|_\infty < 1$ .

It is not difficult to understand the condition as we shall explain below.

Condition 1 part (1) says that there does *not* exist any nonzero  $\Delta x$  satisfying both  $\Psi_J^\top \bar{x} = \Psi_J^\top (\bar{x} + \Delta x)$  and  $\Phi \bar{x} = \Phi (\bar{x} + \Delta x)$ . Otherwise, there exists a nonempty line segment  $\mathcal{I}$  connecting  $\bar{x} - \alpha \Delta x$  and  $\bar{x} + \alpha \Delta x$  for some sufficiently small  $\alpha > 0$  such that  $\Phi x = \Phi \bar{x}$  and  $\|\Psi^\top x\|_1$  is constant for  $x \in \mathcal{I}$ . Hence,  $\bar{x}$  cannot be the unique minimizer. This condition implies that the role of  $\ell_1$  minimization is “limited” to recovering the support set  $I$ ; if an oracle gives us  $I$ , we must solely rely on  $\Psi$ ,  $\Phi$ , and  $b$  to recover  $\bar{x}$ , since  $\|\Psi^\top x\|_1$  is locally linear near  $\bar{x}$  over  $\{x : \text{supp}(\Psi^\top x) = I\}$  and, thus, lacks the ability to pick  $\bar{x}$  out. In addition, part (1) of the condition implies the existence of  $\rho$  and  $\tau$  such that

$$\|\Psi_I^\top x\|_2 \leq \rho \|\Psi_J^\top x\|_1 + \tau \|\Phi x\|_2 \tag{2}$$

for all  $x \in \mathbb{R}^n$ , which will be used later in Theorem 3.

Condition 1 part (2) states the existence of a strictly-complementary *dual certificate*  $y$ . To see this, let us check a part of the optimality condition of Eq. 1a:  $0 \in \Psi \partial \|\cdot\|_1(\Psi^\top x) - \Phi^\top \beta$ , where vector  $\beta$  is the Lagrangian multiplier. We rewrite the condition as  $0 = \Psi y - \Phi^\top \beta$  where  $y \in \partial \|\cdot\|_1(\Psi^\top x)$ , which translates to  $y_I = \text{sign}(\Psi_I^\top x)$  and  $\|y_J\|_\infty \leq 1$ . This  $y$  certifies the optimality of  $\bar{x}$ . For solution uniqueness and/or robustness, we shall later show that the strict inequality  $\|y_J\|_\infty < 1$  is necessary. Similar to the role played by the well-known Karush-Kuhn-Tucker (KKT) condition in convex optimization, Condition 1 serves as a necessary and sufficient condition for a signal not only to be a solution, but also to be the unique solution. This, hence, strengthens the corresponding results deduced by the KKT condition.

It is worth mentioning a variant of Condition 1 as follows.

**Condition 2** Given  $\bar{x} \in \mathbb{R}^n$ , let  $I = \text{supp}(\Psi^\top \bar{x})$ . There exists a nonempty index set  $J \subseteq I^c$  such that the index sets  $I$ ,  $J$ , and  $K = (I \cup J)^c$  satisfy

- (1)  $\text{Ker}(\Psi_J^\top) \cap \text{Ker}(\Phi) = \{0\}$ ;
- (2) There exists  $y \in \mathbb{R}^l$  such that  $\Psi y \in \text{Im}(\Phi^\top)$ ,  $y_I = \text{sign}(\Psi_I^\top \bar{x})$ ,  $\|y_J\|_\infty < 1$ , and  $\|y_K\|_\infty \leq 1$ .

In Condition 2, a smaller  $J$  relaxes part (2) but gives a larger  $\text{Ker}(\Psi_J^\top)$  and thus tightens part (1). Although Condition 2 allows a more flexible  $J$  than Condition 1, we shall show that they are equivalent.

The comparisons between Condition 1 and those in the existing literature are given in Section 4.

### 3.2 Main results

Depending on the specific models in Eq. 1, we need the following assumptions:

**Assumption 1** Matrix  $\Phi$  has full row-rank.

**Assumption 2**  $\lambda_{\max}(\Psi \Psi^\top) = 1$ .

**Assumption 3** Matrix  $\Psi$  has full row-rank.

Assumptions 1 and 3 are standard to avoid redundancy. Assumption 2 is non-essential as we can scale any general matrix  $\Psi$  by multiplying  $\frac{1}{\sqrt{\lambda_{\max}(\Psi \Psi^\top)}}$ . Below we state our main results and delay their proofs to Sections 6–8.

**Theorem 1** (Uniqueness) Under Assumption 1, let  $\hat{x}$  be a solution to problem (1a) or (1b), or under Assumptions 1 and 3, let  $\hat{x}$  be a solution to problem (1c). The following statements are equivalent:

- 1) Solution  $\hat{x}$  is unique;
- 2) Condition 1 holds for  $\bar{x} = \hat{x}$ ;
- 3) Condition 2 holds for  $\bar{x} = \hat{x}$ .

This theorem states that Conditions 1 and 2 are equivalent, and they are necessary and sufficient for a solution  $\hat{x}$  to problem (1a), or to problem (1b), or to problem (1c) to be unique. To state our next result on robustness, we let

$$r(J) := \sup_{u \in \text{Ker}(\Psi_J^\top) \setminus \{0\}} \frac{\|u\|_2}{\|\Phi u\|_2}.$$

Part (1) of Condition 1 ensures that  $0 < r(J) < +\infty$ . If  $\Psi = Id$ , then  $u \in \text{Ker}(\Psi_J^\top) \setminus \{0\}$  is a sparse nonzero vector with maximal support  $J^c$ , and  $r(J)$  is the

inverse of the minimal singular value of the submatrix  $\Phi_{J^c}$ . Below we claim that Condition 1 ensures the robustness of problems (1b) and (1c) to arbitrary noise in  $b$ .

**Theorem 2** (Robustness to measurement noise) *Under Assumptions 1–3, given an original signal  $x^* \in \mathbb{R}^n$ , let  $I = \text{supp}(\Psi^\top x^*)$  and  $J = I^c$ . For arbitrary noise  $w$ , let  $b = \Phi x^* + w$  and  $\delta = \|w\|_2$ . If Condition 1 is met for  $\bar{x} = x^*$ , then*

- 1) *For any  $C_0 > 0$ , there exists a constant  $C_1 > 0$  such that every minimizer  $x_{\delta,\lambda}$  of problem (1b) using parameter  $\lambda = C_0\delta$  satisfies*

$$\|\Psi^\top(x_{\delta,\lambda} - x^*)\|_1 \leq C_1\delta;$$

- 2) *Every minimizer  $x_\delta$  of problem (1c) satisfies*

$$\|\Psi^\top(x_\delta - x^*)\|_1 \leq C_2\delta.$$

The constraints  $C_1$  and  $C_2$  are given as follows. Define

$$\beta = (\Phi\Phi^\top)^{-1}\Phi\Psi y, \quad C_3 = r(J)\sqrt{|I|}, \quad \text{and} \quad C_4 = \frac{1 + \text{Cond}(\Psi)\|\Phi\|C_3}{1 - \|y_J\|_\infty},$$

with which we have

$$C_1 = 2C_3 + C_0\|\beta\|_2 + \frac{(1 + C_0\|\beta\|_2/2)^2C_4}{C_0},$$

$$C_2 = 2C_3 + 2C_4\|\beta\|_2.$$

*Remark 1* From the results of Theorem 2, it is straightforward to derive  $\ell_1$  and  $\ell_2$  bounds for  $x_{\delta,\lambda} - x^*$  and  $x_\delta - x^*$  under Assumption 2.

*Remark 2* Since  $C_0$  is free to choose, one can choose the optimal  $C_0 = \sqrt{\frac{4C_4}{4\|\beta\|_2 + C_4\|\beta\|_2^2}}$  and simplify  $C_1$  to

$$C_1 = 2C_3 + C_4\|\beta\|_2 + \sqrt{C_4^2\|\beta\|_2^2 + 4C_4\|\beta\|_2} \leq 2C_3 + 2C_4\|\beta\|_2 + 2,$$

which becomes very similar to  $C_2$ . This reflects the equivalence between problems (1b) and (1c) in the sense that given  $\lambda$ , one can find  $\delta$  such that they have the same solution, and vice versa.

*Remark 3* Both  $C_1$  and  $C_2$  are the sum of  $2C_3$  and other terms.  $2C_3$  alone bounds the error when  $\Psi^\top x_{\delta,\lambda}$  (or  $\Psi^\top x_\delta$ ) and  $\Psi^\top x^*$  have matching signs. Since  $C_3$  does not depend on  $y_J$ , part (2) of Condition 1 does not play any role, whereas part (1) plays the major role. When the signs of  $\Psi^\top x_{\delta,\lambda}$  (or  $\Psi^\top x_\delta$ ) and  $\Psi^\top x^*$  do not match, the remaining terms in  $C_1$  and  $C_2$  are involved, and they are affected by part (2) of Condition 1; in particular,  $\|y_J\|_\infty < 1$  plays a big role as  $C_4$  is inversely proportional to  $1 - \|y_J\|_\infty$ . Also, since there is no knowledge about the support of  $\Psi^\top x_{\delta,\lambda}$ , which may or may not equal to that of  $\Psi^\top x^*$ ,  $C_4$  inevitably depends on the global properties of  $\Psi$  and  $\Phi$ . In contrast,  $C_3$  only depends on the restricted property of  $\Phi$ .

When  $\Psi^\top x^*$  is only approximately sparse, we need to choose  $I$  as the support of the  $|I|$  largest entries of  $\Psi^\top x^*$  in Condition 1. Then we have the following robustness result of approximately sparse signals for Eq. 1c.

**Theorem 3** (Robustness for approximately sparse signals) *Under Assumption 3, given an original signal  $x^* \in \mathbb{R}^n$  with  $|I|$  largest entries of  $\Psi^\top x^*$  supported on  $I$ , let  $J = I^c$ . For arbitrary noise  $w$ , let  $b = \Phi x^* + w$  and  $\delta = \|w\|_2$ . If Condition 1 is met on  $I$ , i.e.,  $\text{Ker}(\Psi_J^\top) \cap \text{Ker}(\Phi) = \{0\}$ , and there exist  $y \in \mathbb{R}^I$  and  $\beta \in \mathbb{R}^m$  such that  $\Psi y = \Phi^\top \beta$  and*

$$y_I = \text{sign}(\Psi_I^\top x^*), \quad \|y_J\|_\infty < 1, \tag{3}$$

then there exist  $\rho$  and  $\tau$  such that

$$\|\Psi_I^\top x\|_2 \leq \rho \|\Psi_J^\top x\|_1 + \tau \|\Phi x\|_2, \tag{4}$$

for all  $x \in \mathbb{R}^n$ , and every minimizer  $x_\delta$  of  $\|\Psi^\top x\|_1$  subjective to  $\|\Phi x - b\|_2 \leq \delta$  satisfies

$$\|\Psi^\top (x_\delta - x^*)\|_2 \leq \frac{2(1 + \rho)}{1 - \|y_J\|_\infty} \|\Psi_J^\top x^*\|_1 + \left( \frac{2(1 + \rho)\|\beta\|_2}{1 - \|y_J\|_\infty} + 2\tau \right) \delta. \tag{5}$$

*Remark 4* With Assumption 3, part (1) of Condition 1 is equivalent to the existence of  $\rho$  and  $\tau$  in Eq. 4 for all  $x \in \mathbb{R}^n$ . When part (1) of Condition 1 is satisfied, the existence of  $\rho$  and  $\tau$  is easy to show. If part (1) is not satisfied, then there exist  $x \neq 0$  such that  $\Psi_J^\top x = 0$  and  $\Phi x = 0$ . Assumption 3 gives us that  $\Psi_I^\top x \neq 0$ . Thus  $\|\Psi_I^\top x\|_2 > 0 = \rho \|\Psi_J^\top x\|_1 + \tau \|\Phi x\|_2$  for any  $\rho$  and  $\tau$ .

Condition (3) can be relaxed into

$$\|y_I - \text{sign}(\Psi_I^\top x^*)\|_2 \leq \theta_1, \quad \|y_J\|_\infty < 1.$$

In this case, denote  $\mu_1 := \rho\theta_1 + \|y_J\|_\infty$  and  $\mu_2 := \tau\theta_1 + \|\beta\|_2$ . If  $\mu_1 < 1$ , then every minimizer  $x_\delta$  of  $\|\Psi^\top x\|_1$  subjective to  $\|\Phi x - b\|_2 \leq \delta$  satisfies

$$\|\Psi^\top (x_\delta - x^*)\|_2 \leq \frac{2(1 + \rho)}{1 - \mu_1} \|\Psi_J^\top x^*\|_1 + \left( \frac{2(1 + \rho)\mu_2}{1 - \mu_1} + 2\tau \right) \delta.$$

*Remark 5* Theorem 3 is inspired by several existing results in the standard  $\ell_1$ -synthesis case; for example Theorem 3.1 in [19] and Theorem 4.33 and Exercise 4.17 in [20]. The result here can be viewed as an extension from the  $\ell_1$ -synthesis case to the  $\ell_1$ -analysis case.

### 4 Related works

In the case of  $\Psi = Id$ , Condition 1 is well known in the literature for  $\ell_1$ -synthesis models. It is initially proposed in [2] as a sufficient condition for solution uniqueness. For problems (1b) and (1c), [29, 30] present sufficient but non-necessary conditions for solution uniqueness. Later, its necessity is established in [17] for model (1b) and



then in [18] for all models in Eq. 1, assuming  $\Psi = Id$  or an orthogonal basis. The solution robustness of model (1b) is given under the same condition in [17]. Below we restrict our literature review to results for the  $\ell_1$ -analysis models.

### 4.1 Previous uniqueness conditions

Papers [7–9, 21, 31] establish the uniqueness of the  $\ell_1$ -analysis models, and some use stronger conditions than ours. Our purpose in the comparison is restricted to uniqueness and robustness to arbitrary noise. The reader should be aware that some of them also imply stronger results such as sign consistency by Conditions 4 and 6 below; while the rest such as Conditions 3 and 5 below are not used to derive stronger results yet.

The following condition in [21], which was proposed by strengthening the null space condition, guarantees the solution uniqueness for problem (1a):

**Condition 3** Given  $\bar{x}$ , let  $Q$  be a basis matrix of  $\text{Ker}(\Phi)$  and  $I = \text{supp}(\Psi^\top \bar{x})$ . The followings are met:

- (1)  $\Psi_{I^c}^\top Q$  is of full-rank;
- (2)  $\|(Q^\top \Psi_{I^c})^+ Q^\top \Psi_I \text{sign}(\Psi_I^\top \bar{x})\|_\infty < 1$ .

Here  $A^+$  is the pseudo-inverse of matrix  $A$  defined as  $A^+ = A^\top (AA^\top)^{-1}$ .

Paper [8] proposes the following condition, which generalizes the synthesis Identifiability Criterion (IC) proposed in [2], for the solution uniqueness and robustness for problems (1a) and (1b) (the robustness requires the non-zero entries of  $\Psi_I^\top x$  to be sufficiently large compared to noise).

**Condition 4** For a given  $\bar{x}$ , index sets  $I = \text{supp}(\Psi^\top \bar{x})$  and  $J = I^c$  satisfy:

- (1)  $\text{Ker}(\Psi_J^\top) \cap \text{Ker}(\Phi) = \{0\}$ ;
- (2) Let  $A^{|J|} = U(U^\top \Phi^\top \Phi U)^{-1} U^\top$  and  $\Omega^{|J|} = \Psi_J^+ (\Phi^\top \Phi A^{|J|} - Id) \Psi_I$ , where  $U$  is a basis matrix of  $\text{Ker}(\Psi_J^\top)$ . Then

$$IC(\text{sign}(\Psi_I^\top \bar{x})) := \min_{u \in \text{Ker}(\Psi_J)} \|\Omega^{|J|} \text{sign}(\Psi_I^\top \bar{x}) - u\|_\infty < 1.$$

According to [8], Conditions 3 and 4 do not contain each other, and Condition 3 does not reduce to the synthesis IC in the synthesis case. Conditions 3 and 4 are determined by the sign and the support; while Conditions 1 and 2 are not determined quantities, and they depend on the existence of certain dual variables. This might be the main difference between Conditions 3, 4 and Conditions 1, 2.

The following example shows that Conditions 3 and 4 are both stronger than Conditions 1 and 2. Let

$$\Psi = \begin{pmatrix} 10.5 & 1 & 10 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \Phi = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \tilde{x} = \begin{pmatrix} 1 \\ -1 \\ -10 \end{pmatrix}, \quad b = \begin{pmatrix} -1 \\ -10 \end{pmatrix}.$$

It is straightforward to verify that Conditions 1 and 2 hold. However, Conditions 3 and 4 fail to hold. Indeed, we have  $\Psi^\top \tilde{x} = (10.5, 0, 0)^\top$  and  $I = \{1\}$ .  $Q = (1, 0, 0)^\top$  is a basis matrix of  $\text{Ker}(\Phi)$ . Thus

$$\|(Q^\top \Psi_{I^c})^+ Q^\top \Psi_I \text{sign}(\Psi_I^\top \tilde{x})\|_\infty = \left\| \left( \frac{10.5}{101}, \frac{105}{101} \right)^\top \right\|_\infty = \frac{105}{101}.$$

Hence, Condition 3 does not hold. Furthermore,  $U = (1, -1, -10)^\top$  is a basis matrix of  $\text{Ker}(\Psi_J^\top)$ , and the definition of  $\Omega^{[J]}$  gives us  $\Omega^{[J]} = (\frac{10.5}{101}, \frac{105}{101})^\top$ . Therefore,  $IC(\text{sign}(\Psi_I^\top \tilde{x})) = \frac{105}{101} > 1$ , so Condition 4 does not hold either. Paper [8] also presents sufficient conditions for solution uniqueness, which are reviewed in [18] and shown to be not necessary.

During the time we are preparing this manuscript, work [31] gives the same condition as Condition 1 and shows its sufficiency. However, the necessity of Condition 1 or 2 is never discussed in the literature.

### 4.2 Previous Robustness Conditions

Turning to solution robustness, [7, 9, 31] have studied the robustness of problems (1b) and (1c) in the Hilbert-space setting. Translating to the finite dimension, the condition in [7] is equivalent to Condition 2. Under Condition 2, work [7] shows the existence of a constant  $C$  (not explicitly given) such that the solution  $x_{\delta,\lambda}$  to Eq. 1b obeys  $\|\Psi^\top(x_{\delta,\lambda} - x^*)\|_2 \leq C\delta$  when  $\lambda$  is set proportional to the noise level  $\delta$ . [31] gives an explicit formula of  $C$  in  $\|x_{\delta,\lambda} - x^*\|_2 \leq C\delta$  for the solution  $x_{\delta,\lambda}$  to Eq. 1b. In order to obtain an explicit formula for  $C$ , [9] introduces the following condition:

**Condition 5** Assume that  $\Psi\Psi^\top$  is invertible, and let  $\hat{\Psi} = (\Psi\Psi^\top)^{-1}\Psi$ . Given  $\bar{x}$ , the following two statements hold:

- (1) There exists some  $y \in \partial\|\cdot\|_1(\Psi^\top \bar{x})$  such that  $\Psi y \in \text{Im}(\Phi^\top)$ ;
- (2) For some  $t \in (0, 1)$ , let  $I(t) = \{i : |y_i| > t\}$ , and the mapping  $\hat{\Phi} := \Phi|_{\text{Span}\{\hat{\psi}_i : i \in I(t)\}}$  is injective.

Under this condition, the solutions to Eqs. 1b and 1c are subject to error bounds whose constants depend on  $t$ ,  $\hat{\Phi}$ , and other quantities.

**Proposition 1** Condition 5 is stronger than Condition 2.

*Proof* Let  $J = I(t)^c$ ; then we have  $\|y_J\|_\infty \leq t < 1$  from the definition of  $I(t)$ . It remains to show that  $\text{Ker}(\Psi_J^\top) \cap \text{Ker}(\Phi) = \{0\}$ . For any  $x \in \text{Ker}(\Psi_J^\top)$ , we have

$$\begin{aligned} x &= (\Psi\Psi^\top)^{-1}\Psi\Psi^\top x = (\Psi\Psi^\top)^{-1}\Psi_J\Psi_J^\top x + (\Psi\Psi^\top)^{-1}\Psi_{J^c}\Psi_{J^c}^\top x \\ &= (\Psi\Psi^\top)^{-1}\Psi_{I(t)}\Psi_{I(t)}^\top x. \end{aligned}$$

Since  $\Phi$  restricted to  $\text{Span}\{\hat{\psi}_i : i \in I(t)\} = \text{Im}((\Psi\Psi^\top)^{-1}\Psi_{I(t)})$  is injective, we have what we need. □

Definition 5 in paper [8] provides a much stronger condition below, which generalizes the synthesis Exact Recovery Condition (ERC) in [32], and this condition strengthens Condition 4 by dropping the dependence on the  $\Psi$ -support (see the definition of  $RC(I)$  below).

**Condition 6** Given  $\bar{x}$ , index sets  $I = \text{supp}(\Psi^\top \bar{x})$  and  $J = I^c$  satisfy:

- (1)  $\text{Ker}(\Psi_J^\top) \cap \text{Ker}(\Phi) = \{0\}$ ;
- (2) Letting  $\Omega^{[J]}$  be given as in Condition 4,

$$RC(I) := \max_{p \in \mathbb{R}^{|I|}, \|p\|_\infty \leq 1} \min_{u \in \text{Ker}(\Psi_J)} \|\Omega^{[J]}p - u\|_\infty < 1.$$

Under this condition, a nice error bound and a certain kind of “weak” sign consistency (between  $\Psi^\top x_{\delta, \lambda}$  and  $\Psi^\top x^*$ ) are given provided that problem (1b) is solved with the parameter  $\lambda = \frac{\rho \|w\|_{2c_J}}{2(1-RC(I))}$  for some  $\rho > 1$ , where  $c_J = \|\Psi_J^\top \Phi^\top (\Phi A^{[J]} \Phi^\top - Id)\|_{2, \infty}$ . When  $1 - RC(I)$  gets close to 0, this  $\lambda$  can become larger than it should be.

### 5 Verifying the conditions

In this section, we present a method to verify Condition 1. Our method includes two steps:

**(Step 1):** Let  $\Phi = U \Sigma V^\top$  be the singular value decomposition of  $\Phi$ . Assume  $V = [v_1, \dots, v_n]$ . Since  $\Phi$  has full row-rank, we have  $\text{Ker}(\Phi) = \text{Span}\{v_{m+1}, \dots, v_n\}$  and  $Q = [v_{m+1}, \dots, v_n]$  as a basis of  $\text{Ker}(\Phi)$ . We verify that  $\Psi_J^\top Q$  is of full-rank by computing the smallest singular value or QR decomposition and thus ensure part (1) of Condition 1.

**(Step 2):** Let  $u_1 = -Q^\top \Psi_I \text{sign}(\Psi_I^\top \bar{x})$  and  $A = Q^\top \Psi_J$ . Solve the convex problem

$$\underset{u \in \mathbb{R}^{|J|}}{\text{minimize}} \|u\|_\infty, \quad \text{subject to } Au = u_1. \tag{6}$$

If the optimal objective of Eq. 6 is strictly less than 1, then part (2) of Condition 1 holds. In fact, we have:

**Proposition 2** Part (2) of Condition 1 holds if and only if Eq. 6 has an optimal objective  $< 1$ .

*Proof* Let  $\hat{u}$  be a minimizer of Eq. 6. Assume  $\|\hat{u}\|_\infty < 1$ . We consider the vector  $y$  composed by  $y_I = \text{sign}(\Psi_I^\top \bar{x})$  and  $y_J = \hat{u}$ . To show part (2) of Condition 1, it suffices to prove  $\Psi y \in \text{Im}(\Phi^\top)$ , or equivalently,  $Q^\top \Psi y = 0$ . Indeed,

$$Q^\top \Psi y = Q^\top \Psi_J y_J + Q^\top \Psi_I y_I = Q^\top \Psi_J \hat{u} + Q^\top \Psi_I y_I = 0.$$

The other direction is obvious. □

The convex program (6) is similar in form to one in [8, Definition 4], but they are used to verify different conditions.

### 6 Proof of Theorem 1

We establish Theorem 1 in two steps. Our first step proves the theorem for problem (1a) only. The second step proves Theorem 1 for problems (1b) and (1c).

#### 6.1 Proof of Theorem 1 for problem (1a)

The equivalence of the three statements is shown in the following order: 3)  $\implies$  1)  $\implies$  2)  $\implies$  3).

3)  $\implies$  1). Consider any perturbation  $\hat{x} + h$  where  $h \in \text{Ker}(\Phi) \setminus \{0\}$ . Take a subgradient  $g \in \partial \|\cdot\|_1(\Psi^\top \hat{x})$  obeying  $g_I = \text{sign}(\Psi_I^\top \hat{x}) = y_I$ ,  $g_K = y_K$ , and  $\|g_J\|_\infty \leq 1$  such that  $\langle g_J, \Psi_J^\top h \rangle = \|\Psi_J^\top h\|_1$ . Then,

$$\|\Psi^\top(\hat{x} + h)\|_1 \geq \|\Psi^\top \hat{x}\|_1 + \langle \Psi g, h \rangle \tag{7a}$$

$$= \|\Psi^\top \hat{x}\|_1 + \langle \Psi g - \Psi y, h \rangle \tag{7b}$$

$$= \|\Psi^\top \hat{x}\|_1 + \langle g - y, \Psi^\top h \rangle \tag{7c}$$

$$= \|\Psi^\top \hat{x}\|_1 + \langle g_J - y_J, \Psi_J^\top h \rangle \tag{7d}$$

$$\geq \|\Psi^\top \hat{x}\|_1 + \|\Psi_J^\top h\|_1(1 - \|y_J\|_\infty), \tag{7e}$$

where Eq. 7b follows from  $\Psi y \in \text{Im}(\Phi^\top) = \text{Ker}(\Phi)^\perp$  and  $h \in \text{Ker}(\Phi)$ , Eq. 7d follows from the setting of  $g$ , and Eq. 7e is an application of the inequality  $\langle x, y \rangle \leq \|x\|_1 \|y\|_\infty$  and  $\langle g_J, \Psi_J^\top h \rangle = \|\Psi_J^\top h\|_1$ . Since  $h \in \text{Ker}(\Phi) \setminus \{0\}$  and  $\text{Ker}(\Psi_J^\top) \cap \text{Ker}(\Phi) = \{0\}$ , we have  $\|\Psi_J^\top h\|_1 > 0$ . Together with the condition  $\|y_J\|_\infty < 1$ , we have  $\|\Psi^\top(\hat{x} + h)\|_1 > \|\Psi^\top \hat{x}\|_1$  for every  $h \in \text{Ker}(\Phi) \setminus \{0\}$  which implies that  $\hat{x}$  is the unique minimizer of Eq. 1a.

1)  $\implies$  2). For every  $h \in \text{Ker}(\Phi) \setminus \{0\}$ , we have  $\Phi(\hat{x} + th) = \Phi \hat{x}$  and can find  $t$  small enough around 0 such that  $\text{sign}(\Psi_I^\top(\hat{x} + th)) = \text{sign}(\Psi_I^\top \hat{x})$ . Since  $\hat{x}$  is the unique solution, for small and nonzero  $t$ , we have

$$\begin{aligned} \|\Psi^\top(\hat{x})\|_1 &< \|\Psi^\top(\hat{x} + th)\|_1 = \|\Psi_I^\top(\hat{x} + th)\|_1 + \|\Psi_{I^c}^\top(\hat{x} + th)\|_1 \\ &= \langle \Psi_I^\top(\hat{x} + th), \text{sign}(\Psi_I^\top(\hat{x} + th)) \rangle + \|t\Psi_{I^c}^\top h\|_1 \\ &= \langle \Psi_I^\top \hat{x} + t\Psi_I^\top h, \text{sign}(\Psi_I^\top \hat{x}) \rangle + \|t\Psi_{I^c}^\top h\|_1 \\ &= \langle \Psi_I^\top \hat{x}, \text{sign}(\Psi_I^\top \hat{x}) \rangle + t\langle \Psi_I^\top h, \text{sign}(\Psi_I^\top \hat{x}) \rangle + \|t\Psi_{I^c}^\top h\|_1 \\ &= \|\Psi^\top(\hat{x})\|_1 + t\langle \Psi_I^\top h, \text{sign}(\Psi_I^\top \hat{x}) \rangle + \|t\Psi_{I^c}^\top h\|_1. \end{aligned}$$

Therefore, for any  $h \in \text{Ker}(\Phi) \setminus \{0\}$ , we have

$$\langle \Psi_I^\top h, \text{sign}(\Psi_I^\top \hat{x}) \rangle < \|\Psi_{I^c}^\top h\|_1, \tag{8}$$

which can be viewed as an analysis version of the synthesis null space condition [5]. If the condition  $\text{Ker}(\Psi_{I^c}^\top) \cap \text{Ker}(\Phi) = \{0\}$  does not hold, we can choose a nonzero vector  $h \in \text{Ker}(\Psi_{I^c}^\top) \cap \text{Ker}(\Phi)$ . We also have  $-h \in \text{Ker}(\Psi_{I^c}^\top) \cap \text{Ker}(\Phi)$ .

Then we have  $\langle \Psi_I^\top h, \text{sign}(\Psi_I^\top \hat{x}) \rangle < 0$  and  $-\langle \Psi_I^\top h, \text{sign}(\Psi_I^\top \hat{x}) \rangle < 0$ , which is a contradiction.

It remains to show the existence of  $y$  in part (2) of Condition 1. This part is in spirit of the methods in papers [17] and [18], which are based on linear programming strong duality. We take  $\hat{y}$  with restrictions  $\hat{y}_I = \text{sign}(\Psi_I^\top \hat{x})$  and  $\hat{y}_{I^c} = 0$ . If such  $\hat{y}$  satisfies  $\Psi \hat{y} \in \text{Im}(\Phi^\top)$ , then the existence has been shown. If  $\Psi \hat{y} \notin \text{Im}(\Phi^\top) = \text{Ker}(\Phi)^\perp$ , then we shall construct a new vector to satisfy part (2) of Condition 1. Let  $Q$  be a basis matrix of  $\text{Ker}(\Phi)$ . We have that  $a := Q^\top \Psi \hat{y}$  must be a nonzero vector. Consider the following problem

$$\underset{z \in \mathbb{R}^l}{\text{minimize}} \|z\|_\infty \quad \text{subject to } Q^\top \Psi z = -a \text{ and } z_I = 0. \tag{9}$$

For any minimizer  $\hat{z}$  of problem (9), we have  $\Psi(\hat{y} + \hat{z}) \in \text{Ker}(\Phi)^\perp = \text{Im}(\Phi^\top)$  and  $(\hat{y} + \hat{z})_I = \hat{y}_I = \text{sign}(\Psi_I^\top \hat{x})$ . Thus, we shall show that the objective of problem (9) is strictly less than 1. To this end, we rewrite problem (9) in an equivalent form as:

$$\underset{z}{\text{minimize}} \|z_{I^c}\|_\infty \quad \text{subject to } Q^\top \Psi_{I^c} z_{I^c} = -a, \tag{10}$$

whose Lagrange dual problem is

$$\underset{p}{\text{maximize}} \langle p, a \rangle \quad \text{subject to } \|\Psi_{I^c}^\top Q p\|_1 \leq 1. \tag{11}$$

Note that  $Qp \in \text{Ker}(\Phi)$  and  $|\langle p, a \rangle| = |\langle p, Q^\top \Psi \hat{y} \rangle| = |\langle p, Q^\top \Psi_I \text{sign}(\Psi_I^\top \hat{x}) \rangle| = |\langle \Psi_I^\top Q p, \text{sign}(\Psi_I^\top \hat{x}) \rangle|$ . By using Eq. 8, for any  $p$ , we have

$$|\langle p, a \rangle| = \begin{cases} |\langle \Psi_I^\top Q p, \text{sign}(\Psi_I^\top \hat{x}) \rangle| = 0, & \text{if } Qp = 0, \\ |\langle \Psi_I^\top Q p, \text{sign}(\Psi_I^\top \hat{x}) \rangle| < \|\Psi_{I^c}^\top Q p\|_1 \leq 1, & \text{otherwise.} \end{cases}$$

Hence, problem (11) is feasible, and its objective value is strictly less than 1. By the linear programming strong duality property, problems (9) and (10) also have solutions, and their objective values are strictly less than 1, too. This completes the proof.

2)  $\implies$  3). Let  $J = I^c$  and  $K = \emptyset$ , then Condition 2 follows.

The proof of 3)  $\implies$  1) is a standard technique in the compressed sensing community.

### 6.2 Proof of Theorem 1 for problems (1b) and (1c)

**Lemma 1** *Let  $\gamma > 0$ . If  $\gamma \|\Phi x - b\|_2^2 + \|\Psi^\top x\|_1$  is constant on a convex set  $\Omega$ , then both  $\Phi x - b$  and  $\|\Psi^\top x\|_1$  are constant on  $\Omega$ .*

*Proof* It suffices to prove the case where the convex set  $\Omega$  has more than one point. Suppose  $x_1$  and  $x_2$  are different points in  $\Omega$ . Consider the line segment  $L$  connecting  $x_1$  and  $x_2$ . By the convexity of  $\Omega$ , we know  $L \subset \Omega$ . Thus,  $\gamma \|\Phi x - b\|_2^2 + \|\Psi^\top x\|_1$

is constant on  $L$ , and we denote the constant value by  $\hat{c}$ . If  $\Phi x_1 - b \neq \Phi x_2 - b$ , then, for any  $0 < \alpha < 1$ , we have

$$\gamma \|\Phi(\alpha x_1 + (1 - \alpha)x_2) - b\|_2^2 + \|\Psi^\top(\alpha x_1 + (1 - \alpha)x_2)\|_1 \tag{12a}$$

$$= \gamma \|\alpha(\Phi x_1 - b) + (1 - \alpha)(\Phi x_2 - b)\|_2^2 + \|\alpha(\Psi^\top x_1) + (1 - \alpha)(\Psi^\top x_2)\|_1 \tag{12b}$$

$$< \alpha(\gamma \|\Phi x_1 - b\|_2^2 + \|\Psi^\top x_1\|_1) + (1 - \alpha)(\gamma \|\Phi x_2 - b\|_2^2 + \|\Psi^\top x_2\|_1) \tag{12c}$$

$$= \alpha \hat{c} + (1 - \alpha) \hat{c} = \hat{c}, \tag{12d}$$

where the strict inequality follows from the *strict* convexity of  $\gamma \|\cdot\|_2^2$  and the convexity of  $\|\Psi^\top x\|_1$ . This means that  $\alpha x_1 + (1 - \alpha)x_2 \in L \subset \Omega$  attains a smaller value than  $\hat{c}$  for any  $0 < \alpha < 1$ , which is a contradiction to the assumption that all points in  $\Omega$  attain the same value  $\hat{c}$ . Therefore, we have  $\Phi x_1 - b = \Phi x_2 - b$ , from which it is easy to obtain  $\|\Psi^\top x_1\|_1 = \|\Psi^\top x_2\|_1$ .  $\square$

We let  $X_\lambda$  and  $Y_\delta$  denote the sets of solutions to problems (1b) and (1c), respectively. Moreover, we assume that these two sets are nonempty. Then, from Lemma 1, we have the following result.

**Corollary 1** *In problem (1b),  $\Phi x - b$  and  $\|\Psi^\top x\|_1$  are constant on  $X_\lambda$ ; in problem (1c),  $\Phi x - b$  and  $\|\Psi^\top x\|_1$  are constant on  $Y_\delta$ .*

*Proof* Since  $\|\Phi x - b\|_2^2 + \lambda \|\Psi^\top x\|_1$  is constant over  $X_\lambda$ , the result follows directly from Lemma 1 for problem (1b). For problem (1c), if  $0 \in Y_\delta$ , then we have  $Y_\delta = \{0\}$  because of the full row-rankness of  $\Psi$ . The result holds trivially. Suppose  $0 \notin Y_\delta$ . Since the optimal objective  $\|\Psi^\top x\|_1$  is constant for all  $x \in Y_\delta$ , we have to show that  $\|\Phi x - b\|_2^2 = \delta$  for all  $x \in Y_\delta$ . If there exist a nonzero  $\hat{x} \in Y_\delta$  such that  $\|\Phi \hat{x} - b\|_2^2 < \delta$ , we can find a non-empty ball  $\mathcal{B}$  centered at  $\hat{x}$  with a sufficiently small radius  $\rho > 0$  such that  $\|\Phi \tilde{x} - b\|_2^2 < \delta$  for all  $\tilde{x} \in \mathcal{B}$ . Let  $\alpha = \min\{\frac{\rho}{2\|\hat{x}\|_2}, \frac{1}{2}\} \in (0, 1)$ . We have  $(1 - \alpha)\hat{x} \in \mathcal{B}$  and  $\|(1 - \alpha)\Psi^\top \hat{x}\|_1 < \|\Psi^\top \hat{x}\|_1$ , which is a contradiction to the assumption that  $x \in Y_\delta$ .  $\square$

*Proof of Theorem 1 for problems (1b) and (1c)* This proof exploits Corollary 1. Since the results of Corollary 1 are identical for problems (1b) and (1c), we present the proof for problem (1b) only.

By assumption,  $X_\lambda$  is nonempty. We pick  $\hat{x} \in X_\lambda$ . Let  $b^* = \Phi \hat{x}$ , which is independent of the choice of  $\hat{x}$  according to Corollary 1. We introduce the following problem

$$\underset{x}{\text{minimize}} \|\Psi^\top x\|_1, \quad \text{subject to } \Phi x = b^* \tag{13}$$

and denote its solution set by  $X^*$ .

Now, we show that  $X_\lambda = X^*$ . Since  $\Phi x = \Phi \hat{x}$  and  $\|\Psi^\top x\|_1 = \|\Psi^\top \hat{x}\|_1$  hold for all  $x \in X_\lambda$  and any  $x$  obeying  $\Phi x = \Phi \hat{x}$  and  $\|\Psi^\top x\|_1 = \|\Psi^\top \hat{x}\|_1$  belongs to  $X_\lambda$ , it suffices to show that  $\|\Psi^\top x\|_1 = \|\Psi^\top \hat{x}\|_1$  for all  $x \in X^*$ . Assume that this does *not* hold. Then, because  $\hat{x}$  is a feasible solution of problem (13), there exists  $\tilde{x} \in X^*$  satisfying  $\|\Psi^\top \tilde{x}\|_1 < \|\Psi^\top \hat{x}\|_1$ . However,  $\|\Phi \tilde{x} - b\|_2 = \|b^* - b\|_2 = \|\Phi \hat{x} - b\|_2$

and  $\|\Psi^\top \tilde{x}\|_1 < \|\Psi^\top \hat{x}\|_1$  mean that  $\tilde{x}$  is a strictly better solution to problem (1b) than  $\hat{x}$ , contradicting the assumption that  $\hat{x} \in X_\lambda$ .

Since  $X_\lambda = X^*$ ,  $\hat{x}$  is the unique solution to problem (1b) if and only if it is the unique solution to problem (13). Since problem (13) is in the same form of problem (1a), applying the part of Theorem 1 for problem (1a), which is already proved, we conclude that  $\hat{x}$  is the unique solution to problem (1b) if and only if Condition 1 or 2 holds. □

### 7 Proof of Theorem 2

**Lemma 2** *Assume that vectors  $\bar{x}$  and  $y$  satisfy Condition 1. Let  $I = \text{supp}(\Psi^\top \bar{x})$  and  $J = I^c$ . We have*

$$\|\Psi^\top x - \Psi^\top \bar{x}\|_1 \leq C_3 \|\Phi(x - \bar{x})\|_2 + C_4 d_y(x, \bar{x}), \quad \forall x, \tag{14}$$

where  $d_y(x, \bar{x}) := \|\Psi^\top x\|_1 - \|\Psi^\top \bar{x}\|_1 - \langle \Psi y, x - \bar{x} \rangle$  is the Bregman distance of function  $\|\Psi^\top \cdot\|_1$ , the absolute constants  $C_3, C_4$  are given in Theorem 2.

*Proof* This proof is divided into two parts. They are partially inspired by [7].

1. this part shows that for any  $u \in \text{Ker}(\Psi_J^\top)$ ,

$$\|\Psi^\top x - \Psi^\top \bar{x}\|_1 \leq \left(1 + \frac{C_3 \|\Phi\|}{\sqrt{\lambda_{\min}(\Psi \Psi^\top)}}\right) \|\Psi^\top(x - u)\|_1 + C_3 \|\Phi(x - \bar{x})\|_2. \tag{15}$$

2. this part shows that

$$f(x) := \min \left\{ \|\Psi^\top(x - u)\|_1 : u \in \text{Ker}(\Psi_J^\top) \right\} \leq (1 - \|y_J\|_\infty)^{-1} d_y(x, \bar{x}). \tag{16}$$

Using the definition of  $C_4$ , combining Eqs. 15 and 16 gives Eq. 14.

*Part 1.* Let  $u \in \text{Ker}(\Psi_J^\top)$ . By the triangle inequality of norms, we get

$$\|\Psi^\top x - \Psi^\top \bar{x}\|_1 \leq \|\Psi^\top(x - u)\|_1 + \|\Psi^\top(u - \bar{x})\|_1. \tag{17}$$

Since  $\bar{x} \in \text{Ker}(\Psi_J^\top)$ , we have  $u - \bar{x} \in \text{Ker}(\Psi_J^\top)$  and thus  $\|u - \bar{x}\|_2 \leq r(J) \|\Phi(u - \bar{x})\|_2$ , where  $r(J) < +\infty$  follows from part (1) of Condition 1. Using the fact that  $\text{supp}(\Psi^\top(u - \bar{x})) = I$ , we derive that

$$\|\Psi^\top(u - \bar{x})\|_1 \leq \sqrt{|I|} \|\Psi^\top(u - \bar{x})\|_2 \tag{18a}$$

$$\leq \sqrt{|I|} \|u - \bar{x}\|_2 \tag{18b}$$

$$\leq \sqrt{|I|} r(J) \|\Phi(u - \bar{x})\|_2 \tag{18c}$$

$$= C_3 \|\Phi(u - \bar{x})\|_2, \tag{18d}$$

where we have used the assumption  $\lambda_{\max}(\Psi\Psi^\top) = 1$  and the definition  $C_3 = r(J)\sqrt{|I|}$ . Furthermore,

$$\|\Phi(u - \bar{x})\|_2 \leq \|\Phi(x - u)\|_2 + \|\Phi(x - \bar{x})\|_2 \tag{19a}$$

$$\leq \|\Phi\| \|x - u\|_2 + \|\Phi(x - \bar{x})\|_2 \tag{19b}$$

$$\leq \frac{\|\Phi\| \|\Psi^\top(x-u)\|_2}{\sqrt{\lambda_{\min}(\Psi\Psi^\top)}} + \|\Phi(x - \bar{x})\|_2 \tag{19c}$$

$$\leq \frac{\|\Phi\| \|\Psi^\top(x-u)\|_1}{\sqrt{\lambda_{\min}(\Psi\Psi^\top)}} + \|\Phi(x - \bar{x})\|_2. \tag{19d}$$

Therefore, we get Eq. 15 after combining Eqs. 17, 18, and 19.

Part 2. Since  $\langle \Psi y, \bar{x} \rangle = \|\Psi^\top \bar{x}\|_1$  implies  $d_y(x, \bar{x}) = \|\Psi^\top x\|_1 - \langle \Psi y, x \rangle$ , it is equivalent to proving

$$f(x) \leq (1 - \|y_J\|_\infty)^{-1} (\|\Psi^\top x\|_1 - \langle \Psi y, x \rangle).$$

Since  $u \in \text{Ker}(\Psi_J^\top)$  is equivalent to  $\Psi_J^\top u = 0$ , the Lagrangian of the minimization problem in Eq. 16 is

$$L(u, v) = \|\Psi^\top(x - u)\|_1 + \langle v, \Psi_J^\top u \rangle = \|\Psi^\top(x - u)\|_1 + \langle \Psi_J v, u - x \rangle + \langle \Psi_J v, x \rangle.$$

Then  $f(x) = \min_u \max_v L(u, v)$ . Following the minimax theorem, we derive that

$$\begin{aligned} f(x) &= \max_v \min_u L(u, v) \\ &= \max_v \min_u \{ \|\Psi^\top(x - u)\|_1 + \langle \Psi_J v, u - x \rangle + \langle \Psi_J v, x \rangle \} \\ &= \max_w \min_u \{ \|\Psi^\top(x - u)\|_1 + \langle w, u - x \rangle + \langle w, x \rangle : w \in \text{Im}(\Psi_J) \} \\ &= \max_w \{ \langle w, x \rangle : w \in \partial \|\Psi^\top \cdot\|_1(0) \cap \text{Im}(\Psi_J) \} \\ &= \max_w \{ \langle c\Psi y + w, x \rangle : w \in \partial \|\Psi^\top \cdot\|_1(0) \cap \text{Im}(\Psi_J) \} - \langle c\Psi y, x \rangle, \quad \forall c > 0 \\ &= c \max_w \{ \langle \Psi y + w, x \rangle : w \in c^{-1} \partial \|\Psi^\top \cdot\|_1(0) \cap \text{Im}(\Psi_J) \} - c \langle \Psi y, x \rangle, \quad \forall c > 0. \end{aligned}$$

Let

$$c = (1 - \|y_J\|_\infty)^{-1}$$

and  $Z_J = \{z \in \mathbb{R}^l : z_I = 0\}$ . Since  $\|y_J\|_\infty < 1$  from part (2) of Condition 1, we have  $c < +\infty$  and get

$$(y + c^{-1} \partial \|\cdot\|_1(0) \cap Z_J) \subset \partial \|\cdot\|_1(0),$$

from which we conclude

$$(\Psi y + c^{-1} \partial \|\Psi^\top \cdot\|_1(0) \cap \text{Im}(\Psi_J)) \subset \partial \|\Psi^\top \cdot\|_1(0).$$

Hence, for any  $w \in c^{-1} \partial \|\Psi^\top \cdot\|_1(0) \cap \text{Im}(\Psi_J)$ , it holds that  $\Psi y + w \subset \partial \|\Psi^\top \cdot\|_1(0)$ , which implies  $\langle \Psi y + w, x \rangle \leq \|\Psi^\top x\|_1$  by the convexity of  $\|\Psi^\top \cdot\|_1$ . Therefore,  $f(x) \leq c(\|\Psi^\top x\|_1 - \langle \Psi y, x \rangle)$ . □

**Lemma 3** ([16], Theorem 3; [17], Lemma 3.5) *Suppose that  $x^* \in \mathbb{R}^n$  is a fixed vector obeying  $\text{supp}(\Psi^\top x^*) = I$  and that there are vectors satisfying  $y \in \partial \|\cdot\|_1(\Psi^\top x^*)$  and*



$\Psi y = \Phi^\top \beta$ . Then for every  $\delta > 0$  and every data vector  $b$  satisfying  $\|\Phi x^* - b\|_2 \leq \delta$ , the following two statements hold:

- 1) Every minimizer  $x_{\delta,\lambda}$  of problem (1b) satisfies  $d_y(x_{\delta,\lambda}, x^*) \leq \frac{(\delta + \lambda\|\beta\|_2/2)^2}{\lambda}$  and  $\|\Phi x_{\delta,\lambda} - b\|_2 \leq \delta + \lambda\|\beta\|_2$ ;
- 2) Every minimizer  $x_\delta$  of problem (1c) satisfies  $d_y(x_\delta, x^*) \leq 2\delta\|\beta\|_2$ .

From  $\Psi y = \Phi^\top \beta$  and the full-rankness of  $\Phi$ , we have  $\beta = (\Phi\Phi^\top)^{-1}\Phi\Psi y$ .

*Proof of Theorem 2* Firstly, we derive that

$$\|\Psi(x_{\delta,\lambda} - x^*)\|_1 \leq C_3\|\Phi(x_{\delta,\lambda} - x^*)\|_2 + C_4d_y(x_{\delta,\lambda}, x^*) \tag{20a}$$

$$\leq C_3\|\Phi x_{\delta,\lambda} - b\|_2 + C_3\|\Phi x^* - b\|_2 + C_4d_y(x_{\delta,\lambda}, x^*) \tag{20b}$$

$$\leq C_3(\delta + \lambda\|\beta\|_2) + C_3\delta + C_4\frac{(\delta + \lambda\|\beta\|_2/2)^2}{\lambda}, \tag{20c}$$

where the first and the third inequalities follow from Lemmas 2 and 3, respectively. Substituting  $\lambda = C_0\delta$  and collecting like terms in Eq. 20c, we obtain the first part of Theorem 2. The second part can be proved in the same way.  $\square$

### 8 Proof of Theorem 3

Let  $w := x_\delta - x^*$  and  $v := \Psi^\top w$ . Then, by the minimality of  $\|\Psi^\top x_\delta\|_1$ , we derive

$$\begin{aligned} \|\Psi^\top x^*\|_1 &\geq \|\Psi^\top x_\delta\|_1 = \|v + \Psi^\top x^*\|_1 = \|v_I + \Psi_I^\top x^*\|_1 + \|v_J + \Psi_J^\top x^*\|_1 \\ &\geq \langle v_I + \Psi_I^\top x^*, \text{sign}(\Psi_I^\top x^*) \rangle + \|v_J\|_1 - \|\Psi_J^\top x^*\|_1 \\ &= \langle v_I, \text{sign}(\Psi_I^\top x^*) \rangle + \|\Psi_I^\top x^*\|_1 + \|v_J\|_1 - \|\Psi_J^\top x^*\|_1. \end{aligned}$$

Rearranging the above inequality and using the fact that  $\|\Psi^\top x^*\|_1 = \|\Psi_I^\top x^*\|_1 + \|\Psi_J^\top x^*\|_1$  yield

$$\|v_J\|_1 \leq 2\|\Psi_J^\top x^*\|_1 + |\langle v_I, \text{sign}(\Psi_I^\top x^*) \rangle|. \tag{21}$$

In addition, we have

$$|\langle v_I, \text{sign}(\Psi_I^\top x^*) \rangle| = |\langle y_I, v_I \rangle| \leq |\langle y, v \rangle| + |\langle y_J, v_J \rangle|.$$

In what follows, we further bound the right-hand side of the above inequality term by term. First of all, in view of the optimization constraint and the feasibility of both  $x_\delta$  and  $x^*$ , we have

$$\|\Phi w\|_2 = \|\Phi(x_\delta - x^*)\|_2 \leq \|\Phi x_\delta - b\|_2 + \|\Phi x^* - b\|_2 \leq 2\delta.$$

Using the condition  $\Psi y = \Phi^\top \beta$ , we obtain

$$\begin{aligned} |\langle y, v \rangle| &= |\langle y, \Psi^\top w \rangle| = |\langle \Psi y, w \rangle| = |\langle \Phi^\top \beta, w \rangle| \\ &= |\langle \beta, \Phi w \rangle| \leq \|\beta\|_2 \cdot \|\Phi w\|_2 \leq 2\delta\|\beta\|_2. \end{aligned}$$

At last,  $|\langle y_J, v_J \rangle| \leq \|y_J\|_\infty \|v_J\|_1$ . Collecting these upper bounds, we derive

$$|\langle v_I, \text{sign}(\Psi_I^\top x^*) \rangle| \leq \|y_J\|_\infty \|v_J\|_1 + 2\|\beta\|_2\delta.$$

Using Eq. 21 and noting that  $\|y_J\|_\infty < 1$ , we get

$$\|v_J\|_1 \leq \frac{2}{1 - \|y_J\|_\infty} \|\Psi_J^\top x^*\|_1 + \frac{2\|\beta\|_2}{1 - \|y_J\|_\infty} \delta. \quad (22)$$

Now, using the inequality (4), we get

$$\|v_I\|_2 \leq \rho \|v_J\|_1 + \tau \|\Phi w\|_2 \leq \frac{2\rho}{1 - \|y_J\|_\infty} \|\Psi_J^\top x^*\|_1 + \left( \frac{2\rho\|\beta\|_2}{1 - \|y_J\|_\infty} + 2\tau \right) \delta. \quad (23)$$

Finally, combing Eqs. 22 and 23, we obtain

$$\begin{aligned} \|v\|_2 &\leq \|v_I\|_2 + \|v_J\|_2 \leq \|v_I\|_2 + \|v_J\|_1 \\ &\leq \frac{2(1 + \rho)}{1 - \|y_J\|_\infty} \|\Psi_J^\top x^*\|_1 + \left( \frac{2(1 + \rho)\|\beta\|_2}{1 - \|y_J\|_\infty} + 2\tau \right) \delta. \end{aligned}$$

This completes the proof.

**Acknowledgments** The work of H. Zhang is supported by China Scholarship Council during his visit to Rice University, and in part by the National Science Foundation of China (No.11501569 and No.61072118). The work of M. Yan is supported in part by the Center for Domain-Specific Computing (CDSC) funded by NSF grants CCF-0926127 and ARO/ARL MURI grant FA9550-10-1-0567. The work of W. Yin is supported in part by NSF grants DMS-0748839 and ECCS-1028790. They thank Rachel Ward and Xiaoya Zhang for their helpful discussions.

## References

1. Elad, M., Milanfar, P., Rubinstein, R.: Analysis versus synthesis in signal priors. *Inverse Prob.* **23**(3), 947 (2007)
2. Fuchs, J.J.: On sparse representations in arbitrary redundant bases. *IEEE Trans. Inf. Theory* **50**(6), 1341–1344 (2004)
3. Candes, E.J., Plan, Y.: A probabilistic and RIPless theory of compressed sensing. *IEEE Trans. Inf. Theory* **57**(11), 7235–7254 (2011)
4. Candes, E.: The restricted isometry property and its implications for compressed sensing. *Comptes Rendus Mathematique* **346**(9–10), 589–592 (2008)
5. Cohen, A., Dahmen, W., DeVore, R.: Compressed sensing and best k-term approximation. *J. Am. Math. Soc.*, 211–231 (2009)
6. Zhang, Y.: Theory of compressive sensing via  $\ell_1$ -minimization: a non-RIP analysis and extensions. *J. Oper. Res. Soc. China* **1**(1), 79–105 (2013)
7. Grasmair, M.: Linear convergence rates for tikhonov regularization with positively homogeneous functionals. *Inverse Prob.* **27**(7), 075014 (2011)
8. Vaiter, S., Peyre, G., Dossal, C., Fadili, J.: Robust sparse analysis regularization. *IEEE Trans. Inf. Theory* **59**(4), 2001–2016 (2013)
9. Haltmeier, M.: Stable signal reconstruction via  $\ell^1$ -minimization in redundant, non-tight frames. *IEEE Trans. Signal Process.* **61**(2), 420–426 (2013)
10. Needell, D., Ward, R.: Stable image reconstruction using total variation minimization. *SIAM Journal on Imaging Sciences* **6**(2), 1035–1058 (2013)
11. Needell, D., Ward, R.: Near-optimal compressed sensing guarantees for total variation minimization. *IEEE Trans. Image Process.* **22**(10), 3941–3949 (2013)
12. Krahmer, F., Ward, R.: Stable and robust sampling strategies for compressive imaging. *IEEE Trans. Image Process.* **23**(2), 612–622 (2014)
13. Kabanava, M., Rauhut, H., Zhang, H.: Robust analysis  $\ell_1$ -recovery from gaussian measurements and total variation minimization. *Eur. J. Appl. Math.* **26**(6), 917–929 (2015)

14. Vaiteer, S., Golbabaee, M., Fadili, J., Peyré, G.: Model selection with low complexity priors. *Information and Inference* **4**(3), 230–287 (2015)
15. Zhang, H., Cheng, L.: On uniqueness guarantees of solution in convex regularized linear inverse problems. *Linear Algebra Appl.* **486**, 475–483 (2015)
16. Burger, M., Osher, S.: Convergence rates of convex variational regularization. *Inverse Prob.* **20**(5), 1411 (2004)
17. Grasmair, M., Haltmeier, M., Scherzer, O.: Necessary and sufficient conditions for linear convergence of  $\ell_1$ -regularization. *Commun. Pure Appl. Math.* **64**(2), 161–182 (2011)
18. Zhang, H., Yin, W., Cheng, L.: Necessary and sufficient conditions of solution uniqueness in  $\ell_1$ -norm minimization. *J. Optim. Theory Appl.* **164**, 109–122 (2015)
19. Hügel, M., Rauhut, H., Strohme, T.: Remote sensing via  $\ell_1$ -minimization. *Found. Comput. Math.* **14**, 115–150 (2014)
20. Foucart, S., Rauhut, H.: A mathematical introduction to compressive sensing. *Applied and Numerical Harmonic Analysis*. Birkhäuser (2013)
21. Nam, S., Davies, M.E., Elad, M., Gribonval, R.: The cosparsity analysis model and algorithms. *Appl. Comput. Harmon. Anal.* **34**(1), 30–56 (2013)
22. Rudin, L., Osher, S., Fatemi, E.: Nonlinear total variation based noise removal algorithms. *Physica D* **60**, 259–268 (1992)
23. Candès, E.J., Eldar, Y.C., Needell, D., Randall, P.: Compressed sensing with coherent and redundant dictionaries. *Appl. Comput. Harmon. Anal.* **31**(1), 59–73 (2011)
24. Liu, Y., Mi, T., Li, S.: Compressed sensing with general frames via optimal-dual-based  $\ell_1$ -analysis. *IEEE Trans. Inf. Theory* **58**(7), 4201–4214 (2012)
25. Donoho, D.L., Tanner, J.: Sparse nonnegative solution of underdetermined linear equations by linear programming. *Proc. Natl. Acad. Sci. U.S.A.* **102**(27), 9446–9451 (2005)
26. Chandrasekaran, V., Recht, B., Parrilo, P.A., Willsky, A.S.: The convex geometry of linear inverse problems. *Found. Comput. Math.* **12**(6), 805–849 (2012)
27. Kabanava, M., Rauhut, H.: Analysis  $\ell_1$ -recovery with frames and gaussian measurements. *Acta Applicandae Mathematicae* **140**(1), 173–195 (2015)
28. Foucart, S.: Stability and robustness of  $\ell_1$ -minimizations with weibull matrices and redundant dictionaries. *Linear Algebra Appl.* **441**, 4–21 (2014)
29. Fuchs, J.J.: Recovery of exact sparse representations in the presence of bounded noise. *IEEE Trans. Inf. Theory* **51**(10), 3601–3608 (2005)
30. Tibshirani, R.J.: The lasso problem and uniqueness. *Electronic Journal of Statistics* **7**, 1456–1490 (2013)
31. Fadili, J., Peyré, G., Vaiteer, S., Deledalle, C.-A., Salmon, J.: Stable Recovery with Analysis Decomposable Priors. In: *Proceedings of Signal Processing with Adaptive Sparse Structured Representations (SPARS)* (2013)
32. Tropp, J.A.: Greed is good: algorithmic results for sparse approximation. *IEEE Trans. Inf. Theory* **50**(10), 2231–2242 (2004)