

TRIPLE HOMOMORPHISMS OF C*-ALGEBRAS

NGAI-CHING WONG

In memory of our beloved friend, Kosita Beidar (1951.10.21–2004.3.9)

ABSTRACT. In this note, we will discuss what kind of operators between C*-algebras preserves Jordan triple products $\{a, b, c\} = (ab^*c + cb^*a)/2$. These include especially isometries and disjointness preserving operators.

1. INTRODUCTION

Recall that a Banach algebra A is an algebra with a norm $\|\cdot\|$ such that $\|ab\| \leq \|a\|\|b\|$, and every Cauchy sequence converges. A complex Banach algebra A is a C*-algebra if there is an involution $*$ defined on A such that $\|a^*a\| = \|a\|^2$. A special example is $B(H)$, the algebra of all bounded linear operators on a (complex) Hilbert space H . By the Gelfand-Naimark-Segel Theorem, C*-algebras are exactly those norm closed $*$ -subalgebras of $B(H)$. An abelian C*-algebra A can also be represented as the algebra $C_0(X)$ of continuous functions on a locally compact Hausdorff space X vanishing at infinity. X is compact if and only if A is unital.

It is well known that the algebraic structure determines the geometric (norm) structure of a C*-algebra A . Indeed, the norm of a self-adjoint element a of A coincides with the spectral radius of a , and the latter is a pure algebraic object. In general, the norm of an arbitrary element a of A is equal to $\|a^*a\|^{1/2}$, and a^*a is self-adjoint. For an abelian C*-algebra $A = C_0(X)$, we note that the underlying space X can be considered as the maximal ideal space of A consisting of nonzero complex homomorphisms (= linear and multiplicative functionals) of A . The topology of X is the hull-kernel topology, and thus be solely determined by the algebraic structure of A .

In this note, we will discuss how much the algebraic structure can be recovered if we know the norm, or other, structure of a C*-algebra. In particular, isometries and disjointness preserving operators of C*-algebras preserve triple products $\{a, b, c\} = (ab^*c + cb^*a)/2$.

The author is very grateful to our late friend, Kosita Beidar, from whom he learned how to look at a seemingly pure analytic problem from the point of view of an algebraist.

Date: August 30, 2004; to appear in Southeast Asian Bulletin of Mathematics.

2000 Mathematics Subject Classification. 46L05, 46B04, 47B48, 17C65.

Key words and phrases. C*-algebras, Jordan triples, isometries, disjointness preserving operators.

Partially supported by Taiwan National Science Council.

2. THE GEOMETRIC STRUCTURE DETERMINES THE ALGEBRAIC STRUCTURE

Suppose $T : A \rightarrow B$ is an isometric linear embedding between C^* -algebras. That is, $\|Tx\| = \|x\|$ for all x in A . We are interested in knowing what kind of algebraic structure T inherits from A to its range, which is in general just a Banach subspace of B . We begin with two famous results.

Theorem 2.1. (Banach and Stone; see, e.g., [6]) *Let X and Y be locally compact Hausdorff spaces. Let $T : C_0(X) \rightarrow C_0(Y)$ be a surjective linear isometry. Then T is a weighted composition operator*

$$Tf = h \cdot f \circ \varphi, \quad \forall f \in C_0(X),$$

where h is a continuous scalar function on Y with $|h(y)| \equiv 1$, and φ is a homeomorphism from Y onto X . Consequently, two abelian C^* -algebras are isomorphic as Banach spaces if, and only if, they are isomorphic as $*$ -algebras.

Here is a sketch of the proof. Let $T^* : M(Y) \rightarrow M(X)$ be the dual map of T , which is again a surjective linear isometry from the Banach space $M(Y) = C_0(Y)^*$ of all bounded Radon measures on Y onto that on X . Restricting T^* to the dual unit balls, which are weak* compact and convex, we get an affine homeomorphism. Since the nonzero extreme points of the dual unit balls are exactly unimodular scalar multiples of point masses, T^* sends a point mass δ_y to $\lambda\delta_x$. Here $y \in Y$, $x \in X$ and $|\lambda| = 1$. We write $x = \varphi(y)$ and $\lambda = h(y)$ to indicate that x and λ depend on y . It follows that

$$Tf(y) = T^*(\delta_y)(f) = h(y)\delta_{\varphi(y)}(f) = h(y)f(\varphi(y)).$$

It is then routine to see that h is unimodular and continuous on Y , and that φ is a homeomorphism from Y onto X .

Theorem 2.2 (Kadison [7]). *Let A and B be C^* -algebras. Let $T : A \rightarrow B$ be a surjective linear isometry. Then there is a unitary element u in $\tilde{B} = B \oplus \mathbb{C}1$, the unitization of B , and a Jordan $*$ -isomorphism $J : A \rightarrow B$ such that*

$$Ta = uJ(a), \quad \forall a \in A.$$

Consequently, two C^* -algebras are isomorphic as Banach spaces if, and only if, they are isomorphic as Jordan $*$ -algebras.

Recall that a Jordan $*$ -homomorphism J preserves linear sums, involutions and Jordan products: $a \circ b = (ab + ba)/2$. It is easy to see that the abelian case can also be written in this form with $u = h$ and $Jf = f \circ \varphi$. In general, the product of a pair of elements in A can be decomposed into two parts $ab = a \circ b + [a, b]$, the sum of the Jordan product and the Lie product $[a, b] = (ab - ba)/2$. It is plain that $a \circ b = b \circ a$ is commutative and $[a, b] = -[b, a]$ is anti-commutative. However they are not associative. The Kadison theorem states that the norm structure of a C^* -algebra determines completely its Jordan structure.

It is interesting to note that Jordan products are determined by squares:

$$a \circ b = \frac{(a+b)^2 - a^2 - b^2}{2}, \quad \forall a, b \in A.$$

A similar algebraic structure exists in C*-algebras, namely, the Jordan triple products:

$$\{a, b, c\} = \frac{ab^*c + cb^*a}{2}.$$

There is also a polar identity for triples:

$$\{a, b, c\} = \frac{1}{8} \sum_{\alpha^2=1} \sum_{\beta^4=1} \alpha\beta \{a + \alpha b + \beta c\}^{(3)},$$

Hence, a linear map T between C*-algebras preserves triple products if and only if it preserves cubes $a^{(3)} = \{a, a, a\} = aa^*a$.

Kaup [8] generalizes Kadison theorem: a linear surjection between C*-algebras $T : A \rightarrow B$ is an isometry if and only if it preserves triple products. A geometric proof of the Kadison Theorem is given by Dang, Friedman and Russo [3]. It goes first to note that a norm exposed face of the dual unit ball U_{B^*} is of the form $F_u = \{\varphi \in B^* : \|\varphi\| = \varphi(u) \leq 1\}$ for a unique partial isometry u in B^{**} . For two φ, ψ in B^* , they are said to be orthogonal to each other if they have polar decompositions $\varphi = u|\varphi|, \psi = v|\psi|$ such that $u \perp v$, i.e., $u^*v = uv^* = 0$. This amounts to say that $\|\varphi \pm \psi\| = \|\varphi\| + \|\psi\|$. Two faces F_u, F_v are orthogonal if and only if $u \perp v$. Then they verify that the dual map T^* of the surjective linear isometry T maps faces to faces and preserves orthogonality. Consequently, $T^{**} : A^{**} \rightarrow B^{**}$ sends orthogonal partial isometries to orthogonal partial isometries. By the spectral theory, every element a in $A \subset A^{**}$ can be approximated in norm by a finite linear sum of orthogonal partial isometries $\sum_j \lambda_j u_j$. Then its cube $a^{(3)}$ can also be approximated by $\sum_j \lambda_j^{(3)} u_j$. It follows that $T(a^{(3)})$ and $(Ta)^{(3)}$ can both be approximated by $\sum_j \lambda_j^{(3)} T^{**} u_j$. Hence $T(a^{(3)}) = (Ta)^{(3)}$, and thus T preserves triple products by the polar identity.

We note that the above (geometric) proof of the Kadison theorem depends very much on the fact the range of the isometry is again a C*-algebra. Extending the Holsztynski theorem [4, 6], Chu and Wong [2] studied non-surjective linear isometries between C*-algebras.

Theorem 2.3 (Chu and Wong [2]). *Let A and B be C*-algebras and let T be a linear isometry from A into B . There is a largest closed projection p in B^{**} such that $T(\cdot)p : A \rightarrow B^{**}$ is a Jordan triple homomorphism and*

$$T(ab^*c + cb^*a)p = T(a)T(b)^*T(c)p + T(c)T(b)^*T(a)p, \quad \forall a, b, c \in A.$$

When A is abelian, we have $\|T(a)p\| = \|a\|$ for all a in A . In particular, T reduces locally to a Jordan triple isomorphism on the JB-triple generated by any a in A , also an abelian C*-algebra, by a closed projection p_a .*

Besides the triple technique, the proof of the above theorem makes use of the concept of representing elements of a C^* -algebra as special sections of a continuous field of Hilbert spaces developed in [10]. It is still geometric.

3. DISJOINTNESS PRESERVING OPERATORS ARE TRIPLE HOMOMORPHISMS

In this section, we do not assume the operator T is isometric. Although the following statement might have been known to experts, we provide a new and short proof here as we do not find any in the literature.

Theorem 3.1. *Let $T : A \rightarrow B$ be a bounded linear map between C^* -algebras. Then T is a triple homomorphism if and only if T^{**} sends partial isometries of A^{**} to partial isometries of B^{**} .*

Proof. One direction is trivial. Suppose T^{**} sends partial isometries of A^{**} to partial isometries of B^{**} . Let u, v be two partial isometries in A . Observe that they are orthogonal to each other, namely, $u^*v = uv^* = 0$, if and only if they have orthogonal initial spaces and orthogonal range spaces. This amounts to saying that $u + \lambda v$ is a partial isometry for all scalar λ with $|\lambda| = 1$. Consequently, T sends orthogonal partial isometries to orthogonal partial isometries. For every a in $A \subset A^{**}$, approximate a in norm by a finite linear sum $\sum_n \lambda_n u_n$ of orthogonal partial isometries in A^{**} . Then its cube $a^{(3)} = aa^*a$ can also be approximated in norm by $\sum_n \lambda_n^{(3)} u_n$. It follows that Ta and $T(a^{(3)})$ can be approximated in norm by $\sum_n \lambda_n T^{**} u_n$ and $\sum_n \lambda_n^{(3)} T^{**} u_n$, respectively. This gives $T(a^{(3)}) = (Ta)^{(3)}$, $\forall a \in A$. By the polar identity, we see that T is a triple homomorphism. \square

We say that a linear map $T : A \rightarrow B$ between C^* -algebras is *disjointness preserving* if

$$a^*b = ab^* = 0 \quad \text{implies} \quad (Ta)^*(Tb) = (Ta)(Tb)^* = 0, \quad \forall a, b \in A.$$

Clearly, T is disjointness preserving if and only if it preserves disjointness of partial isometries. It is clear that every triple homomorphism preserves disjointness. Looking at the well-known result of Jarosz [5, 6] in the abelian case, we see that not every disjointness preserving map is a triple homomorphism. Indeed, let $T : C_0(X) \rightarrow C_0(Y)$ be a bounded disjointness preserving linear map between abelian C^* -algebras. Then there is a closed subset Y_0 of Y on which every Tf vanishes. On $Y_1 = Y \setminus Y_0$ there is a bounded continuous function h and a continuous map φ from Y_1 into X such that $Tf|_{Y_1} = h \cdot f \circ \varphi$ for all f in $C_0(X)$. Hence, T is a triple homomorphism if and only if $T^{**}1$ is a partial isometry in $C_0(Y)^{**}$. We end this note with a proof of this fact for the non-abelian case.

Theorem 3.2. *Let $T : A \rightarrow B$ be a bounded linear map between C^* -algebras. Then T is a triple homomorphism if and only if T is disjointness preserving and $T^{**}1$ is a partial isometry in B^{**} .*

Proof. We verify the sufficiency only. For simplicity of notations, denote again by T the bidual map of T from A^{**} into B^{**} .

Let a be a self-adjoint element of A . Identify the commutative C*-subalgebra of A generated by 1 and a with $C(X)$, where $X \subseteq [0, \|a\|]$ is the spectrum of a . Let $0 = \alpha_0 < \alpha_1 < \cdots < \alpha_{n-1} < \alpha_n = \|a\| + 1$ such that $X = \cup_k X_k$ is a partition of X with $X_k = X \cap [\alpha_{k-1}, \alpha_k) \neq \emptyset$, and pick an arbitrary point x_k from X_k for each $k = 1, 2, \dots, n$. In particular,

$$1 = \sum_k 1_{X_k},$$

where 1_{X_k} is the characteristic function of the set X_k . For $1 < j < k$, we can find two sequences $\{f_m\}_m$ and $\{g_m\}_m$ in $C(X) \cap A$ such that $f_m g_{m+p} = 0$ for $m, p = 0, 1, \dots$, $f_m \rightarrow 1_{X_j}$ and $g_m \rightarrow 1_{X_k}$ pointwisely on X . By the weak* continuity of T , we see that

$$T(1_{X_k})T(f_m)^* = \lim_{p \rightarrow \infty} T(g_{m+p})T(f_m)^* = 0 \quad \text{for all } m = 1, 2, \dots$$

Thus

$$T(1_{X_k})T(1_{X_j})^* = \lim_{m \rightarrow \infty} T(1_{X_k})T(f_m)^* = 0.$$

Similarly, we have

$$T(1_{X_k})^*T(1_{X_j}) = 0.$$

When $j = 1$, we note that 1_{X_1} is an open projection in A^{**} . Hence there is an increasing net $\{a_\lambda\}_\lambda$ in A converges to 1_{X_1} weakly (see [9, Proposition 3.11.9], and also [1]). Using an argument similar to above, we still get

$$T(1_{X_k})T(1_{X_1})^* = T(1_{X_k})^*T(1_{X_1}) = 0.$$

Consequently, for each $j = 1, 2, \dots, n$, we have

$$T(1)T(1_{X_j})^*T(1) = \sum_{m,n} T(1_{X_n})T(1_{X_j})^*T(1_{X_m}) = (T(1_{X_j}))^{(3)}.$$

This gives

$$\sum_n T(1_{X_n}) = T1 = (T1)^{(3)} = \sum_n (T(1_{X_n}))^{(3)}.$$

Multiplying the above identity on the left by $T(1_{X_n})^*$ and $((T(1_{X_n}))^{(3)})^*$ respectively, we see that

$$(T(1_{X_n}) - ((T(1_{X_n}))^{(3)})^*(T(1_{X_n}) - ((T(1_{X_n}))^{(3)})) = 0.$$

Hence $T(1_{X_n})$ is a partial isometry for each n and orthogonal to the others. It follows that

$$\begin{aligned} (T(f))^{(3)} &= \lim \left(\sum_n f(x_n)T(1_{X_n}) \right)^{(3)} = \lim \sum_n f(x_n)^{(3)}(T(1_{X_n}))^{(3)} \\ &= \lim \sum_n f(x_n)^{(3)}T(1_{X_n}) = T(f^{(3)}), \end{aligned}$$

for all f in $C(X)$. By the polar identity, T preserves triple products in $C(X)$. Let $p = T1^*T1$ and $q = T1T1^*$ be the initial and range projections of the partial isometry $T1$, respectively. We have

$$2Tf = 2T\{1, 1, f\} = qTf + Tfp, \quad \forall f \in C(X).$$

It then follows that

$$Tf = qTf = Tfp,$$

for all f in $C(X)$. Thus for each self-adjoint element a of A ,

$$(1) \quad T(a^{(3)}) = (Ta)^{(3)} \quad \text{and} \quad Ta = qTa = Tap.$$

At this point, we assume further that $T1 = p = q$ is a projection. It then follows from (1) that for every self-adjoint element a of A ,

$$Ta = T(a^*) = T\{1, a, 1\} = \{T1, a, T1\} = T1(Ta)^*T1 = (Ta)^*.$$

Therefore, T also preserves self-adjointness. For any self-adjoint elements a, b of A , by observing the identities $T(a \pm b)^{(3)} = (T(a \pm b))^{(3)}$, we also see that $T(aba) = TaTbTa$. Altogether, it follows that $T(a + ib)^{(3)} = (T(a + ib))^{(3)}$. In other words, T is a triple homomorphism from A into B .

In general, we consider the map $J = T1^*T$ from A into $T1^*TA \subseteq B^{**}$. It follows from (1) that J is also disjointness preserving. Since $J1 = T1^*T1 = p$ is a projection, J is a triple homomorphism. Hence $J(a^{(3)}) = (Ja)^{(3)}$, and thus

$$T1^*T(a^{(3)}) = T1^*Ta(Ta)^*T1T1^*Ta = T1^*Ta(Ta)^*qTa = T1^*(Ta)^{(3)},$$

for all a in A by (1). Therefore,

$$qT(a^{(3)}) = T1T1^*T(a^{(3)}) = T1T1^*(Ta)^{(3)} = q(Ta)^{(3)},$$

and then by (1) again,

$$T(a^{(3)}) = (Ta)^{(3)}, \quad \forall a \in A.$$

This says T is a triple homomorphism from A into B . □

REFERENCES

- [1] L. G. Brown, "Semicontinuity and multipliers of C^* -algebras", *Can. J. Math.*, vol. XL, no. 4 (1988), 865–988.
- [2] C.-H. Chu and N.-C. Wong, "Isometries between C^* -algebras", *Revista Matematica Iberoamericana*, **20** (2004), no. 1, 87–105.
- [3] T. Dang, Y. Friedman and B. Russo, "Affine geometric proofs of the Banach Stone theorems of Kadison and Kaup", *Rocky Mountain J. Math.*, **20** (1990), no. 2, 409–428.
- [4] W. Holsztynski, "Continuous mappings induced by isometries of spaces of continuous functions", *Studia Math.* **26** (1966), 133–136.
- [5] K. Jarosz, "Automatic continuity of separating linear isomorphisms", *Canad. Math. Bull.* **33** (1990), 139–144.
- [6] J.-S. Jeang and N.-C. Wong, "Weighted composition operators of $C_0(X)$'s", *J. Math. Anal. Appl.* **201** (1996), 981–993.
- [7] R. V. Kadison, "Isometries of operator algebras", *Ann. of Math.* **54** (1951), 325–338.
- [8] W. Kaup, "A Riemann mapping theorem for bounded symmetric domains in complex Banach spaces", *Math. Z.* **138** (1983), 503–529.
- [9] G. K. Pedersen, "C*-algebras and their automorphism groups", Academic Press, 1979, London.
- [10] N.-C. Wong, "Left quotients of a C^* -algebra, I: Representation via vector sections", *J. Operator Theory* **32** (1994), 185–201.

DEPARTMENT OF APPLIED MATHEMATICS, NATIONAL SUN YAT-SEN UNIVERSITY, AND NATIONAL CENTER FOR THEORETICAL SCIENCES, KAOHSIUNG, 80424, TAIWAN, R.O.C.

E-mail address: wong@math.nsysu.edu.tw