TRANSITION PROBABILITIES OF NORMAL STATES DETERMINE THE JORDAN STRUCTURE OF A QUANTUM SYSTEM

CHI-WAI LEUNG, CHI-KEUNG NG, AND NGAI-CHING WONG

ABSTRACT. Let $\Phi : \mathfrak{S}(M_1) \to \mathfrak{S}(M_2)$ be a bijection (not assumed affine nor continuous) between the sets of normal states of two quantum systems, modelled on the self-adjoint parts of von Neumann algebras M_1 and M_2 , respectively. This paper concerns with the situation when Φ preserves (or partially preserves) one of the following three notions of "transition probability" on the normal state spaces: the Uhlmann transition probability P_U , the Raggio transition probability P_B and an "asymmetric transition probability" P_0 as defined in this article.

It is shown that the two systems are isomorphic, i.e. M_1 and M_2 are Jordan *isomorphic, if Φ preserves all pairs with zero Uhlmann (respectively, Raggio or asymmetric) transition probability, i.e., for any normal states μ and ν , we have

$$P(\Phi(\mu), \Phi(\nu)) = 0$$
 if and only if $P(\mu, \nu) = 0$,

where P stands for P_U (respectively, P_R or P_0). Furthermore, as an extension of Wigner's theorem, it is shown that there is a Jordan *-isomorphism $\Theta : M_2 \to M_1$ with

$$\Phi = \Theta^*|_{\mathfrak{S}(M_1)}$$

if and only if Φ preserves the "asymmetric transition probability". This is also equivalent to Φ preserving the Raggio transition probability. Consequently, if Φ preserves the Raggio transition probability, it will preserve the Uhlmann transition probability as well. As another application, the sets of normal states equipped with either the usual metric, the Bures metric or "the metric induced by the self-dual cone" are complete Jordan *-invariants for the underlying von Neumann algebras.

1. INTRODUCTION

Let \mathfrak{H}_1 and \mathfrak{H}_2 be two (complex) Hilbert spaces and $T : \mathfrak{H}_1 \to \mathfrak{H}_2$ be a bijective map (not assumed linear nor continuous). Wigner's theorem states that if T preserves the transition probability, in the sense that

$$|\langle T(\xi), T(\eta) \rangle|^2 = |\langle \xi, \eta \rangle|^2 \qquad (\xi, \eta \in \mathfrak{H}_1),$$

then there exist a unitary or an anti-unitary $S : \mathfrak{H}_1 \to \mathfrak{H}_2$ and a function $f : \mathfrak{H}_2 \to \mathbb{T}$ such that $T(\xi) = f(\xi)S(\xi)$ ($\xi \in \mathfrak{H}_1$). Uhlhorn's theorem, as a generalization of Wigner's theorem, states that if dim $\mathfrak{H}_1 \geq 3$ and T preserves pairs with zero transition probability,

Date: July 21, 2018.

Key words and phrases. von Neumann algebras; normal states; transition probability; metric spaces; Jordan *-isomorphisms.

in the sense that

$$\langle T(\xi), T(\eta) \rangle = 0$$
 if and only if $\langle \xi, \eta \rangle = 0$ $(\xi, \eta \in \mathfrak{H}_1)$,

then there exist a unitary or an anti-unitary S and a function $g: \mathfrak{H}_2 \to \mathbb{C} \setminus \{0\}$ such that $T(\xi) = g(\xi)S(\xi) \ (\xi \in \mathfrak{H}_1).$

Let A be a (complex) C^{*}-algebra and $\mu, \nu \in A^*$ be pure states of A. The transition probability between μ and ν is defined to be the quantity

$$P(\mu,\nu) := \mu(\mathbf{s}_{\nu}),$$

where \mathbf{s}_{ν} is the support projection of ν in A^{**} . It is well-known that $P(\mu, \nu) = P(\nu, \mu)$, i.e., $\mu(\mathbf{s}_{\nu}) = \nu(\mathbf{s}_{\mu})$, for pure states μ and ν (see e.g. [4]). Suppose that $\pi : A \to \mathcal{L}(\mathfrak{H})$ is a *-representation of A and $\xi \in \mathfrak{H}$, we denote, as usual,

(1.1)
$$\omega_{\xi}(x) := \langle \pi(x)\xi, \xi \rangle \qquad (x \in A).$$

In the case when $A = \mathcal{L}(\mathfrak{H})$ and $\pi : \mathcal{L}(\mathfrak{H}) \to \mathcal{L}(\mathfrak{H})$ is the default representation, the functionals ω_{ξ} and ω_{η} are pure normal states of $\mathcal{L}(\mathfrak{H})$ (where $\xi, \eta \in \mathfrak{H}$) and we have

$$P(\omega_{\xi}, \omega_{\eta}) = |\langle \xi, \eta \rangle|^2.$$

In this setting, Wigner's (respectively, Uhlhorn's) theorem can be interpreted as structural results concerning bijections between pure normal state spaces of $\mathcal{L}(\mathfrak{H}_1)$ and $\mathcal{L}(\mathfrak{H}_2)$ that preserve (respectively, partially preserve) the transition probability, and several proofs are given (see e.g. [11] or [23, Theorem 1]). They have also been extended to the setting of indefinite inner product spaces by Molnár (see [18, Theorem 1] and [19, Corollary 1]). Through our study we will also give another proof for Wigner's theorem (see Corollary 3.3).

On the other hand, Shultz provided a throughout study of transition probability preserving bijections between pure state spaces of general C^* -algebras. Under some extra conditions, such maps are induced by the dual maps of algebraic or Jordan *-isomorphisms of the C^* -algebras (see e.g., [3, 4, 23] for details). Related considerations of maps between pure state spaces of C^* -algebras preserving transition probability or other properties can also be found in, e.g., [5, 16, 25].

However, the pure state setting of transition probability is inappropriate to be adapted to the case of von Neumann algebras. Unlike $\mathcal{L}(\mathfrak{H})$, a general von Neumann algebra may not have any pure normal state at all. Therefore, people are looking for suitable notions of transition probability on the space $\mathfrak{S}(M)$ of all normal states on a von Neumann algebra M (see e.g. [2, 6, 21, 24, 28]). Here, by a normal state on M, we means a norm one positive normal linear functional on M, and it is different from the notion of "physical states" as introduced in [9].

Let $\mathbf{R}(M)$ denote the collection of all (unitary equivalence classes of) faithful unital *representations of a von Neumann algebra M. For any $\mu, \nu \in \mathfrak{S}(M)$ and $(\mathfrak{H}, \pi) \in \mathbf{R}(M)$, we set $\mathfrak{H}(\mu) := \{\xi \in \mathfrak{H} : \omega_{\xi} = \mu\}$ (could be empty). The quantity

$$P_U(\mu,\nu) := \sup\{|\langle \xi,\eta \rangle|^2 : \xi \in \mathfrak{H}(\mu), \eta \in \mathfrak{H}(\nu), (\mathfrak{H},\pi) \in \mathbf{R}(M)\}$$

is well-defined and is called the *Uhlmann transition probability* of μ and ν ([27]). The Uhlmann transition probability is related to the so-called *Bures distance* d_B through the formula

(1.2)
$$d_B(\mu,\nu) := \sqrt{2 - 2\sqrt{P_U(\mu,\nu)}}.$$

This metric d_B is in general different from the usual distance d_1 on $\mathfrak{S}(M)$ given by

$$d_1(\mu, \nu) := \|\mu - \nu\|.$$

In [20], Raggio defined another transition probability. Suppose that $(M, \mathfrak{H}, \mathfrak{P}, \mathfrak{P}, J)$ is the standard form for M as in [12] (see Section 2 below for a brief exploration). By [12, Lemma 2.10], for any $\mu \in \mathfrak{S}(M)$, there is a unique $\xi_{\mu} \in \mathfrak{P}$ satisfying

(1.3)
$$\mu = \omega_{\xi_{\mu}}.$$

If $\mu, \nu \in \mathfrak{S}(M)$, the positive real number

$$P_R(\mu,\nu) := \langle \xi_\mu, \xi_\nu \rangle$$

is called the *Raggio transition probability* of μ and ν . As in the Uhlmann case, the Raggio transition probability induces a metric on $\mathfrak{S}(M)$ by

(1.4)
$$d_2(\mu,\nu) := \sqrt{2 - 2\sqrt{P_R(\mu,\nu)}} \qquad (\mu,\nu \in \mathfrak{S}(M)).$$

This metric coincides with the one induced from \mathfrak{H} , namely,

(1.5)
$$d_2(\mu,\nu) = \|\xi_{\mu} - \xi_{\nu}\| \qquad (\mu,\nu \in \mathfrak{S}(M)).$$

In [20, Corollary 1], the following relation between the Raggio and the Uhlmann transition probabilities was presented:

(1.6)
$$P_U(\mu,\nu) \le P_R(\mu,\nu) \le P_U(\mu,\nu)^{1/2} \quad (\mu,\nu\in\mathfrak{S}(M)).$$

In addition, there is a more naïve extension of the "transition probability":

(1.7)
$$P_0(\mu,\nu) := \mu(\mathbf{s}_{\nu}) \qquad (\mu,\nu \in \mathfrak{S}(M)).$$

Strictly speaking, P_0 is not a transition probability, because unlike the two extensions above, P_0 is asymmetric, and $P_0(\mu, \nu) = 1$ is equivalent to $\mathbf{s}_{\mu} \leq \mathbf{s}_{\nu}$ instead of $\mu = \nu$ (c.f. [21, p.325]). Nevertheless, abusing the language, we still call P_0 the "asymmetric transition probability". It seems to be conceptual clear and technically easier to work with it.

Notice that two normal states $\mu, \nu \in \mathfrak{S}(M)$ are orthogonal, i.e., having orthogonal support projections, exactly when they have zero transition probability in any (and equivalently, all) of the above three settings (see (3.10) in Section 3).

The main concern of this article is on those bijections (not assumed affine nor continuous) from the normal state space of one von Neumann algebra to that of another preserving either one of the three transition probabilities above. We obtain two analogues of Wigner's theorem for bijections between normal state spaces of general quantum systems (which are modelled on self-adjoint elements of von Neumann algebras). Furthermore, several weak analogues of Uhlhorn's theorem for normal state spaces of general quantum systems were also obtained.

More precisely, it is shown that a bijection between normal state spaces preserving either the "asymmetric transition probability" (as defined in (1.7)) or the Raggio transition probability was shown to be induced by a Jordan *-isomorphism (see Theorems 3.2(b) and 3.8). The result concerning the "asymmetric transition probability" can be regarded as an extension of the original Wigner's theorem because of Corollary 3.3. Moreover, we verified that the normal state space equipped with either the Uhlmann transition probability, the Raggio transition probability or the "asymmetric transition probability", completely identifies the underlying quantum system (see Theorems 3.2 and 3.4). Consequently, bijections between normal state spaces preserving the Raggio transition probability will preserve the Uhlmann transition probability (see Corollary 3.9).

This study highlighted the importance of the Raggio and the Uhlmann transition probability in quantum mechanics and it also established a strong relation between these two notions of transition probability. On the other hand, the notion of "asymmetric transition probability" that defined in (1.7) seems to be conceptually clearer and easier to implement in physics, although it is not strictly speaking a transition probability. Actually, Theorem 3.2(b) implies that the datum of measurements of observables associated with support projections of states at all other states is sufficient to determine the quantum system completely.

In developing our main results, we also obtained that several metric spaces associated with the sets of normal states of von Neumann algebras (without any algebraic structure) are complete Jordan *-invariants for the underlying algebras (see Corollary 3.11).

2. NOTATIONS AND PRELIMINARIES

Throughout this article, M, M_1 and M_2 are (complex) von Neumann algebras. We denote by $\mathfrak{S}(M)$ and $\mathcal{P}(M)$ the normal state space of M and the set of all projections in M, respectively.

Let $(M, \mathfrak{H}, \mathfrak{P}, J)$ denote the (unique) standard form of a von Neumann algebra M (see [12]). In other words, \mathfrak{H} is a (complex) Hilbert space with M being a (unital) von Neumann subalgebra of $\mathcal{L}(\mathfrak{H})$, J is a conjugate linear isometric involution on \mathfrak{H} and $\mathfrak{P} \subseteq \mathfrak{H}$ is a cone which is *self-dual*, in the sense that

$$\mathfrak{P} = \{\eta \in \mathfrak{H} : \langle \eta, \xi \rangle \geq 0, \text{ for any } \xi \in \mathfrak{P} \},$$

such that the following conditions hold:

(1) JMJ = M', (2) $JcJ = c^*$ for any $c \in \mathfrak{Z}(M)$, (3) $J\xi = \xi$ for any $\xi \in \mathfrak{P}$, (4) $aa^{\mathbf{t}}(\mathfrak{P}) \subseteq \mathfrak{P}$ for any $a \in M$, where M' is the commutant of M in $\mathcal{L}(\mathfrak{H}), \mathcal{Z}(M) := M \cap M'$ and $a^{\mathbf{t}} := JaJ$. We put

$$S_{\mathfrak{H}} := \{\xi \in \mathfrak{H} : \|\xi\| = 1\}$$
 and $S_{\mathfrak{P}} := \mathfrak{P} \cap S_{\mathfrak{H}}.$

Remark 2.1. Suppose that $\{x_i\}_{i\in\mathfrak{I}}$ is a net in M that WOT-converges to $x \in M$, when considered as operators in $\mathcal{L}(H)$. As $\omega_{\xi}(x_i) \to \omega_{\xi}(x)$ ($\xi \in \mathfrak{P}$) and $\{\omega_{\xi} : \xi \in \mathfrak{P}\} = M_*^+$, we know that $\{x_i\}_{i\in\mathfrak{I}}$ weak*-converges to x. Thus, the WOT on $M \subseteq \mathcal{L}(\mathfrak{H})$ coincides with the weak*-topology. In particular, if $\{e_i\}_{i\in\mathfrak{I}}$ is an increasing net in $\mathcal{P}(M)$ with $e_i \uparrow e_0 \in \mathcal{P}(M)$, then $e_i y e_i \xrightarrow{w^*} e_0 y e_0$ ($y \in M$), because $\omega_{\xi}(e_i y e_i) = \langle y e_i \xi, e_i \xi \rangle \to \omega_{\xi}(e_0 y e_0)$ ($\xi \in \mathfrak{P}$).

The following proposition contains some known results. Note that Part (b) of it inherits from [7, Theorem 2.2], while part (a) can be regarded as a result of Dye, because all the ingredients for its proof are already in [10] (and a similar discussion can be found in [22], although it is not explicitly stated there). We give a simple proof for part (b) here, so that we have a complete elementary proof for Corollary 3.3 below.

Proposition 2.2. Let M_1 and M_2 be two von Neumann algebras. Suppose that there is an orthoisomorphism $\Gamma : \mathcal{P}(M_1) \to \mathcal{P}(M_2)$, i.e. Γ is bijective, and for any $p, q \in \mathcal{P}(M_1)$,

$$p q = 0$$
 if and only if $\Gamma(p)\Gamma(q) = 0$.

(a) (Dye) M_1 and M_2 are Jordan *-isomorphic.

(b) If Γ extends to a bounded linear map $\tilde{\Gamma}: M_1^{\mathrm{sa}} \to M_2^{\mathrm{sa}}$, then $\tilde{\Gamma}$ is a Jordan isomorphism.

Proof: (a) By [10, Lemma 1], the bijection Γ is an order isomorphism that sends central projections to central projections. Let e_k be the central projection in M_k such that $e_k M_k$ is the type I_2 part of M_k (k = 1, 2).

We first show that $\Gamma(e_1) = e_2$ and $(1 - e_1)M_1$ is Jordan *-isomorphic to $(1 - e_2)M_2$. In fact, as Γ is an order isomorphism with $\Gamma(1 - e_1) \in \mathcal{Z}(M_2)$, it restricts to an orthoisomorphism from $\mathcal{P}((1 - e_1)M_1)$ onto $\mathcal{P}(\Gamma(1 - e_1)M_2)$. The absence of nonzero type I_2 summand in $(1 - e_1)M_1$ and the Corollary in [10, p. 83] ensure that $\Gamma|_{\mathcal{P}((1-e_1)M_1)}$ extends to a Jordan *-isomorphism from $(1 - e_1)M_1$ onto $\Gamma(1 - e_1)M_2$. Hence, $\Gamma(1 - e_1)M_2$ does not have a nonzero type I_2 summand neither. This means $\Gamma(1-e_1)e_2 = 0$, or equivalently,

$$\Gamma(1-e_1) \le 1-e_2.$$

Similarly, $\Gamma^{-1}(1-e_2) \leq 1-e_1$. By [10, Lemma 1], one has $1-\Gamma(e_1) = \Gamma(1-e_1) = 1-e_2$.

It remains to show that e_1M_1 is Jordan *-isomorphic to e_2M_2 . Indeed, because $\Gamma(e_1) = e_2$, the map Γ restricts to an orthoisomorphism from $\mathcal{P}(e_1M_1)$ onto $\mathcal{P}(e_2M_2)$. Since

$$\Gamma(\mathfrak{Z}(\mathbf{e}_1 M_1) \cap \mathcal{P}(\mathbf{e}_1 M_1)) = \mathfrak{Z}(\mathbf{e}_2 M_2) \cap \mathcal{P}(\mathbf{e}_2 M_2),$$

 Γ induces an orthoisomorphism from $\mathcal{P}(\mathcal{Z}(\mathbf{e}_1 M_1))$ onto $\mathcal{P}(\mathcal{Z}(\mathbf{e}_2 M_2))$, and the Corollary in [10, p. 83] implies that $\mathcal{Z}(\mathbf{e}_1 M_1)$ is *-isomorphic to $\mathcal{Z}(\mathbf{e}_2 M_2)$. The conclusion now follows from the fact that $\mathbf{e}_k M_k = \mathcal{Z}(\mathbf{e}_k M_k) \otimes M_2(\mathbb{C})$ (k = 1, 2).

(b) Let $n \in \mathbb{N}$ and $r_1, ..., r_n \in \mathbb{R}$. Consider $\{p_1, ..., p_n\}$ to be a set of orthogonal elements in $\mathcal{P}(M_1)$. If we denote $a := \sum_{k=1}^n r_k p_k$, then we have

$$\tilde{\Gamma}(a^2) = \sum_{k=1}^n r_k^2 \Gamma(p_k) = \tilde{\Gamma} \left(\sum_{k=1}^n r_k p_k\right)^2$$

(the last equality uses the fact that Γ preserves orthogonality). Since elements with finite spectrum is norm-dense in M_1^{sa} , the continuity of $\tilde{\Gamma}$ ensures that $\tilde{\Gamma}(x^2) = \tilde{\Gamma}(x)^2$ $(x \in M_1^{\text{sa}})$, as claimed.

Our next proposition is also well-known (although we do not find the second conclusion explicitly stated anywhere), and we will only give a brief account for it.

Proposition 2.3. Let $\Theta : M_2 \to M_1$ be a Jordan *-isomorphism. Then $\Theta^*(\mathfrak{S}(M_1)) = \mathfrak{S}(M_2)$ and $\Theta^*|_{\mathfrak{S}(M_1)}$ preserves both the Raggio and the Uhlmann transition probabilities.

Proof: By [14, Theorem 10], there is a central projection $f_2 \in \mathcal{P}(M_2) \cap \mathcal{Z}(M_2)$ such that Θ restricts to a *-isomorphism $\Theta_i : f_2M_2 \to f_1M_1$ as well as a *-anti-isomorphism $\Theta_a : (1 - f_2)M_2 \to (1 - f_1)M_1$, where $f_1 := \Theta(f_2)$. For $k \in \{1, 2\}$, one has

$$\mathfrak{S}(M_k) = \left\{ \left(t\mu_k, (1-t)\nu_k \right) : t \in [0,1]; \mu_k \in \mathfrak{S}(f_k M_k); \nu_k \in \mathfrak{S}\left((1-f_k)M_k \right) \right\}$$

and $\mathbf{R}(M_k) = \mathbf{R}(f_k M_k) \times \mathbf{R}((1 - f_k) M_k)$ in the canonical way.

Since all *-isomorphisms and *-anti-isomorphisms between von Neumann algebras are isometric and weak*-continuous, the same is true for Jordan *-isomorphisms and we have $\Theta^*(\mathfrak{S}(M_1)) = \mathfrak{S}(M_2).$

Note that Θ_i and Θ_a induce canonical bijections from $\mathbf{R}(f_1M_1)$ onto $\mathbf{R}(f_2M_2)$ and from $\mathbf{R}((1-f_1)M_1)$ onto $\mathbf{R}((1-f_2)M_2)$, respectively. Indeed, if (π, \mathfrak{H}) is a *-representation of $(1-f_1)M_1$ and $(\bar{\pi}, \bar{\mathfrak{H}})$ is its conjugate representation, then $(\bar{\pi} \circ \Theta_a, \bar{\mathfrak{H}})$ is a *-representation of $(1-f_2)M_2$. Now, it follows from the definition of P_U that $\Theta^*|_{\mathfrak{S}(M_1)}$ preserves P_U .

On the other hand, if $(M_k, \mathfrak{H}_k, \mathfrak{P}_k, J_k)$ is the standard form of M_k (k = 1, 2), then the uniqueness of the standard form ensures that $(M_k, \mathfrak{H}_k, \mathfrak{P}_k, J_k)$ decomposes canonically with respect to f_k and $1 - f_k$. Moreover, if $(M, \mathfrak{H}, \mathfrak{P}, J)$ is the standard form of a von Neumann algebra M, then $x \mapsto Jx^*J$ is a *-anti-isomorphism from M to M', and $(M', \mathfrak{H}, \mathfrak{P}, J)$ is the standard form of M'. From these, one can check easily that $\Theta^*|_{\mathfrak{S}(M_1)}$ preserves P_R .

3. The main results

Set $\mathcal{P}_{\sigma}(M) := \{\mathbf{s}_{\mu} : \mu \in \mathfrak{S}(M)\}$. For any $p \in \mathcal{P}(M)$, it follows from Zorn's Lemma that there is an orthogonal family $\{p_i\}_{i \in \mathfrak{I}}$ in $\mathcal{P}_{\sigma}(M)$ satisfying

$$(3.1) p = \sum_{i \in \mathfrak{I}} p_i$$

(the convergence is taken in the weak*-topology). We write

$$F_0(p) := \{\nu \in \mathfrak{S}(M) : \nu(p) = 0\}$$

Obviously, $F_0(p)$ coincides with the closed face $(1-p)\mathfrak{S}(M)(1-p)\cap\mathfrak{S}(M)$ of $\mathfrak{S}(M)$. Moreover, since

(3.2)
$$\nu(\mathbf{s}_{\mu}) = 0$$
 if and only if $\mathbf{s}_{\nu}\mathbf{s}_{\mu} = 0$,

we have

(3.3)
$$F_0(\mathbf{s}_{\mu}) = \{ \nu \in \mathfrak{S}(M) : \mathbf{s}_{\nu}\mathbf{s}_{\mu} = 0 \} \qquad (\mu \in \mathfrak{S}(M))$$

If $p = \sum_{i \in \mathfrak{I}} p_i$ is as in (3.1), then

(3.4)
$$F_0(p) = \bigcap_{i \in \mathfrak{I}} F_0(p_i).$$

See, e.g., [26], for more explorations between projections and their associated faces.

We say that a map $\Phi : \mathfrak{S}(M_1) \to \mathfrak{S}(M_2)$ is biorthogonality preserving if for any $\mu, \nu \in \mathfrak{S}(M_1)$, one has

$$\mathbf{s}_{\mu}\mathbf{s}_{\nu} = 0$$
 if and only if $\mathbf{s}_{\Phi(\mu)}\mathbf{s}_{\Phi(\nu)} = 0.$

Lemma 3.1. Let M_1 and M_2 be two von Neumann algebras. Suppose that $\Phi : \mathfrak{S}(M_1) \to \mathfrak{S}(M_2)$ is a biorthogonality preserving bijection.

(a) There exists an orthoisomorphism $\check{\Phi} : \mathfrak{P}(M_1) \to \mathfrak{P}(M_2)$ such that

(3.5)
$$\Phi(F_0(p)) = F_0(\check{\Phi}(p)) \quad (p \in \mathcal{P}(M_1)) \quad and \quad \check{\Phi}(\mathbf{s}_{\mu}) = \mathbf{s}_{\Phi(\mu)} \quad (\mu \in \mathfrak{S}(M_1)).$$

(b) If M_1 or M_2 does not have a type I_2 summand, then Φ extends to a Jordan isomorphism from M_1^{sa} to M_2^{sa} .

(c) If
$$\Theta: M_2 \to M_1$$
 is a Jordan *-isomorphism satisfying $\Phi = \Theta^*|_{\mathfrak{S}(M_1)}$, then

$$\Phi(\nu)(\mathbf{s}_{\Phi(\mu)}) = \nu(\mathbf{s}_{\mu}) \quad (\mu, \nu \in \mathfrak{S}(M_1)), \quad and \quad \Phi = \Theta^{-1}|_{\mathfrak{P}(M_1)}.$$

Proof. (a) We denote by $\mathcal{F}(M_k)$ the set of all closed faces of $\mathfrak{S}(M_k)$ (k = 1, 2). The bijectivity of Φ and (3.3) tell us that Φ is biorthogonality preserving if and only if

(3.6)
$$\Phi(F_0(\mathbf{s}_{\mu})) = F_0(\mathbf{s}_{\Phi(\mu)}) \qquad (\mu \in \mathfrak{S}(M_1)).$$

Let $p \in \mathcal{P}(M_1)$, and $p := \sum_{i \in \mathfrak{I}} \mathbf{s}_{\mu_i}$ be a decomposition as in (3.1) for a family $\{\mu_i\}_{i \in \mathfrak{I}}$ in $\mathfrak{S}(M_1)$ with its elements having disjoint support projections. By the hypothesis, elements

in $\{\Phi(\mu_i)\}_{i\in\mathfrak{I}}$ have disjoint support projections, and hence $\sum_{i\in\mathfrak{I}} \mathbf{s}_{\Phi(\mu_i)}$ converges in the weak*-topology to a projection $\check{\Phi}(p) \in \mathcal{P}(M_2)$. Since Φ is injective, (3.4) and (3.6) imply

$$\Phi(F_0(p)) = \bigcap_{i \in \mathfrak{I}} \Phi(F_0(\mathbf{s}_{\mu_i})) = \bigcap_{i \in \mathfrak{I}} F_0(\mathbf{s}_{\Phi(\mu_i)}) = F_0\left(\sum_{i \in \mathfrak{I}} \mathbf{s}_{\Phi(\mu_i)}\right) = F_0(\check{\Phi}(p)).$$

Moreover, the map $F_0: p \mapsto F_0(p)$ is a bijection from $\mathcal{P}(M_k)$ onto $\mathcal{F}(M_k)$ for k = 1, 2 ([4, Theorem 3.35]). These show that $\check{\Phi}(p)$ is independent of the choice of $\{\mu_i\}_{i\in\mathfrak{I}}$, and that Φ induces a map $\Phi^{\mathfrak{F}}: \mathcal{F}(M_1) \to \mathcal{F}(M_2)$.

In the same way, Φ^{-1} induces a map from $\mathcal{F}(M_2)$ to $\mathcal{F}(M_1)$ which is clearly the inverse of $\Phi^{\mathcal{F}}$. Therefore, $\Phi^{\mathcal{F}}$ is a bijection, and the bijectivity of the map $\check{\Phi} : \mathcal{P}(M_1) \to \mathcal{P}(M_2)$ follows from the bijectivity of F_0 .

Suppose now that $p, q \in \mathcal{P}(M_1)$ satisfying pq = 0. Then for any $p', q' \in \mathcal{P}_{\sigma}(M_1)$ with $p' \leq p$ and $q' \leq q$, one has p'q' = 0. Hence, from the hypothesis concerning Φ and the definition of $\check{\Phi}$, we conclude that $\check{\Phi}(p)\check{\Phi}(q) = 0$. Again, by considering Φ^{-1} , we know that $\check{\Phi}$ is an orthoisomorphism.

- (b) This follows from the Corollary in [10, p. 83].
- (c) By second equality in (3.5),

$$\mu(\Theta(\Phi(\mathbf{s}_{\mu}))) = \mu(\Theta(\mathbf{s}_{\Phi(\mu)})) = \Theta^*(\mu)(\mathbf{s}_{\Theta^*(\mu)})) = 1$$

Thus, $\mathbf{s}_{\mu} \leq \Theta(\check{\Phi}(\mathbf{s}_{\mu})) = \Theta(\mathbf{s}_{\Theta^{*}(\mu)})$. Conversely, as $\mu(\mathbf{s}_{\mu}) = 1$, one has $\Theta^{*}(\mu)(\Theta^{-1}(\mathbf{s}_{\mu})) = 1$, which means that $\mathbf{s}_{\Theta^{*}(\mu)} \leq \Theta^{-1}(\mathbf{s}_{\mu})$. These give

(3.7)
$$\mathbf{s}_{\mu} = \Theta(\check{\Phi}(\mathbf{s}_{\mu})) = \Theta(\mathbf{s}_{\Theta^{*}(\mu)}),$$

and hence $\Phi(\nu)(\mathbf{s}_{\Phi(\mu)}) = \nu(\mathbf{s}_{\mu})$. On the other hand, due to the construction of $\check{\Phi}$ in the argument for part (a), Equality (3.7) also produces the second conclusion.

Theorem 3.2. Let M_1 and M_2 be von Neumann algebras, and let $\Phi : \mathfrak{S}(M_1) \to \mathfrak{S}(M_2)$ be a bijection.

(a) If Φ is biorthogonality preserving, then M_1 and M_2 are Jordan *-isomorphic.

(b) There is a Jordan *-isomorphism $\Theta: M_2 \to M_1$ satisfying $\Phi = \Theta_*|_{\mathfrak{S}(M_1)}$ if and only if Φ preserves the "asymmetric transition probability" P_0 , i.e.,

$$P_0(\Phi(\mu), \Phi(\nu)) = P_0(\mu, \nu) \qquad (\mu, \nu \in \mathfrak{S}(M_1)).$$

Proof: (a) This follows directly from Lemma 3.1(a) and Proposition 2.2(a).

(b) Suppose that such a Jordan *-isomorphism Θ exists. Then Lemma 3.1(c) tells us that $P_0(\Phi(\nu), \Phi(\mu)) = P_0(\nu, \mu) \ (\mu, \nu \in \mathfrak{S}(M_1)).$

For the converse implication, we first note that because $\mu(\mathbf{s}_{\nu}) = \Phi(\mu)(\mathbf{s}_{\Phi(\nu)}) \ (\mu, \nu \in \mathfrak{S}(M))$, the map Φ is biorthogonality preserving (see (3.2)). Consider $\check{\Phi} : \mathcal{P}(M_1) \to \mathfrak{S}(M)$

 $\mathcal{P}(M_2)$ to be the map as in Lemma 3.1. Let M_1^0 be the real linear span of $\mathcal{P}(M_1)$ in M_1 . We want to extend $\check{\Phi}$ to M_1^0 by setting

$$\check{\Phi}(x) := \sum_{k=1}^{n} r_k \check{\Phi}(p_k),$$

when $x = \sum_{k=1}^{n} r_k p_k$ for some $n \in \mathbb{N}$, $r_1, ..., r_n \in \mathbb{R}$ and $p_1, ..., p_n \in \mathcal{P}(M_1)$. To show that this extension is well-defined, let us consider $p_k = \sum_{i \in \mathfrak{I}_k} \mathbf{s}_{\mu_{k,i}}$ to be a decomposition as in (3.1) (k = 1, ..., n). By the construction of $\check{\Phi}$ in the proof of Lemma 3.1(a), for any $\mu \in \mathfrak{S}(M_1)$,

$$\Phi(\mu) \Big(\sum_{k=1}^{n} r_k \check{\Phi}(p_k) \Big) = \sum_{k=1}^{n} r_k \sum_{i \in \mathfrak{I}_k} \Phi(\mu) \big(\mathbf{s}_{\Phi(\mu_{k,i})} \big) = \sum_{k=1}^{n} r_k \sum_{i \in \mathfrak{I}_k} \mu \big(\mathbf{s}_{\mu_{k,i}} \big) = \mu(x).$$

Thus, the surjectivity of Φ implies that $\check{\Phi}(x)$ is independent of the choices of $r_1, ..., r_n$ nor $p_1, ..., p_n$. Obviously, $\check{\Phi}$ is a linear map on M_1^0 satisfying

(3.8)
$$\nu(\check{\Phi}(x)) = \Phi^{-1}(\nu)(x) \qquad (x \in M_1^0; \nu \in \mathfrak{S}(M_2)).$$

This implies $\|\check{\Phi}(x)\| = \|x\|$ $(x \in M_1^0)$, and $\check{\Phi}$ extends to a linear isometry from M_1^{sa} onto M_2^{sa} . By Proposition 2.2(b), this extension, and hence its inverse $\Theta : M_2^{\text{sa}} \to M_1^{\text{sa}}$ is a Jordan isomorphism. Furthermore, Relation (3.8) tells us that $\Phi = \Theta_*|_{\mathfrak{S}(M_2)}$.

Note that one may use, for example, [15, Theorem 2] instead of Proposition 2.2(b) to conclude the proof of part (b) above. However, the current proof enables us to get an elementary proof of our next corollary (note that the results in [10] are only needed in Theorem 3.2(a)).

Through the argument of this corollary, one may also regard Theorem 3.2(b) as an extension of Wigner's theorem, and hence we get another proof for Wigner's theorem. Notice that the first part of the proof of this corollary is similar to that of [23, Theorem 1], but instead of showing the extension to be affine and employing [15, Corollary 5], we show that the extension preserves P_0 and use Theorem 3.2(b) to obtain the conclusion.

Corollary 3.3. (Wigner) Let \mathfrak{H}_1 and \mathfrak{H}_2 be two Hilbert spaces and let

$$\mathfrak{S}_{\mathbf{p}}(\mathcal{L}(\mathfrak{H}_k)) := \{ \omega_{\xi} : \xi \in S_{\mathfrak{H}_k} \} \qquad (k = 1, 2).$$

If $\Phi : \mathfrak{S}_{p}(\mathcal{L}(\mathfrak{H}_{1})) \to \mathfrak{S}_{p}(\mathcal{L}(\mathfrak{H}_{2}))$ is a bijection that preserves the transition probability, there is a Jordan *-isomorphism $\Theta : \mathcal{L}(\mathfrak{H}_{2}) \to \mathcal{L}(\mathfrak{H}_{1})$ with $\Phi = \Theta_{*}|_{\mathfrak{S}_{p}(\mathcal{L}(\mathfrak{H}_{1}))}$.

Proof: For any k = 1, 2 and $\xi \in S_{\mathfrak{H}_k}$, we know that $\mathbf{s}_{\omega_{\xi}}$ is the projection e_{ξ} from \mathfrak{H}_k onto $\mathbb{C} \cdot \xi$. Through diagonalisation of positive trace-class operators, we know that for each $\mu \in \mathfrak{S}(\mathcal{L}(\mathfrak{H}_1))$, there exist $n \in \mathbb{N} \cup \{\infty\}$, an orthonormal sequence $\{\xi_i\}_{i=1}^n$ in $S_{\mathfrak{H}_i}$ and a sequence $\{t_i\}_{i=1}^n$ in (0,1] with $\sum_{i=1}^n t_i = 1$ such that $\mu = \sum_{i=1}^n t_i \omega_{\xi_i}$ (converges in

norm). In this case, we propose to set

$$\bar{\Phi}(\mu) := \sum_{i=1}^{n} t_i \Phi(\omega_{\xi_i}).$$

For any finite orthonormal sequence $\{\zeta_j\}_{j=1}^N$ in $S_{\mathfrak{H}_1}$, one has, by the hypothesis,

(3.9)
$$\sum_{i=1}^{n} t_i \Phi(\omega_{\xi_i}) \left(\sum_{j=1}^{N} \mathbf{s}_{\Phi(\omega_{\zeta_j})} \right) = \sum_{j=1}^{N} \sum_{i=1}^{n} t_i \omega_{\xi_i}(\mathbf{s}_{\omega_{\zeta_j}}) = \mu \left(\sum_{j=1}^{N} \mathbf{s}_{\omega_{\zeta_j}} \right)$$

Since Φ is surjective, the above tells us that the value of $\sum_{i=1}^{n} t_i \Phi(\omega_{\xi_i})$ on any finite rank projection in $\mathcal{P}(\mathcal{L}(\mathfrak{H}_2))$ is independent of the decomposition $\mu = \sum_{i=1}^{n} t_i \omega_{\xi_i}$. Thus, $\bar{\Phi}(\mu)$ is well-defined.

On the other hand, Relation (3.9) also tells us that $\Phi : \mathfrak{S}(\mathcal{L}(\mathfrak{H}_1)) \to \mathfrak{S}(\mathcal{L}(\mathfrak{H}_2))$ is an injection, and the surjectivity of $\overline{\Phi}$ follows from the surjectivity of Φ . Furthermore, (3.9) implies that $\overline{\Phi}(\mu)(\mathbf{s}_{\Phi(\omega_{\eta})}) = \mu(\mathbf{s}_{\omega_{\eta}}) \ (\eta \in S_{\mathfrak{H}_1}).$

Let $\nu \in \mathfrak{S}(\mathcal{L}(\mathfrak{H}_1))$ and $\nu = \sum_{l=1}^m r_l \omega_{\eta_l}$ be a decomposition of ν similar to that of μ in the above. As $\{\mathbf{s}_{\omega_{\eta_l}}\}_{l=1}^m$ (and hence $\{\mathbf{s}_{\Phi(\omega_{\eta_l})}\}_{l=1}^m$) is an orthogonal sequence and $r_l > 0$ (for all l = 1, 2, ..., m), we have

$$\mathbf{s}_{\nu} = \sum_{l=1}^{m} \mathbf{s}_{\omega_{\eta_l}}$$
 as well as $\mathbf{s}_{\bar{\Phi}(\nu)} = \sum_{l=1}^{m} \mathbf{s}_{\Phi(\omega_{\eta_l})}$

(the convergences are in the weak*-topology). Hence,

$$\bar{\Phi}(\mu)(\mathbf{s}_{\bar{\Phi}(\nu)}) = \bar{\Phi}(\mu) \Big(\sum_{l=1}^m \mathbf{s}_{\Phi(\omega_{\eta_l})}\Big) = \sum_{l=1}^m \mu(\mathbf{s}_{\omega_{\eta_l}}) = \mu(\mathbf{s}_{\nu}).$$

Finally, Theorem 3.2(b) gives a Jordan *-isomorphism $\Theta : \mathcal{L}(\mathfrak{H}_1) \to \mathcal{L}(\mathfrak{H}_2)$ satisfying $\Phi = \Theta_* \mid_{\mathfrak{S}(\mathcal{L}(\mathfrak{H}_1))}$.

On the other hand, Theorem 3.2(a) can be regarded as an extension of a weak form of Uhlhorn's theorem for the normal state space of von Neumann algebras. In particular, we have the following application of it.

Theorem 3.4. Let M_1 and M_2 be von Neumann algebras. Then M_1 and M_2 are Jordan *isomorphic if there is a bijection $\Phi : \mathfrak{S}(M_1) \to \mathfrak{S}(M_2)$ satisfying any one of the following conditions:

(1) Φ preserves the usual metric d₁;

(2) Φ preserves pairs with zero Raggio transition probabilities, i.e. for any $\mu, \nu \in \mathfrak{S}(M_1)$,

$$P_R(\Phi(\mu), \Phi(\nu)) = 0$$
 if and only if $P_R(\mu, \nu) = 0$;

- (3) Φ preserves pairs with zero Uhlmann transition probabilities;
- (4) Φ preserves pairs with zero "asymmetric transition probabilities".

Proof: We claim that in each of the four cases, Φ is biorthogonality preserving, and thus Theorem 3.2(a) applies.

Indeed, the assertion for the case of Φ preserving the usual metric d_1 follows from the well-known fact that $\mathbf{s}_{\mu}\mathbf{s}_{\nu} = 0$ if and only if $\|\mu - \nu\| = 2$.

Suppose that Φ preserves pairs with zero Raggio transition probabilities. By (1.4) and (1.5), we know that $P_R(\mu,\nu) = 0$ if and only if $\|\xi_{\mu} - \xi_{\nu}\|^2 = 2$. On the other hand, it follows from [12, Lemma 2.10(2)] that $\|\xi_{\mu} - \xi_{\nu}\|^2 = 2$ if and only if $\|\mu - \nu\| = 2$, because $\|\xi_{\mu} - \xi_{\nu}\| \|\xi_{\mu} + \xi_{\nu}\| = \sqrt{4 - 4\langle\xi_{\mu},\xi_{\nu}\rangle^2}$. Thus, the assertion for the second case follows from that of the first case.

Finally, the assertions for the third (respectively, the fourth) case, follows from (1.6) (respectively, (3.2)) and the second case.

As seen in the above, for any $\mu, \nu \in \mathfrak{S}(M_1)$, one has

 $(3.10) \quad \mathbf{s}_{\mu}\mathbf{s}_{\nu} = 0 \quad \Longleftrightarrow \quad P_{U}(\mu,\nu) = 0 \quad \Longleftrightarrow \quad P_{R}(\mu,\nu) = 0 \quad \Longleftrightarrow \quad P_{0}(\mu,\nu) = 0.$

In particular, we obtained an alternative proof of [1, Lemma 1.8].

One may wonder if it is possible to get a stronger conclusion for Theorem 3.2(a) (and hence a stronger conclusion for Theorem 3.4) similar to that of Theorem 3.2(b). However, the following example shows that it is impossible even in the case when $M_1 = M_2 = \mathcal{L}(\ell^2)$.

Example 3.5. Let $M = \mathcal{L}(\mathfrak{H})$ with dim $\mathfrak{H} \geq 2$. Let \sim be an equivalence relation in $\mathfrak{S}(M)$ defined by

$$\mu \sim \nu$$
 if and only if $\mathbf{s}_{\mu} = \mathbf{s}_{\nu}$.

Let \mathfrak{C} be the set of equivalence classes of $\mathfrak{S}(M)$ under \sim . Suppose that ζ_1 and ζ_2 are two orthogonal elements in $S_{\mathfrak{H}}$, and $e_{\zeta_k} \in \mathfrak{P}(M)$ is the orthogonal projection onto $\mathbb{C} \cdot \zeta_k$ (k = 1, 2). For any $t \in (0, 1)$, if we set $\mu_t := t\omega_{\zeta_1} + (1 - t)\omega_{\zeta_2}$, then $\mathbf{s}_{\mu_t} = e_{\zeta_1} + e_{\zeta_2}$. Hence, $\{\mu_t : t \in (0, 1)\} \subseteq C_0$ for an element $C_0 \in \mathfrak{C}$. Consider any bijection $\Phi_0 : C_0 \to C_0$ with

$$\Phi_0(\mu_t) = (1 - t)\omega_{\zeta_1} + t\omega_{\zeta_2} \qquad (t \in (0, 1)),$$

and define a bijection $\Phi: \mathfrak{S}(M) \to \mathfrak{S}(M)$ by setting $\Phi|_{C_0} = \Phi_0$ as well as

$$\Phi(\mu) = \mu \qquad (\mu \in \mathfrak{S}(M) \setminus C_0).$$

From the definition of Φ , we know that $\mathbf{s}_{\Phi(\mu)} = \mathbf{s}_{\mu}$ ($\mu \in \mathfrak{S}(M)$), and Φ is biorthogonality preserving. However, since

$$\|\omega_{\zeta_1} - \mu_t\| = 2 - 2t$$
 and $\|\Phi(\omega_{\zeta_1}) - \Phi(\mu_t)\| = 2t$ $(t \in (0, 1)),$

one concludes that Φ cannot be induced by any continuous map from M_* to itself.

Nevertheless, in the case when the bijection Φ actually preserves the Raggio transition probability, we will see in Theorem 3.8 below that the conclusion as in Theorem 3.2(b) holds. In order to obtain this result, we need some more preparations. A normed space X is said to be *strictly convex* if for any $x, y \in X$, the condition ||x + y|| = ||x|| + ||y|| implies that x and y are linearly dependent. Clearly, any Hilbert space is strictly convex. Let us recall the following well-known fact in Banach spaces theory.

Lemma 3.6. Suppose that X_1 and X_2 are real Banach spaces such that X_2 is strictly convex. If K is a convex subset of X_1 and $f : K \to X_2$ is a metric preserving map, then f is automatically an affine map.

For the benefit of the readers, we sketch a proof here. In fact, in order to show

$$f(tx + (1-t)y) = tf(x) + (1-t)f(y)$$

for any $x \neq y$ in K and $t \in (0, 1)$, we may assume (by "shifting" K and f if necessary) that y = 0 and that f(0) = 0. In this case, we have

$$(3.11) ||f(x) - f(tx)|| = ||x - tx|| = (1 - t)||f(x)|| = ||f(x)|| - t||x|| = ||f(x)|| - ||f(tx)||,$$

and the strict convexity gives $f(x) - f(tx) \in \mathbb{R} \cdot f(tx)$. This, together with (3.11), establishes the required relation: f(tx) = tf(x).

Proposition 3.7. Let $(M_k, \mathfrak{H}_k, J_k, \mathfrak{P}_k)$ be a von Neumann algebra in its standard form (k = 1, 2). There are canonical bijective correspondences (through restrictions) amongst the following:

- the set $\mathfrak{I}_{\mathfrak{H}}$ of complex linear isometries from \mathfrak{H}_1 to \mathfrak{H}_2 sending \mathfrak{P}_1 onto \mathfrak{P}_2 ;
- the set $\mathbb{J}_{\mathfrak{P}}$ of metric preserving surjections from \mathfrak{P}_1 onto \mathfrak{P}_2 ;
- the set \mathfrak{I}_S of metric preserving surjections from $S_{\mathfrak{P}_1}$ onto $S_{\mathfrak{P}_2}$.

Proof: For every $\rho \in \mathfrak{I}_{\mathfrak{H}}$, one clearly has $\rho|_{S_{\mathfrak{P}_1}} \in \mathfrak{I}_S$. The assignment $\rho \mapsto \rho|_{S_{\mathfrak{P}_1}}$ defines an injection $R: \mathfrak{I}_{\mathfrak{H}} \to \mathfrak{I}_S$, because $S_{\mathfrak{P}_1}$ generates \mathfrak{H}_1 . Secondly, if $\chi \in \mathfrak{I}_S$, then

$$\langle \chi(\xi), \chi(\eta) \rangle = \langle \xi, \eta \rangle \in \mathbb{R}_+ \qquad (\xi, \eta \in S_{\mathfrak{P}_1}),$$

and hence the extension $\tilde{\chi} : t\xi \mapsto t\chi(\xi)$ $(\xi \in S_{\mathfrak{P}_1}, t \in \mathbb{R}_+)$ belongs to $\mathfrak{I}_{\mathfrak{P}}$. This gives an injection $E : \mathfrak{I}_S \to \mathfrak{I}_{\mathfrak{P}}$. Furthermore, as elements in $\mathfrak{I}_{\mathfrak{H}}$ are affine, the composition $E \circ R : \mathfrak{I}_{\mathfrak{H}} \to \mathfrak{I}_{\mathfrak{P}}$ coincides with the restriction map $\rho \mapsto \rho|_{\mathfrak{P}_1}$. Thus, it remains to show that $E \circ R$ is surjective.

Let us now consider $\varphi \in \mathcal{I}_{\mathfrak{P}}$. Since the only extreme point in \mathfrak{P}_k is the zero element, we know from Lemma 3.6 that $\varphi(0) = 0$. The metric preserving assumption now implies

(3.12)
$$\|\varphi(\xi)\| = \|\xi\|$$
 and $\langle\varphi(\xi),\varphi(\eta)\rangle = \langle\xi,\eta\rangle \in \mathbb{R}_+$ $(\xi,\eta\in\mathfrak{P}_1).$

For k = 1, 2, we denote by $\mathfrak{H}_k^{\mathrm{sa}}$ the real Hilbert space $\mathfrak{P}_k - \mathfrak{P}_k$. As \mathfrak{P}_k is a self-dual cone, if $\eta \in \mathfrak{H}_k^{\mathrm{sa}}$, there exist unique elements $\xi^+, \xi^- \in \mathfrak{P}_k$ with $\xi = \xi^+ - \xi^-$ and $\|\xi\|^2 = \|\xi^+\|^2 + \|\xi^-\|^2$ (see e.g. [13, Lemme I.1.2]).

Define $\tilde{\varphi}: \mathfrak{H}_1^{\mathrm{sa}} \to \mathfrak{H}_2^{\mathrm{sa}}$ by

$$\tilde{\varphi}(\xi) := \varphi(\xi^+) - \varphi(\xi^-) \qquad (\xi \in \mathfrak{H}_1^{\mathrm{sa}}).$$

For every $\xi, \eta \in \mathfrak{H}_1^{sa}$, one knows from (3.12) that

$$\|\tilde{\varphi}(\xi) - \tilde{\varphi}(\eta)\|^2 = \|\varphi(\xi^+) - \varphi(\xi^-) - \varphi(\eta^+) + \varphi(\eta^-)\|^2 = \|\xi^+ - \xi^- - \eta^+ + \eta^-\|^2,$$

which means that $\tilde{\varphi}$ preserves metric. Hence,

$$\|\tilde{\varphi}(\xi)\|^{2} = \|\xi\|^{2} = \|\xi^{+}\|^{2} + \|\xi^{-}\|^{2} = \|\varphi(\xi^{+})\|^{2} + \|\varphi(\xi^{-})\|^{2},$$

and the uniqueness of $\tilde{\varphi}(\xi)^{\pm}$ produces

(3.13)
$$\tilde{\varphi}(\xi)^{\pm} = \varphi(\xi^{\pm}) \qquad (\xi \in \mathfrak{H}_1^{\mathrm{sa}}).$$

If $\psi := \varphi^{-1} : \mathfrak{P}_2 \to \mathfrak{P}_1$ and $\tilde{\psi}$ is defined in the same way as $\tilde{\varphi}$, then, by a similar property as (3.13) for $\tilde{\psi}$, we obtain that, for each $\zeta \in \mathfrak{H}_1^{\mathrm{sa}}$,

$$\tilde{\varphi}(\tilde{\psi}(\zeta)) = \tilde{\varphi}(\tilde{\psi}(\zeta)^+ - \tilde{\psi}(\zeta)^-) = \varphi(\tilde{\psi}(\zeta)^+) - \varphi(\tilde{\psi}(\zeta)^-) = \varphi(\psi(\zeta^+)) - \varphi(\psi(\zeta^-)) = \zeta.$$

Consequently, $\tilde{\varphi}$ is surjective. It now follows from the Mazur-Ulam theorem that $\tilde{\varphi}$ is a linear isometry from \mathfrak{H}_1^{sa} onto \mathfrak{H}_2^{sa} . Finally, the complexification, $\bar{\varphi}$, of $\tilde{\varphi}$ is an element in $\mathcal{J}_{\mathfrak{H}}$ (note that linear isometries preserve inner products) satisfying $\bar{\varphi}|_{\mathfrak{P}_1} = \varphi$.

Recall that a projection $p \in \mathcal{P}(M) \setminus \{0\}$ is said to be σ -finite if any family of non-zero orthogonal subprojections of p is countable. It is easy to check that $\mathcal{P}_{\sigma}(M)$ consists exactly of σ -finite projections and the sum of a finite number of orthogonal σ -finite projections is again σ -finite. We also recall that a von Neumann algebra is said to be σ -finite if its identity is a σ -finite projection.

Theorem 3.8. Let M_1 and M_2 be two von Neumann algebras, and let $\Phi : \mathfrak{S}(M_1) \to \mathfrak{S}(M_2)$ be a bijection. Then Φ preserves the Raggio transition probability if and only if one can find a (necessarily unique) Jordan *-isomorphism $\Theta : M_2 \to M_1$ satisfying $\Phi = \Theta_*|_{\mathfrak{S}(M_1)}$.

Proof: One direction of the equivalence follows from Proposition 2.3. For the opposite direction, we assume in the following that Φ preserves the Raggio transition probability.

Notice that because of Relation (3.10), the map Φ is biorthogonality preserving, and Lemma 3.1 gives an orthoisomorphism $\check{\Phi} : \mathcal{P}(M_1) \to \mathcal{P}(M_2)$. Moreover, by Relations (1.4) as well as (1.5), the map $\varphi : S_{\mathfrak{P}_1} \to S_{\mathfrak{P}_2}$ given by

$$\varphi(\xi_{\mu}) := \xi_{\Phi(\mu)} \qquad (\mu \in \mathfrak{S}(M_1))$$

is a metric preserving surjection, and Proposition 3.7 tells us that it extends to a complex linear isometry $\bar{\varphi} : \mathfrak{H}_1 \to \mathfrak{H}_2$ satisfying $\bar{\varphi}(\mathfrak{P}_1) = \mathfrak{P}_2$.

By considering finite sums of elements in $\mathcal{P}_{\sigma}(M_1)$, one obtains, through (3.1), an increasing net $\{e_i\}_{i\in\mathfrak{I}}$ of σ -finite projections such that $e_i \uparrow 1$. Let us put $f_i := \check{\Phi}(e_i)$ $(i \in \mathfrak{I})$. Then all f_i are σ -finite and $f_i \uparrow 1$ (because $\check{\Phi}$ is an orthoisomorphism).

By Corollary 2.5 and Lemma 2.6 of [12], the standard form for $e_i M_1 e_i$ is

$$(e_i M_1 e_i, e_i e_i^{\mathsf{t}} \mathfrak{H}_1, e_i e_i^{\mathsf{t}} \mathfrak{P}_1, e_i e_i^{\mathsf{t}} J_1 e_i e_i^{\mathsf{t}})$$

(observe that $e_i^{\mathbf{t}} x e_i^{\mathbf{t}} \eta = x e_i^{\mathbf{t}} \eta = x \eta$, whenever $x \in e_i M_1 e_i, \eta \in e_i e_i^{\mathbf{t}} \mathfrak{H}_1$). In a similar way, $(f_i M_2 f_i, f_i f_i^{\mathbf{t}} \mathfrak{H}_2, f_i f_i^{\mathbf{t}} \mathfrak{H}_2, f_i f_i^{\mathbf{t}} \mathfrak{H}_2 f_i f_i^{\mathbf{t}})$ is the standard from of $f_i M_2 f_i$.

We identify, as usual,

$$\mathfrak{S}(e_i M_1 e_i) \cong e_i \mathfrak{S}(M_1) e_i \cap \mathfrak{S}(M_1) = F_0(1 - e_i)$$

and $\mathfrak{S}(f_i M_2 f_i) \cong F_0(1-f_i)$ in the canonical ways. From this, the map Φ induces, through Lemma 3.1(a), a bijection $\Phi_i : \mathfrak{S}(e_i M_1 e_i) \to \mathfrak{S}(f_i M_2 f_i)$. For each $\mu \in \mathfrak{S}(e_i M_1 e_i)$, let $\xi^i_{\mu} \in S_{e_i e_i^{\dagger} \mathfrak{P}_1}$ be the element with $\mu(x) = \langle x \xi^i_{\mu}, \xi^i_{\mu} \rangle$ ($x \in e_i M_1 e_i$). Then

$$\mu(y) = \mu(e_i y e_i) = \langle y \xi^i_\mu, \xi^i_\mu \rangle \qquad (y \in M_1),$$

and the uniqueness of the element ξ_{μ} in \mathfrak{P}_1 satisfying (1.3) implies that $\xi^i_{\mu} = \xi_{\mu}$. Hence, if $\varphi_i : S_{e_i e_i^{t} \mathfrak{P}_1} \to S_{f_i f_i^{t} \mathfrak{P}_2}$ is the bijection defined by $\varphi_i(\xi^i_{\mu}) := \xi^i_{\Phi_i(\mu)}$ ($\mu \in \mathfrak{S}(e_i M_1 e_i)$), we have $\varphi_i = \varphi|_{S_{e_i e_i^{t} \mathfrak{P}_1}}$.

The above shows that $\psi_i := \bar{\varphi}|_{e_i e_i^{\mathsf{t}} \mathfrak{H}_1}$ is a bijective isometry from $e_i e_i^{\mathsf{t}} \mathfrak{H}_1$ to $f_i f_i^{\mathsf{t}} \mathfrak{H}_2$ with $\psi_i(e_i e_i^{\mathsf{t}} \mathfrak{P}_1) = f_i f_i^{\mathsf{t}} \mathfrak{P}_2$. Since both $e_i M_1 e_i$ and $f_i M_2 f_i$ are σ -finite, [8, Théorème 3.3] gives a Jordan *-isomorphism $\Lambda_i : e_i M_1 e_i \to f_i M_2 f_i$ such that for every $x \in e_i M_1 e_i$ and $\xi \in S_{e_i e_i^{\mathsf{t}} \mathfrak{P}_1}$,

(3.14)
$$\Phi(\omega_{\xi})(\Lambda_{i}(x)) = \omega_{\varphi(\xi)}(\Lambda_{i}(x)) = \langle \Lambda_{i}(x)\varphi_{i}(\xi), \varphi_{i}(\xi) \rangle = \langle x\xi, \xi \rangle.$$

In particular, one has $\Phi_i = (\Lambda_i^{-1})^*|_{\mathfrak{S}(e_i M_1 e_i)}$.

As in the beginning of the proof, Φ_i is biorthogonality preserving and induces an orthoisomorphism $\check{\Phi}_i : \mathcal{P}(e_i M_1 e_i) \to \mathcal{P}(f_i M_2 f_i)$ satisfying Relation (3.5). It then follows from

$$\Phi(F_0(p) \cap \mathfrak{S}(e_i M_1 e_i)) = F_0(\check{\Phi}(p)) \cap \mathfrak{S}(f_i M_2 f_i) \qquad (p \in \mathfrak{P}(e_i M_1 e_i))$$

that $\check{\Phi}_i = \check{\Phi}|_{\mathcal{P}(e_i M_1 e_i)}$. Thus, Lemma 3.1(c) implies $\Lambda_i|_{\mathcal{P}(e_i M_1 e_i)} = \check{\Phi}|_{\mathcal{P}(e_i M_1 e_i)}$. From this, we know that whenever $i \leq j$, one has $\Lambda_j|_{\mathcal{P}(e_i M_1 e_i)} = \Lambda_i|_{\mathcal{P}(e_i M_1 e_i)}$, which ensures that

$$\Lambda_j|_{e_i M_1 e_i} = \Lambda_i$$

Set $M_1^e := \bigcup_{i \in \mathfrak{I}} e_i M_1 e_i$ and $M_2^f := \bigcup_{i \in \mathfrak{I}} f_i M_2 f_i$. The above allows us to define a Jordan *-isomorphism $\Lambda_0 : M_1^e \to M_2^f$ satisfying $\Lambda_0|_{e_i M_1 e_i} = \Lambda_i$, and (3.14) gives

(3.15)
$$\omega_{\varphi(\xi)}(\Lambda_0(x)) = \omega_{\xi}(x) \qquad (x \in M_1^e, \xi \in \mathfrak{P}_1)$$

because φ is an isometry and $\bigcup_{i\in\mathfrak{I}}e_ie_i^{\mathsf{t}}\mathfrak{P}_1$ is norm-dense in \mathfrak{P}_1 . We thus know from $\{\omega_{\varphi(\xi)}:\xi\in\mathfrak{P}_1\}=(M_2)^+_*$ that Λ_0 is weak*-continuous.

On the other hand, since $e_i y e_i \xrightarrow{w^*} y$ for any $y \in M_1$ (see e.g. Remark 2.1), M_1^e is weak*-dense in M_1 . Hence, Λ_0 extends to a weak*-continuous complex linear map $\Lambda: M_1 \to M_2$ such that $\Lambda(M_1^+) \subseteq M_2^+$, $\Lambda(1) = 1$ and, because of (3.15),

(3.16)
$$\Phi(\mu)(\Lambda(x)) = \mu(x) \qquad (x \in M_1, \mu \in \mathfrak{S}(M_1)).$$

Similarly, Φ^{-1} induces a positive linear map $\Upsilon : M_2 \to M_1$ satisfying the corresponding property as (3.16). Clearly, Υ is the inverse of Λ , and Λ is an order isomorphism. By [15, Corollary 5], Λ is a Jordan *-isomorphism, and $\Theta := \Lambda^{-1}$ is the required map.

The proof above can be shorten quite a bit if [8, Théorème 3.3] holds for the non- σ -finite case. However, this seems to be unknown. Note that even in the later work of [13, Theorem VII.1.1], which generalised [8, Théorème 3.3] to the case of JBW^* -algebras, the σ -finite assumption was still imposed.

The following corollary is a direct consequence of Theorems 3.4(b) and 3.8 as well as Proposition 2.3.

Corollary 3.9. If $\Phi : \mathfrak{S}(M_1) \to \mathfrak{S}(M_2)$ is a bijection preserving either the Raggio transition probability or the "asymmetric transition probability", then it preserves the Uhlmann transition probability as well.

It is natural to ask if the converse of the above holds. This lead to the following question.

Question 3.10. If $\Phi : \mathfrak{S}(M_1) \to \mathfrak{S}(M_2)$ is a bijection preserving the Uhlmann transition probability, can one find a Jordan *-isomorphism $\Theta : M_2 \to M_1$ satisfying $\Phi = \Theta_*|_{\mathfrak{S}(M_1)}$?

Let us end this section with another application of our main results. Here, $d_{\|\cdot\|}$ denotes the metric on \mathfrak{P} defined by the norm on \mathfrak{H} .

Corollary 3.11. Let M be a von Neumann algebra. Each one of the following metric spaces: $(\mathfrak{P}, d_{\parallel \cdot \parallel})$, $(\mathfrak{S}(M), d_B)$, $(\mathfrak{S}(M), d_1)$ and $(\mathfrak{S}(M), d_2)$ is a complete Jordan *-invariant for M.

Proof: The fact that $(\mathfrak{S}(M), d_1)$ is a complete Jordan *-invariant for M is already proved in Theorem 3.4. Moreover, it follows from Theorem 3.4 and Relation (1.4) (respectively, (1.2)) that $(\mathfrak{S}(M), d_2)$ (respectively, $(\mathfrak{S}(M), d_B)$) is a complete Jordan *-invariant. Consequently, $(\mathfrak{P}, d_{\parallel \cdot \parallel})$ is also a complete Jordan *-invariant because of Proposition 3.7. \Box

Furthermore, if dim $M \geq 2$, then the metric space $B_{\mathfrak{P}} := \{\xi \in \mathfrak{P} : \|\xi\| \leq 1\}$ (under the metric induced by the norm on \mathfrak{H}) is also a complete Jordan *-invariant for M. In fact, it is not hard to see that the set of extreme points of $B_{\mathfrak{P}}$ is $S_{\mathfrak{P}} \cup \{0\}$. Thus, using the fact that $S_{\mathfrak{P}_2}$ is not a singleton set, a continuity argument will verify that if $\phi : B_{\mathfrak{P}_1} \to B_{\mathfrak{P}_2}$ is a distance preserving bijection, then $\phi(0) = 0$. One can find the details of this, as well as its generalization to all non-commutative L_p -spaces $(p \in (1, \infty))$, in our further work on this subject ([17]).

4. Acknowledgement

The authors are supported by National Natural Science Foundation of China (11471168) and Taiwan MOST grant (102-2115-M-110-002-MY2).

References

- P.M. Alberti and G. Peltri, On Bures distance over standard form vN-algebras, preprint (arXiv:math/ 00081643v3).
- [2] P.M. Alberti and A. Uhlmann, On Bures distance and *-algebraic transition probability between inner derived positive linear forms over W*-algebras, Acta Appl. Math. 60 (2000), 1-37.
- [3] E.M. Alfsen and F.W. Shultz, State spaces of operator algebras, Birkhüser (2001).
- [4] E.M. Alfsen and F.W. Shultz, Geometry of state spaces of operator algebras, Birkhüser (2003).
- [5] L.G. Brown, Complements to Various Stone-Weierstrass Theorems for C^{*}-algebras and a Theorem of Shultz, Commun. Math. Phys. 143 (1992), 405-413.
- [6] D. Buchholz and E. Størmer, Superposition, transition probabilities and primitive observables in infinite quantum systems, to appear in Commun. Math. Phys. (arXiv:1411.2100v3)
- [7] M.A. Chebotar, W.F. Ke, P.H. Lee and N.C. Wong, Mappings preserving zero products, Stud. Math. 155 (2003), 77-94.
- [8] A. Connes, Caractérisation des espaces vectoriels ordonnés sous-jacents aux algèbres de von Neumann, Ann. Inst. Fourier 24 (1974), 121-155.
- [9] E. Christensen, Measures on projections and physical states, Commun. Math. Phys. 86 (1982), 529-538.
- [10] H.A. Dye, On the geometry of projections in certain operator algebras, Ann. Math. 61 (1955), 73-89.
- [11] Gy.P. Gehér, An elementary proof for the non-bijective version of Wigner's theorem, Physics Letters A 378 (2014) 2054-2057.
- [12] U. Haagerup, The standard form of von Neumann algebras, Math. Scand. **37** (1975), 271-283.
- [13] B. Iochum, Cônes autopolaires et algèbres de Jordan, Lecture Notes in Mathematics 1049, Springer-Verlag, Berlin, (1984).
- [14] R.V. Kadison, Isometries of operator algebras, Ann. Math. 54 (1951), 325-338.
- [15] R.V. Kadison, A generalized Schwarz inequality and algebraic invariants for operator algebras, Ann. Math. 564 (1952), 494-503.
- [16] L.E. Labuschagne, Pure state transformations induced by linear operators, Expo. Math. 23 (2005), 319-348.
- [17] C.W. Leung, C.K. Ng and N.C. Wong, Isometries on the cones of non-commutative L^p , in preparation.
- [18] L. Molnár, Generalization of Wigners unitary-antiunitary theorem for indefinite inner product spaces, Commun. Math. Phys. 210 (2000), 785-791.
- [19] L. Molnár, Orthogonality preserving transformations on indefinite inner product spaces: generalization of Uhlhorns version of Wigners theorem, J. Funct. Anal. 194 (2002), 248-262.
- [20] G.A. Raggio, Comparison of Uhlmann's transition probability with the one induced by the natural positive cone of von Neumann algebras in standard form, Lett. Math. Phys. 6 (1982), 233-236.
- [21] G.A. Raggio, Generalized transition probabilities and application, in Quantum Probability and Applications to the Quantum Theory of Irreversible Processes, Lect. Note in Math. 1055, Springer-Verlag (1984).
- [22] D. Sherman, Noncommutative L^p structure encodes exactly Jordan structure, J. Funct. Anal. **221** (2005), 150-166.
- [23] F.W. Shultz, Pure states as a dual object for C*-algebras, Commun. Math. Phys. 82 (1982), 497-509.

- [24] K. Schmüdgen, Transition probabilities of positive linear functionals on *-algebras, preprint (arXiv:1304.6322v1).
- [25] E. Størmer, Positive linear maps of operator algebras, Acta Math. 110 (1963), 233-278.
- [26] E. Størmer, On partially ordered vector spaces and their duals with applications to simplexes and C*-algebras, Proc. London Math. Soc. 18 (1968), 245-265.
- [27] A. Uhlmann, The "transition probability" in the state space of a *-algebra, Rep. Math. Phys. 9 (1976), 273-279.
- [28] A. Uhlmann, Transition probability (fidelity) and its relatives, Found. Phys. 41 (2011), 288-298.

(Chi-Wai Leung) DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, HONG KONG.

E-mail address: cwleung@math.cuhk.edu.hk

(Chi-Keung Ng) Chern Institute of Mathematics and LPMC, Nankai University, Tianjin 300071, China.

E-mail address: ckng@nankai.edu.cn

(Ngai-Ching Wong) DEPARTMENT OF APPLIED MATHEMATICS, NATIONAL SUN YAT-SEN UNIVER-SITY, KAOHSIUNG, 80424, TAIWAN.

E-mail address: wong@math.nsysu.edu.tw