

Estimation of Functionals of Sparse Covariance Matrices

JIANQING FAN^{*}, PHILIPPE RIGOLLET[†], AND WEICHEN WANG[‡]

Princeton University

Abstract. High-dimensional statistical tests often ignore correlations to gain simplicity and stability leading to null distributions that depend on functionals of correlation matrices such as their Frobenius norm and other ℓ_r norms. Motivated by the computation of critical values of such tests, we investigate the difficulty of estimation the functionals of sparse correlation matrices. Specifically, we show that simple plug-in procedures based on thresholded estimators of correlation matrices are sparsity-adaptive and minimax optimal over a large class of correlation matrices. Akin to previous results on functional estimation, the minimax rates exhibit an elbow phenomenon. Our results are further illustrated in simulated data as well as an empirical study of data arising in financial econometrics.

AMS 2000 subject classifications: Primary 62H12; secondary 62H15, 62C20, 62H25.

Key words and phrases: Covariance matrix, Functional estimation, High-dimensional testing, Minimax, Elbow effect.

1. INTRODUCTION

Covariance matrices are at the core of many statistical procedures such as principal component analysis or linear discriminant analysis. Moreover, not only do they arise as natural quantities to capture interactions between variables but, as we illustrate below, they often characterize the asymptotic variance of commonly used estimators. Following the original papers of Bickel and Levina [BL08a, BL08b], much work has focused on the inference of high-dimensional covariance matrices under sparsity [CL11, CRZ13, CY12, CZZ10, CZ12, Kar08, LF09, RWRY11] and other structural assumptions related to sparse principal component analysis [AW09, BR13b, BR13a, BJNP13, CMW13, CMW14, JL09, LV12, RLZ09, Ma13, OMH13, PJ12, FFL08, FLM11, FLM13, SSM09, VL12, ZHT06]. This area of research is very active and, as a result, this list of references is illustrative rather than comprehensive. This line of work can be split into two main themes: estimation and detection. The former is the main focus of the present paper. However, while most of the literature has focused on estimating

^{*}Supported in part by NSF Grants DMS-1206464 and DMS-1406266 and NIH grant R01-GM072611-9.

[†]Supported in part by NSF grants DMS-1317308, CAREER-DMS-1053987 and by the Howard B. Wentz Jr. Junior Faculty award.

[‡]Supported in part by NSF grant DMS-1206464.

the covariance matrix itself, under various performance measures, we depart from this line of work by focusing on functionals of the covariance matrix rather than the covariance matrix itself.

Estimation of functionals of unknown signals such as regression functions or densities is known to be different in nature from estimation of the signal itself. This problem has received most attention in nonparametric estimation, originally in the Gaussian white noise model [INK87, NK87, Fan91, EL96] (see also [Nem00] for a survey of results in the Gaussian white noise model) and later extended to density estimation [HM87, BR88] and various other models such as regression [DN90, CL05, CL06, Kle06] and inverse problems [But07, BM11]. Most of these papers study the estimation of quadratic functionals and, interestingly, exhibit an elbow in the rates of convergence: there exists a critical regularity parameter below which the rate of estimation is nonparametric and above which, it becomes parametric. As we will see below the phenomenon also arises when regularity is measured by sparsity.

Over the past decade, sparsity has become the prime measure of regularity, both for its flexibility and generality. In particular, smooth functions can be viewed as functions with a sparse expansion in an appropriate basis. At a high level, sparsity assumes that many of the unknown parameters are equal to zero or nearly so, so that the few nonzero parameters can be consistently estimated using a small number of observations relative to the apparent dimensionality of the problem. Moreover, sparsity acts not only as a regularity parameter that stabilizes statistical procedures but also as key feature for interpretability. Indeed, it is often the case that setting many parameters to zero simply corresponds to a simpler sub-model. The main idea is to let data select the correct sub-model. This is the case in particular for covariance matrix estimation where zeros in the matrix correspond to uncorrelated variables. Yet, while the value of sparsity for covariance matrix estimation has been well established, to the best of our knowledge, this paper provides the first analysis for the estimation of functionals of sparse covariance matrix. Indeed, the actual performance of many estimators critically depends on such functionals. Therefore accurate functional estimation leads to a better understanding the performance of many estimators and can ultimately serve as a guide to selecting the best estimator. Applications of our results are illustrated in Section 2.

Our work is not only motivated by real applications, but also by a natural extension of the theoretical analysis carried out in the sparse Gaussian sequence model [CL05]. In that paper, Cai and Low assume that the unknown parameter θ belong to an ℓ_q -ball, where $q > 0$ can be arbitrarily close to 0. Such balls are known to emulate sparsity and actually correspond to a more accurate notion of sparsity for signal θ that is encountered in applications (see e.g. [FR13]). They also show that a nonquadratic estimator can be fully efficient to estimate quadratic functionals. We extend some of these results to covariance matrix estimation. Such an extension is not trivial since, unlike the Gaussian sequence model, covariance matrix lies at high-dimensional manifolds and its estimation exhibits complicated dependencies in the structure of the noise.

We also compare our results for optimal rates of estimating matrix functionals with that of estimating matrix itself. Many methods have been proposed to estimate covariance matrix in different sense of sparsity using different techniques

including thresholding [BL08a], tapering [BL08b, CZZ10, CZ12] and penalized likelihood [LF09] to name only a few. These methods often lead to minimax optimal rates in various classes and under several metrics [CZZ10, CZ12, RT12]. However, the optimal rates of estimating matrix functionals has not yet been covered by much literature. Intuitively, it should have faster rates of convergence on estimating a matrix functional than itself since it is just a one-dimensional estimating problem and the estimating error cancel with each other when we sum those elements together. We will see this is indeed the case when we compare the minimax rates of estimating matrix functionals with those of estimating matrices.

The rest of the paper is organized as follows. We begin in Section 2 by two motivating examples of high-dimensional hypothesis testing problems: a two-sample testing problem of Gaussian means that arises in genomics and validating the efficiency of markets based on the Capital Asset Pricing Model (CAPM). Next, in Section 3, we introduce an estimator of the quadratic functional of interest that is based on the thresholding estimator introduced in [BL08a]. We also prove its optimality in a minimax sense over a large class of sparse covariance matrices. The study is further extended to estimating other measures of sparsity of covariance matrix. Finally, we study the numerical performance of our estimator in Section 5 on simulated experiments as well as in the framework of the two applications described in Section 2. All proofs are relegated to the appendix.

NOTATION: Let d be a positive integer. The space of $d \times d$ positive semi-definite matrices is denoted by \mathbf{S}_d^+ . For any two integers $c < d$, define $[c : d] = \{c, c + 1, \dots, d\}$ to be the sequence of contiguous integers between c and d , and we simply write $[d] = \{1, \dots, d\}$. I_d denotes the identity matrix of \mathbb{R}^d . Moreover, for any subset $S \subset [d]$, denote by $\mathbf{1}_S \in \{0, 1\}^d$ the column vector with j th coordinate equal to one iff $j \in S$. In particular, $\mathbf{1}_{[d]}$ denotes the d dimensional vector of all ones.

We denote by tr the trace operator on square matrices and by diag the linear operator that sets to 0 all the off diagonal elements of a square matrix. The Frobenius norm of a real matrix M is denoted by $\|M\|_F$ and is defined by $\|M\|_F = \sqrt{\text{tr}(M^\top M)}$. Note that $\|M\|_F$ is a the Hilbert-Schmidt norm associated with the inner product $\langle A, B \rangle = \text{tr}(A^\top B)$ defined on the space of real rectangular matrices of the same size. Moreover, $|A|$ denotes the determinant of a square matrix A . The variance of a random variable X is denote by $\text{var}(X)$.

In the proofs, we often employ C to denote a generic positive constant that may change from line to line.

2. TWO MOTIVATING EXAMPLES

In this section, we describe our main motivation for estimating quadratic functionals of a high-dimensional covariance matrix in the light of two applications to high-dimensional testing problems. The first one is a high-dimensional two-sample hypothesis testing with applications in Gene-set testing. The second example is about testing the validity of the *capital asset pricing model (CAPM)* from financial economics.

2.1 Two-sample Hypothesis Testing in high dimensions

In various statistical applications, in particular in genomics, the dimensionality of the problems is so large that statistical procedures involving inverse covariance

matrices are not viable due to its lack of stability both from a statistical and numerical point of view. This limitation can be well illustrated on a showcase example: two-sample hypothesis testing [BS96] in high dimensions.

Suppose that we observe two independent samples $X_1^{(1)}, \dots, X_{n_1}^{(1)} \in \mathbb{R}^p$ that are i.i.d. $\mathcal{N}(\mu_1, \Sigma_1)$ and $X_1^{(2)}, \dots, X_{n_2}^{(2)} \in \mathbb{R}^p$ that are i.i.d. $\mathcal{N}(\mu_2, \Sigma_2)$. The goal is to test $H_0 : \mu_1 = \mu_2$ vs $H_1 : \mu_1 \neq \mu_2$.

Assume first that $\Sigma_1 = \Sigma_2 = \Sigma$. In this case, the canonical test for this problem is based on Hotelling's T^2 statistic defined by

$$T^2 = \frac{n_1 n_2}{n} (\bar{X}^{(1)} - \bar{X}^{(2)})^\top \hat{\Sigma}^{-1} (\bar{X}^{(1)} - \bar{X}^{(2)})$$

where

$$\bar{X}^{(j)} = \frac{1}{n_j} \sum_{i=1}^{n_j} X_i^{(j)}, \quad j = 1, 2, \quad \hat{\Sigma} = \frac{1}{n} \sum_{j=1}^2 \sum_{i=1}^{n_j} (X_i^{(j)} - \bar{X}^{(j)})(X_i^{(j)} - \bar{X}^{(j)})^\top.$$

When dimensionality p is large, Bai and Saranadasa [BS96] showed that the test based on Hotelling's T^2 has low power and suggest the statistics M defined as

$$M = (\bar{X}^{(1)} - \bar{X}^{(2)})^\top (\bar{X}^{(1)} - \bar{X}^{(2)}) - \frac{n}{n_1 n_2} \text{tr}(\hat{\Sigma}).$$

This avoids the stability and invertability of the estimated sample covariance matrix $\hat{\Sigma}$ when p is large. This statistics leads to a provably more powerful test asymptotically [BS96]. It is not hard to see that the statistic M implements the Naive Bayes rule which ignores the dependence structure.

It may be shown [BS96, Appendix A.1] under mild moment and scaling conditions that the statistic M under the null hypothesis $H_0 : \mu_1 = \mu_2$ is asymptotically Gaussian:

$$Z = \frac{M}{\sqrt{\text{var}(M)}} \rightarrow \mathcal{N}(0, 1) \quad \text{as } n, p \rightarrow \infty, \quad \frac{n}{p} \rightarrow \gamma > 0, \quad \frac{n_1}{n_1 + n_2} \rightarrow \kappa \in (0, 1).$$

Moreover, it holds

$$\text{var}(M) = 2 \frac{n(n-1)}{(n_1 n_2)^2} \|\Sigma\|_{\mathbb{F}}^2 (1 + o(1)).$$

Clearly the asymptotic variance of M depends on the unknown covariance matrix Σ through its quadratic functional, and in order to compute the pivotal test statistic Z , Bai and Saranadasa suggest to estimate $\|\Sigma\|_{\mathbb{F}}^2$ by the quantity

$$B^2 = \frac{n^2}{(n+2)(n-1)} \left[\|\hat{\Sigma}\|_{\mathbb{F}}^2 - \frac{1}{n} (\text{tr}(\hat{\Sigma}))^2 \right].$$

They show that B^2 is a consistent estimator of $\|\Sigma\|_{\mathbb{F}}^2$ in the sense that $B^2 = (1 + o_P(1)) \|\Sigma\|_{\mathbb{F}}^2$. Clearly this solution does not leverage any sparsity assumption and may suffer from power deficiency if the matrix Σ is indeed sparse. Rather, if the covariance matrix Σ is believed to be sparse, one may prefer to use a thresholded estimator for Σ as in [BL08a] rather than the empirical covariance

matrix $\hat{\Sigma}$. In this case, the same analysis as [BS96] leads to the following estimator of $\|\Sigma\|_{\mathbb{F}}^2$:

$$\widehat{\|\Sigma\|_{\mathbb{F}}^2} = \sum_{i,j=1}^p \hat{\sigma}_{ij}^2 1\{|\hat{\sigma}_{ij}| > \tau\}.$$

where $\{\hat{\sigma}_{ij}, i, j \in [p]\}$ could be either the elements of empirical covariance matrix $\hat{\Sigma}$ or some other consistent estimator of σ_{ij} and $\tau > 0$ is a threshold parameter.

More recently, Chen and Qin [CQ10] took into account the case $\Sigma_1 \neq \Sigma_2$ and proposed a statistic which has good properties under weaker assumptions. Especially, only very weak assumption on the relationship between p and n is needed. They consider the statistic

$$T = \sum_{k=1}^2 \frac{1}{n_k(n_k - 1)} \sum_{i \neq j}^{n_k} (X_i^{(k)})^\top X_j^{(k)} - \frac{2}{n_1 n_2} \sum_{i=1}^{n_1} \sum_{j=1}^{n_2} (X_i^{(1)})^\top X_j^{(2)},$$

and showed that under H_0 ,

$$Z = \frac{T}{\sqrt{\text{var}(T)}} \rightarrow \mathcal{N}(0, 1) \quad \text{as } n, p \rightarrow \infty, \quad n^{-1/3}(\log p) \rightarrow 0, \quad \frac{n_1}{n_1 + n_2} \rightarrow \kappa \in (0, 1).$$

Moreover,

$$\text{var}(T) = \left\{ \frac{2}{n_1(n_1 - 1)} \|\Sigma_1\|_{\mathbb{F}}^2 + \frac{2}{n_2(n_2 - 1)} \|\Sigma_2\|_{\mathbb{F}}^2 + \frac{4}{n_1 n_2} \langle \Sigma_1, \Sigma_2 \rangle \right\} (1 + o(1)).$$

They used the following estimator of $\|\Sigma_i\|_{\mathbb{F}}^2$, $i = 1, 2$, and $\text{tr}(\Sigma_1 \Sigma_2)$:

$$\widehat{\|\Sigma_i\|_{\mathbb{F}}^2} = \frac{1}{n_i(n_i - 1)} \text{tr} \left\{ \sum_{j \neq k}^{n_i} (X_j^{(i)} - \bar{X}_{-(j,k)}^{(i)}) (X_j^{(i)})^\top (X_k^{(i)} - \bar{X}_{-(j,k)}^{(i)}) (X_k^{(i)})^\top \right\},$$

and

$$\widehat{\langle \Sigma_1, \Sigma_2 \rangle} = \frac{1}{n_1 n_2} \text{tr} \left\{ \sum_{l=1}^{n_1} \sum_{k=1}^{n_2} (X_l^{(1)} - \bar{X}_{-l}^{(1)}) (X_l^{(1)})^\top (X_k^{(2)} - \bar{X}_{-k}^{(2)}) (X_k^{(2)})^\top \right\},$$

where $\bar{X}_{-(j,k)}^{(i)}$ denotes the i th sample mean when both $X_j^{(i)}$ and $X_k^{(i)}$ are excluded, and $\bar{X}_{-l}^{(i)}$ denotes the i th sample mean when $X_l^{(i)}$ is excluded.

Again, these estimators do not leverage any sparsity assumption. Akin to the above reasoning, we can also estimate $\|\Sigma_i\|_{\mathbb{F}}^2$, $i = 1, 2$ and $\langle \Sigma_1, \Sigma_2 \rangle$ using thresholding. It is not hard to derive a theory for estimating quadratic functionals involving two covariance matrices but the details of this procedure are beyond the scope of the present paper.

To conclude this section, observe that both approaches in [BS96] and [CQ10] circumvent the difficulty of estimating the inverse covariance matrix by simply taking it equal to the identity. Instead, a hybrid approach would consist of estimating Σ_i^{-1} by D_i^{-1} , where $\hat{D}_i = \text{diag}(\hat{\Sigma}_i)$, $i = 1, 2$. This essentially corresponds to normalizing the empirical variance of each coordinate to 1 and put variables

into the same scale. To illustrate our point, consider for simplicity the case where $\Sigma_1 = \Sigma_2 = \Sigma$ and let

$$(2.1) \quad \tilde{T} = \sum_{i=1}^2 \frac{1}{n_i(n_i - 1)} \sum_{j \neq k}^{n_i} (X_j^{(i)})^\top \hat{D}^{-1} X_k^{(i)} - \frac{2}{n_1 n_2} \sum_{i=1}^{n_1} \sum_{j=1}^{n_2} (X_i^{(1)})^\top \hat{D}^{-1} X_j^{(2)},$$

where $\hat{D} = \text{diag}(\hat{\Sigma})$. Then under H_0 , using the same analysis as [BS96, CQ10], it is not hard to prove that \tilde{T} is asymptotically Gaussian and that under similar conditions as above,

$$\text{var}(\tilde{T}) = \frac{2n^2}{(n_1 n_2)^2} \|R\|_{\mathbb{F}}^2 (1 + o(1)),$$

where $R = \text{diag}(\Sigma)^{-1/2} \Sigma \text{diag}(\Sigma)^{-1/2}$ is the correlation matrix associated to Σ . Therefore, in order to estimate the distribution of \tilde{T} under the null hypothesis, and thus critical values, it suffices to estimate the quadratic functional $\|R\|_{\mathbb{F}}^2$.

2.2 Testing High Dimensional CAPM Model

The capital asset pricing model (CAPM) is a simple financial model that postulates how individual asset returns are related to the market risks. Specifically, the individual excessive return $Y_t^{(i)}$ of asset $i \in [N]$ over the riskfree rate at time $t \in [T]$ can be expressed as an affine function of the vector of K risk factors $f_t \in \mathbb{R}^K$, according to the following regression model

$$(2.2) \quad Y_t^{(i)} = \alpha_i + \beta_i^\top f_t + \epsilon_t^{(i)},$$

where we assume that for any $t \in [T]$, the excessive returns of K risk factors $f_t \in \mathbb{R}^K$ are observed. The case $K = 1$ with f_t being the excessive return of the market portfolio corresponds to the original CAPM [Sha64, Lin65, Mos66]. It is nowadays more common to employ the Fama-French three-factor model (see [FF93] for their definition) for the US equity market, corresponding to $K = 3$.

For simplicity, let us rewrite the model (2.2) in the following vectorial form

$$Y_t = \alpha + B f_t + \epsilon_t, \quad t \in [T]$$

where $Y_t = (Y_t^{(1)}, \dots, Y_t^{(N)})^\top$, $\alpha = (\alpha_1, \dots, \alpha_N)^\top$, $B = (\beta_1, \dots, \beta_N)^\top$ and $\epsilon_t = (\epsilon_t^{(1)}, \dots, \epsilon_t^{(N)})^\top$. Assume further that $(\epsilon_t)_{t \in [T]}$ is a stationary random process with expectation $\mathbb{E}[\epsilon_t] = 0$ and covariance matrix $\mathbb{E}[\epsilon_t \epsilon_t^\top] = \Sigma$, unknown. Under this model, the multi-factor pricing model, which includes CAPM as a specific example, postulates that $\alpha = 0$, namely, all returns are fully compensated by their risks: No extra returns are possible and the market is efficient. This leads us to naturally consider the following hypothesis testing problem $H_0 : \alpha = 0$ vs $H_1 : \alpha \neq 0$.

Traditionally, the number of portfolios N is assumed to be smaller than the number T of observations and, in this case, the GRS [GRS89] is commonly employed. However, the number of portfolios in the market is often very large and the assumption $N < T$ is enforced by grouping individual stocks into testing portfolios, leading to a potential loss of information and problems of data-snooping and endogeneity. Recently, Pesaran and Yamagata [PY12] pointed out to this issue and provided a solution by ignoring the correlation in the Wald statistics.

Let $\hat{\alpha}_i \in \mathbb{R}, \hat{\epsilon}^{(i)} \in \mathbb{R}^T, i \in [N]$ denote respectively the estimated intercepts and residual vectors for each of the N linear regression problems (2.2) that are obtained by ordinary least squares. Write $\hat{\alpha} = (\hat{\alpha}_1, \dots, \hat{\alpha}_N)^\top$ and for any $t \in T$, define $\hat{\epsilon}_t = (\hat{\epsilon}_t^{(1)}, \dots, \hat{\epsilon}_t^{(N)})^\top$. Then, an unbiased estimator of $\Sigma = \text{var}(\epsilon_t)$ is given by the covariance matrices of residuals

$$\tilde{\Sigma} = \frac{1}{T - K - 1} \sum_{t=1}^T \hat{\epsilon}_t \hat{\epsilon}_t^\top.$$

Let $\hat{D} = \text{diag}(\tilde{\Sigma})$ and $M_F = I_T - F(F^\top F)^{-1}F^\top$ denotes the projection matrix onto the null space of F where $F = (f_1, \dots, f_T)^\top$. Define $W_d = (\mathbf{1}_{[T]}^\top M_F \mathbf{1}_{[T]}) \hat{\alpha}^\top \hat{D}^{-1} \hat{\alpha}$ the Wald-type of test statistics with correlation ignored, whose normalized version is given by

$$(2.3) \quad J_\alpha = \frac{W_d - \mathbb{E}(W_d)}{\sqrt{\text{var}(W_d)}}.$$

Under some conditions, it can be shown [PY12] that $J_\alpha \rightarrow \mathcal{N}(0, 1)$ as $N \rightarrow \infty$ under the null hypothesis. Moreover, if the $\epsilon_t^{(i)}$ are i.i.d. Gaussian, it holds that

$$\mathbb{E}(W_d) = \frac{\nu N}{\nu - 2},$$

and

$$\text{var}(W_d) = \frac{2N(\nu - 1)}{\nu - 4} \left(\frac{\nu}{\nu - 2} \right)^2 [1 + (N - 1)\bar{\rho}^2 + O(\nu^{-1/2})],$$

where $\nu = T - K - 1$ is the degrees of freedom and

$$\bar{\rho}^2 = \frac{2}{N(N - 1)} \sum_{i=2}^N \sum_{j=1}^{i-1} \rho_{ij}^2,$$

where $\rho = D^{-1/2} \Sigma D^{-1/2}$ with $D = \text{diag}(\Sigma)$ is the correlation matrix of the stationary process $(\epsilon_t)_{t \in [T]}$. The authors go on to propose an estimator of the quadratic functional $\bar{\rho}^2$ by replacing the correlation coefficients ρ_{ij} in the above expression by $\hat{\rho}_{i,j} \mathbb{1}(|\hat{\rho}_{i,j}| > \tau)$ where $(\hat{\rho}_{i,j})_{i,j \in [N]} = \hat{D}^{-1/2} \tilde{\Sigma} \hat{D}^{-1/2}$ and $\tau > 0$ is a threshold parameter. However, they did not provide any analysis of this method, nor any guidance to chose τ .

3. OPTIMAL ESTIMATION OF QUADRATIC FUNCTIONALS

In the previous section, we have described rather general questions involving the estimation of quadratic functions of covariance or correlation matrices. We begin by observing that consistent estimation of $\|\Sigma\|_{\mathbb{F}}^2$ is impossible unless $p = o(n)$. This precludes in particular the high-dimensional framework that motivates our study. In fact, for non-degenerates case in which $\min_j \sigma_{jj} > c$, the functional $\|\Sigma\|_{\mathbb{F}}^2 \geq c^2 p$, which diverges rapidly.

Our goal is to estimate the Frobenius norm $\|\Sigma\|_{\mathbb{F}}^2$ of a sparse $p \times p$ covariance matrix Σ using n i.i.d. observations $X_1, \dots, X_n \sim \mathcal{N}(0, \Sigma)$. Observe that $\|\Sigma\|_{\mathbb{F}}^2$ can be decomposed as $\|\Sigma\|_{\mathbb{F}}^2 = Q(\Sigma) + D(\Sigma)$ where $Q(\Sigma) = \sum_{i \neq j} \sigma_{ij}^2$ corresponds to the off-diagonal elements and $D(\Sigma) = \sum_j \sigma_{jj}^2$ corresponds to the diagonal

elements. The following theorem, implies that even if $\Sigma = \text{diag}(\Sigma)$ is diagonal, the quadratic functional $\|\Sigma\|_{\mathbb{F}}^2$ cannot be estimated consistently if $p \geq n$. This makes sense intuitively as the diagonal of Σ consists of p unknown parameters while we have only n observations. While usually such a situation is overcome by assuming sparsity, such an assumption is unrealistic for variance parameters. Indeed, if $\sigma_{jj} = 0$ for some $j \in [p]$ then, all the X_i s will have their j th coordinate equal to zero and the dimension of the problem can be reduced trivially by simply removing this coordinate.

PROPOSITION 3.1. *Fix $n, p \geq 1$ and let*

$$\mathcal{D}_p = \{\Sigma \in \mathbf{S}_p^+ : \Sigma = \text{diag}(\Sigma), \Sigma_{ii} \leq 1\}$$

be the class of diagonal covariance matrices with diagonal elements bounded by 1. Then there exists a universal constant $C > 0$ such that

$$\inf_{\hat{D}} \sup_{\Sigma \in \mathcal{D}_p} \mathbb{E}[\hat{D} - D(\Sigma)]^2 \geq C \frac{p}{n}.$$

In particular, it implies that

$$\inf_{\hat{F}} \sup_{\Sigma \in \mathcal{D}_p} \mathbb{E}[\hat{F} - \|\Sigma\|_{\mathbb{F}}^2]^2 \geq C \frac{p}{n},$$

where the infima are taken with over all real valued measurable functions of the observations.

The proof is postponed to the appendix, Section B.1.

To overcome the above limitation, we consider the following class of sparse covariance matrices (indeed correlation matrices). For any $q \in [0, 2)$, $R > 0$ let $\mathcal{F}_q(R)$ denote the set of $p \times p$ covariance matrices defined by

$$(3.1) \quad \mathcal{F}_q(R) = \left\{ \Sigma \in \mathbf{S}_p^+ : \sum_{i \neq j} |\sigma_{ij}|^q \leq R, \text{diag}(\Sigma) = I_p \right\}.$$

Note that for this class of functions, we assume that the variance along each coordinate is normalized to 1. This normalization is frequently obtained by sample estimates, as shown in the previous section. This simplified assumption is motivated also by Proposition 3.1 above which implies that $\|\Sigma\|_{\mathbb{F}}^2$ for general covariance matrix cannot be estimated accurately in the large p small n regime since sparsity assumptions on the diagonal elements are implausible. Note that the condition $\text{diag}(\Sigma) = I_p$ implies that diagonal elements $D(\Sigma)$ of matrices in $\mathcal{F}_q(R)$ can be estimated without error so that we could possibly achieve consistency even if the case of large p small n .

Matrices in $\mathcal{F}_q(R)$ have many small coefficients for small values of q and R . In particular, when $q = 0$, there are no more than R entries of non-vanishing correlations in each row. Following a major trend in the estimation of sparse covariance matrices [BL08a, BL08b, CL11, CY12, CZZ10, CZ12, Kar08, LF09] we employ a thresholding estimator of the covariance matrix as a running horse to

estimate the quadratic functionals. From the n i.i.d. observations $X_1, \dots, X_n \sim \mathcal{N}(0, \Sigma)$, we form the empirical covariance matrix $\hat{\Sigma}$ that is defined by

$$(3.2) \quad \hat{\Sigma} = \frac{1}{n} \sum_{k=1}^n X_k X_k^\top$$

with elements $\hat{\Sigma} = \{\hat{\sigma}_{ij}\}_{ij}$ and for any threshold $\tau > 0$, let $\tilde{\Sigma}_\tau = \{\tilde{\sigma}_{ij}\}_{ij}$ denote the thresholding estimator of Σ defined by $\tilde{\sigma}_{ij} = \hat{\sigma}_{ij} \mathbb{1}\{|\hat{\sigma}_{ij}| > \tau\}$ if $i \neq j$ and $\tilde{\sigma}_{ii} = \hat{\sigma}_{ii}$.

Next, we employ a simple plug-in estimator for $Q(\Sigma)$:

$$(3.3) \quad \widehat{Q(\Sigma)} = Q(\tilde{\Sigma}_\tau) = \sum_{i \neq j} \hat{\sigma}_{ij}^2 \mathbb{1}\{|\hat{\sigma}_{ij}| > \tau\},$$

Note that no value of the diagonal elements is used to estimate $Q(\Sigma)$.

In the rest of this section, we establish that $\widehat{Q(\Sigma)}$ is minimax adaptive over the scale $\{\mathcal{F}_q(R), q \in [0, 2), R > 0\}$. Interestingly, we will see that the minimax rate presents an elbow as often in quadratic functional estimation.

THEOREM 3.1. *Assume that $\gamma \log(p) < n$ for some constant $\gamma > 8$ and fix $C_0 \geq 4$. Consider the threshold*

$$\tau = 2C_0 \sqrt{\frac{\gamma \log p}{n}},$$

and assume that $\tau \leq 1$. Then, for any $q \in [0, 2), R > 0$, the plug-in estimator $Q(\tilde{\Sigma}_\tau)$ satisfies

$$\mathbb{E}[(Q(\tilde{\Sigma}_\tau) - Q(\Sigma))^2] \leq C_1 \psi_{n,p}(q, R) + C_2 p^{4-\gamma/2},$$

where

$$\psi_{n,p}(q, R) = \frac{R^2}{n} \vee R^2 \left(\frac{\log p}{n}\right)^{2-q},$$

and C_1, C_2 are positive constants depending on γ, C_0, q . T

The proof is postponed to the appendix, Section A.1

Note that the rates $\psi_{n,p}(q, R)$ present an elbow at $q = 1 - \log \log p / \log n$ as usually the case in functional estimation. We now argue that the rates $\psi_{n,p}(q, R)$ are optimal in a minimax sense for a wide range of settings. In particular, the elbow effect arising from the maximum in the definition of ψ is not an artifact. In the following theorem, we emphasize the dependence on Σ by using the notation \mathbb{E}_Σ for the expectation with respect to the distribution of the sample X_1, \dots, X_n , where $X_i \sim \mathcal{N}(0, \Sigma)$.

THEOREM 3.2. *Fix $q \in [0, 2), R > 0$ and assume $2 \log p < n$ and $R^2 < (p-1)n^{-q}/2$. Then, there exists a positive constant $C_3 > 0$ such that,*

$$\inf_{\hat{Q}} \sup_{\Sigma \in \mathcal{F}_q} \mathbb{E}_\Sigma[(\hat{Q} - Q(\Sigma))^2] \geq C_3 \phi_{n,p}(q, R),$$

where $\phi_{n,p}(q, R)$ is defined by

$$(3.4) \quad \phi_{n,p}(q, R) = \frac{R^2}{n} \vee \left\{ R^2 \left(\frac{\log \left(\frac{p-1}{R^2 n^q} + 1 \right)}{2n} \right)^{2-q} \wedge R^{4/q} \wedge 1 \right\}$$

and the infimum is taken over all measurable functions \hat{Q} of the sample X_1, \dots, X_n .

The proof is postponed to the appendix, subsection B.2. A few remarks are in order.

1. The additional term of order $p^{4-\gamma/2}$ in Theorem 3.1 can be made negligible by taking γ large enough. To show this tradeoff explicitly, we decided keep this term.
2. When $\delta < R^2 < p^\delta n^{-q}$ for some constant $\delta < 1$, a slightly stronger requirement than Theorem 3.2 trivially satisfied for constant R and $q = 0$, the lower bound there can be written as

$$(3.5) \quad \phi_{n,p}(q, R) = \frac{R^2}{n} \vee \left\{ R^2 \left(\frac{\log p}{n} \right)^{2-q} \wedge 1 \right\}.$$

Observe that the above lower bound matches the upper bound presented in Theorem 3.1 when $R^{\frac{2}{2-q}} \log p \leq n$. Arguably, this is the most interesting range as it characterizes rates of convergence (to zero) as opposed to rates of divergence, that may be of different nature (see, e.g., [Ver12]). In other words, the rates given in (3.5) are minimax adaptive with respect to n , R , p , and q . In our formulation, we allow $R = R_{n,p}$ to depend on other parameters of the problem. We choose here to keep the notation light.

3. The reason we choose correlation matrix class to present the elbow effect is just for simplicity. Actually, we can replace the constraint $\text{diag}(\Sigma) = I_p$ in the definition of $\mathcal{F}_q(R)$ by boundedness of diagonal elements of Σ . Then for estimating off-diagonal elements $Q(\Sigma)$, following exactly the same derivation, the same elbow phenomenon has been noticed. Meanwhile the optimal rate for estimating diagonal elements $D(\Sigma)$ is again of the order p/n . This optimal rate can be attained by the estimator:

$$(3.6) \quad \widehat{D(\Sigma)} = \frac{1}{n(n-1)} \sum_{i=1}^p \sum_{k \neq j} X_{k,i}^2 X_{j,i}^2.$$

We omitted the proof here. Thus if we do not have prior information about diagonal elements, we could still estimate optimally the quadratic functional of a covariance matrix by applying the thresholding method (3.3) for off-diagonal elements, together with (3.6) for diagonal elements.

4. The rate $\phi_{n,p}(q, R)$ presents the same elbow phenomenon at $q = 1$ observed in the estimation of functionals, starting with work of Bickel and Ritov [BR88] and Fan [Fan91]. Closer to the present setup is the the work of Cai and Low [CL05] who study the estimation of functionals of “sparse” sequences in the infinite Gaussian sequence model. There, a parameter $\alpha > 0$ controls the speed of decay of the unknown coefficients. Note that while smaller values q lead to sparser matrices Σ , no estimator can benefit further from sparsity below $q = 1$, unlike in the case of estimation of Σ . Again, this is inherent to estimating functionals.
5. The condition $R^2 < (p-1)n^{-q}/2$ corresponds to the high-dimensional regime and allows us to keep clean terms in the logarithm. Similar assumptions are made in related literature (see, e.g., [CZ12]).

4. EXTENSION TO NON-QUADRATIC FUNCTIONALS

Closely related to quadratic functional is the ℓ_r functional of covariance matrices, which is defined by

$$(4.1) \quad \ell_r(\Sigma) = \max_{i \leq p} \sum_{j \leq p} |\sigma_{ij}|^r.$$

It is often used to measure the sparsity of a covariance matrix. Note that $\ell_1(\Sigma)$ functional is indeed the ℓ_1 -norm of the covariance matrix Σ , whereas when $r = 2$, ℓ_r functional is the maximal row-wise quadratic functional. Thus the non-quadratic ℓ_r functional is just a natural extension of quadratic functional with an outside maximum, whose optimal estimation problem will be the main focus of this section.

We consider a class of matrix with row-wise sparsity structure as follows:

$$(4.2) \quad \mathcal{G}_q(R) = \left\{ \Sigma \in \mathbf{S}_p^+ : \max_{i \leq p} \sum_{j \leq p} |\sigma_{ij}|^q \leq R, \Sigma_{ii} \leq 1 \right\},$$

for $q \in [0, r)$ and $R > 0$ which can depend on n and p . Note that $\Sigma_{ii} \leq 1$ is not essential, we could replace it by boundedness of diagonal elements of Σ . Similar class of parametric spaces for covariance matrices has been considered by [BL08a] with $q = 1$. The space is also more general than weak ℓ_q ball considered by [CZ12].

We should also expect an elbow effect in the minimax lower bound, which is shown in the following theorem.

THEOREM 4.1. *Fix $q \in [0, r)$, $R > 0$ and assume that $2 \log p < n$ and $R^2 < (p-1)n^{-q}/2$. Then, there exists a positive constant $C_4 > 0$ such that,*

$$\inf_{\hat{L}} \sup_{\Sigma \in \mathcal{G}_q} \mathbb{E}_{\Sigma} [(\hat{L} - \ell_r(\Sigma))^2] \geq C_4 \tilde{\phi}_{n,p}(q, R),$$

where $\tilde{\phi}_{n,p}(q, R)$ is defined by

$$(4.3) \quad \tilde{\phi}_{n,p}(q, R) = R^2 \frac{\log p}{n} \vee \left\{ R^2 \left(\frac{\log \left(\frac{p-1}{R^2 n^q} + 1 \right)}{2n} \right)^{r-q} \wedge R^{2r/q} \wedge 1 \right\}$$

and the infimum is taken over all measurable functions \hat{L} of the sample X_1, \dots, X_n .

The proof is similar to that of Theorem 3.2 and is relegated to the appendix, subsection B.4. As in (3.5), when $\delta < R^2 < p^\delta n^{-q}$ for some $\delta < 1$, the lower bound in Theorem 4.1 can be written as

$$(4.4) \quad \tilde{\phi}_{n,p}(q, R) = R^2 \frac{\log p}{n} \vee \left\{ R^2 \left(\frac{\log p}{n} \right)^{r-q} \wedge 1 \right\}.$$

To establish the upper bound, we consider again a thresholding estimator. Naturally, we estimate ℓ_r functional of each single row, denoted by $\ell_r^{(i)}(\Sigma) = \sum_j |\sigma_{ij}|^r$, using the thresholding technique. Following the same notation as the previous section, the estimator is defined by

$$(4.5) \quad \widehat{\ell_r(\Sigma)} = \ell_r(\tilde{\Sigma}_\tau) = \max_i \ell_r^{(i)}(\tilde{\Sigma}_\tau) = \max_i \sum_{j \leq p} |\hat{\sigma}_{ij}|^r \mathbb{1}_{\{|\hat{\sigma}_{ij}| > \tau\}},$$

for a threshold $\tau > 0$. We will see in the next theorem that this estimator achieves the adaptive minimax optimal rate.

THEOREM 4.2. Assume that $\gamma \log(p) < n$ for some constant $\gamma > 8$ and fix $C_0 \geq 4$. Consider the threshold

$$\tau = 2C_0 \sqrt{\frac{\gamma \log p}{n}}$$

and assume that $\tau \leq 1$. Then, for any $q \in [0, r)$, $R > 0$, the plug-in estimator $\ell_r(\tilde{\Sigma}_\tau)$ satisfies

$$\mathbb{E}[(\ell_r(\tilde{\Sigma}_\tau) - \ell_r(\Sigma))^2] \leq C_5 \tilde{\psi}_{n,p}(q, R) + C_6 p^{4-\gamma/2},$$

where

$$\tilde{\psi}_{n,p}(q, R) = \begin{cases} \frac{R^2 \log p}{n} & \text{if } q < \max\{r-1, 0\}, \\ R^2 \left(\frac{\gamma \log p}{n}\right)^{r-q} & \text{if } q \geq \max\{r-1, 0\}. \end{cases}$$

and C_5 and C_6 are positive constants.

The proof is indeed a generalization to that of Theorem 3.1, yet is of more general interests. In proving Theorem 3.1, we used the decomposition

$$\hat{\sigma}_{ij}^2 - \sigma_{ij}^2 = 2\sigma_{ij}(\hat{\sigma}_{ij} - \sigma_{ij}) + (\hat{\sigma}_{ij} - \sigma_{ij})^2,$$

which is actually the Taylor expansion of $\hat{\sigma}_{i,j}^2$ at $\sigma_{i,j}$. Carefully scrutinizing the proof, we find that the first term has the parametric rate $O(R^2/n)$ whereas the second term contributes to the rate $O(R^2(\log p/n)^{2-q})$. This phenomenon can be generalized to the ℓ_r -functional. In the latter case, we will apply Taylor expansion of $|\hat{\sigma}_{ij}|^r$ at $|\sigma_{ij}|$, and the first-order term will contribute to the parametric rate of $O(R^2 \log p/n)$ while the second-order term has the rate $O(R^2(\log p/n)^{r-q})$. The elbow effect stems from the dominance of estimation errors of the first- and second-order terms of Taylor's expansion. We relegate the proof to the appendix, subsection A.2.

A few remarks should be mentioned:

1. The combination of the two theorems imply that the estimator $\ell_r(\tilde{\Sigma}_\tau)$ is minimax adaptive over the space $\{\mathcal{G}_q(R), q \in [0, r), R > 0\}$ under very mild conditions. The adaptive minimax optimal rate of convergence is given by (4.4). The term $p^{4-\gamma/2}$ can be made as small as possible by choosing large enough γ .
2. The ℓ_r functional involves the maxima of the row sums. Compared it with estimating the quadratic functional, we need to pay the price of an extra $\log p$ term in the parametric rate.
3. The rate $\tilde{\psi}_{n,p}(q, R)$ presents the elbow phenomenon at $q = r - 1$ if $r > 1$. So quadratic row-wise functional $\ell_2(\tilde{\Sigma}_\tau)$ bears the same elbow behavior as the quadratic functional $Q(\tilde{\Sigma}_\tau)$.

5. NUMERICAL EXPERIMENTS

Simulations are conducted in this section to evaluate the numerical performance of our plug-in estimator for quadratic functionals. Then the proposed method is applied to two high dimensional testing problems: simulated two-sample data and real financial equity market data.

5.1 Quadratic Functional Estimation

We first study the behavior of estimators $\widehat{Q}(\widehat{\Sigma}) + \widehat{D}(\widehat{\Sigma})$ for the total quadratic functional and $\widehat{Q}(\widehat{\Sigma}) = Q(\widehat{\Sigma}_\tau)$ for its off-diagonal part. To that end, four sparse covariance matrix structures were used in the simulations:

- (M1) auto-correlation AR(1) covariance matrix $\sigma_{ij} = 0.25^{|i-j|}$;
- (M2) banded correlation matrix with $\sigma_{ij} = 0.3$ if $|i - j| = 1$ and 0 otherwise;
- (M3) sparse matrix with a block, size $p/20$ by $p/20$, of correlation 0.3;
- (M4) identity matrix with maximal level of sparsity.

We chose $p = 500$ and let n vary from 30 to 100. For estimating the total quadratic functional, our proposed thresholding estimator, Bai and Saranadasa (BS, [BS96]) estimator, and Chen and Qin (CQ, [CQ10]) estimator were applied to each setting for repetition of 500 times. Their mean absolute estimation errors were reported in log scale (base 2) in Figure 1 with their standard deviations omitted here. BS and CQ cannot be directly used for off-diagonal quadratic functional estimation, so we deducted $\widehat{D}(\widehat{\Sigma})$ from both of them to serve as an estimator for only the off-diagonal part. The mean absolute estimation errors, compared with our proposed estimator $Q(\widehat{\Sigma}_\tau)$, are depicted in log scale (base 2) in Figure 2.




Fig 1: Performance of estimating $\|\Sigma\|_F^2$ using thresholded estimator $\widehat{Q} + \widehat{D}$ (blue dotted), CQ (orange solid) and BS (red dashed). The mean of absolute errors over 500 repetitions in log scale (base 2) versus the sample size were reported for matrix M1 (top left), M2 (top right), M3 (bottom left), M4 (bottom right).




Fig 2: Performance of estimating $Q(\Sigma)$ using thresholded estimator \hat{Q} (blue dotted), $CQ\text{-}\hat{D}$ (orange solid) and $BS\text{-}\hat{D}$ (red dashed). The mean of absolute errors over 500 repetitions in log scale (base 2) versus the sample size were reported for matrix M1 (top left), M2 (top right), M3 (bottom left), M4 (bottom right).

The four plots correspond to the aforementioned four covariance structures. We did not report the estimation error of directly using the naive plug-in which is an obvious disaster. In all the four cases, the BS (red dashed line) method does not perform well in the “large p small n ” regime. The method CQ (orange solid line) exhibits a relatively small estimation error in general, but it can still be improved using the thresholding method. As theory shows, the method CQ is ratio-consistent [CQ10], so our method (blue dotted line) is better only to a second order, which was captured by the small gap between blue and orange curves. When estimating only off-diagonal quadratic functionals (Figure 2), the advantage of the thresholding method is even sharper since the error caused by non-sparse diagonal elements has been eliminated. The improved performance comes from the prior knowledge of sparsity, thus our method works best for very sparse matrix, especially well for identity matrix as seen in Figure 1.

A practical question is how to choose a proper threshold, as this is important to the performance of the thresholding estimator. In the above simulations, we chose $\tau = C\sqrt{\log p/n}$ with constant C slightly different for the four cases but all close to 1.5. In the next two applications to hypothesis testing, we employ the cross validation to choose a proper thresholding. The procedure consists of the following steps:

- (1) The data is split into training data $D_S^{(v)}$ of sample size n_1 and testing data $D_{S^c}^{(v)}$ of sample size $n - n_1$ for m times, $v = 1, 2, \dots, m$;
- (2) The training data $D_S^{(v)}$ is used to construct the thresholding estimator $Q(\tilde{\Sigma}_\tau^{(v)})$ under a grid of thresholds while the testing data $D_{S^c}^{(v)}$ constructs the non-thresholded ratio-consistent estimator $\hat{Q}^{(v)}$ e.g. using CQ estimator of $\|\Sigma\|_F^2$;
- (3) The candidates of thresholds are $\tau_j = j\Delta\sqrt{\log(p)/n_1}$ for $j = 1, 2, \dots, J$ where J is chosen to be a reasonably large number, say 50, and Δ is such that $J\Delta\sqrt{\log(p)/n_1} \leq \hat{M} := \max_i \hat{\sigma}_{ii}$;
- (4) The final j^* is taken to be the minimizer of the following optimization problem:

$$\min_{j \in \{1, 2, \dots, J\}} \frac{1}{m} \sum_{v=1}^m \left| Q(\tilde{\Sigma}_{\tau_j}^{(v)}) - \hat{Q}^{(v)} \right|.$$

- (5) The final estimator $Q(\tilde{\Sigma}_{\tau_{j^*}})$ is obtained by applying threshold $\tau_{j^*} = j^*\Delta\sqrt{\log(p)/n}$ to the empirical covariance matrix of the n observations.

Bickel and Levina [BL08a] suggested to use $n_1 = n/\log n$ for estimating covariance matrices. This is consistent with our experience for estimating functionals when no prior knowledge about the covariance matrix structure is provided. We will apply this splitting rule in the later simulation studies on high dimensional hypothesis testing.

5.2 Application to High Dimensional Two-sample Testing

In this section, we apply the thresholding estimator of quadratic functionals to the high dimensional two-sample testing problem. Two groups of data are simulated from the Gaussian models:

$$X_{i,j} \sim \mathcal{N}(\mu_i, \Sigma), \quad \text{for } i = 1, 2 \text{ and } j = 1, \dots, n/2.$$

The dimensions considered for this problem are $(p, n) \in \{(500, 100), (1000, 150), (2000, 200)\}$. We choose the sparse covariance structure to be 2 by 2 block diagonal matrices with 250 of them having correlations 0.3 and the rest having correlations 0. So the off-diagonal quadratic functional is always 45, which does not increase with p in our setting. The mean vectors μ_1 and μ_2 are chosen as follows. Let $\mu_1 = 0$ and the percentage of $\mu_{1,k} = \mu_{2,k}$ to be in $\{0\%, 50\%, 95\%, 100\%\}$. The 100% proportion corresponds to the case where the two groups are identical, thus gives information about accuracy of the size of the tests. The 95% proportion represents the situation where the alternative hypotheses are sparse. For those k such that $\mu_{1,k} \neq \mu_{2,k}$, we simply chose the value of each $\mu_{2,k}$ equally. To make the power comparable among different configurations, we use a constant signal-to-noise ratio $\eta = \|\mu_1 - \mu_2\| / \sqrt{\text{tr}(\Sigma^2)} = 0.1$ across experiments.

Table 1 reports the empirical power and size of six testing methods based on 500 repetitions:

TABLE 1
Empirical testing power and size of 6 testing methods based on 500 simulations.

Prop. of Equalities	BS	newBS	CQ	newCQ	Bonf	BH
$p = 500, n = 100$						
0%	0.408	0.422	0.428	0.432	0.104	0.110
50%	0.396	0.422	0.418	0.428	0.110	0.116
95%	0.422	0.440	0.438	0.442	0.208	0.214
100% (size)	0.030	0.036	0.036	0.038	0.042	0.042
$p = 1000, n = 150$						
0%	0.696	0.710	0.718	0.718	0.082	0.086
50%	0.698	0.712	0.712	0.714	0.106	0.112
95%	0.702	0.716	0.718	0.722	0.308	0.328
100% (size)	0.040	0.044	0.048	0.046	0.050	0.050
$p = 2000, n = 200$						
0%	0.930	0.938	0.940	0.940	0.138	0.146
50%	0.918	0.922	0.924	0.928	0.104	0.106
95%	0.922	0.928	0.930	0.930	0.324	0.338
100% (size)	0.046	0.050	0.050	0.050	0.046	0.046

TABLE 2
Mean and SD of relative errors for estimating quadratic functionals (in percentage).

	$p = 500, n = 100$	$p = 1000, n = 150$	$p = 2000, n = 200$
BS	4.93 (2.48)	4.47 (1.56)	5.05 (1.10)
newBS	2.12 (1.43)	0.74 (0.56)	0.54 (0.40)
CQ	3.72 (1.97)	2.32 (1.24)	1.70 (0.91)
newCQ	2.77 (1.38)	1.27 (0.64)	0.62 (0.33)

- (BS) Bai and Saranadasa's original test.
- (newBS) Bai and Saranadasa's modified test where $\text{tr}(\Sigma^2)$ is estimated by thresholding the sample covariance matrix.
- (CQ) Chen and Qin's original test.
- (newCQ) Chen and Qin's modified test where $\text{tr}(\Sigma_i^2)$ and $\text{tr}(\Sigma_1 \Sigma_2)$ are estimated by thresholding their empirical counterparts.
- (Bonf) Bonferroni correction: This method regards the high-dimensional testing problem as p univariate testing problems. If there is a p-value that is less than $0.05/p$, the null hypothesis is rejected.
- (BH) Benjamini-Hochberg method. The method is similar to the Bonferroni correction, but employs the Benjamini-Hochberg method in decision making.

For estimating quadratic functionals, the cross-validation is employed using $n/\log(n)$ splitting rule. The first four methods are evaluated at the 5% significance level while Bonferroni correction and Benjamini-Hochberg correction are evaluated at 5% family-wise error rate or FDR. We also list the average relative estimation errors for the quadratic functionals of the first four methods in Table 2. Here the average is taken over four different proportions of equalities and the average for CQ and newCQ is also taken over errors in estimating $\text{tr}(\Sigma_1^2)$ and $\text{tr}(\Sigma_2^2)$.

Several comments are in order. First, the first four methods based on Wald-type of statistic with correlation ignored perform much better, in terms of the power, than the last two methods which combines individual tests. Even in the




Fig 3: P-values of testing $H_0 : \alpha = 0$ in the CAPM and Fama-French 3 factor models based on S & P 500 monthly returns from January 1995 to December 2012.

case that proportional of equalities is 95% where the individual difference is large for non-identical means, aggregating the signals together in Wald type of statistic still outperforms. However, in the case of 0% identical means, the power of Bonferroni or FDR method is extremely small, due to small individual differences. Second, the method newCQ, which combines CQ and thresholding estimator of the quadratic functional, has the highest power and performs the best among all methods. The corrected BS method also improves the performance by estimating the quadratic functionals better compared with original BS. CQ indeed is more powerful than BS as claimed by Chen and Qin [CQ10], but we can even improve the performance of those two methods more by leveraging the sparsity structure of covariance matrices.

5.3 Application to Testing Multifactor Pricing Model

In this section, we test the validity of the Capital Asset Pricing Model (CAPM) and Fama-French models using Pesaran and Yamagata's method (2.3) for the securities in the Standard & Poor 500 (S&P 500) index. Following the literature, we used 60 monthly stock returns to construct test statistics since monthly returns are nearly independent. The composition of index keeps changing annually, so we selected only 276 large stocks. The monthly returns (adjusted for dividend) between January 1990 and December 2012 are downloaded from the Wharton Research Data Services (WRDS) database. The time series on the risk-free rates and Fama-French three factors are obtained from Ken French's data library. If only the first factor i.e. the excessive return of the market portfolio is used, Fama-French model reduces to the CAPM model. We tested the null hypothesis $H_0 : \alpha = 0$ for both models. The p-values of the tests are depicted in Figure 3, which are computed based on running windows of previous 60 months.

The results suggest that market efficiency is time dependent and the Fama-French model are rejected less frequently than the CAPM. Before 1998, the evidence that $\alpha \neq 0$ is very strong. After 1998, the Fama-French 3-factor model holds most of the time except the period 2007-2009 that contains the financial

crisis. On the other hand, the CAPM is rejected for an extended period of time during this period.

APPENDIX A: PROOF OF UPPER BOUNDS

A.1 Proof of Theorem 3.1

Denote by $\mathcal{E} = \mathbb{E}[(Q(\tilde{\Sigma}_\tau) - Q(\Sigma))^2]$ the quantity of interest and first decompose it into three parts:

$$\mathcal{E} = \mathbb{E} \left[\left(\sum_{i \neq j} (\hat{\sigma}_{ij}^2 \mathbb{1}(|\hat{\sigma}_{ij}| > \tau) - \sigma_{ij}^2) \right)^2 \right] \leq 3(L_1 + L_2 + L_3),$$

where

$$\begin{aligned} L_1 &= \left(\sum_{i \neq j} \sigma_{ij}^2 \mathbb{1}(|\sigma_{ij}| \leq \tau) \right)^2, \\ L_2 &= \mathbb{E} \left[\left(\sum_{i \neq j} (\hat{\sigma}_{ij}^2 - \sigma_{ij}^2) \mathbb{1}(|\sigma_{ij}| > \tau) \right)^2 \right], \\ L_3 &= \mathbb{E} \left[\left(\sum_{i \neq j} \hat{\sigma}_{ij}^2 (\mathbb{1}(|\hat{\sigma}_{ij}| > \tau) - \mathbb{1}(|\sigma_{ij}| > \tau)) \right)^2 \right]. \end{aligned}$$

We bound each term separately, beginning with L_1 :

$$L_1 = \left(\sum_{i \neq j} |\sigma_{ij}|^q |\sigma_{ij}|^{2-q} \mathbb{1}(|\sigma_{ij}| \leq \tau) \right)^2 \leq R^2 \tau^{4-2q}.$$

Next, to control L_2 , observe that

$$\hat{\sigma}_{ij}^2 - \sigma_{ij}^2 = (\hat{\sigma}_{ij} - \sigma_{ij})^2 + 2\sigma_{ij}(\hat{\sigma}_{ij} - \sigma_{ij}).$$

It yields $L_2 \leq 2L_4 + 8L_5$, where

$$\begin{aligned} L_4 &= \mathbb{E} \left[\left(\sum_{i \neq j} (\hat{\sigma}_{ij} - \sigma_{ij})^2 \mathbb{1}(|\sigma_{ij}| > \tau) \right)^2 \right], \\ L_5 &= \mathbb{E} \left[\left(\sum_{i \neq j} \sigma_{ij} (\hat{\sigma}_{ij} - \sigma_{ij}) \mathbb{1}(|\sigma_{ij}| > \tau) \right)^2 \right]. \end{aligned}$$

Using respectively the Cauchy-Schwarz inequality, Lemma A.3 and the simple inequality

$$(A.1) \quad \sum_{i \neq j} \mathbb{1}(|\sigma_{ij}| > \tau) \leq \sum_{i \neq j} |\sigma_{ij}|^q \tau^{-q} \leq R\tau^{-q},$$

we get

$$\begin{aligned} L_4 &\leq \mathbb{E} \left[\left(\sum_{i \neq j} (\hat{\sigma}_{ij} - \sigma_{ij})^4 \mathbb{1}(|\sigma_{ij}| > \tau) \right) \left(\sum_{i \neq j} \mathbb{1}(|\sigma_{ij}| > \tau) \right) \right] \\ &\leq Cn^{-2} \left(\sum_{i \neq j} \mathbb{1}(|\sigma_{ij}| > \tau) \right)^2 \\ &\leq CR^2 \tau^{-2q} n^{-2}. \end{aligned}$$

Lemma A.3 together with the Jensen inequality yield $\mathbb{E}[(\hat{\sigma}_{ij} - \sigma_{ij})^2] \leq C/n$. Thus, by the Cauchy-Schwarz inequality, we get $\mathbb{E}[(\hat{\sigma}_{ij} - \sigma_{ij})(\hat{\sigma}_{st} - \sigma_{st})] = C/n$. This yields the following control on L_5 :

$$L_5 \leq \frac{C}{n} \left(\sum_{i \neq j} |\sigma_{ij}| \mathbb{1}(|\sigma_{ij}| > \tau) \right)^2.$$

The cases $q \leq 1$ and $q > 1$ are treated separately leading to the elbow effect in Theorem 3.1. If $q \leq 1$, since $|\sigma_{ij}| \leq 1$, we have

$$L_5 \leq \frac{C}{n} \left(\sum_{i \neq j} |\sigma_{ij}| \right)^2 \leq \frac{C}{n} \left(\sum_{i \neq j} |\sigma_{ij}|^q \right)^2 \leq CR^2 n^{-1}.$$

If $q > 1$, we get from Hölder's inequality that

$$L_5 \leq \frac{C}{n} \left(\sum_{i \neq j} |\sigma_{ij}|^q \right)^{2/q} \left(\sum_{i \neq j} \mathbb{1}(|\sigma_{ij}| > \tau) \right)^{2-2/q} \leq CR^2 \tau^{2-2q} n^{-1}.$$

Finally, we bound L_3 as $L_3 \leq 3(L_6 + L_7 + L_8)$, where

$$L_6 = \mathbb{E} \left[\left(\sum_{i \neq j} \hat{\sigma}_{ij}^2 \mathbb{1}(|\hat{\sigma}_{ij}| \leq \tau, |\sigma_{ij}| > \tau) \right)^2 \right],$$

$$L_7 = \mathbb{E} \left[\left(\sum_{i \neq j} \hat{\sigma}_{ij}^2 \mathbb{1}(|\hat{\sigma}_{ij}| > \tau, \frac{\tau}{2} < |\sigma_{ij}| \leq \tau) \right)^2 \right],$$

$$L_8 = \mathbb{E} \left[\left(\sum_{i \neq j} \hat{\sigma}_{ij}^2 \mathbb{1}(|\hat{\sigma}_{ij}| > \tau, |\sigma_{ij}| \leq \frac{\tau}{2}) \right)^2 \right].$$

We now proceed to controlling each term L_6, L_7 and L_8 as follows. Using (A.1), we get

$$L_6 \leq \tau^4 \left(\sum_{i \neq j} \mathbb{1}(|\sigma_{ij}| > \tau) \right)^2 \leq R^2 \tau^{4-2q}.$$

Next,

$$\begin{aligned} L_7 &\leq \mathbb{E} \left(\sum_{i \neq j} \hat{\sigma}_{ij}^2 \mathbb{1}(\frac{\tau}{2} < |\sigma_{ij}| \leq \tau) \right)^2 \\ &\leq 2\mathbb{E} \left(\sum_{i \neq j} (\hat{\sigma}_{ij}^2 - \sigma_{ij}^2) \mathbb{1}(\frac{\tau}{2} < |\sigma_{ij}| \leq \tau) \right)^2 + 2 \left(\sum_{i \neq j} \sigma_{ij}^2 \mathbb{1}(\frac{\tau}{2} < |\sigma_{ij}| \leq \tau) \right)^2 \end{aligned}$$

The first term can be treated in the same way as that of L_2 and has the same rate as L_2 . The second term is bounded by $2L_1$ and has the same rate as L_1 .

Finally, we control L_8 as follows:

$$\begin{aligned} L_8 &\leq \mathbb{E} \left[\left(\sum_{i \neq j} \hat{\sigma}_{ij}^2 \mathbb{1}(|\hat{\sigma}_{ij} - \sigma_{ij}| > \tau/2) \right)^2 \right] \\ &= \sum_{i \neq j} \sum_{s \neq t} \mathbb{E} \left[\hat{\sigma}_{ij}^2 \hat{\sigma}_{st}^2 \mathbb{1}(|\hat{\sigma}_{ij} - \sigma_{ij}| > \frac{\tau}{2}) \mathbb{1}(|\hat{\sigma}_{st} - \sigma_{st}| > \frac{\tau}{2}) \right] \\ &\leq \sum_{i \neq j} \sum_{s \neq t} \left(\mathbb{E} \left[\hat{\sigma}_{ij}^4 \hat{\sigma}_{st}^4 \right] \mathbb{E} \left[\mathbb{1}(|\hat{\sigma}_{ij} - \sigma_{ij}| > \frac{\tau}{2}) \mathbb{1}(|\hat{\sigma}_{st} - \sigma_{st}| > \frac{\tau}{2}) \right] \right)^{1/2} \\ &\leq C \sum_{i \neq j} \sum_{s \neq t} \left(\mathbb{P} \left(|\hat{\sigma}_{ij} - \sigma_{ij}| > \frac{\tau}{2} \right) \mathbb{P} \left(|\hat{\sigma}_{st} - \sigma_{st}| > \frac{\tau}{2} \right) \right)^{1/4}. \end{aligned}$$

Taking now $\tau = 2C_0(\frac{\gamma \log p}{n})^{\frac{1}{2}}$ for $\gamma > 8$ large enough, by Lemma A.2, we get

$$L_8 \leq C \sum_{i \neq j} \sum_{s \neq t} p^{-\gamma/2} \leq Cp^{4-\gamma/2}$$

Hence combining all the inequalities together, we conclude that for such τ , we get

$$\mathcal{E} \leq \begin{cases} C \left[R^2 \left(\tau^{4-2q} + \frac{\tau^{-2q}}{n^2} + \frac{1}{n} \right) + p^{4-\gamma/2} \right] & \text{if } 0 \leq q \leq 1, \\ C \left[R^2 \left(\tau^{4-2q} + \frac{\tau^{-2q}}{n^2} + \frac{\tau^{2-2q}}{n} \right) + p^{4-\gamma/2} \right] & \text{if } 1 < q < 2. \end{cases}$$

Plugging the value of τ completes the proof.

A.2 Proof of Theorem 4.2

First of all, note that

$$\mathbb{E}[(\ell_r(\tilde{\Sigma}_\tau) - \ell_r(\Sigma))^2] \leq \mathbb{E}[\max_i (\ell_r^{(i)}(\tilde{\Sigma}_\tau) - \ell_r^{(i)}(\Sigma))^2].$$

Therefore, we need only to bound the right hand side in the above inequality.

As in the proof of Theorem 3.1, we decompose $\max_i (\ell_r^{(i)}(\tilde{\Sigma}_\tau) - \ell_r^{(i)}(\Sigma))^2$ into three terms.

$$\begin{aligned} \max_i \left(\ell_r^{(i)}(\hat{\Sigma}^\top) - \ell_r^{(i)}(\Sigma) \right)^2 &= \max_i \left(\sum_j |\hat{\sigma}_{ij}|^r \mathbb{1}\{|\hat{\sigma}_{ij}| > \tau\} - |\sigma_{ij}|^r \right)^2 \\ &\leq 3(L_1 + L_2 + L_3), \end{aligned}$$

where

$$\begin{aligned} L_1 &= \max_i \left(\sum_j |\sigma_{ij}|^r \mathbb{1}\{|\sigma_{ij}| \leq \tau\} \right)^2, \\ L_2 &= \max_i \left(\sum_j (|\hat{\sigma}_{ij}|^r - |\sigma_{ij}|^r) \mathbb{1}\{|\sigma_{ij}| > \tau\} \right)^2, \\ L_3 &= \max_i \left(\sum_j |\hat{\sigma}_{ij}|^r (\mathbb{1}\{|\hat{\sigma}_{ij}| > \tau\} - \mathbb{1}\{|\sigma_{ij}| > \tau\}) \right)^2. \end{aligned}$$

We can bound L_1 and L_3 with similar techniques as in the proof of Theorem 3.1. Therefore, we focus only on bounding L_2 .

First notice if $r \leq 1$, $|\hat{\sigma}_{ij}|^r - |\sigma_{ij}|^r \leq |\hat{\sigma}_{ij} - \sigma_{ij}|^r$ by the subadditivity of x^r for $x > 0$. With this fact,

$$\begin{aligned} \mathbb{E}L_2 &\leq \mathbb{E} \left[\max_i \left(\sum_j |\hat{\sigma}_{ij} - \sigma_{ij}|^r \mathbb{1}\{|\sigma_{ij}| > \tau\} \right)^2 \right] \\ &\leq \left(\max_i \sum_j \mathbb{1}\{|\sigma_{ij}| > \tau\} \right)^2 \mathbb{E} \left[\max_{i,j} |\hat{\sigma}_{ij} - \sigma_{ij}|^{2r} \right]. \end{aligned}$$

The right hand side is further bounded by $CR^2(\log p/n)^{r-q}$ due to Lemma A.3 and simple inequality (A.1) with sum over only j in this case.

For any positive integers a and real number $b \geq a$, define

$$C_b^a = \frac{b(b-1)\cdots(b-a+1)}{a!}.$$

If $r > 1$, by Taylor's expansion of $|\hat{\sigma}_{ij}|^r$ at the point $|\sigma_{ij}|$, there exists a θ_{ij} that lies between $|\sigma_{ij}|$ and $|\hat{\sigma}_{ij}|$ such that

$$\begin{aligned} L_2 = \max_i \left(\sum_{m=1}^{\lfloor r \rfloor - 1} C_r^m \sum_{j=1}^p |\sigma_{ij}|^{r-m} (|\hat{\sigma}_{ij}| - |\sigma_{ij}|)^m \mathbb{1}\{|\sigma_{ij}| > \tau\} \right. \\ \left. + C_r^{\lfloor r \rfloor} \sum_{j=1}^p |\theta_{ij}|^{r-\lfloor r \rfloor} (|\hat{\sigma}_{ij}| - |\sigma_{ij}|)^{\lfloor r \rfloor} \mathbb{1}\{|\sigma_{ij}| > \tau\} \right)^2, \end{aligned}$$

where $\lfloor r \rfloor$ represents the largest integer that is smaller than or equal to r . Using the fact that $\| |\hat{\sigma}_{ij}| - |\sigma_{ij}| \| \leq |\hat{\sigma}_{ij} - \sigma_{ij}|$ and $C_r^m \leq e^r$ for $m = 1, \dots, \lfloor r \rfloor$, the above square summation is bounded by

$$C \max_i \left[\sum_{m=1}^{\lfloor r \rfloor - 1} h_{im}^2 + R_{i\lfloor r \rfloor}^2 \right],$$

where C depends on r and

$$\begin{aligned} h_{im} &= \sum_{j=1}^p |\sigma_{ij}|^{r-m} |\hat{\sigma}_{ij} - \sigma_{ij}|^m \mathbb{1}\{|\sigma_{ij}| > \tau\}, \\ R_{i\lfloor r \rfloor} &= \sum_{j=1}^p |\theta_{ij}|^{r-\lfloor r \rfloor} |\hat{\sigma}_{ij} - \sigma_{ij}|^{\lfloor r \rfloor} \mathbb{1}\{|\sigma_{ij}| > \tau\}. \end{aligned}$$

We need to bound each $\mathbb{E}[\max_i h_{im}^2]$ and $\mathbb{E}[\max_i R_{i\lfloor r \rfloor}^2]$. Let $M_0 = \lceil r - q \rceil$ be the smallest integer that is no smaller than $r - q$. Then, $M_0 \geq r - q$ and $M_0 < r - q + 1$. We bound h_{im} terms separately for low-order and high-order regimes: $1 \leq m < M_0$ and $M_0 \leq m \leq \lfloor r \rfloor - 1$. Notice when $q \geq r - 1$, low-order terms do not exist; and high-order terms may or may not exist which depends on how large r is.

For low-order terms $1 \leq m < M_0$, due to the fact $r - m \geq r - M_0 + 1 > q$,

$$\begin{aligned} \mathbb{E} \left[\max_i h_{im}^2 \right] &\leq \left(\max_i \sum_{j,k=1}^p (|\sigma_{ij}| |\sigma_{ik}|)^{r-m} \mathbb{1}\{|\sigma_{ij}| > \tau, |\sigma_{ik}| > \tau\} \right) \\ &\quad \times \mathbb{E} \left[\max_{i,j,k} |\hat{\sigma}_{ij} - \sigma_{ij}|^m |\hat{\sigma}_{ik} - \sigma_{ik}|^m \right] \\ &\leq \left(\max_i \sum_{j=1}^p |\sigma_{ij}|^{r-m} \right)^2 \mathbb{E} \left[\max_{i,j} |\hat{\sigma}_{ij} - \sigma_{ij}|^{2m} \right] \\ &\leq C \left(\max_i \sum_{j=1}^p |\sigma_{ij}|^q \right)^2 \left(\frac{\log p}{n} \right)^m, \end{aligned}$$

where the last inequality utilizes Lemma A.3. Thus, the sum of the low-order terms is dominated by the first term with rate $O(R^2 \log p/n)$. For high-order terms $M_0 \leq m \leq \lfloor r \rfloor - 1$, since $r - m - q \leq r - M_0 - q \leq 0$, we have

$$\begin{aligned} \mathbb{E} \left[\max_i h_{im}^2 \right] &\leq \mathbb{E} \left[\max_i \left(\sum_{j=1}^p |\sigma_{ij}|^q \tau^{r-m-q} |\hat{\sigma}_{ij} - \sigma_{ij}|^m \mathbb{1}_{\{|\sigma_{ij}| > \tau\}} \right)^2 \right] \\ &\leq C \tau^{2(r-m-q)} \left(\max_i \sum_{j=1}^p |\sigma_{ij}|^q \right)^2 \left(\frac{\log p}{n} \right)^m \leq C R^2 \left(\frac{\log p}{n} \right)^{r-q}. \end{aligned}$$

Now, it remains to bound the remaining term. Similar to the case $r \leq 1$, we have

$$|\theta_{ij}|^{r-\lfloor r \rfloor} \leq |\sigma_{ij}|^{r-\lfloor r \rfloor} + |\hat{\sigma}_{ij} - \sigma_{ij}|^{r-\lfloor r \rfloor} \leq 1 + |\hat{\sigma}_{ij} - \sigma_{ij}|^{r-\lfloor r \rfloor},$$

by again the subadditivity of function $x^{r-\lfloor r \rfloor}$, $x > 0$. Using same tricks as before, Lemma A.3 and inequality (A.1),

$$\mathbb{E}[\max_i R_{i\lfloor r \rfloor}^2] \leq C R^2 \left(\frac{\log p}{n} \right)^{r-q}.$$

The remaining term obviously disappears if r is integer. Combining the bounds that we have obtained so far, $L_2 \leq C \tilde{\psi}_{n,p}(q, R)$. It attains the parametric rate only when $r > 1$ and $q < r - 1$. Otherwise, the nonparametric rate shows up. This completes the proof.

A.3 Technical Lemmas

LEMMA A.1 (Laurent and Massart, 2000 [LM00]). *Suppose Y_1, Y_2, \dots, Y_D are i.i.d. standard Gaussian random variables and $Z = \sum_{i=1}^D a_i (Y_i^2 - 1)$ for some constants a_1, a_2, \dots, a_D . Then,*

$$(A.2) \quad \mathbb{P}(|Z| \geq 2|a|_2 \sqrt{x} + 2|a|_\infty x) \leq 2e^{-x},$$

where $|a|_\infty = \sup_i |a_i|$ and $|a|_2^2 = \sum a_i^2$.

Similar results to the following lemma have been obtained by various authors, including [BL04]. Below, we give a non-asymptotic bound to facilitate our technical proof.

LEMMA A.2. *For $\Sigma \in \mathcal{F}_q$, $X_k \sim \mathcal{N}(0, \Sigma)$ and any constant $C_0 \geq 4$, we have*

$$(A.3) \quad \mathbb{P}(|\hat{\sigma}_{i,j} - \sigma_{i,j}| > t) \leq 4e^{-nt^2/C_0^2},$$

for all $t \leq C_0$, where $\hat{\sigma}_{ij} = \frac{1}{n} \sum_{k=1}^n X_{ki} X_{kj}$. Consequently, if $\gamma \log(p) < n$, we have

$$(A.4) \quad \mathbb{P} \left(|\hat{\sigma}_{i,j} - \sigma_{i,j}| > 4 \sqrt{\frac{\gamma \log(p)}{n}} \right) \leq 4p^{-\gamma}.$$

PROOF. Let $Y_{kij} = \frac{1}{2}(X_{ki} + X_{kj})$ and $Z_{kij} = \frac{1}{2}(X_{ki} - X_{kj})$. Then, they are jointly normal with marginal distributions $Y_{kij} \sim \mathcal{N}(0, \sigma_{Y_{ij}}^2)$ and $Z_{kij} \sim \mathcal{N}(0, \sigma_{Z_{ij}}^2)$, where

$$\sigma_{Y_{ij}}^2 = \frac{1}{4}(\sigma_{ii} + 2\sigma_{ij} + \sigma_{jj}), \quad \text{and} \quad \sigma_{Z_{ij}}^2 = \frac{1}{4}(\sigma_{ii} - 2\sigma_{ij} + \sigma_{jj}).$$

Define $\Delta_{Y_{ij}} = \frac{1}{n} \sum_{k=1}^n (Y_{kij}^2 - \sigma_{Y_{ij}}^2)$ and $\Delta_{Z_{ij}} = \frac{1}{n} \sum_{k=1}^n (Z_{kij}^2 - \sigma_{Z_{ij}}^2)$. It is easily seen that $\hat{\sigma}_{ij} - \sigma_{ij} = \Delta_{Y_{ij}} - \Delta_{Z_{ij}}$.

By Lemma A.1, $\mathbb{P}\left(|\Delta_{Y_{ij}}| \geq 2\frac{\sigma_{Y_{ij}}^2}{\sqrt{n}}\sqrt{x} + 2\frac{\sigma_{Y_{ij}}^2}{n}x\right) \leq 2e^{-x}$. Choosing $x = \gamma \log(p)$ and using $\gamma \log(p) < n$, we have

$$\begin{aligned} \mathbb{P}\left(|\Delta_{Y_{ij}}| \geq 4\sigma_{Y_{ij}}^2\sqrt{\frac{\gamma \log p}{n}}\right) &\leq \mathbb{P}\left(|\Delta_{Y_{ij}}| \geq 2\sigma_{Y_{ij}}^2\left(\sqrt{\frac{\gamma \log p}{n}} + \frac{\gamma \log p}{n}\right)\right) \\ &\leq 2p^{-\gamma}. \end{aligned}$$

The similar result holds for $\Delta_{Z_{ij}}$. Therefore, by using $\sigma_{ii} \leq 1$, we have

$$\begin{aligned} &\mathbb{P}\left(|\hat{\sigma}_{ij} - \sigma_{ij}| > 4\sqrt{\frac{\gamma \log p}{n}}\right) \\ &\leq \mathbb{P}\left(|\Delta_{Y_{ij}}| + |\Delta_{Z_{ij}}| > 4\sqrt{\frac{\gamma \log p}{n}}\right) \\ &\leq \mathbb{P}\left(|\Delta_{Y_{ij}}| + |\Delta_{Z_{ij}}| > 4\sqrt{\frac{\gamma \log p}{n}}(\sigma_{Y_{ij}}^2 + \sigma_{Z_{ij}}^2)\right) \\ &\leq \mathbb{P}\left(|\Delta_{Y_{ij}}| > 4\sqrt{\frac{\gamma \log p}{n}}\sigma_{Y_{ij}}^2\right) + \mathbb{P}\left(|\Delta_{Z_{ij}}| > 4\sqrt{\frac{\gamma \log p}{n}}\sigma_{Z_{ij}}^2\right) \\ &\leq 4p^{-\gamma}. \end{aligned}$$

This proves the second part of the lemma. For any constant $C_0 > 4$, we get

$$\mathbb{P}\left(|\hat{\sigma}_{ij} - \sigma_{ij}| > C_0\sqrt{\frac{\gamma \log p}{n}}\right) \leq 4p^{-\gamma}.$$

Taking $t = C_0\sqrt{\frac{\gamma \log p}{n}} \leq C_0$, we obtain the first part of the lemma. \square

LEMMA A.3. *Let X_1, \dots, X_n be n i.i.d random vectors such that $X_1 \sim \mathcal{N}_p(0, \Sigma)$, $\Sigma = \{\sigma_{ij}\}$ and $\hat{\Sigma} = \{\hat{\sigma}_{ij}\}$ be the sample covariance matrix. Then, there exists a positive constant C (which depends on r) such that*

$$\mathbb{E}(\hat{\sigma}_{ij} - \sigma_{ij})^4 \leq \frac{C}{n^2} \quad \text{and} \quad \mathbb{E}\left[\max_{i,j}(\hat{\sigma}_{ij} - \sigma_{ij})^{2m}\right] \leq C\left(\frac{\log p}{n}\right)^m$$

for all positive $m \leq r$, provided that $\gamma \log p < n$ for constant $\gamma > 8$.

PROOF. The first part of the lemma is easily derived by direct computation. In fact, more general results for the sample mean, which include our case, have already been derived in the literature. Now let us derive the bound for the second conclusion. Define function $f_t(x) = \exp(tx^{1/(2m)})$ for a pre-chosen $t > 0$. Then,

$f_t(x)$ itself is convex and monotonically increasing for $x > 0$ if $m \leq 1/2$. However, it is not necessarily convex if $m > 1/2$. In this case we employ another function

$$g_t(x) = \begin{cases} \frac{e^{2m-1}}{2m} \left(1 + tx \left(\frac{2m-1}{t}\right)^{1-2m}\right), & \text{if } 0 < x \leq \left((2m-1)/t\right)^{2m}, \\ f_t(x), & \text{otherwise.} \end{cases}$$

It is easy to see $f_t(x) \leq g_t(x) \leq e^{2m-1}f_t(x)$ and $g_t(x)$ is convex and monotonically increasing for $x > 0$. We will apply $g_t(x)$ to bound the expected maxima (use $f_t(x)$ instead for $m \leq 1/2$). Denote $\eta_{ij} = \hat{\sigma}_{ij} - \sigma_{ij}$, then

$$\mathbb{E} \left[\max_{i,j} (\hat{\sigma}_{ij} - \sigma_{ij})^{2m} \right] = \mathbb{E} \left[g_t^{-1} \left(\max_{i,j} g_t(\eta_{ij}^{2m}) \right) \right] \leq g_t^{-1} \left(\mathbb{E} \left[\max_{i,j} g_t(\eta_{ij}^{2m}) \right] \right),$$

where the last inequality is due to the concavity of g_t^{-1} and Jensen's inequality. By monotonicity and $f_t(x) \leq g_t(x) \leq e^{2m-1}f_t(x)$, the right hand side is bounded by

$$f_t^{-1} \left(e^{2m-1} \mathbb{E} \left[\max_{i,j} f_t(\eta_{ij}^{2m}) \right] \right) \leq \left(\frac{\log(e^{2m-1} \mathbb{E} \max_{i,j,\delta} e^{t\delta\eta_{ij}})}{t} \right)^{2m},$$

where $\delta \in \{-1, 1\}$. The max in the right hand side is further bounded by replacing max with sum over i, j and δ , that is

$$\left(\frac{\log(e^{2m-1} \mathbb{E} \sum_{i,j,\delta} e^{t\delta\eta_{ij}})}{t} \right)^{2m} \leq t^{-2m} \left(2r - 1 + \log \sum_{i,j} \mathbb{E}[e^{t\eta_{ij}} + e^{-t\eta_{ij}}] \right)^{2m}.$$

We will show that

$$(A.5) \quad \mathbb{E} \exp(\lambda(X_{ki}X_{kj} - \sigma_{ij})) \leq \exp(16\lambda^2),$$

for $|\lambda| \leq 3/16$. Therefore, for $t/n \leq 3/16$, we have

$$\mathbb{E} \left[\max_{i,j} (\hat{\sigma}_{ij} - \sigma_{ij})^{2m} \right] \leq \left((2r+2) \frac{\log p}{t} + 16 \frac{t}{n} \right)^{2m}.$$

Choosing $t = \sqrt{n \log(p)}/8$, we get the optimal bound of $C \left(\frac{\log p}{n}\right)^m$ where C only depends on r . And the corresponding $\lambda = t/n = \sqrt{\log(p)}/8n$ is indeed smaller than $3/16$ due to the assumption $\gamma \log p < n$ for $\gamma > 8$.

So it left only to prove the claim (A.5), which actually says $X_{ki}X_{kj} - \sigma_{ij}$ is sub-exponential. We omit index k for simplicity in the following. By the same notation in proof of Lemma A.2, we have $X_iX_j - \sigma_{ij} = (Y_{ij}^2 - \sigma_{Y_{ij}}^2) - (Z_{ij}^2 - \sigma_{Z_{ij}}^2)$. Therefore, by Holder's inequality,

$$\mathbb{E} \left[e^{\lambda(X_iX_j - \sigma_{ij})} \right] \leq \left(\mathbb{E} \left[e^{2\lambda(Y_{ij}^2 - \sigma_{Y_{ij}}^2)} \right] \mathbb{E} \left[e^{-2\lambda(Z_{ij}^2 - \sigma_{Z_{ij}}^2)} \right] \right)^{1/2},$$

where $Y_{ij} \sim N(0, \sigma_{Y_{ij}}^2)$ and $Z_{ij} \sim N(0, \sigma_{Z_{ij}}^2)$. We first bound the Y_{ij} term:

$$\begin{aligned} \mathbb{E} \left[e^{2\lambda(Y_{ij}^2 - \sigma_{Y_{ij}}^2)} \right] &= \exp \left(-\frac{1}{2} \log(1 - 4\lambda\sigma_{Y_{ij}}^2) - 2\lambda\sigma_{Y_{ij}}^2 \right) \\ &\leq \exp \left((4\lambda\sigma_{Y_{ij}}^2)^2 \right) \leq e^{16\lambda^2}. \end{aligned}$$

The first inequality is due to the fact $-\log(1-x) - x \leq 2x^2$ for $x \leq 3/4$ with $x = 4\lambda\sigma_{Y_{ij}}^2$ and the last inequality is because $\sigma_{Y_{ij}}^2 \leq 1$. Similarly the same bound holds for Z_{ij} term for $4\lambda\sigma_{Z_{ij}}^2 \geq -3/4$. Combining them together, we completes the proof for claim (A.5) and thus for the lemma. \square

APPENDIX B: MINIMAX LOWER BOUNDS

Our lower bounds rely on standard arguments from minimax theory. We refer to [Tsy09, Chapter 2] for more details. In the sequel, let $\text{KL}(P, \bar{P})$ denote the Kullback-Leibler divergence between two distributions P and \bar{P} , where $P \ll \bar{P}$. It is defined by

$$\text{KL}(P, \bar{P}) = \int \log \left(\frac{dP}{d\bar{P}} \right) dP.$$

B.1 Proof of Proposition 3.1

We are going to employ a simple two-point lower bound. Fix $\epsilon \in (0, 1/2)$ and let P_p^n (respectively \bar{P}_p^n) denote the distribution of a sample X_1, \dots, X_n where $X_1 \sim \mathcal{N}(0, I_p)$ (respectively $X_1 \sim \mathcal{N}(0, (1 - \epsilon)I_p)$). Next, observe that $I_p, (1 - \epsilon)I_p \subset \mathcal{D}_p$ so that

$$(B.1) \quad \sup_{\Sigma \in \mathcal{D}_p} \mathbb{E} |\hat{D} - D(\Sigma)| \geq \max_{\Sigma \in \{I_p, (1-\epsilon)I_p\}} \mathbb{E} |\hat{D} - D(\Sigma)|$$

Moreover, $|D(I_p) - D((1 - \epsilon)I_p)| = p(2\epsilon - \epsilon^2) > p\epsilon$. Then, it follows from the Markov inequality that

$$(B.2) \quad \begin{aligned} \frac{1}{p\epsilon} \max_{\Sigma \in \{I_p, (1-\epsilon)I_p\}} \mathbb{E} |\hat{D} - D(\Sigma)| &\geq \max_{\Sigma \in \{I_p, (1-\epsilon)I_p\}} \mathbb{P}[|\hat{D} - D(\Sigma)| > p\epsilon] \\ &\geq \frac{1}{4} \exp[-\text{KL}(P_p^n, \bar{P}_p^n)], \end{aligned}$$

where the last inequality follows from [Tsy09, Theorem 2.2, (iii)].

Completion of the proof requires an upper bound on $\text{KL}(P_p^n, \bar{P}_p^n)$. To that end, note that it follows from the chain rule and simple algebra that

$$\text{KL}(P_p^n, \bar{P}_p^n) = np \text{KL}(P_1^1, \bar{P}_1^1) = \frac{np}{2} \left[\log(1 - \epsilon) + \frac{\epsilon}{1 - \epsilon} \right] \leq \frac{np}{2} \frac{\epsilon^2}{1 - \epsilon} \leq np\epsilon^2.$$

Taking now $\epsilon = 1/(2\sqrt{np}) \leq 1/2$ yields $\text{KL}(P_p^n, \bar{P}_p^n) \leq 1/4$. Together with (B.1) and (B.2), it yields

$$\inf_{\hat{D}} \sup_{\Sigma \in \mathcal{D}_p} \mathbb{E} |\hat{D} - D(\Sigma)| \geq \frac{1}{8e^{1/4}} \sqrt{\frac{p}{n}}.$$

To conclude the proof, we square the above inequality and employ Jensen's inequality.

B.2 Proof of Theorem 3.2

To prove minimax lower bounds, we employ a standard technique that consists of reducing the estimation problem to a testing problem. We split this proof into two parts and begin by proving

$$\inf_{\hat{Q}} \sup_{\Sigma \in \mathcal{F}_q} \mathbb{E}_{\Sigma} [\hat{Q} - Q(\Sigma)]^2 \geq C \frac{R^2}{n},$$

for some positive constant $C > 0$. To that end, for any $A \in \mathbf{S}_p^+$, let \mathbb{P}_A denote the distribution of $X \sim \mathcal{N}(0, A)$. It is not hard to show if $|A| > 0$ and $|B| > 0$, $A, B \in \mathbf{S}_p^+$, then the Kullback-Leibler divergence between \mathbb{P}_A and \mathbb{P}_B is given by

$$(B.3) \quad \text{KL}(\mathbb{P}_A, \mathbb{P}_B) = \frac{1}{2} \left[\log \left(\frac{|B|}{|A|} \right) + \text{tr}(B^{-1}A) - p \right].$$

Next, take A and B to be of the form

$$A^{(k)} = \begin{pmatrix} \mathbf{1}\mathbf{1}^\top & a\mathbf{1}\mathbf{1}^\top & 0 \\ a\mathbf{1}\mathbf{1}^\top & \mathbf{1}\mathbf{1}^\top & 0 \\ 0 & 0 & I_{p-k} \end{pmatrix}, \quad B^{(k)} = \begin{pmatrix} \mathbf{1}\mathbf{1}^\top & b\mathbf{1}\mathbf{1}^\top & 0 \\ b\mathbf{1}\mathbf{1}^\top & \mathbf{1}\mathbf{1}^\top & 0 \\ 0 & 0 & I_{p-k} \end{pmatrix}$$

where $a, b \in (0, 1/2)$, 0 is a generic symbol to indicate that the missing space is filled with zeros, and $\mathbf{1}$ denotes a vector of ones of length $k/2$. Note that if we have random variables $(X, Y, Z_1, \dots, Z_{p-2})$ chosen from distribution $\mathcal{N}(0, A^{(2)})$ meaning that Z_k 's are independent with X, Y but the correlation between X and Y is a , then random vector $(X, \dots, X, Y, \dots, Y, Z_1, \dots, Z_{p-k})$ with $k/2$ X 's and Y 's in it follows $\mathcal{N}(0, A^{(k)})$. It is obvious that these two matrices are degenerate and comes from perfectly correlated random variables. Since perfectly correlated random variables do not add new information, for such matrices, an application of (B.3) yields

$$\text{KL}(\mathbb{P}_{A^{(k)}}, \mathbb{P}_{B^{(k)}}) = \text{KL}(\mathbb{P}_{A^{(2)}}, \mathbb{P}_{B^{(2)}}) = \frac{1-ab}{1-b^2} - \frac{1}{2} \log \left(\frac{1-a^2}{1-b^2} \right) - 1.$$

Next, using the convexity inequality $\log(1+x) \geq x - x^2/2$ for all $x > 0$, we get that

$$\text{KL}(\mathbb{P}_{A^{(k)}}, \mathbb{P}_{B^{(k)}}) \leq \frac{(a-b)^2}{2(1-b^2)} \left[1 + \frac{(a+b)^2}{2(1-b^2)} \right] \leq 2(a-b)^2,$$

using the fact that $a, b \in (0, 1/2)$. Take now if $R > 4$

$$a = \frac{1}{4}, \quad b = a + \frac{1}{4\sqrt{n}}, \quad k = \sqrt{R}$$

so that we indeed have $a, b \in (0, 1/2)$ and also $A^{(k)}, B^{(k)} \in \mathcal{F}_q$ obviously. If $R < 4$, take $k = 2, a = \sqrt{R}/8, b = a + \sqrt{R}/64n$ instead. Moreover, this choice leads to $n\text{KL}(\mathbb{P}_A, \mathbb{P}_B) \leq 1/5$. Using standard techniques to reduce estimation problems to testing problems (see, e.g., [Tsy09, Theorem 2.5]), we find that

$$\inf_{\hat{Q}} \max_{\Sigma \in \{A, B\}} \mathbb{E}_\Sigma [(\hat{Q} - Q(\Sigma))^2] \geq C(Q(A) - Q(B))^2.$$

For the above choice of A and B , we have

$$(Q(A^{(k)}) - Q(B^{(k)}))^2 = \frac{k^4}{4}(a^2 - b^2)^2 \geq C \frac{R^2}{n}.$$

Since $A^{(k)}, B^{(k)} \in \mathcal{F}_q$, the above two displays imply that

$$\inf_{\hat{Q}} \max_{\Sigma \in \mathcal{F}_q} \mathbb{E}_\Sigma [(\hat{Q} - Q(\Sigma))^2] \geq C \frac{R^2}{n},$$

which completes the proof of the first part of the lower bound.

For the second part of the lower bound, we reduce our problem to a testing problem of the same flavor as [ACBL14, BR13b]. Note however, that our construction is different because the covariance matrices considered in these papers do not yield large enough lower bounds. We use the following construction.

Fix an integer $k \in [p-1]$ and let $\mathcal{S} = \{S \subset [p-1] : |S| = k\}$ denote the set of subsets of $[p-1]$ that have cardinality k . Fix $a \in (0, 1)$ to be chosen later and for any $S \in \mathcal{S}$, recall that $\mathbf{1}_S$ is the column vector in $\{0, 1\}^{p-1}$ with support given by S . For each $S \in \mathcal{S}$, we define the following $p \times p$ covariance matrix:

$$(B.4) \quad \Sigma_S = \begin{pmatrix} 1 & a\mathbf{1}_S^\top \\ a\mathbf{1}_S & I_{p-1} \end{pmatrix}$$

Let \mathbb{P}_0 denote the distribution of $X \sim \mathcal{N}_p(0, I_p)$ and \mathbb{P}_S denote the distribution of $X \sim \mathcal{N}_p(0, \Sigma_S)$. Let \mathbb{P}_0^n (resp. \mathbb{P}_S^n) denote the distribution of $\mathbf{X} = (X_1, \dots, X_n)$ of a collection n i.i.d random variables drawn from \mathbb{P}_0 (resp. \mathbb{P}_S). Moreover, let $\bar{\mathbb{P}}^n$ denote the distribution of \mathbf{X} where the X_i 's are drawn as follows: first draw S uniformly at random from \mathcal{S} and then, conditionally on S , draw X_1, \dots, X_n independently from \mathbb{P}_S . Note that $\bar{\mathbb{P}}^n$ is the mixture of n independent samples rather the distribution of n independent random vectors drawn from a mixture distribution. Consider the following testing problem:

$$H_0 : \mathbf{X} \sim \mathbb{P}_0^n \quad \text{vs.} \quad H_1 : \mathbf{X} \sim \bar{\mathbb{P}}^n.$$

Using [Tsy09, Theorem 2.2, part (iii)], we get that for any test $\psi = \psi(\mathbf{X})$, we have

$$\mathbb{P}_0^n(\psi = 0) \vee \max_{S \in \mathcal{S}} \mathbb{P}_S^n(\psi = 1) \geq \mathbb{P}_0^n(\psi = 0) \vee \bar{\mathbb{P}}^n(\psi = 1) \geq \frac{1}{4} \exp(-\chi^2(\bar{\mathbb{P}}^n, \mathbb{P}_0)),$$

where we recall that the χ^2 -divergence between two probability distributions P and Q is defined by

$$\chi^2(P, Q) = \begin{cases} \int \left(\frac{dP}{dQ} - 1 \right)^2 dQ, & \text{if } P \ll Q \\ \infty, & \text{otherwise.} \end{cases}$$

Lemma B.1 implies that for suitable choices of the parameters a and k , we have $\chi^2(\bar{\mathbb{P}}^n, \mathbb{P}_0) \leq 2$ so that the test errors are bounded below by a constant $C = e^{-2}/4$. Since $Q(\Sigma_S) = 2ka^2$ for any $S \in \mathcal{S}$, it follows from a standard reduction from hypothesis testing to estimation (see, e.g., [Tsy09, Theorem 2.5]) that the above result implies the following lower bound:

$$(B.5) \quad \inf_{\hat{Q}} \max_{\Sigma \in \mathcal{H}} \mathbb{E}_\Sigma [(\hat{Q} - Q(\Sigma))^2] \geq Ck^2a^4,$$

for some positive constant C , where the infimum is taken over all estimators \hat{Q} of $Q(\Sigma)$ based on n observations and \mathcal{H} is the class of covariance matrices defined by

$$\mathcal{H} = \{I_p\} \cup \{\Sigma_S : S \in \mathcal{S}\}.$$

To complete the proof, observe that the values of a and k prescribed in Lemma B.1 imply that $\mathcal{H} \subset \mathcal{F}_q$ and give the desired lower bound. Note first that, for any choice of a and k , the following holds trivially: $I_p \in \mathcal{F}_q$ and $\text{diag}(\Sigma_S) = I_p$ for any $S \in \mathcal{S}$. Write $\Sigma_S = (\sigma_{ij})$ and observe that

$$\sum_{i \neq j} |\sigma_{ij}|^q = 2ka^q.$$

Next, we treat each case of Lemma B.1 separately.

Case 1. Note first that $2ka^q = R/2 < R$ so that $\Sigma_S \in \mathcal{F}_q$. Moreover, $k^2a^4 = CR^{4/q}$.

Case 2. Note first that $2ka^q \leq R/2 < R$ so that $\Sigma_S \in \mathcal{F}_q$. Since $k \geq 2$ and $k^2 \leq R^2n^q$, we have

$$k \geq \frac{R}{4} \left(\frac{\log\left(\frac{p-1}{k^2} + 1\right)}{2n} \right)^{-q/2}.$$

Therefore,

$$k^2a^4 \geq \frac{R^2}{16} \left(\frac{\log\left(\frac{p-1}{R^2n^q} + 1\right)}{2n} \right)^{2-q} \wedge \frac{1}{4}.$$

Combining the two cases, we get

$$k^2a^4 \geq C \left[R^2 \left(\frac{\log\left(\frac{p-1}{R^2n^q} + 1\right)}{2n} \right)^{2-q} \wedge R^{4/q} \wedge 1 \right].$$

Together with (B.5), this completes the proof of the second part of the lower bound.

B.3 A technical lemma on χ^2 divergences

LEMMA B.1. *Consider a mixture of Gaussian product distributions*

$$\tilde{\mathbb{P}}^n = \frac{1}{m} \sum_{j=1}^m \mathbb{P}_j^n$$

where $\mathbb{P}_j \sim \mathcal{N}_p(0, \Sigma_j)$ such that $\mathbb{P}_j \ll \mathbb{P}_0$. Then

$$(B.6) \quad \chi^2(\tilde{\mathbb{P}}^n, \mathbb{P}_0^n) = \frac{1}{m^2} \sum_{j,k=1}^m |I_p - (\Sigma_j - I)(\Sigma_k - I)|^{-n/2} - 1.$$

Furthermore, assume $2(\log p) \leq n$. Consider the mixture $\tilde{\mathbb{P}}^n$ defined in subsection B.2 where k and a are defined as follows.

1. If $R < 4\left(\frac{\log p}{n}\right)^{q/2}$, then take $k = 1$ and $a = (R/4)^{1/q}$.
2. If $R \geq 4\left(\frac{\log p}{n}\right)^{q/2}$, then take k to be the largest integer such that

$$(B.7) \quad k \leq \frac{R}{2} \left(\frac{\log\left(\frac{p-1}{k^2} + 1\right)}{2n} \right)^{-q/2}.$$

and

$$(B.8) \quad a = \left(\frac{\log\left(\frac{p-1}{k^2} + 1\right)}{2n} \right)^{1/2} \wedge (2k)^{-1/2}.$$

Such choices yield in both cases

$$(B.9) \quad \chi^2(\tilde{\mathbb{P}}^n, \mathbb{P}_0^n) \leq e - 1.$$

Moreover, in case 2 we have that (i) $k \geq 2$ and (ii) under the assumption that $R^2 < (p-1)n^{-q}/2$, we also have $k^2 \leq R^2n^q < (p-1)/2$.

PROOF. To unify the notation, we will work directly with $\mathbb{P}_S, S \in \mathcal{S}$ rather than $\mathbb{P}_j, j \in [m]$. However, in the first part of the proof, we will not use the specific form Σ_S nor that of \mathcal{S} . For now, we simply assume that Σ_S is invertible (we will check this later on). Recall that

$$\chi^2(\bar{\mathbb{P}}^n, \mathbb{P}_0^n) = \mathbb{E}_0 \left[\left(\frac{d\bar{\mathbb{P}}^n}{d\mathbb{P}_0^n} - 1 \right)^2 \right] = \frac{1}{|\mathcal{S}|^2} \sum_{S, T \in \mathcal{S}} \left(\mathbb{E}_0 \left[\frac{d\mathbb{P}_S}{d\mathbb{P}_0} \frac{d\mathbb{P}_T}{d\mathbb{P}_0} \right] \right)^n - 1,$$

where \mathbb{E}_0 denotes the expectation with respect to \mathbb{P}_0 . Furthermore,

$$\mathbb{E}_0 \left[\frac{d\mathbb{P}_S}{d\mathbb{P}_0} \frac{d\mathbb{P}_T}{d\mathbb{P}_0} \right] = \frac{1}{(|\Sigma_S| |\Sigma_T|)^{1/2}} \mathbb{E}_0 \left[\exp \left(-\frac{1}{2} X^\top (\Sigma_S^{-1} + \Sigma_T^{-1} - 2I_p) X \right) \right].$$

Consider the spectral decomposition of $\Sigma_S^{-1} + \Sigma_T^{-1} - 2I_p = U \Lambda U^\top$, where U is an orthogonal matrix and Λ is a diagonal matrix with eigenvalues $\lambda_1, \dots, \lambda_p$ on its diagonal. Then, by rotational invariance of the Gaussian distribution, it holds

$$\begin{aligned} \mathbb{E}_0 \left[\exp \left(-\frac{1}{2} X^\top (\Sigma_S^{-1} + \Sigma_T^{-1} - 2I_p) X \right) \right] &= \mathbb{E}_0 \left[\exp \left(-\frac{1}{2} X^\top \Lambda X \right) \right] \\ &= \prod_{j=1}^p \mathbb{E}_0 \left[\exp \left(-\frac{1}{2} \lambda_j X_j^2 \right) \right] \\ &= \begin{cases} \prod_{j=1}^p (1 + \lambda_j)^{-1/2} = |I + \Lambda|^{-1/2} & \text{if } \max_j \lambda_j < 1 \\ \infty & \text{otherwise} \end{cases} \end{aligned}$$

To ensure that the above expression is finite, note that the Cauchy-Schwarz inequality yields

$$\begin{aligned} \left(\mathbb{E}_0 \left[\frac{d\mathbb{P}_S}{d\mathbb{P}_0} \frac{d\mathbb{P}_T}{d\mathbb{P}_0} \right] \right)^2 &\leq \mathbb{E}_0 \left[\left(\frac{d\mathbb{P}_S}{d\mathbb{P}_0} \right)^2 \right] \mathbb{E}_0 \left[\left(\frac{d\mathbb{P}_T}{d\mathbb{P}_0} \right)^2 \right] \\ &= (\chi^2(\mathbb{P}_S, \mathbb{P}_0) + 1)(\chi^2(\mathbb{P}_T, \mathbb{P}_0) + 1) < \infty \end{aligned}$$

where the two χ^2 divergences are finite because $\mathbb{P}_S \ll \mathbb{P}_0$ for any $S \in \mathcal{S}$. Therefore,

$$\mathbb{E}_0 \left[\frac{d\mathbb{P}_S}{d\mathbb{P}_0} \frac{d\mathbb{P}_T}{d\mathbb{P}_0} \right] = \frac{|I + \Lambda|^{-1/2}}{(|\Sigma_S| |\Sigma_T|)^{1/2}} = \frac{|\Sigma_S^{-1} + \Sigma_T^{-1} - I_p|^{-1/2}}{(|\Sigma_S| |\Sigma_T|)^{1/2}}.$$

Next, observe that

$$\begin{aligned} (|\Sigma_S| |\Sigma_T| |\Sigma_S^{-1} + \Sigma_T^{-1} - I_p|)^{-1/2} &= (|\Sigma_S (\Sigma_S^{-1} + \Sigma_T^{-1} - I_p) \Sigma_T|)^{-1/2} \\ &= |I - (\Sigma_S - I)(\Sigma_T - I)|^{-1/2}. \end{aligned}$$

Since we have not used the specific form of $\Sigma_S, S \in \mathcal{S}$, this bound is valid for any mixture and completes the proof of (B.6).

Next we apply this bound to the specific choice for Σ_S of (B.4). Note that the minimal eigenvalue of the matrices $\Sigma_S, S \in \mathcal{S}$ is $1 - \sqrt{ka^2}$. Later we will show $2a^2k \leq 1$, which implies that Σ_S is always positive definite. In particular, this implies that $\mathbb{P}_S \ll \mathbb{P}_0$ for any $S \in \mathcal{S}$. Moreover, it follows from definition (B.4) that

$$I - (\Sigma_S - I)(\Sigma_T - I) = \begin{pmatrix} 1 - a^2 \mathbf{1}_S^\top \mathbf{1}_T & 0 \\ 0 & I - a^2 \mathbf{1}_S \mathbf{1}_T^\top \end{pmatrix},$$

where 0 is a generic symbol to indicate space filled by zeros. Expanding the determinant along the first row (or column), we get

$$|I - (\Sigma_S - I)(\Sigma_T - I)|^{-1/2} = (1 - a^2 \mathbf{1}_S^\top \mathbf{1}_T)^{-1/2} |I - a^2 \mathbf{1}_S \mathbf{1}_T^\top|^{-1/2} = (1 - a^2 \mathbf{1}_S^\top \mathbf{1}_T)^{-1},$$

where in the second equality, we used Sylvester's determinant theorem. By (B.6), we have

$$\chi^2(\bar{\mathbb{P}}^n, \mathbb{P}_0^n) = \frac{1}{|\mathcal{S}|^2} \sum_{S, T \in \mathcal{S}} (1 - a^2 \mathbf{1}_S^\top \mathbf{1}_T)^{-n} - 1$$

As to be verified later, $2a^2k \leq 1$. Using the fact that $(1 - x)^{-1} \leq \exp(2x)$ for $x \in [0, 1/2]$ and the symmetry, we have

$$\chi^2(\bar{\mathbb{P}}^n, \mathbb{P}_0^n) \leq \frac{1}{|\mathcal{S}|} \sum_{S \in \mathcal{S}} \exp(2na^2 |S \cap [k]|) - 1 = \mathcal{E}[\exp(2na^2 |S \cap [k]|) - 1],$$

where \mathcal{E} denotes the expectation with respect to the distribution of S randomly chosen from \mathcal{S} . In particular, $|S \cap [k]| = \sum_{i=1}^k \mathbb{1}(i \in S)$ is the sum of k negatively associated random variables. Therefore, the above expectation is further bounded by, using negative association,

$$\prod_{i=1}^k \mathcal{E}[e^{2na^2 \mathbb{1}(i \in S)}].$$

Next, for a given by (B.8), we have

$$\prod_{i=1}^k \mathcal{E}[e^{2na^2 \mathbb{1}(i \in S)}] = [(e^{2na^2} - 1) \frac{k}{p-1} + 1]^k \leq [1 + \frac{1}{k}]^k \leq e.$$

We now show for both cases of the lemma, we have $2a^2k \leq 1$. Indeed for case 1, we get $2a^2k = 2(R/4)^{2/q} < 2(\log p)/n \leq 1$. For case 2, $2a^2k \leq 1$ follows trivially from the definition of a . Also observe that $k \geq 2$ since,

$$\frac{R}{2} \left(\frac{\log(\frac{p-1}{4} + 1)}{n} \right)^{-q/2} \geq 2 \left(\frac{\log p}{n} \right)^{q/2} \left(\frac{\log(\frac{p-1}{4} + 1)}{n} \right)^{-q/2} \geq 2.$$

This proves part (i) of the statement on k . To prove part (ii), observe that $R^2 < (p-1)n^{-q}/2$ implies that

$$2^{1-2/q} < 1 < \log \left(\frac{p-1}{R^2 n^q} + 1 \right),$$

which is equivalent to

$$Rn^{q/2} > \frac{R}{2} \left(\frac{\log(\frac{p-1}{R^2 n^q} + 1)}{2n} \right)^{-q/2},$$

Therefore, $k^2 \leq R^2 n^q < (p-1)/2$. □

B.4 Proof of Theorem 4.1

The proof follows a similar idea to that of Theorem 3.2. For the second part of the lower bound, we use exactly the same construction of two hypotheses as in Lemma B.1. Then, it follows that for the ℓ_r functional,

$$\inf_{\hat{L}} \max_{\Sigma \in \mathcal{H}} \mathbb{E}_{\Sigma} [(\hat{L} - \ell_r(\Sigma))^2] \geq Ck^2a^{2r},$$

for some positive constant C , where the infimum is taken over all estimators \hat{L} of $\ell_r(\Sigma)$ based on n observations. In case 1, $k^2a^{2r} = CR^{2r/q}$ while in case 2, following the same arguments as before,

$$k^2a^{2r} \geq \frac{R^2}{16} \left(\frac{\log\left(\frac{p-1}{R^{2n^q}} + 1\right)}{2n} \right)^{r-q} \wedge \frac{1}{2}.$$

This completes the second part of the lower bound.

The first part of the result is a little bit more complicated than the construction of $A^{(k)}$ and $B^{(k)}$ in subsection B.2 due to the extra $\log p$ term in the lower bound. We need to consider a mixture of measures in order to capture the complexity of the problem. With a slight abuse of notation, we redefine $(2k) \times (2k)$ matrices $A^{(k)}, B^{(k)}$ as follows:

$$A^{(k)} = \begin{pmatrix} \mathbf{1}\mathbf{1}^\top & a\mathbf{1}\mathbf{1}^\top \\ a\mathbf{1}\mathbf{1}^\top & \mathbf{1}\mathbf{1}^\top \end{pmatrix}, \quad B^{(k)} = \begin{pmatrix} \mathbf{1}\mathbf{1}^\top & b\mathbf{1}\mathbf{1}^\top \\ b\mathbf{1}\mathbf{1}^\top & \mathbf{1}\mathbf{1}^\top \end{pmatrix},$$

where $a, b \in (0, 1/2)$ and $\mathbf{1}$ denotes a vector of ones of length k . Since $R^2 < p$, we now construct the block diagonal covariance matrices

$$\Sigma_m^{(k)} = \text{diag}(C_1, C_2, \dots, C_M, I_{p-2kM}), \quad m = 1, 2, \dots, M$$

where the m -th diagonal block is chosen to be $C_m = B^{(k)}$ while others are $C_i = A^{(k)}$ for $i \neq m$ and $M = \lfloor p/R \rfloor$. Also define $\Sigma_0^{(k)}$ to be of the same structure with $C_i = A^{(k)}$ for all i . Then, we have $\Sigma_m^{(R/2)} \in \mathcal{G}_q(R)$ for $m = 0, 1, \dots, M$, since each row of $\Sigma_m^{(R/2)}$ only contains at most R nonzero elements that are bounded by 1.

Let \mathbb{P}_0 denote the distribution of $X \sim \mathcal{N}_p(0, \Sigma_0^{(R/2)})$ and \mathbb{P}_m denote the distribution of $X \sim \mathcal{N}_p(0, \Sigma_m^{(R/2)})$. Let \mathbb{P}_0^n (resp. \mathbb{P}_m^n) denote the distribution of $\mathbf{X} = (X_1, \dots, X_n)$ of n i.i.d random variables drawn from \mathbb{P}_0 (resp. \mathbb{P}_m). Moreover, let $\bar{\mathbb{P}}^n$ denote the uniform mixture of \mathbb{P}_m^n over $m \in [M]$. Consider the testing problem:

$$\begin{aligned} H_0 &: \mathbf{X} \sim \mathbb{P}_0^n \\ H_1 &: \mathbf{X} \sim \bar{\mathbb{P}}^n. \end{aligned}$$

Using [Tsy09, Theorem 2.2, part (iii)] as before, we need to show χ^2 -divergence can be bounded by a constant. By the same calculation as in Lemma B.1, we have

$$\chi^2(\bar{\mathbb{P}}^n, \mathbb{P}_0^n) = \frac{1}{M^2} \sum_{1 \leq i, j \leq M} |I - [(\Sigma_0^{(1)})^{-1} \Sigma_i^{(1)} - I][(\Sigma_0^{(1)})^{-1} \Sigma_j^{(1)} - I]|^{-n/2} - 1.$$

Note that χ^2 -divergence here depends on $\Sigma_m^{(1)}$ instead of $\Sigma_m^{(R/2)}$ since perfectly correlated random variables do not add additional information and hence do not affect χ^2 -divergence (see Section B.2). Using the definition of $\Sigma_m^{(1)}$'s, we obtain

$$\begin{aligned} & |I - [(\Sigma_0^{(1)})^{-1}\Sigma_i^{(1)} - I][(\Sigma_0^{(1)})^{-1}\Sigma_j^{(1)} - I]| \\ &= \begin{cases} 1 - 2(1+a^2)\left(\frac{a-b}{1-a^2}\right)^2 + \frac{(a-b)^4}{(1-a^2)^2}, & \text{if } i = j \\ 1, & \text{otherwise.} \end{cases} \end{aligned}$$

Therefore,

$$\chi^2(\bar{\mathbb{P}}^n, \mathbb{P}_0^n) = \frac{1}{M} \left\{ \left(1 - 2(1+a^2)\left(\frac{a-b}{1-a^2}\right)^2 + \frac{(a-b)^4}{(1-a^2)^2} \right)^{-n/2} - 1 \right\},$$

which is bounded by $((1-5(a-b)^2)^{-n/2}-1)/M$ due to the fact $2(1+a^2)/(1-a^2)^2 \leq 5$ for $a \leq 1/2$. Now choose

$$a = \frac{1}{4}, b = a + \frac{1}{4}\sqrt{\frac{\log p}{n}}.$$

By assumption, there exists a constant $c_0 > 1$ such that $R^2 \leq c_0 p$. Thus,

$$\chi^2(\bar{\mathbb{P}}^n, \mathbb{P}_0^n) \leq \frac{1}{M} \left\{ \left(1 - \frac{\log p}{2n} \right)^{-n/2} - 1 \right\} \leq \frac{R}{p} e^{\log p/2} \leq \sqrt{c_0}.$$

Using standard techniques to reduce estimation problems to testing problems as before, we find

$$\inf_{\hat{L}} \max_{\Sigma \in \{\Sigma_m^{(R/2)} : m=0, \dots, M\}} \mathbb{E}_{\Sigma} [(\hat{L} - \ell_r(\Sigma))^2] \geq C(\ell_r(\Sigma_0^{(R/2)}) - \ell_r(\Sigma_1^{(R/2)}))^2.$$

For the above choice of $\Sigma_m^{(k)}$, we have

$$(\ell_r(\Sigma_0^{(R/2)}) - \ell_r(\Sigma_1^{(R/2)}))^2 = \frac{R^2}{4}(b^r - a^r)^2 \geq CR^2 \frac{\log p}{n}.$$

Since $\Sigma_m^{(R/2)} \in \mathcal{G}_q$, the above two displays imply that

$$\inf_{\hat{L}} \max_{\Sigma \in \mathcal{G}_q} \mathbb{E}_{\Sigma} [(\hat{L} - \ell_r(\Sigma))^2] \geq C \frac{R^2 \log p}{n},$$

which together with the other part of the lower bound, completes the proof of the theorem.

REFERENCES

- [ACBL14] Ery Arias-Castro, Sébastien Bubeck, and Gábor Lugosi, *Detecting positive correlations in a multivariate sample*, Bernoulli (to appear) (2014).
- [AW09] Arash A. Amini and Martin J. Wainwright, *High-dimensional analysis of semidefinite relaxations for sparse principal components*, Annals of Statistics **37** (2009), no. 5B, 2877–2921.

- [BJNP13] Aharon Birnbaum, Iain M. Johnstone, Boaz Nadler, and Debashis Paul, *Minimax bounds for sparse PCA with noisy high-dimensional data*, *Ann. Statist.* **41** (2013), no. 3, 1055–1084.
- [BL04] Bickel, P.J. and Levina, E. (2004). Some theory for Fisher’s linear discriminant, ‘naive Bayes’, and some alternatives when there are many more variables than observations. *Bernoulli*, **10**, 989–1010.
- [BL08a] Peter J. Bickel and Elizaveta Levina, *Covariance regularization by thresholding*, *Annals of Statistics* **36** (2008), no. 6, 2577–2604.
- [BL08b] Peter J. Bickel and Elizaveta Levina, *Regularized estimation of large covariance matrices*, *Ann. Statist.* **36** (2008), no. 1, 199–227. [MR2387969 \(2009a:62255\)](#)
- [BM11] Cristina Butucea and Katia Meziani, *Quadratic functional estimation in inverse problems*, *Stat. Methodol.* **8** (2011), no. 1, 31–41. [MR2741507](#)
- [BR88] P. J. Bickel and Y. Ritov, *Estimating integrated squared density derivatives: sharp best order of convergence estimates*, *Sankhyā Ser. A* **50** (1988), no. 3, 381–393. [MR1065550 \(91e:62079\)](#)
- [BR13a] Quentin Berthet and Philippe Rigollet, *Complexity theoretic lower bounds for sparse principal component detection*, COLT 2013 - The 26th Conference on Learning Theory, Princeton, NJ, June 12–14, 2013 (Shai Shalev-Shwartz and Ingo Steinwart, eds.), *JMLR W&CP*, vol. 30, 2013, pp. 1046–1066.
- [BR13b] Quentin Berthet and Philippe Rigollet, *Optimal detection of sparse principal components in high dimension*, *Ann. Statist.* **41** (2013), no. 1, 1780–1815.
- [BS96] Zhidong Bai and Hewa Saranadasa, *Effect of high dimension: by an example of a two sample problem*, *Statist. Sinica* **6** (1996), no. 2, 311–329. [MR1399305 \(97i:62062\)](#)
- [But07] Cristina Butucea, *Goodness-of-fit testing and quadratic functional estimation from indirect observations*, *Ann. Statist.* **35** (2007), no. 5, 1907–1930. [MR2363957 \(2009e:62182\)](#)
- [CL05] T. Tony Cai and Mark G. Low, *Nonquadratic estimators of a quadratic functional*, *Ann. Statist.* **33** (2005), no. 6, 2930–2956. [MR2253108 \(2007k:62058\)](#)
- [CL06] ———, *Optimal adaptive estimation of a quadratic functional*, *Ann. Statist.* **34** (2006), no. 5, 2298–2325. [MR2291501 \(2008m:62054\)](#)
- [CL11] T. Tony Cai and Weidong Liu, *Adaptive thresholding for sparse covariance matrix estimation*, *J. Amer. Statist. Assoc.* **106** (2011), no. 494, 672–684. [MR2847949 \(2012i:62143\)](#)
- [CMW13] T. Tony Cai, Zongming Ma, and Yihong Wu, *Sparse PCA: Optimal rates and adaptive estimation*, *The Annals of Statistics* **41** (2013), no. 6, 3074–3110.
- [CMW14] T. Tony Cai, Zongming Ma, and Yihong Wu, *Optimal estimation and rank detection for sparse spiked covariance matrices*, *Probab. Theory Related Fields* (to appear) (2014).
- [CQ10] Song Xi Chen and Ying-Li Qin, *A two-sample test for high-dimensional data with applications to gene-set testing*, *Ann. Statist.* **38** (2010), no. 2, 808–835. [MR2604697 \(2011c:62187\)](#)

- [CRZ13] T. Tony Cai, Zhao Ren, and Harrison H. Zhou, *Optimal rates of convergence for estimating Toeplitz covariance matrices*, Probab. Theory Related Fields **156** (2013), no. 1-2, 101–143. [MR3055254](#)
- [CY12] T. Tony Cai and Ming Yuan, *Adaptive covariance matrix estimation through block thresholding*, Ann. Statist. **40** (2012), no. 4, 2014–2042. [MR3059075](#)
- [CZ12] T. Tony Cai and Harrison H. Zhou, *Minimax estimation of large covariance matrices under ℓ_1 -norm*, Statist. Sinica **22** (2012), no. 4, 1319–1349. [MR3027084](#)
- [CZZ10] T. Tony Cai, Cun-Hui Zhang, and Harrison H. Zhou, *Optimal rates of convergence for covariance matrix estimation*, Ann. Statist. **38** (2010), no. 4, 2118–2144. [MR2676885](#) ([2011h:62215](#))
- [DN90] David L Donoho and Michael Nussbaum, *Minimax quadratic estimation of a quadratic functional*, Journal of Complexity **6** (1990), no. 3, 290 – 323.
- [EL96] Sam Efromovich and Mark Low, *On optimal adaptive estimation of a quadratic functional*, Ann. Statist. **24** (1996), no. 3, 1106–1125. [MR1401840](#) ([97k:62078](#))
- [Fan91] Jianqing Fan, *On the estimation of quadratic functionals*, Ann. Statist. **19** (1991), no. 3, 1273–1294. [MR1126325](#) ([92j:62006](#))
- [FF93] Eugene F. Fama and Kenneth R. French, *Common risk factors in the returns on stocks and bonds*, Journal of Financial Economics **33** (1993), no. 1, 3 – 56.
- [FFL08] Jianqing Fan, Yingying Fan, and Jinchi Lv, *High dimensional covariance matrix estimation using a factor model*, J. Econometrics **147** (2008), no. 1, 186–197. [MR2472991](#) ([2010b:62199](#))
- [FLM11] Jianqing Fan, Yuan Liao, and Martina Mincheva, *High-dimensional covariance matrix estimation in approximate factor models*, Ann. Statist. **39** (2011), no. 6, 3320–3356. [MR3012410](#)
- [FLM13] ———, *Large covariance estimation by thresholding principal orthogonal complements*, J. R. Stat. Soc. Ser. B. Stat. Methodol. **75** (2013), no. 4, 603–680, With 33 discussions by 57 authors and a reply by Fan, Liao and Mincheva. [MR3091653](#)
- [FR13] Simon Foucart and Holger Rauhut, *A mathematical introduction to compressive sensing*, Applied and Numerical Harmonic Analysis, Birkhäuser/Springer, New York, 2013. [MR3100033](#)
- [GRS89] Michael R. Gibbons, Stephen A. Ross, and Jay Shanken, *A test of the efficiency of a given portfolio*, Econometrica **57** (1989), no. 5, pp. 1121–1152.
- [HM87] Peter Hall and J. S. Marron, *Estimation of integrated squared density derivatives*, Statist. Probab. Lett. **6** (1987), no. 2, 109–115. [MR907270](#) ([88h:62061](#))
- [INK87] I. Ibragimov, A. Nemirovskii, and R. Khasminskii, *Some problems on nonparametric estimation in gaussian white noise*, Theory of Probability & Its Applications **31** (1987), no. 3, 391–406.
- [JL09] Iain M. Johnstone and Arthur Yu Lu, *On consistency and sparsity for principal components analysis in high dimensions*, J. Amer. Statist. Assoc. **104** (2009), no. 486, 682–693. [MR2751448](#)

- [Kar08] Noureddine El Karoui, *Operator norm consistent estimation of large-dimensional sparse covariance matrices*, *Annals of Statistics* **36** (2008), no. 6, 2717–2756.
- [Kle06] Jussi Klemelä, *Sharp adaptive estimation of quadratic functionals*, *Probab. Theory Related Fields* **134** (2006), no. 4, 539–564. [MR2214904 \(2007b:62039\)](#)
- [LF09] Clifford Lam and Jianqing Fan, *Sparsistency and rates of convergence in large covariance matrix estimation*, *Ann. Statist.* **37** (2009), no. 6B, 4254–4278. [MR2572459 \(2011d:62064\)](#)
- [Lin65] John Lintner, *The valuation of risk assets and the selection of risky investments in stock portfolios and capital budgets*, *The Review of Economics and Statistics* **47** (1965), no. 1, pp. 13–37.
- [LM00] B. Laurent and P. Massart, *Adaptive estimation of a quadratic functional by model selection*, *Ann. Statist.* **28** (2000), no. 5, 1302–1338. [MR1805785 \(2002c:62052\)](#)
- [LV12] Elizaveta Levina and Roman Vershynin, *Partial estimation of covariance matrices*, *Probab. Theory Related Fields* **153** (2012), no. 3-4, 405–419. [MR2948681](#)
- [Ma13] Zongming Ma, *Sparse principal component analysis and iterative thresholding*, *The Annals of Statistics* **41** (2013), no. 2, 772–801.
- [Mos66] Jan Mossin, *Equilibrium in a capital asset market*, *Econometrica* **34** (1966), no. 4, pp. 768–783.
- [Nem00] Arkadi Nemirovski, *Topics in non-parametric statistics*, *Lectures on probability theory and statistics (Saint-Flour, 1998)*, *Lecture Notes in Math.*, vol. 1738, Springer, Berlin, 2000, pp. 85–277. [MR1775640 \(2001h:62074\)](#)
- [NK87] A. S. Nemirovskii and R. Z. Khasminskii, *Nonparametric estimation of the functionals of the products of a signal observed in white noise*, *Problemy Peredachi Informatsii* **23** (1987), no. 3, 27–38. [MR914348 \(90b:62046\)](#)
- [OMH13] Alexei Onatski, Marcelo J. Moreira, and Marc Hallin, *Asymptotic power of sphericity tests for high-dimensional data*, *The Annals of Statistics* **41** (2013), no. 3, 1204–1231.
- [PJ12] Debashis Paul and Iain M. Johnstone, *Augmented sparse principal component analysis for high dimensional data*, *Arxiv:1202.1242v1* (2012).
- [PY12] M Hashem Pesaran and Takashi Yamagata, *Testing capm with a large number of assets*, *IZA Discussion Papers 6469*, Institute for the Study of Labor (IZA), 2012.
- [RLZ09] Adam J. Rothman, Elizaveta Levina, and Ji Zhu, *Generalized thresholding of large covariance matrices*, *J. Amer. Statist. Assoc.* **104** (2009), no. 485, 177–186. [MR2504372 \(2010g:62186\)](#)
- [RT12] Philippe Rigollet and Alexandre Tsybakov, *Comment: “Minimax estimation of large covariance matrices under ℓ_1 -norm”*, *Statist. Sinica* **22** (2012), no. 4, 1358–1367. [MR3027087](#)
- [RWR11] Pradeep Ravikumar, Martin J. Wainwright, Garvesh Raskutti, and Bin Yu, *High-dimensional covariance estimation by minimizing ℓ_1 -penalized log-determinant divergence*, *Electron. J. Stat.* **5** (2011), 935–

980. [MR2836766](#)
- [Sha64] William F. Sharpe, *Capital asset prices: A theory of market equilibrium under conditions of risk*, The Journal of Finance **19** (1964), no. 3, pp. 425–442.
- [SSM09] Dan Shen, Haipeng Shen, and J.S. Marron, *Consistency of sparse PCA in high dimension, low sample size contexts*, Annals of Statistics **37** (2009), no. 6B, 4104–4130.
- [Tsy09] Alexandre B. Tsybakov, *Introduction to nonparametric estimation*, Springer Series in Statistics, Springer, New York, 2009, Revised and extended from the 2004 French original, Translated by Vladimir Zaiats. [MR2724359 \(2011g:62006\)](#)
- [Ver12] Nicolas Verzelen, *Minimax risks for sparse regressions: Ultra-high-dimensional phenomena*, Electron. J. Stat. **6** (2012), 38–90. [MR2879672](#)
- [VL12] Vincent Vu and Jing Lei, *Minimax rates of estimation for sparse PCA in high dimensions*, Proceedings of the Fifteenth International Conference on Artificial Intelligence and Statistics April 21-23, 2012 La Palma, Canary Islands., JMLR W&CP, vol. 22, 2012, pp. 1278–1286.
- [ZHT06] Hui Zou, Trevor Hastie, and Robert Tibshirani, *Sparse principal component analysis*, Journal of Computational and Graphical Statistics **15** (2006), no. 2, 265–286 (English).

DEPARTMENT OF OPERATIONS RESEARCH
AND FINANCIAL ENGINEERING
PRINCETON UNIVERSITY
PRINCETON, NJ 08544, USA
{JQFAN, RIGOLLET, WEICHENW}@PRINCETON.EDU