

A Combinatorial Curvature Flow for Compact 3-Manifolds with Boundary

Feng Luo

abstract

We introduce a combinatorial curvature flow for PL metrics on compact triangulated 3-manifolds with boundary consisting of surfaces of negative Euler characteristic. The flow tends to find the complete hyperbolic metric with totally geodesic boundary on a manifold. Some of the basic properties of the combinatorial flow are established. The most important ones is that the evolution of the combinatorial curvature satisfies a combinatorial heat equation. It implies that the total curvature decreases along the flow. The local convergence of the flow to the hyperbolic metric is also established if the triangulation is isotopic to a totally geodesic triangulation.

§1. Introduction

1.1. The purpose of this paper is to announce the construction of a combinatorial curvature flow which is a 3-dimensional analogy of the work of [CL]. In [CL], we introduced a 2-dimensional combinatorial curvature flow for triangulated surfaces of non-positive Euler characteristic. It is shown that for any initial choice of PL metric of circle packing type, the flow exists for all time and converges exponentially fast to the Andreev-Koebe-Thurston's circle packing metrics.


The basic building blocks for the 2-dimensional flow are hyperbolic and Euclidean triangles. In the 3-dimensional case, the basic building blocks are the *hyperideal tetrahedra* discovered by Bao and Bonahon [BB]. Given an ideal triangulation of a compact 3-manifold with boundary consisting of surfaces of negative Euler characteristic, we replace each (truncated) tetrahedron by a hyperideal tetrahedron by assigning the edge lengths. The isometric gluing of these tetrahedra gives a hyperbolic cone metric on the 3-manifold. The *PL curvature* of the cone metric at an edge is 2π less than the sum of dihedral angles at the edge. The combinatorial curvature flow that we propose is the following system of ordinary differential equations,

$$(1.1) \quad dx_i/dt = K_i$$

where x_i is the length of the i -th edge and K_i is the curvature of the cone metric (x_1, \dots, x_n) at the i -th edge. The equation (1.1) captures the essential features of the 2-dimensional combinatorial Ricci flow in [CL]. The most important of all is that the PL curvature evolves according to a combinatorial heat equation. Thus the corresponding maximal principle applies. The flow has the tendency of finding the complete hyperbolic metric of totally geodesic boundary on the manifold. By analyzing the singularity formations in equation (1.1), it is conceivable that one could give a new proof of Thurston's geometrization theorem for these manifolds using (1.1). Furthermore, the flow (1.1) will be a useful tool to find algorithmically the complete hyperbolic metric.

1.2. We now provide some details of the approach. Suppose M is a compact 3-manifold with boundary consisting of surfaces of negative Euler characteristic. An *ideal triangulation* (or truncated triangulation) of M is the following finite set of data. Take a finite collection of 3-simplexes and identify their faces in pairs by homeomorphisms. The quotient space with a small regular neighborhood of each vertex removed is homeomorphic to the 3-manifold M . By abuse the use of language, we will call (the homeomorphism images of) i -dimensional cells ($i \geq 1$)

of the ideal triangulation (truncated triangulation) in the interior of M , the *edges, triangles and tetrahedra*. A *hyperideal tetrahedron* in the 3-dimensional hyperbolic space is a compact convex polyhedron so that it is diffeomorphic to a truncated tetrahedron in the 3-dimensional Euclidean space and its four hexagonal faces are right-angled hyperbolic hexagons. (Note that two compact subsets of \mathbf{R}^n are diffeomorphic if there is a diffeomorphism between two open neighborhoods of them sending one compact set to the other). See the figure below. In the beautiful work of [BB], Bao and Bonahon give a complete characterization of hyperideal convex polyhedra. As a consequence of their work, hyperideal tetrahedra are completely characterized by their six dihedral angles at the six edges so that the sum of three dihedral angles associated to edges adjacent to each vertex is less than π .



The six edges of a hyperideal tetrahedron are the intersections of its hexagonal faces

figure

One of our main technical observations is the following,

Theorem 1. *The volume of a hyperideal tetrahedron is a strictly convex function of its dihedral angles.*

Note that each hyperideal tetrahedron is also determined by its six edge lengths. By the Schlaefli formula, an equivalent statement to theorem 1 is that the Hessian matrix of the map sending six edge lengths to the six dihedral angle is strictly positive definite. This positive definite matrix provides a basis for constructing the combinatorial Laplacian operator for the curvature evolution equation.

Now given an ideal triangulated 3-manifold (M, T) , let E be the set of edges in the triangulation and let n be the number of edges in E . An assignment $x : E \rightarrow \mathbf{R}_{>0}$ is called a *hyperbolic cone metric associated to the triangulation T* if for each tetrahedron t in T with edges e_1, \dots, e_6 , the six numbers $x_i = x(e_i)$ ($i=1, \dots, 6$) are the edge lengths of a hyperideal tetrahedron in \mathbf{H}^3 . The set of all hyperbolic cone metrics associated to T is denoted by $L(M, T)$ which will be considered as an open subset of $\mathbf{R}^n = \mathbf{R}^E$. The PL curvature of a cone metric $x \in L(M, T)$ is the map $K(x) : E \rightarrow \mathbf{R}$ sending an edge e to the PL curvature of x at the edge e . Again we identify the set of all PL curvatures $\{K(x) | x \in L(M, T)\}$ with a subset of \mathbf{R}^n . The combinatorial curvature flow is the vector field in $L(M, T)$ defined by equation (1.1) where K_i in the right-hand side is the PL curvature of the metric $x = (x_1, \dots, x_n)$ at time t at the i -th edge.

Theorem 2. *For any ideal triangulated 3-manifold, under the combinatorial curvature flow*

(1.1), the PL curvature $K_i(t)$ evolves according to a combinatorial heat equation,

$$(1.2) \quad dK_i(t)/dt = \sum_{j=1}^n a_{ij}K_j(t)$$

where the matrix $[a_{ij}]_{n \times n}$ is symmetric negative definite. In particular, the total curvature $\sum_{i=1}^n K_i^2(t)$ is strictly decreasing along the flow unless $K_i(t) = 0$ for all i .

Theorem 3. *For any ideal triangulated 3-manifold (M, T) , the equilibrium points of the combinatorial curvature flow (1.1) are the complete hyperbolic metric with totally geodesic boundary. Furthermore, each equilibrium point is a local attractor of the flow.*

Another consequence of the convexity is the following local rigidity result for hyperbolic cone metrics without constraints on cone angles. Note that by [HK], hyperbolic cone metrics with cone angles at most 2π are locally rigid.

Theorem 4. *For any ideal triangulated 3-manifold (M, T) , the curvature map $\Pi : L(M, T) \rightarrow \mathbf{R}^n$ sending a metric x to its curvature $K(x)$ is a local diffeomorphism. In particular, a hyperbolic cone metric associated to an ideal triangulation is locally determined by its cone angles.*

In the rest of the note, we will sketch briefly the ideas of the proofs of the results. In the last section, we propose several questions related to the combinatorial curvature flow whose resolution will lead to a new proof of Thurston's geometrization theorem for this class of 3-manifolds.

1.3. Acknowledgment. We thank Ben Chow for conversations on the topic of Ricci flow and X.S. Lin for discussions. The work is supported in part by the NSF and a research grant from the Rutgers University.

§2. Sketch of the Proofs

2.1. The key step in the proof of all theorems is to establish the convexity theorem 1. Here is a quick way to see it. Suppose x_1, \dots, x_6 are the lengths of the six edges of a hyperideal tetrahedron so that the corresponding dihedral angles are a_1, \dots, a_6 . Let $a = (a_1, \dots, a_6)$, $x = (x_1, \dots, x_6)$ and V be the volume of the hyperideal tetrahedron. Then by using cosine laws for hyperbolic triangles and hyperbolic right-angled hexagons, we see that both functions $x = x(a)$ and $a = a(x)$ are smooth in a and x respectively. In particular, they are diffeomorphisms. By the Schläfli formula $\partial V / \partial a_i = -x_i/2$, the Jacobian matrix $[\partial x_i / \partial a_j]_{6 \times 6}$ is symmetric. The Jacobian matrix is non-singular due to the fact that the map $a = a(x)$ is a diffeomorphism. Thus its signature is independent of the choice of hyperideal tetrahedra. One checks directly that in the case the hyperideal tetrahedron is regular (i.e., all x_i 's are the same and all a_i 's are the same), the Jacobian matrix is positive definite. Thus theorem 1 follows. Since the inverse of a symmetric positive definite matrix is again symmetric and positive definite, we see that the matrix $[\partial a_i / \partial x_j]_{6 \times 6}$ is also symmetric and positive definite.

2.2. To prove theorem 2, one uses the equation (1.1). Namely, to understand the evolution of the curvature, it suffices to understand the evolution of the individual dihedral angle $a_i(t)$. By the chain rule, we have,

$$(2.1) \quad da_i/dt = \sum_{j=1}^6 (\partial a_i / \partial x_j) dx_j/dt = \sum_{j=1}^6 (\partial a_i / \partial x_j) K_j.$$

By the discussion above, the matrix $[\partial a_i / \partial x_j]_{6 \times 6}$ is positive definite. By summing over all dihedral angles adjacent to a fixed edge, we obtain theorem 2 since the sum of semi-negative definite matrices are still semi-negative definite. A further study shows that the resulting matrix is strictly negative definite.

2.3. Both theorems 3 and 4 follow from the fact that the combinatorial curvature flow (1.1) is the negative gradient flow of a strictly convex function defined on the space $L(M, T)$. Indeed, by the Schlaefli formula $dV = -1/2 \sum_i x_i da_i$, if we form $H = 2V + \sum_{i=1}^6 a_i x_i$, then $\partial H / \partial x_i = a_i$. Thus the function H is a strictly convex function of the edge lengths x_1, \dots, x_6 . For any hyperbolic cone metric x associated to the ideal triangulation T , we define

$$(2.2) \quad H(x) = 2\text{vol}(x) - \sum_i K_i x_i$$

where $\text{vol}(x)$ is the volume of the metric. One checks easily that $H(x)$ is a strictly convex function of x and the gradient of $H(x)$ is $-(K_1, \dots, K_n)$. Thus theorem 3 follows from the standard theory of ordinary differential equation. To see theorem 4, one uses the fact that the map sending a point to the gradient of a strictly smooth convex function is a local diffeomorphism.

§3. Some Remarks and Questions

This work and [CL] are motivated by the work of Richard Hamilton [Hi] on Ricci flow. The strategy of [Hi] is to find a flow deforming metrics so that its curvature evolves according to a heat-type equation. The main focus of study then shifts from the evolution of the metrics to the evolution of its curvature using the maximal principle for heat equations. As long as one has control of the curvature evolution, then one gets some control of the metric evolutions by study either the singularity formation or the long time convergence. Theorem 2 above seems to indicate that the combinatorial flow (1.1) deforms the cone metrics in the "right" direction. There remains the task of understanding the singularity formations in (1.1) which corresponds to the degeneration of the hyperideal tetrahedra. This is being investigated. Below are some thoughts on this topic. The motivations come from the work of [Th], [CV], [Riv2], [Lei] and [CL].

3.1. To understand the singularity formation, we will focus our attention to the function H in (2.2). Following [CV] and [Riv2], our goal is to find the (linear) conditions on the ideal triangulation so that it will guarantee the existence of critical points for H . The existence of the ideal triangulation satisfying the (linear) condition will be related to the topology of the 3-manifold and will be resolved by topological arguments.

Suppose (M, T) is an ideal triangulated compact 3-manifold so that each boundary component of M has negative Euler characteristic. A pair (e, t) where e is an edge and t is a tetrahedron containing e is called a *corner* in T . Following Rivin [Riv1], Casson and Lackenby [La1], we say the triangulated manifold (M, T) supports a *linear hyperbolic structure* if one can assign each corner of T a positive number called the *dihedral angle* so that (1) the sum of dihedral angles of all corners adjacent to each fixed edge is 2π , and (2) the sum of dihedral angles of every triple of corners $(e_1, t), (e_2, t), (e_3, t)$ where e_1, e_2, e_3 are adjacent to a fixed vertex is less than π . By the work of [BB], given a linear hyperbolic structure on (M, T) , we can realize each individual tetrahedron by a hyperideal tetrahedron whose dihedral angles are the assigned numbers so that the sum of the dihedral angles at each edge is 2π . It can be shown that if (M, T) supports a linear hyperbolic structure, then the manifold M is irreducible

without incompressible tori. One would ask if the converse is also true. The work of Lackenby [La2] gives some evidences that the following may have a positive answer. See also the work of [KR].

Question 1. *Suppose M is a compact irreducible 3-manifold with incompressible boundary consisting of surfaces of negative Euler characteristic. If M contains no incompressible tori and annuli, is there any ideal triangulation of M which supports a linear hyperbolic structure?*

The next question is the 3-dimensional analogy of the 2-dimensional singularity formation analysis in the work of [CV], [Riv2] and [Le].

Question 2. *Suppose (M, T) is an ideal triangulated 3-manifold which supports a linear hyperbolic structure. Does H have a local minimal point in the space $L(M, T)$ of all cone metrics associated to (M, T) ?*

Positive resolutions of above two questions will produce a new proof of Thurston's geometrization theorem for this class of 3-manifolds.

3.2. We remark that the moduli space of all hyperideal tetrahedra parametrized by their edge lengths x_1, \dots, x_6 is not a convex subset of \mathbf{R}^6 . This is the main reason that we have only local convergence and local rigidity in theorems 3 and 4. However, it is conceivable that theorem 4 may still be true globally. We do not know yet if the space $L(M, T)$ of all cone metrics associated to the ideal triangulated manifold is homeomorphic to a Euclidean space. It is likely to be the case. In fact, one would hope that there is a diffeomorphism $h : \mathbf{R}_{>0} \rightarrow \mathbf{R}_{>0}$ so that if we parameterize the space of all hyperideal tetrahedra by $(t_1, \dots, t_6) = (h(x_1), \dots, h(x_6))$, then the space becomes convex in t -coordinate. Evidently, if this holds, it implies the space $L(M, T)$ is convex in the t -coordinate.

References

- [BB] Bao, Xiliang; Bonahon, Francis: Hyperideal polyhedra in hyperbolic 3-space. *Bull. Soc. Math. France* 130 (2002), no. 3, 457–491.
- [CL] Chow, Bennett; Luo, Feng: Combinatorial Ricci flows on surfaces. *J. Differential Geom.* 63 (2003), no. 1, 97–129.
- [CV] Colin de Verdière, Yves: Un principe variationnel pour les empilements de cercles. *Invent. Math.* 104 (1991), no. 3, 655–669.
- [Hi] Hamilton, Richard S. Three-manifolds with positive Ricci curvature. *J. Differential Geom.* 17 (1982), no. 2, 255–306.
- [HK] Hodgson, Craig D.; Kerckhoff, Steven P: Rigidity of hyperbolic cone-manifolds and hyperbolic Dehn surgery. *J. Differential Geom.* 48 (1998), no. 1, 1–59.
- [KR] Kang, Ensil and JH Rubinstein, J. H: ideal triangulations of 3-manifolds I, preprint.
- [La1] Lackenby, Marc: Word hyperbolic Dehn surgery. *Invent. Math.* 140 (2000), no. 2, 243–282.
- [La2] Lackenby, Marc: Taut ideal triangulations of 3-manifolds. *Geom. Topol.* 4 (2000), 369–395 (electronic).
- [Le] Leimon, Gregory: Characterizing the Delaunay decompositions of compact hyperbolic surfaces. *Geom. Topol.* 6 (2002), 361–391 (electronic).
- [Ri1] Rivin, Igor: Combinatorial optimization in geometry. *Adv. in Appl. Math.* 31 (2003), no. 1, 242–271.

[Ri2] Rivin, Igor: Euclidean structures on simplicial surfaces and hyperbolic volume. *Ann. of Math.* (2) 139 (1994), no. 3, 553–580.

[Th] Thurston, William: *Topology and geometry of 3-manifolds*, Lecture notes, Princeton University, 1978.

Department of Mathematics, Rutgers University, Piscataway, NJ 07059

Email: fluo@math.rutgers.edu