

EVOLUTION OF ANGULAR MOMENTUM AND CENTER OF MASS AT NULL INFINITY

PO-NING CHEN, JORDAN KELLER,
MU-TAO WANG, YE-KAI WANG, AND SHING-TUNG YAU

ABSTRACT. We study how conserved quantities such as angular momentum and center of mass evolve with respect to the retarded time at null infinity, which is described in terms of a Bondi-Sachs coordinate system. These evolution formulae complement the classical Bondi mass loss formula for gravitational radiation. They are further expressed in terms of the potentials of the shear and news tensors. The consequences that follow from these formulae are (1) Supertranslation invariance of the fluxes of the CWY conserved quantities. (2) A conservation law of angular momentum à la Christodoulou. (3) A duality paradigm for null infinity. In particular, the supertranslation invariance distinguishes the CWY angular momentum and center of mass from the classical definitions.

1. INTRODUCTION

In this article, we study the evolution of angular momentum and center of mass at null infinity of asymptotically flat vacuum spacetimes. These evolution formulae complement the classical Bondi mass loss formula for gravitational radiations. We are particularly interested in the total flux of angular momentum and center of mass.

For a good notion of conserved quantities, one expects that the total flux is independent of the choice of coordinate systems. However, as indicated by Penrose [19], the notion of “angular momentum carried away by gravitational radiation” can be shifted by supertranslations, an infinite dimensional symmetry at null infinity. Such ambiguity has been a crucial obstacle to a clear understanding of conserved quantities at null infinity. In this article, we consider both the classical and the Chen-Wang-Yau (CWY) [4] definitions for angular momentum and center of mass at null infinity. A key result

P.-N. Chen is supported by NSF grant DMS-1308164 and Simons Foundation collaboration grant #584785, M.-T. Wang is supported by NSF grant DMS-1810856, Y.-K. Wang is supported by MOST Taiwan grant 107-2115-M-006-001-MY2, 109-2628-M-006-001 -MY3 and S.-T. Yau is supported by NSF grants PHY-0714648 and DMS-1308244. The authors would like to thank the National Center for Theoretical Sciences at National Taiwan University where part of this research was carried out. This material is based upon work supported by the National Science Foundation under Grant No. DMS-1810856.

is the supertranslation invariance of the flux of the CWY angular momentum and center of mass. This invariance distinguishes the CWY definitions from the classical definitions.

Consider the future null infinity \mathcal{I}^+ of an asymptotically flat spacetime, which is described in terms of a Bondi-Sachs coordinate system. \mathcal{I}^+ is identified with $I \times S^2$, where $I \subset (-\infty, +\infty)$ is an interval parametrized by the retarded time u and S^2 is the standard unit 2-sphere equipped with the standard round metric σ_{AB} . Let m denote the mass aspect, N_A the angular momentum aspect, C_{AB} the shear tensor, and N_{AB} the news tensor of \mathcal{I}^+ . One can view m as a smooth function, N_A a smooth one-form, and C_{AB} and N_{AB} smooth symmetric traceless 2-tensors (with respect to σ_{AB}) on S^2 that depend on u . In particular, $\partial_u C_{AB} = N_{AB}$. See a brief description of \mathcal{I}^+ in the Bondi-Sachs coordinates and the definitions of these quantities in Section 2.

All integrals in this paper on the sphere are taken over the standard two-sphere S^2 with the standard round metric σ_{AB} . We take the standard formulae for energy and linear momentum:

$$\begin{aligned} E &= \int_{S^2} 2m \\ P^k &= \int_{S^2} 2m \tilde{X}^k, \quad k = 1, 2, 3 \end{aligned} \tag{1.1}$$

where $\tilde{X}^k, k = 1, 2, 3$ are the standard coordinate functions on \mathbb{R}^3 restricted to the unit sphere S^2 .

Furthermore, we consider the classical angular momentum

$$\tilde{J}^k = \int_{S^2} \epsilon^{AB} \nabla_B \tilde{X}^k [N_A - \frac{1}{4} C_A^D \nabla^B C_{DB}], \tag{1.2}$$

and the classical center of mass

$$\tilde{C}^k = \int_{S^2} \nabla^A \tilde{X}^k [N_A - u \nabla_A m - \frac{1}{4} C_A^D \nabla^B C_{DB} - \frac{1}{16} \nabla_A (C_{DE} C^{DE})], \tag{1.3}$$

where ∇_A denotes the covariant derivative with respect to σ_{AB} , and ϵ_{AB} denotes the volume form of σ_{AB} and $k = 1, 2, 3$. The indexes are raised, lowered, and contracted with respect to σ_{AB} . Our definition is that of Dray-Streubel [12]. See Section III.B of Flanagan-Nichols [13] for details.

Remark 1.1. In the above definitions of conserved quantities, we omit the constant $\frac{1}{8\pi}$.

Furthermore, we consider the CWY angular momentum J^k and center of mass C^k as the limits of the CWY quasi-local angular momentum and center of mass [4, 5] on \mathcal{I}^+ evaluated in [15].

$$J^k = \int_{S^2} \epsilon^{AB} \nabla_B \tilde{X}^k \left(N_A - \frac{1}{4} C_{AB} \nabla_D C^{DB} - c \nabla_A m \right)$$

$$\begin{aligned}
C^k = & \int_{S^2} \nabla^A \tilde{X}^k \left[N_A - u \nabla_A m - \frac{1}{4} C_{AB} \nabla_D C^{DB} - \frac{1}{16} \nabla_A (C^{DE} C_{DE}) \right. \\
& \left. - c \nabla_A m + 2 \epsilon_{AB} (\nabla^B \underline{c}) m \right] \\
& + \int_{S^2} 3 \tilde{X}^k c m - \frac{1}{4} \tilde{X}^k \nabla_A \underline{F}^{AB} \nabla^D \underline{F}_{DB}
\end{aligned}$$

where c and \underline{c} are the potentials of C_{AB} , as given in (2.9) and $\underline{F}_{AB} = \frac{1}{2}(\epsilon_{AD} \nabla_B \nabla^D \underline{c} + \epsilon_{BD} \nabla_A \nabla^D \underline{c})$. For definiteness, the potentials are assumed to be supported in the $\ell \geq 2$ modes.

In Theorem 11 and Theorem 16 of [15], it is shown that J^k and $\frac{C^k}{E}$ are the limit of the Chen-Wang-Yau quasi-local angular momentum and center of mass (omitting constant $1/8\pi$) under the zero linear momentum assumption

$$\int_{S^2} m(u, x) \tilde{X}^i = 0. \quad (1.4)$$

The CWY angular momentum and center of mass modify the classical definitions as follows:

$$\begin{aligned}
J^k &= \tilde{J}^k - \int_{S^2} \epsilon^{AB} \nabla_B \tilde{X}^k c \nabla_A m \\
C^k &= \tilde{C}^k + \int_{S^2} \nabla^A \tilde{X}^k (-c \nabla_A m + 2 \epsilon_{AB} (\nabla^B \underline{c}) m) \\
&+ \int_{S^2} 3 \tilde{X}^k c m - \frac{1}{4} \tilde{X}^k \nabla_A \underline{F}^{AB} \nabla^D \underline{F}_{DB}
\end{aligned} \quad (1.5)$$

The correction terms come from solving the optimal isometric embedding equation in the theory of Wang-Yau quasilocal mass [25, 26] and are non-local. They provide the reference terms that are critical in the Hamiltonian approach of defining conserved quantities. See [16] for a definition of angular momentum in the context of perturbations of Kerr, in which the referencing is achieved by the uniformization theorem.

The ten conserved quantities $(E, P^k, \tilde{J}^k, \tilde{C}^k)$, or (E, P^k, J^k, C^k) , are functions on I that depend on the retarded time u . We compute the derivatives of these conserved quantities with respect to u . In particular, for the classical angular momentum and center of mass, we obtain

Theorem 1.2. *The classical angular momentum \tilde{J}^k and center of mass \tilde{C}^k , $k = 1, 2, 3$ evolve according to the following:*

$$\partial_u \tilde{J}^k = \frac{1}{4} \int_{S^2} \left[\epsilon^{AE} \nabla_E \tilde{X}^k (C_{AB} \nabla_D N^{BD} - N_{AB} \nabla_D C^{BD}) + \tilde{X}^k \epsilon^{AB} (C_A{}^D N_{DB}) \right], \quad (1.6)$$

$$\partial_u \tilde{C}^k = \frac{1}{4} \int_{S^2} \left[\nabla^A \tilde{X}^k \left(\frac{u}{2} \nabla_A |N|^2 + C_{AB} \nabla_D N^{BD} - N_{AB} \nabla_D C^{BD} \right) \right]. \quad (1.7)$$

The evolution formulae (1.6) and (1.7) can be further expressed in terms of the potentials of C_{AB} and N_{AB} :

Theorem 1.3. *Suppose c and \underline{c} are the potentials of C_{AB} and n and \underline{n} are the potentials of N_{AB} , as given in (2.9) and (2.10), then*

$$\begin{aligned}\partial_u \tilde{J}^k &= \frac{1}{8} \int_{S^2} \tilde{X}^k ([c, \Delta(\Delta + 2)n]_1 + [\underline{c}, \Delta(\Delta + 2)\underline{n}]_1) \\ \partial_u \tilde{C}^k &= \frac{1}{8} \int_{S^2} \tilde{X}^k \left(u [((\Delta + 2)n)^2 + ((\Delta + 2)\underline{n})^2 - 4\epsilon^{AB} \nabla_A n \nabla_B (\Delta + 2)\underline{n}] \right. \\ &\quad \left. + [(\Delta + 2)c, (\Delta + 2)n]_2 + [(\Delta + 2)\underline{c}, (\Delta + 2)\underline{n}]_2 \right),\end{aligned}\tag{1.8}$$

where $[\cdot, \cdot]_1$ is the Poisson bracket on S^2 defined in (4.1) and $[\cdot, \cdot]_2$ is another bracket on S^2 defined in (4.2).

The Bondi-Metzner-Sachs (BMS) group acts on \mathcal{I}^+ . It includes supertranslations which we will review in further details in Section 5. The ambiguity of supertranslations has presented an essential difficulty to understanding the structure of \mathcal{I}^+ since the 1960s. Among (m, N_A, C_{AB}, N_{AB}) , only N_{AB} is a supertranslation invariant quantity. It is natural to ask whether total flux of angular momentum is invariant under a supertranslation. For the classical angular momentum, we prove that

Corollary 1.4 (Theorem 5.1). *Suppose \mathcal{I}^+ extends from $u = -\infty$ to $u = +\infty$ and the news tensor decays as*

$$N_{AB}(u, x) = O(|u|^{-1-\varepsilon}) \text{ as } u \rightarrow \pm\infty,$$

then the total flux of the classical angular momentum \tilde{J}^k is supertranslation invariant if and only if

$$\lim_{u \rightarrow +\infty} m(u, x) - \lim_{u \rightarrow -\infty} m(u, x)\tag{1.9}$$

is supported in the $l \leq 1$ modes.

In particular, if $\lim_{u \rightarrow +\infty} m(u, x) - \lim_{u \rightarrow -\infty} m(u, x)$ contains $l \geq 2$ modes, the total flux of the classical angular momentum will depend on the supertranslation. This demonstrates how the total flux of the classical angular momentum can be shifted by a supertranslation. On the other hand, we show that the CWY angular momentum is free of such supertranslation ambiguity.

Theorem 1.5 (Theorem 5.4). *Suppose the news tensor decays as*

$$N_{AB}(u, x) = O(|u|^{-1-\varepsilon}) \text{ as } u \rightarrow \pm\infty.$$

Then the total flux of J^k is supertranslation invariant.

Remark 1.6. In the above statement, supertranslation invariant means that it is equivariant under ordinary ($l = 1$) translation and is invariant

under higher mode ($l \geq 2$) of the supertranslation. See the statement of Theorem 5.4 for further details.

We also show that the invariance under supertranslation distinguishes the CWY center of mass from the classical center of mass. Indeed, the total flux of the classical center of mass is invariant under supertranslation if and only if $\lim_{u \rightarrow +\infty} m(u, x) - \lim_{u \rightarrow -\infty} m(u, x)$ is a constant function on S^2 . On the other hand, the total flux of the CWY center of mass is always supertranslation invariant. See the statement of Theorem 5.5.

Next, we show that if a spacetime admits a Bondi-Sachs coordinate system with vanishing news tensor, then (E, P^k, J^k, C^k) are constant (independent of the retarded time u) and supertranslation invariant. See the statement of Theorem 6.2 for further details.

While our focus is on the study of angular momentum and center of mass in a Bondi-Sachs coordinate system, we show that the evolution formulae for the classical angular momentum can be carried over to the framework of the stability of Minkowski spacetime [9] if we take Rizzi's definition of angular momentum [20, 21]. This provides a conservation law of angular momentum that complements the conservation law for linear momentum of Christodoulou [7, Equation (13)].

Another natural consequence of (1.8) is a duality paradigm among sets of null infinity data (m, N_A, C_{AB}, N_{AB}) , through replacing the potentials $(c, \underline{c}, n, \underline{n})$ by $(-\underline{c}, c, -\underline{n}, n)$.

Corollary 1.7. *Given a set of null infinity data (m, N_A, C_{AB}, N_{AB}) defined on $[u_1, u_2] \times S^2$, there exists a dual set of null infinity data $(m^*, N_A^*, C_{AB}^*, N_{AB}^*)$ that has the same (classical) energy, linear momentum, angular momentum, and center-of-mass.*

These are dual sets of null infinity data that are indistinguishable in terms of the classical conserved quantities.

The paper is organized as follows. In Section 2, we introduce the definitions and integration by parts formulae used throughout the paper. The flux of classical conserved quantities is computed in Section 3 and is rewritten in terms of the potentials in Section 4. The aforementioned consequences of flux formulae are presented in Section 5 to Section 7. In the last section, we consider the case of quadrupole moment radiation. With the future theoretical and numerical investigation in mind, we express the flux formulae in terms of the spherical harmonics expansion of potentials explicitly.

2. BACKGROUND INFORMATION

In this section, we describe the Bondi-Sachs coordinate system and recall several useful formulae for functions and tensors on S^2 .

2.1. Bondi-Sachs coordinates. In terms of a Bondi-Sachs coordinate system (u, r, x^2, x^3) , near \mathcal{I}^+ of a vacuum spacetime, the metric takes the form

$$g_{\alpha\beta}dx^\alpha dx^\beta = -UVdu^2 - 2Ududr + r^2h_{AB}(dx^A + W^A du)(dx^B + W^B du). \quad (2.1)$$

The index conventions here are $\alpha, \beta = 0, 1, 2, 3$, $A, B = 2, 3$, and $u = x^0, r = x^1$. See [2, 17] for more details of the construction of the coordinate system.

The metric coefficients U, V, h_{AB}, W^A of (2.1) depend on u, r, θ, ϕ , but $\det h_{AB}$ is independent of u and r . These gauge conditions thus reduce the number of metric coefficients of a Bondi-Sachs coordinate system to six (there are only two independent components in h_{AB}). On the other hand, the boundary conditions $U \rightarrow 1, V \rightarrow 1, W^A \rightarrow 0, h_{AB} \rightarrow \sigma_{AB}$ are imposed as $r \rightarrow \infty$ (such boundary conditions may not be satisfied in a radiative spacetime). Here σ_{AB} denotes a standard round metric on S^2 . The special gauge choice implies a hierarchy among the vacuum Einstein equations, see [17, 14].

Assuming the outgoing radiation condition [2, 22, 17], the boundary condition and the vacuum Einstein equation imply that as $r \rightarrow \infty$, all metric coefficients can be expanded in inverse integral powers of r .¹ In particular (see Chrusciel-Jeziarski-Kijowski [10, (5.98)-(5.100)] for example),

$$\begin{aligned} U &= 1 - \frac{1}{16r^2}|C|^2 + O(r^{-3}), \\ V &= 1 - \frac{2m}{r} + \frac{1}{r^2} \left(\frac{1}{3}\nabla^A N_A + \frac{1}{4}\nabla^A C_{AB}\nabla_D C^{BD} + \frac{1}{16}|C|^2 \right) + O(r^{-3}), \\ W^A &= \frac{1}{2r^2}\nabla_B C^{AB} + \frac{1}{r^3} \left(\frac{2}{3}N^A - \frac{1}{16}\nabla^A |C|^2 - \frac{1}{2}C^{AB}\nabla^D C_{BD} \right) + O(r^{-4}), \\ h_{AB} &= \sigma_{AB} + \frac{C_{AB}}{r} + \frac{1}{4r^2}|C|^2\sigma_{AB} + O(r^{-3}) \end{aligned}$$

where $m = m(u, x^A)$ is the mass aspect, $N_A = N_A(u, x^A)$ is the angular aspect and $C_{AB} = C_{AB}(u, x^A)$ is the shear tensor of this Bondi-Sachs coordinate system. Note that our convention of angular momentum aspect differs from that of Chrusciel-Jeziarski-Kijowski [10], $N_A = -3N_{A(CJK)}$. Here we take norm, raise and lower indices of tensors with respect to the metric σ_{AB} . We also define the *news tensor* $N_{AB} = \partial_u C_{AB}$.

2.2. Integral formulae on 2-sphere. Let σ_{AB} be the standard round metric on S^2 with respect to which the indexes of tensors are raised or

¹The outgoing radiation condition assumes the traceless part of the r^{-2} term in the expansion of h_{AB} is zero. The presence of this traceless term will lead to a logarithmic term in the expansions of W^A and V . Spacetimes with metrics which admit an expansion in terms of $r^{-j} \log^i r$ are called ‘‘polyhomogeneous’’ and are studied in [11]. They do not obey the outgoing radiation condition or the peeling theorem [23], but they do appear as perturbations of the Minkowski spacetime by the work of Christodoulou-Klainerman [9].

lowered. Let ∇_A be covariant derivative with respect to σ_{AB} . Let ϵ_{AB} be the volume form. The following identity

$$\epsilon_{AB}\epsilon_{CD} = \sigma_{AC}\sigma_{BD} - \sigma_{AD}\sigma_{BC} \quad (2.2)$$

and its contraction

$$\epsilon_{AB}\epsilon^A{}_C = \sigma_{BC} \quad (2.3)$$

will be used frequently.

The curvature formula on S^2 gives

$$\nabla_A\nabla_B\nabla_C u - \nabla_B\nabla_A\nabla_C u = \sigma_{AC}\nabla_B u - \sigma_{BC}\nabla_A u$$

for a smooth function u on S^2 . In particular, we have

$$\begin{aligned} \nabla_D\nabla^D\nabla_A u &= \nabla_A(\Delta + 1)u \\ \epsilon^{AB}\nabla_A\nabla_B\nabla_C u &= \epsilon_C{}^B\nabla_B u. \end{aligned} \quad (2.4)$$

Let $\tilde{X}^k, k = 1, 2, 3$ be the restriction to S^2 of the standard coordinate functions in \mathbb{R}^3 . It is well-known that they are eigenfunctions for σ_{AB} :

$$\Delta\tilde{X}^k = -2\tilde{X}^k.$$

\tilde{X}^k also satisfies the Hessian equation

$$\nabla_A\nabla_B\tilde{X}^k = -\tilde{X}^k\sigma_{AB}. \quad (2.5)$$

In general, an eigenfunction f with

$$\Delta f = -\ell(\ell + 1)f \quad (2.6)$$

is said to be of mode ℓ . We need the following integration by parts lemma:

Lemma 2.1. *Suppose u and v are smooth functions on S^2 of mode m and n respectively. Then*

$$\int_{S^2} \tilde{X}^k \epsilon^{AB} \nabla_A u \nabla_B v = 0$$

unless $m = n$.

Proof. Integrating by parts, we obtain

$$\int_{S^2} \tilde{X}^k \epsilon^{AB} \nabla_A u \nabla_B v = \int_{S^2} (Y^A \nabla_A v) u,$$

where $Y^A = \epsilon^{AB} \nabla_B \tilde{X}^k$ is a rotation Killing field. Since Δ commutes with $Y^A \nabla_A$, $Y^A \nabla_A v$ is of the same mode as v . \square

The following integrating by parts formulae will be useful in the later sections.

Lemma 2.2. *For any smooth functions u, v on S^2 , we have*

$$\int_{S^2} \tilde{X}^k \epsilon^{AB} \nabla_A (\Delta u) \nabla_B v = \int_{S^2} \tilde{X}^k \epsilon^{AB} \nabla_A u \nabla_B (\Delta v) \quad (2.7)$$

$$\int_{S^2} \tilde{X}^k \epsilon^{AB} \nabla_A \nabla_D u \nabla_B \nabla^D v = - \int_{S^2} \tilde{X}^k \epsilon^{AB} \nabla_A u \nabla_B (\Delta + 2)v. \quad (2.8)$$

Proof. We prove the second formula and the first formula follows similarly. Integrating by parts the left hand side, we obtain

$$-\int_{S^2} \nabla_D \tilde{X}^k \epsilon^{AB} \nabla_A u \nabla_B \nabla^D v - \int_{S^2} \tilde{X}^k \epsilon^{AB} \nabla_A u \nabla_D \nabla_B \nabla^D v$$

Integrating the first term by parts again, we obtain

$$\int_{S^2} \nabla_B \nabla_D \tilde{X}^k \epsilon^{AB} \nabla_A u \nabla^D v - \int_{S^2} \tilde{X}^k \epsilon^{AB} \nabla_A u \nabla_D \nabla_B \nabla^D v$$

By (2.4), this is equal to

$$-\int_{S^2} \tilde{X}^k \epsilon^{AB} \nabla_A u \nabla_B v - \int_{S^2} \tilde{X}^k \epsilon^{AB} \nabla_A u \nabla_B (\Delta + 1)v.$$

□

Lemma 2.3. *For any smooth function u on S^2 , we have*

$$\begin{aligned} \int_{S^2} [2\nabla_A \nabla_B u \nabla^A \nabla^B u - (\Delta u)^2] &= \int_{S^2} u \Delta (\Delta + 2)u \\ \int_{S^2} \tilde{X}^i [2\nabla_A \nabla_B u \nabla^A \nabla^B u - (\Delta u)^2] &= \int_{S^2} \tilde{X}^i [(\Delta + 2)u]^2. \end{aligned}$$

Proof. We use the following formulae in the derivation

$$\begin{aligned} \Delta |\nabla u|^2 &= 2|\nabla^2 u|^2 + 2\nabla u \cdot \nabla (\Delta + 1)u \\ \Delta (u^2) &= 2|\nabla u|^2 + 2u\Delta u \\ \Delta (u\Delta u) &= (\Delta u)^2 + 2\nabla u \cdot \nabla (\Delta u) + u\Delta^2 u. \end{aligned}$$

We prove the second formula and the first one follows similarly. Integrating by parts twice gives

$$\int_{S^2} \tilde{X}^i \nabla_A \nabla_B u \nabla^A \nabla^B u = \int_{S^2} u \nabla^A \nabla^B (\tilde{X}^i \nabla_A \nabla_B u)$$

We compute

$$\begin{aligned} &\nabla^A \nabla^B (\tilde{X}^i \nabla_A \nabla_B u) \\ &= (\nabla^A \nabla^B \tilde{X}^i) \nabla_A \nabla_B u + 2\nabla^B \tilde{X}^i \nabla^A \nabla_A \nabla_B u + \tilde{X}^i \nabla^A \nabla^B \nabla_A \nabla_B u \\ &= -\tilde{X}^i \Delta u + 2\nabla^B \tilde{X}^i \nabla_B (\Delta + 1)u + \tilde{X}^i \Delta (\Delta + 1)u \\ &= \tilde{X}^i \Delta^2 u + 2\nabla^B \tilde{X}^i \nabla_B (\Delta + 1)u \end{aligned}$$

where we use $\nabla^A \nabla_A \nabla_B u = \nabla_B (\Delta + 1)u$.

On the other hand, we have the identity:

$$2\nabla^B u \nabla_B v = \Delta(uv) - u\Delta v - v\Delta u$$

and thus

$$2\nabla^B \tilde{X}^i \nabla_B (\Delta + 1)u = \Delta(\tilde{X}^i (\Delta + 1)u) - \tilde{X}^i \Delta (\Delta + 1)u + 2\tilde{X}^i (\Delta + 1)u.$$

Putting all together gives:

$$\begin{aligned} & \int_{S^2} \tilde{X}^i \nabla_A \nabla_B u \nabla^A \nabla^B u \\ &= \int_{S^2} \tilde{X}^i \Delta^2 u + \int u [\Delta(\tilde{X}^i(\Delta+1)u) - \tilde{X}^i \Delta(\Delta+1)u + 2\tilde{X}^i(\Delta+1)u]. \\ &= \int_{S^2} \tilde{X}^i [(\Delta u)^2 + 2u\Delta u + 2u^2] \end{aligned}$$

Therefore,

$$\int_{S^2} \tilde{X}^i [2\nabla_A \nabla_B u \nabla^A \nabla^B u - (\Delta u)^2] = \int_{S^2} \tilde{X}^i [(\Delta u)^2 + 4u\Delta u + 4u^2] = \int_{S^2} \tilde{X}^i [(\Delta+2)u]^2.$$

□

2.3. Closed and Co-closed Decomposition. In this subsection, we consider symmetric traceless 2-tensors C_{AB} and N_{AB} on S^2 with the decomposition (see [15, Appendix B] for a derivation)

$$C_{AB} = \nabla_A \nabla_B c - \frac{1}{2} \sigma_{AB} \Delta c + \frac{1}{2} (\epsilon_A^E \nabla_E \nabla_B \underline{c} + \epsilon_B^E \nabla_E \nabla_A \underline{c}) \quad (2.9)$$

$$N_{AB} = \nabla_A \nabla_B n - \frac{1}{2} \sigma_{AB} \Delta n + \frac{1}{2} (\epsilon_A^E \nabla_E \nabla_B \underline{n} + \epsilon_B^E \nabla_E \nabla_A \underline{n}) \quad (2.10)$$

for smooth functions $c, \underline{c}, n, \underline{n}$ on S^2 that are referred as potentials of C_{AB} and N_{AB} . The potentials are unique up to their 0 and 1 mode. In the case we consider when C_{AB} and N_{AB} depend on u , all $c, \underline{c}, n, \underline{n}$ depend on u as well.

Proposition 2.4. *Closed and co-closed parts of a symmetric traceless 2-tensors on S^2 are dual to each other in the following sense.*

- (1) Denote the space of symmetric traceless 2-tensors on S^2 by \widehat{Sym} . Then the map $\varepsilon_2 : \widehat{Sym} \rightarrow \widehat{Sym}$, $\varepsilon_2(C_{AB}) = \epsilon_A^D C_{DB}$ satisfies

$$\varepsilon_2(\nabla_A \nabla_B c - \frac{1}{2} \sigma_{AB} \Delta c) = \frac{1}{2} (\epsilon_A^E \nabla_E \nabla_B \underline{c} + \epsilon_B^E \nabla_E \nabla_A \underline{c}), \quad (2.11)$$

$$\varepsilon_2 \left(\frac{1}{2} (\epsilon_A^E \nabla_E \nabla_B \underline{c} + \epsilon_B^E \nabla_E \nabla_A \underline{c}) \right) = -\nabla_A \nabla_B \underline{c} + \frac{1}{2} \sigma_{AB} \Delta \underline{c}. \quad (2.12)$$

- (2) The following identity holds for symmetric traceless 2-tensors

$$\epsilon_D^B \nabla^D C_{BA} = \epsilon_A^D \nabla^B C_{BD}. \quad (2.13)$$

In other words, we have a commutative diagram of isomorphisms

$$\begin{array}{ccc} \widehat{Sym} & \xrightarrow{\varepsilon_2} & \widehat{Sym} \\ \downarrow \text{div} & & \downarrow \text{div} \\ \Lambda^1 & \xrightarrow{*} & \Lambda^1, \end{array}$$

where Λ^1 denotes the space of 1-forms and $(*\omega)_A = \epsilon_A^B \omega_B$ is the Hodge star on 1-forms.

Proof. We use (2.2) and (2.3) in the derivation. Since $\epsilon^{AB}\varepsilon_2(C_{AB}) = 0$ and $\sigma^{AB}\varepsilon_2(C_{AB}) = 0$, $\varepsilon_2(C_{AB})$ is symmetric and traceless. In particular,

$$\epsilon_A^D C_{DB} = \frac{1}{2}(\epsilon_A^D C_{DB} + \epsilon_B^D C_{DA}) \quad (2.14)$$

and (2.11) and (2.12) follow by direct computation.

To verify (2.13), note that both sides are equal to $\nabla^D C_{DE}$ after contracted with ϵ^A_E . \square

In the following two lemmas, we express several integrals involving the shear tensor and the news tensor in terms of their potentials. These formulae will help us to derive Theorem 1.3 from Theorem 1.2.

Lemma 2.5. *Suppose Y^A is either $\nabla^A \tilde{X}^k$ or $\epsilon^{AB} \nabla_B \tilde{X}^k$, and C_{AB} and N_{AB} are given by (2.9) and (2.10), then*

$$\begin{aligned} & \int_{S^2} Y^A C_{AB} \nabla_D N^{BD} \\ &= -\frac{1}{4} \int_{S^2} Y^A [(\Delta + 2)n \nabla_A (\Delta + 2)c + (\Delta + 2)\underline{n} \nabla_A (\Delta + 2)\underline{c}] \\ & \quad + \frac{1}{4} \int_{S^2} Y^A \epsilon_A^D [\nabla_D ((\Delta + 2)c) (\Delta + 2)\underline{n} - \nabla_D ((\Delta + 2)\underline{c}) (\Delta + 2)n] \end{aligned} \quad (2.15)$$

Proof. First of all, note that

$$\nabla^B Y^A C_{AB} = 0, \epsilon^{BD} \nabla_D Y^A C_{AB} = 0.$$

From

$$\nabla_D N^{BD} = \frac{1}{2} \nabla^B (\Delta + 2)n + \frac{1}{2} \epsilon^{BD} \nabla_D (\Delta + 2)\underline{n},$$

we integrate by parts to get

$$\int_{S^2} Y^A C_{AB} \nabla_D N^{BD} = -\frac{1}{2} \int_{S^2} Y^A (\nabla^B C_{AB} (\Delta + 2)n + \epsilon^{BD} \nabla_D C_{AB} (\Delta + 2)\underline{n}).$$

By (2.13)

$$\epsilon^{DB} \nabla_D C_{BA} = \epsilon_{AD} \nabla_B C^{BD}$$

and $\nabla^B C_{BD} = \frac{1}{2} \nabla^D (\Delta + 2)c + \frac{1}{2} \epsilon^{BD} \nabla_B (\Delta + 2)\underline{c}$, we obtain the desired formula. \square

The above generalizes the integral identities derived in [15, (65), (66)]:

$$\begin{aligned} & \int_{S^2} Y^A F_A^B \nabla^D F_{DB} = 0, \\ & \int_{S^2} Y^A \underline{F}_A^B \nabla^D \underline{F}_{DB} = 0 \end{aligned}$$

for $Y^A = \epsilon^{AB} \nabla_B \tilde{X}^k$.

Skew-symmetrizing (2.15), we obtain:

$$\begin{aligned}
& \int_{S^2} Y^A (C_{AB} \nabla_D N^{BD} - N_{AB} \nabla_D C^{BD}) \\
&= \frac{1}{4} \int_{S^2} Y^A [(\Delta + 2)c \nabla_A (\Delta + 2)n - (\Delta + 2)n \nabla_A (\Delta + 2)c] \\
& \quad + \frac{1}{4} \int_{S^2} Y^A [(\Delta + 2)\underline{c} \nabla_A (\Delta + 2)\underline{n} - (\Delta + 2)\underline{n} \nabla_A (\Delta + 2)\underline{c}] \\
& \quad + \frac{1}{4} \int_{S^2} Y^A \epsilon_A^D \nabla_D [(\Delta + 2)c(\Delta + 2)\underline{n} - (\Delta + 2)\underline{c}(\Delta + 2)n].
\end{aligned} \tag{2.16}$$

Next we prove

Lemma 2.6.

$$\begin{aligned}
\int_{S^2} N_{AB} N^{AB} &= \frac{1}{2} \int_{S^2} n \Delta (\Delta + 2)n + \underline{n} \Delta (\Delta + 2)\underline{n} \\
\int_{S^2} \tilde{X}^k N_{AB} N^{AB} &= \frac{1}{2} \int_{S^2} \tilde{X}^k [((\Delta + 2)n)^2 + ((\Delta + 2)\underline{n})^2 - 4\epsilon^{AB} \nabla_A n \nabla_B (\Delta + 2)\underline{n}].
\end{aligned}$$

Proof. Using the formula $\epsilon_A^C \epsilon^{BD} = \delta_A^B \sigma^{CD} - \delta_A^D \sigma^{CB}$ and $\epsilon^{AB} \epsilon_{AE} = \sigma^{BE}$, we compute that

$$\begin{aligned}
N_{AB} N^{AB} &= \nabla_A \nabla_B n \nabla^A \nabla^B n - \frac{1}{2} (\Delta n)^2 + \nabla_A \nabla_B \underline{n} \nabla^A \nabla^B \underline{n} - \frac{1}{2} (\Delta \underline{n})^2 \\
& \quad + 2\epsilon^{AC} \nabla_A \nabla_B n \nabla_C \nabla^B \underline{n}
\end{aligned} \tag{2.17}$$

Integrating by parts yields

$$\begin{aligned}
\int_{S^2} \epsilon^{AC} \nabla_A \nabla_B n \nabla_C \nabla^B \underline{n} &= - \int_{S^2} \epsilon^{AC} \nabla^B \nabla_A \nabla_B n \nabla_C \underline{n} \\
&= - \int_{S^2} \epsilon^{AC} \nabla_A (\Delta + 1)n \nabla_C \underline{n} = 0
\end{aligned}$$

The first formula now follows from the first formula in Lemma 2.3. The second formula follows from the second and third formula in Lemma 2.3. \square

The second formula of Lemma 2.6 can be polarized and we obtain

$$\begin{aligned}
\int_{S^2} \tilde{X}^k C_{AB} N^{AB} &= \frac{1}{2} \int_{S^2} \tilde{X}^k [(\Delta + 2)c(\Delta + 2)n + (\Delta + 2)\underline{c}(\Delta + 2)\underline{n} \\
& \quad - 2\epsilon^{AB} (\nabla_A c \nabla_B (\Delta + 2)\underline{n} + \nabla_A n \nabla_B (\Delta + 2)\underline{c})]
\end{aligned} \tag{2.18}$$

3. EVOLUTION OF CONSERVED QUANTITIES

In this section, we compute the evolution of the classical angular momentum and center of mass. These formulae will be used to calculate the total flux of the conserved quantities.

Let's first review the evolution of the metric under the Einstein equation. It is well-known (see [10, (5.102)] for example) that the evolution of the mass aspect function is given by

$$\partial_u m = -\frac{1}{8}N_{AB}N^{AB} + \frac{1}{4}\nabla^A\nabla^B N_{AB}. \quad (3.1)$$

The modified mass aspect function \hat{m} is defined to be [24]

$$\hat{m} = m - \frac{1}{4}\nabla^A\nabla^B C_{AB} = m - \frac{1}{8}\Delta(\Delta + 2)c \quad (3.2)$$

and satisfies

$$\partial_u \hat{m} = -\frac{1}{8}N_{AB}N^{AB}. \quad (3.3)$$

Therefore,

$$\begin{aligned} \partial_u E &= -\frac{1}{4}\int_{S^2} N_{AB}N^{AB} \\ \partial_u P^k &= -\frac{1}{4}\int_{S^2} \tilde{X}^k N_{AB}N^{AB}, \quad k = 1, 2, 3. \end{aligned}$$

We also recall the evolution of N_A (see [10, (5.103)] for example):

$$\begin{aligned} \partial_u N_A &= \nabla_A m - \frac{1}{4}\nabla^D(\nabla_D\nabla^E C_{EA} - \nabla_A\nabla^E C_{ED}) \\ &\quad + \frac{1}{4}\nabla_A(C_{BE}N^{BE}) - \frac{1}{4}\nabla_B(C^{BD}N_{DA}) + \frac{1}{2}C_{AB}\nabla_D N^{DB}. \end{aligned}$$

The formula can be rewritten in the following form:

Proposition 3.1. *The angular momentum aspect N_A evolves according to*

$$\begin{aligned} \partial_u N_A &= \nabla_A m + \frac{1}{4}\epsilon_{AB}\nabla^B(\epsilon^{PQ}\nabla_P\nabla^E C_{EQ}) + \frac{1}{8}\nabla_A(C_{BE}N^{BE}) \\ &\quad + \frac{1}{8}\epsilon_{AB}\nabla^B(\epsilon^{PQ}C_P^E N_{EQ}) + \frac{1}{2}C_{AB}\nabla_D N^{DB}. \end{aligned} \quad (3.4)$$

Proof. We rewrite the terms $-\frac{1}{4}\nabla^D(\nabla_D\nabla^E C_{EA} - \nabla_A\nabla^E C_{ED})$ and $-\frac{1}{4}\nabla_B(C^{BD}N_{DA})$. First we check the following identity directly:

$$\epsilon_{AB}\nabla^B(\epsilon^{PQ}\nabla_P\nabla^E C_{EQ}) = -\nabla^D(\nabla_D\nabla^E C_{EA} - \nabla_A\nabla^E C_{ED}).$$

As for the term $C_B^D N_{DA}$, we use the following general formulae for symmetric traceless 2-tensors on the 2-sphere:

$$\begin{aligned} C_B^D N_{DA} + N_B^D C_{DA} &= (C_{DE}N^{DE})\sigma_{AB} \\ C_B^D N_{DA} - N_B^D C_{DA} &= -(\epsilon^{PQ}C_P^E N_{EQ})\epsilon_{AB} \end{aligned}$$

Therefore,

$$2C_B^D N_{DA} = (C_{DE}N^{DE})\sigma_{AB} - (\epsilon^{PQ}C_P^E N_{EQ})\epsilon_{AB}.$$

□

Equation (3.4) is indeed equivalent to equation (4) on page 48 of [8]. We apply (3.4) to derive the evolution of the classical angular momentum and center of mass.

Theorem 3.2 (Theorem 1.2). *The classical angular momentum and center of mass evolve according to the following:*

$$\partial_u \tilde{J}^k = \frac{1}{4} \int_{S^2} \left[\epsilon^{AE} \nabla_E \tilde{X}^k (C_{AB} \nabla_D N^{BD} - N_{AB} \nabla_D C^{BD}) + \tilde{X}^k \epsilon^{AB} (C_A{}^D N_{DB}) \right], \quad (3.5)$$

$$\partial_u \tilde{C}^k = \frac{1}{4} \int_{S^2} \left[\nabla^A \tilde{X}^k \left(\frac{u}{2} \nabla_A |N|^2 + C_{AB} \nabla_D N^{BD} - N_{AB} \nabla_D C^{BD} \right) \right], \quad (3.6)$$

where $k = 1, 2, 3$.

Proof. By (1.2),

$$\partial_u \tilde{J}^k = \int_{S^2} \epsilon^{AB} \nabla_B \tilde{X}^k \left[\partial_u N_A - \frac{1}{4} \partial_u (C_A{}^D \nabla^B C_{DB}) \right].$$

First, we deal with the term $\frac{1}{4} \epsilon_{AB} \nabla^B (\epsilon^{PQ} \nabla_P \nabla^E C_{EQ})$ on the right hand side of (3.4) and claim that

$$\int_{S^2} Y^A \epsilon_{AB} \nabla^B (\epsilon^{PQ} \nabla_P \nabla^E C_{EQ}) = 0 \quad (3.7)$$

for $Y^A = \nabla^A \tilde{X}^k$ or $\epsilon^{AB} \nabla_B \tilde{X}^k$. Integrating by parts, the integral becomes

$$\int_{S^2} \epsilon_{AB} \nabla^A Y^B (\epsilon^{PQ} \nabla_P \nabla^E C_{EQ}).$$

Since $\epsilon^{PQ} \nabla_P \nabla^E C_{EQ} = -\frac{1}{2} \Delta (\Delta + 2) \underline{c}$ and $\epsilon_{AB} \nabla^A Y^B$ is zero or $2\tilde{X}^k$, the integral vanishes.

Hence, we obtain

$$\begin{aligned} & \partial_u \tilde{J}^k \\ &= \int_{S^2} \epsilon^{AB} \nabla_B \tilde{X}^k \left[\frac{1}{8} \epsilon_{AE} \nabla^E (\epsilon^{PQ} C_P{}^E N_{EQ}) + \frac{1}{2} C_{AB} \nabla_D N^{DB} - \frac{1}{4} \partial_u (C_A{}^D \nabla^B C_{DB}) \right] \end{aligned}$$

since the integral of $\nabla_A m + \frac{1}{8} \nabla_A (C_{BE} N^{BE})$ against $\epsilon^{AB} \nabla_B \tilde{X}^k$ vanishes. Integrating by parts the first term and use $\epsilon^{AB} \epsilon_{AE} = \delta_E^B$, we obtain the desired formula.

We now turn to the formula for \tilde{C}^k . By (1.3) and (3.7),

$$\begin{aligned} & \partial_u \tilde{C}^k \\ &= \int_{S^2} \nabla^A \tilde{X}^k \left[\partial_u N_A - \nabla_A m + \frac{u}{8} \nabla_A |N|^2 - \frac{1}{4} \partial_u (C_A{}^D \nabla^B C_{DB}) - \frac{1}{16} \nabla_A \partial_u (C_{DE} C^{DE}) \right] \\ &= \int_{S^2} \nabla^A \tilde{X}^k \left[\frac{u}{8} \nabla_A |N|^2 + \frac{1}{8} \nabla_A (C_{BE} N^{BE}) + \frac{1}{2} C_{AB} \nabla_D N^{DB} \right. \\ & \quad \left. - \frac{1}{4} \partial_u (C_A{}^D \nabla^B C_{DB}) - \frac{1}{16} \nabla_A \partial_u (C_{DE} C^{DE}) \right] \end{aligned}$$

since the integral of $\frac{1}{8}\epsilon_{AB}\nabla^B(\epsilon^{PQ}C_P^E N_{EQ})$ against $\nabla^A\tilde{X}^k$ vanishes. We arrive at the desired formula since $\partial_u(C_{DE}C^{DE}) = 2(C_{BE}N^{BE})$. \square

4. EVOLUTION FORMULAE IN TERMS OF POTENTIALS

In this section, we rewrite the evolution formulae in terms of the potentials of the shear and the news tensor.

4.1. Energy and linear momentum. First we recall the formulae for the energy and linear momentum.

Proposition 4.1. *Suppose C_{AB} and N_{AB} are given as in (2.9) and (2.10), we have*

$$\begin{aligned}\partial_u E &= -\frac{1}{8}\int_{S^2} [n\Delta(\Delta+2)n + \underline{n}\Delta(\Delta+2)\underline{n}] \\ \partial_u P^k &= -\frac{1}{8}\int_{S^2} \tilde{X}^k [((\Delta+2)n)^2 + ((\Delta+2)\underline{n})^2 - 4\epsilon^{AB}\nabla_A n \nabla_B(\Delta+2)\underline{n}].\end{aligned}$$

Proof. These follow from Lemma 2.6. \square

4.2. Proof of Theorem 1.3. We first prove the following Proposition:

Proposition 4.2. *The evolution formulae of the conserved quantities can be written as*

$$\begin{aligned}\partial_u \tilde{J}^k &= \frac{1}{8}\int_{S^2} \tilde{X}^k \epsilon^{AB} [\nabla_A c \nabla_B \Delta(\Delta+2)n + \nabla_A \underline{c} \nabla_B \Delta(\Delta+2)\underline{n}] \\ \partial_u \tilde{C}^k &= \frac{1}{8}\int_{S^2} u \tilde{X}^k [((\Delta+2)n)^2 + ((\Delta+2)\underline{n})^2 - 4\epsilon^{AB}\nabla_A n \nabla_B(\Delta+2)\underline{n}] \\ &\quad + \frac{1}{16}\int_{S^2} [\tilde{X}^k (\Delta(\Delta+2)\underline{c}(\Delta+2)\underline{n} - \Delta(\Delta+2)\underline{n}(\Delta+2)\underline{c})] \\ &\quad + \frac{1}{16}\int_{S^2} [\tilde{X}^k (\Delta(\Delta+2)c(\Delta+2)n - \Delta(\Delta+2)n(\Delta+2)c)].\end{aligned}$$

Proof. We write

$$4\partial_u \tilde{J}^k = \int_{S^2} -\tilde{X}^k \epsilon^{AB} C_B^D N_{DA} + \int_{S^2} Y_k^A (C_{AB} \nabla_D N^{BD} - N_{AB} \nabla_D C^{BD}) = (1)+(2)$$

and compute (1) and (2) separately.

Note that (1) = $-\int_{S^2} \tilde{X}^k \varepsilon_2(C_{AB}) N^{AB}$ and recall that $\varepsilon_2(C_{AB})$ has potentials $-\underline{c}$ and c . Applying (2.18), we get

$$\begin{aligned}
(1) &= -\frac{1}{2} \int_{S^2} \tilde{X}^k [-(\Delta + 2)\underline{c}(\Delta + 2)n + (\Delta + 2)c(\Delta + 2)\underline{n}] \\
&\quad + \int_{S^2} \epsilon^{AB} (\nabla_A \underline{c} \nabla_B (\Delta + 2)\underline{n} - \nabla_A n \nabla_B (\Delta + 2)c) \\
&= -\frac{1}{2} \int_{S^2} \tilde{X}^k [-(\Delta + 2)\underline{c}(\Delta + 2)n + (\Delta + 2)c(\Delta + 2)\underline{n}] \\
&\quad - \int_{S^2} \tilde{X}^k \epsilon^{AB} [\nabla_A \underline{c} \nabla_B (\Delta + 2)\underline{n} - \nabla_A n \nabla_B (\Delta + 2)c] \\
&= -\frac{1}{2} \int_{S^2} \tilde{X}^k [-(\Delta + 2)\underline{c}(\Delta + 2)n + (\Delta + 2)c(\Delta + 2)\underline{n}] \\
&\quad - \int_{S^2} \tilde{X}^k \epsilon^{AB} [\nabla_A c \nabla_B (\Delta + 2)n + \nabla_A \underline{c} \nabla_B (\Delta + 2)\underline{n}]
\end{aligned}$$

where we used (2.7) in the last equality.

Applying (2.16) to $Y^A = Y_k^A$, we have

$$\begin{aligned}
(2) &= \frac{1}{2} \int_{S^2} \tilde{X}^k \epsilon^{AB} [\nabla_A (\Delta + 2)c \nabla_B (\Delta + 2)n + \nabla_A (\Delta + 2)\underline{c} \nabla_B (\Delta + 2)\underline{n}] \\
&\quad + \frac{1}{2} \int_{S^2} \tilde{X}^k [(\Delta + 2)c(\Delta + 2)\underline{n} - (\Delta + 2)\underline{c}(\Delta + 2)n]
\end{aligned}$$

Therefore,

$$\begin{aligned}
(1) + (2) &= -\int_{S^2} \tilde{X}^k \epsilon^{AB} [\nabla_A c \nabla_B (\Delta + 2)n + \nabla_A \underline{c} \nabla_B (\Delta + 2)\underline{n}] \\
&\quad + \frac{1}{2} \int_{S^2} \tilde{X}^k \epsilon^{AB} [\nabla_A (\Delta + 2)c \nabla_B (\Delta + 2)n + \nabla_A (\Delta + 2)\underline{c} \nabla_B (\Delta + 2)\underline{n}] \\
&= \frac{1}{2} \int_{S^2} \tilde{X}^k [\epsilon^{AB} \nabla_A \Delta c \nabla_B (\Delta + 2)n + \nabla_A \Delta \underline{c} \nabla_B (\Delta + 2)\underline{n}],
\end{aligned}$$

and the desired formula follows by (2.7).

As for the evolution of the center of mass, we apply (2.15) and note that

$$\int_{S^2} \nabla^A \tilde{X}^k \epsilon_A^D \nabla_D [(\Delta + 2)c(\Delta + 2)\underline{n} + (\Delta + 2)\underline{c}(\Delta + 2)n] = 0.$$

Therefore,

$$\begin{aligned}
\partial_u \tilde{C}^k &= \frac{1}{8} \int_{S^2} u \tilde{X}^k [((\Delta + 2)n)^2 + ((\Delta + 2)\underline{n})^2 - 4\epsilon^{AB} \nabla_A n \nabla_B (\Delta + 2)\underline{n}] \\
&\quad - \frac{1}{16} \int_{S^2} [\nabla^A \tilde{X}^k (\nabla_A (\Delta + 2)\underline{c} (\Delta + 2)\underline{n} - \nabla_A (\Delta + 2)\underline{n} (\Delta + 2)\underline{c})] \\
&\quad - \frac{1}{16} \int_{S^2} [\nabla^A \tilde{X}^k (\nabla_A (\Delta + 2)c (\Delta + 2)n - \nabla_A (\Delta + 2)n (\Delta + 2)c)] \\
&= \frac{1}{8} \int_{S^2} u \tilde{X}^k [((\Delta + 2)n)^2 + ((\Delta + 2)\underline{n})^2 - 4\epsilon^{AB} \nabla_A n \nabla_B (\Delta + 2)\underline{n}] \\
&\quad + \frac{1}{16} \int_{S^2} \tilde{X}^k [\Delta (\Delta + 2)\underline{c} (\Delta + 2)\underline{n} - \Delta (\Delta + 2)\underline{n} (\Delta + 2)\underline{c}] \\
&\quad + \frac{1}{16} \int_{S^2} \tilde{X}^k [\Delta (\Delta + 2)c (\Delta + 2)n - \Delta (\Delta + 2)n (\Delta + 2)c]
\end{aligned}$$

□

To obtain the formulae given in Theorem 1.3, we rewrite the above formulae in terms of bracket operators on S^2 .

Definition 4.3. For two smooth functions u and v on S^2 , denote

$$[u, v]_1 = \epsilon^{AB} \nabla_A u \nabla_B v \quad (4.1)$$

and

$$[u, v]_2 = \frac{1}{2} ((\Delta u)v - (\Delta v)u). \quad (4.2)$$

In view of Definition 4.3, we can write

$$\partial_u \tilde{J}^k = \frac{1}{8} \int_{S^2} \tilde{X}^k ([c, \Delta (\Delta + 2)n]_1 + [\underline{c}, \Delta (\Delta + 2)\underline{n}]_1)$$

and similarly for the center of mass. This proves Theorem 1.3.

5. SUPERTRANSLATION INVARIANCE OF THE TOTAL FLUX

5.1. Total flux of classical conserved quantities. We study the effect of supertranslation on the total flux of conserved quantities along null infinity or, equivalently, the difference of conserved quantities at timelike infinity and spatial infinity. As in the previous section, suppose $I = (-\infty, \infty)$ and \mathcal{I}^+ is complete extending from spatial infinity ($u = -\infty$) to timelike infinity ($u = +\infty$). A supertranslation is a change of coordinates $(\bar{u}, \bar{x}^A) \rightarrow (u, x^A)$ such that $u = \bar{u} + f(x)$, $x^A = \bar{x}^A$ on \mathcal{I}^+ . Let m , C_{AB} , and N_{AB} denote the mass aspect, the shear, and the news, respectively, in the (u, x^A) coordinate system. Since the spherical coordinate is unchanged, we use x to denote either x^A or \bar{x}^A throughout this section. It is well-known (see [10, (C.117) and (C.119)] for example) that the shear $\bar{C}_{AB}(\bar{u}, x)$, and the news $\bar{N}_{AB}(\bar{u}, x)$

in the (\bar{u}, x) coordinate system are given by

$$\bar{C}_{AB}(\bar{u}, x) = C_{AB}(\bar{u} + f(x), x) - 2\nabla_A \nabla_B f + \Delta f \sigma_{AB} \quad (5.1)$$

$$\bar{N}_{AB}(\bar{u}, x) = N_{AB}(\bar{u} + f(x), x) \quad (5.2)$$

We assume that there exists a constant $\varepsilon > 0$ such that

$$N_{AB}(u, x) = O(|u|^{-1-\varepsilon}) \text{ as } u \rightarrow \pm\infty. \quad (5.3)$$

Note that the limits of the shear tensor exist

$$\lim_{u \rightarrow \pm\infty} C_{AB}(u, x) = C_{AB}(\pm)$$

as a result of (5.3).

Similarly, (5.3) implies that the limits of the angular momentum exist

$$\lim_{u \rightarrow \pm\infty} \tilde{J}^k(u) = \tilde{J}^k(\pm).$$

Denote the corresponding quantities after supertranslation by $\tilde{J}_f^k(\pm)$.

Let $Y^A = \epsilon^{AB} \nabla_B \tilde{X}^k$. By (3.5), the total angular momentum flux is

$$\begin{aligned} & \tilde{J}^k(+)-\tilde{J}^k(-) \\ &= \frac{1}{4} \int_{-\infty}^{+\infty} \int_{S^2} \left[Y^A (C_{AB} \nabla_D N^{BD} - N_{AB} \nabla_D C^{BD}) + \tilde{X}^k \epsilon^{AB} C_A{}^D N_{DB} \right] (u, x) dS^2 du \\ &= \frac{1}{4} \int_{-\infty}^{+\infty} \int_{S^2} \left[-\nabla_D Y^A C_{AB} N^{BD} + Y^A (-\nabla_D C_{AB} N^{BD} - N_{AB} \nabla_D C^{BD}) \right] (u, x) dS^2 du \\ &+ \frac{1}{4} \int_{-\infty}^{+\infty} \int_{S^2} \tilde{X}^k \epsilon^{AB} (C_A{}^D N_{DB})(u, x) dS^2 du \end{aligned} \quad (5.4)$$

in the (u, x) coordinates and

$$\begin{aligned} & \tilde{J}_f^k(+)-\tilde{J}_f^k(-) \\ &= \frac{1}{4} \int_{-\infty}^{+\infty} \int_{S^2} \left[-\nabla_D Y^A \bar{C}_{AB} \bar{N}^{BD} + Y^A (-\nabla_D \bar{C}_{AB} \bar{N}^{BD} - \bar{N}_{AB} \nabla_D \bar{C}^{BD}) \right] (\bar{u}, x) dS^2 d\bar{u} \\ &+ \frac{1}{4} \int_{-\infty}^{+\infty} \int_{S^2} \tilde{X}^k \epsilon^{AB} (\bar{C}_A{}^D \bar{N}_{DB})(\bar{u}, x) dS^2 d\bar{u} \end{aligned} \quad (5.5)$$

in the (\bar{u}, x) coordinates.

Applying the chain rule on (5.1) yields

$$\begin{aligned} \nabla_D \bar{C}_{AB}(\bar{u}, x) &= N_{AB}(\bar{u} + f, x) \nabla_D f + (\nabla_D C_{AB})(\bar{u} + f, x) - \nabla_D F_{AB}, \\ \nabla_D \bar{C}^{BD}(\bar{u}, x) &= N^{BD}(\bar{u} + f, x) \nabla_D f + (\nabla_D C^{BD})(\bar{u} + f, x) - \nabla^B (\Delta + 2)f. \end{aligned}$$

To simplify notation, we introduce the u independent symmetric traceless 2-tensor

$$F_{AB} = 2\nabla_A \nabla_B f - \Delta f \sigma_{AB}$$

and thus $\nabla_D F^{BD} = \nabla^B (\Delta + 2)f$.

Equation (5.5) can be rewritten as

$$\begin{aligned} & \tilde{J}_f^k(+)-\tilde{J}_f^k(-) \\ &= \frac{1}{4} \int_{-\infty}^{\infty} \int_{S^2} \left[-\nabla_D Y^A (C_{AB} - F_{AB}) N^{BD} + Y^A \omega_A \right] (\bar{u} + f, x) dS^2 d\bar{u} \\ & \quad + \frac{1}{4} \int_{-\infty}^{+\infty} \int_{S^2} \left[\tilde{X}^k \epsilon^{AB} (C_A^D - F_A^D) N_{DB} \right] (\bar{u} + f, x) dS^2 d\bar{u} \end{aligned} \quad (5.6)$$

where

$$\begin{aligned} \omega_A(u, x) &= \left(-N_{AB}(u, x) \nabla_D f - \nabla_D C_{AB} + \nabla_D F_{AB} \right) N^{BD}(u, x) \\ & \quad - N_{AB}(u, x) \left(N^{BD}(u, x) \nabla_D f + \nabla_D C^{BD}(u, x) - \nabla_D F^{BD}(x) \right). \end{aligned}$$

Note that the integrand is evaluated at $(\bar{u} + f, x)$ in equation (5.6), to which the change of variable will be applied.

By the decaying assumption of the news (5.3), we can apply change of variable $u = \bar{u} + f$ to (5.6) and rewrite it as

$$\begin{aligned} & \tilde{J}_f^k(+)-\tilde{J}_f^k(-) \\ &= \frac{1}{4} \int_{-\infty}^{+\infty} \int_{S^2} \left[-\nabla_D Y^A (C_{AB} - F_{AB}) N^{BD} + Y^A \omega_A + \tilde{X}^k \epsilon^{AB} (C_A^D - F_A^D) N_{DB} \right] (u, x) dS^2 du \end{aligned} \quad (5.7)$$

Combining (5.4) and (5.7), we obtain

$$\begin{aligned} & \left(\tilde{J}_f^k(+)-\tilde{J}_f^k(-) \right) - \left(\tilde{J}^k(+)-\tilde{J}^k(-) \right) \\ &= \frac{1}{4} \int_{-\infty}^{\infty} \int_{S^2} -Y^A |N|^2 \nabla_A f dS^2 du \\ & \quad + \frac{1}{4} \int_{-\infty}^{\infty} \int_{S^2} \left[-Y^A F_{AB} \nabla_D N^{BD} + Y^A N_{AB} \nabla_D F^{BD} - \tilde{X}^k \epsilon^{AB} F_A^D N_{DB} \right] dS^2 du \end{aligned}$$

where we used the identity $2N_{AB}N^{BD} = |N|^2 \delta_A^D$.

Observe that the second integral is of the same form as $\partial_u \tilde{J}$ given in (3.5) and one can thus simplify it as in the proof of Proposition 4.2 to get

$$\begin{aligned} & \left(J_f^k(+)-J_f^k(-) \right) - \left(J^k(+)-J^k(-) \right) \\ &= \frac{1}{4} \int_{-\infty}^{\infty} \int_{S^2} f Y^A \nabla_A |N|^2 dS^2 du + \frac{1}{4} \int_{-\infty}^{\infty} \int_{S^2} \tilde{X}^k \epsilon^{AB} \nabla_A n \nabla_B \Delta (\Delta + 2) f dS^2 du \end{aligned}$$

Integrating by parts, we arrive at

$$\begin{aligned} & \left(\tilde{J}_f^k(+)-\tilde{J}_f^k(-) \right) - \left(\tilde{J}^k(+)-\tilde{J}^k(-) \right) \\ &= \frac{1}{4} \int_{-\infty}^{+\infty} \int_{S^2} f Y^A \nabla_A (|N|^2 - \Delta (\Delta + 2) n) dS^2 du \\ &= \int_{S^2} -2f Y^A \nabla_A (m(+)-m(-)) dS^2 \end{aligned} \quad (5.8)$$

where

$$m(\pm) = \lim_{u \rightarrow \pm\infty} m(u, x).$$

Here we used the mass loss formula (3.1) in the form $\partial_u m = \frac{1}{8}\Delta(\Delta + 2)n - \frac{1}{8}|N|^2$. Note that $m(+)-m(-)$ is of the same mode as $Y^A \nabla_A(m(+)-m(-))$ because Y^A is a Killing field.

In summary, we obtain a necessary and sufficient condition for the total flux of the classical angular momentum to be supertranslation invariant.

Theorem 5.1. *Suppose the news tensor decays as*

$$N_{AB}(u, x) = O(|u|^{-1-\varepsilon}) \text{ as } u \rightarrow \pm\infty.$$

The total flux of the classical angular momentum \tilde{J}^k is supertranslation invariant if and only if

$$m(+)-m(-)$$

(as a function on S^2) is supported in the $l \leq 1$ modes.

Moreover, the above condition holds when the rescaled curvature components P (see Definition A.5) at \mathcal{I}^+ satisfy

$$\lim_{u \rightarrow \infty} P - \lim_{u \rightarrow -\infty} P \tag{5.9}$$

is supported in the $l \leq 1$ modes.

Remark 5.2. Theorem 5.1 is motivated by the investigation in [7], which is built on the framework of stability of Minkowski spacetime. Indeed, equation (11) and (12) of [7]

$$Z^+ - Z^- = \nabla\Phi \quad \text{div}(\Sigma^+ - \Sigma^-) = Z^+ - Z^-$$

imply $\lim_{u \rightarrow \infty} \underline{c}(u, x) = \lim_{u \rightarrow -\infty} \underline{c}(u, x)$. Using moreover (10) of [7]

$$\Delta\Phi = -2(F - \bar{F}),$$

we get

$$\nabla_A(8F - \Delta(\Delta + 2)c|_{-\infty}^{+\infty}) = 0.$$

Moreover, the total flux of the classical center of mass is supertranslation invariant under the same condition

Theorem 5.3. *Suppose the news tensor decays as*

$$N_{AB}(u, x) = O(|u|^{-1-\varepsilon}) \text{ as } u \rightarrow \pm\infty,$$

The total flux of the classical center of mass \tilde{C}^k is supertranslation invariant if and only if

$$m(+)-m(-)$$

is a constant function on S^2 .

Proof. Denoting $\tilde{C}^k(\pm) = \lim_{u \rightarrow \pm\infty} \tilde{C}^k(u)$, by (3.6) we have

$$\begin{aligned} & \tilde{C}^k(+)-\tilde{C}^k(-) \\ &= \frac{1}{4} \int_{-\infty}^{+\infty} \int_{S^2} u|N|^2(u, x) \tilde{X}^k + \nabla^A \tilde{X}^k [C_{AB} \nabla_D N^{BD} - N_{AB} \nabla_D C^{BD}] (u, x) dS^2 du. \end{aligned} \quad (5.10)$$

On the other hand,

$$\begin{aligned} & \tilde{C}_f^k(+)-\tilde{C}_f^k(-) \\ &= \frac{1}{4} \int_{-\infty}^{+\infty} \int_{S^2} \bar{u}|\bar{N}|^2(\bar{u}, x) \tilde{X}^k + \nabla^A \tilde{X}^k [\bar{C}_{AB} \nabla_D \bar{N}^{BD} - \bar{N}_{AB} \nabla_D \bar{C}^{BD}] (\bar{u}, x) dS^2 d\bar{u}. \end{aligned}$$

Proceed in the same way as in the case of angular momentum, we obtain

$$\begin{aligned} & \left(\tilde{C}_f^k(+)-\tilde{C}_f^k(-) \right) - \left(\tilde{C}^k(+)-\tilde{C}^k(-) \right) \\ &= \frac{1}{4} \int_{-\infty}^{+\infty} \int_{S^2} -\tilde{X}^k |N|^2 f - \nabla^A \tilde{X}^k |N|^2 \nabla_A f dS^2 du \\ & \quad + \frac{1}{4} \int_{-\infty}^{+\infty} \int_{S^2} \nabla^A \tilde{X}^k (-2 \nabla_A \nabla_B f + \Delta f \sigma_{AB}) \nabla_D N^{BD} + \nabla^A \tilde{X}^k N_{AB} \nabla^B (\Delta + 2) f dS^2 du \end{aligned}$$

We simplify the second integral as

$$\begin{aligned} & \int_{S^2} -2 \tilde{X}^k \nabla_A f \nabla_D N^{AD} + 2 \nabla^A \tilde{X}^k \nabla_A f \nabla_B \nabla_D N^{BD} - 2 \nabla^A \tilde{X}^k \nabla^B N_{AB} \cdot f \\ &= \int_{S^2} 2 \left(\nabla^A \tilde{X}^k \nabla_A f \nabla_B \nabla_D N^{BD} + \tilde{X}^k \nabla^A \nabla^B N_{AB} \cdot f \right) \end{aligned}$$

and the mass loss formula $\partial_u m = \frac{1}{4} \nabla^A \nabla^B N_{AB} - \frac{1}{8} |N|^2$ implies that

$$\begin{aligned} & \left(\tilde{C}_f^k(+)-\tilde{C}_f^k(-) \right) - \left(\tilde{C}^k(+)-\tilde{C}^k(-) \right) \\ &= \int_{S^2} 2 \tilde{X}^k f (m(+)-m(-)) + 2 \nabla^A \tilde{X}^k (m(+)-m(-)) \nabla_A f \\ &= \int_{S^2} \left(6 \tilde{X}^k (m(+)-m(-)) - 2 \nabla^A \tilde{X}^k \nabla_A (m(+)-m(-)) \right) f. \end{aligned}$$

Hence, $\tilde{C}^k(+)-\tilde{C}^k(-)$ is invariant under arbitrary supertranslation if and only if $6 \tilde{X}^k (m(+)-m(-)) - 2 \nabla^A \tilde{X}^k \nabla_A (m(+)-m(-))$ is supported in the $l \leq 1$ modes.

Multiplying the expression by \tilde{X}^k and summing over $k = 1, 2, 3$, we get $m(+)-m(-)$ is supported in the $l \leq 2$ modes. However, a direct computations shows that if $m(+)-m(-)$ contains a $l = 2$ mode, then $6 \tilde{X}^k (m(+)-m(-)) - 2 \nabla^A \tilde{X}^k \nabla_A (m(+)-m(-))$ contains a $l = 3$ mode. Similarly, if $m(+)-m(-)$ contains a $l = 1$ mode, then $6 \tilde{X}^k (m(+)-m(-)) - 2 \nabla^A \tilde{X}^k \nabla_A (m(+)-m(-))$ contains a $l = 2$ mode. Thus, $m(+)-m(-)$ is

constant if and only if $\tilde{C}^k(+)$ – $\tilde{C}^k(-)$ is invariant under arbitrary supertranslation \square

5.2. Total flux of the CWY conserved quantities. In this subsection, we show that the total flux of the CWY angular momentum and center of mass is supertranslation invariant. We decompose f into its modes:

$$f = \alpha_0 + \alpha_i \tilde{X}^i + f_{l \geq 2}$$

and let $J^k(\pm)$ be the limits of the CWY angular momentum in the u coordinate and $J_f^k(\pm)$ be the limits of the CWY angular momentum in the \bar{u} coordinate. We have

Theorem 5.4. *Suppose the news tensor decays as*

$$N_{AB}(u, x) = O(|u|^{-1-\varepsilon}) \text{ as } u \rightarrow \pm\infty.$$

Then the total flux of J^k is supertranslation invariant. Namely,

$$\left(J_f^k(+) - J_f^k(-) \right) - \left(J^k(+) - J^k(-) \right) = \alpha_i \varepsilon^{ik}_j (P^j(+) - P^j(-)).$$

Proof. Note that

$$J^k = \tilde{J}^k - \int_{S^2} Y^A c \nabla_A m \quad (5.11)$$

where $Y^A = \varepsilon^{AB} \nabla_B \tilde{X}^k$.

The assumption (5.3) on the decay of news tensor implies that the limit of mass aspect function is invariant of supertranslation

$$\lim_{\bar{u} \rightarrow \pm\infty} \bar{m}(\bar{u}, x) = \lim_{u \rightarrow \pm\infty} m(u, x) \text{ or } \bar{m}(\pm) = m(\pm). \quad (5.12)$$

Moreover, we have

$$\lim_{\bar{u} \rightarrow \pm\infty} \bar{C}_{AB}(\bar{u}, x) = \lim_{u \rightarrow \pm\infty} C_{AB}(u, x) - 2\nabla_A \nabla_B f + \Delta f \sigma_{AB}. \quad (5.13)$$

If we denote the closed potential of $\lim_{\bar{u} \rightarrow +\infty} \bar{C}_{AB}$ and $\lim_{\bar{u} \rightarrow +\infty} C_{AB}$ by $\bar{c}(+)$ and $c(+)$ respectively, we have

$$\bar{c}(+) = c(+) - 2f_{\ell \geq 2} \quad (5.14)$$

as functions on S^2 . Evaluating the definition of the CWY angular momentum (5.11) at $+\infty$ gives

$$J^k(+) = \tilde{J}^k(+) - \int_{S^2} Y^A c(+) \nabla_A m(+),$$

and

$$J_f^k(+) = \tilde{J}_f^k(+) - \int_{S^2} Y^A \bar{c}(+) \nabla_A \bar{m}(+).$$

Taking the difference and applying (5.12) and (5.14), we derive

$$J_f^k(+) - J^k(+) = \tilde{J}_f^k(+) - \tilde{J}^k(+) + 2 \int_{S^2} f_{\ell \geq 2} Y^A \nabla_A m(+).$$

We derive a similar equation at $-\infty$ and thus

$$\begin{aligned} & \left(J_f^k(+)-J_f^k(-) \right) - \left(J^k(+)-J^k(-) \right) \\ &= \left(\tilde{J}_f^k(+)-\tilde{J}_f^k(-) \right) - \left(\tilde{J}^k(+)-\tilde{J}^k(-) \right) + 2 \int_{S^2} f_{\ell \geq 2} Y^A \nabla_A (m(+)-m(-)) \\ &= -2 \int_{S^2} f_{\ell \leq 1} Y^A \nabla_A (m(+)-m(-)) \end{aligned}$$

by (5.8). It follows that

$$\left(J_f^k(+)-J_f^k(-) \right) - \left(J^k(+)-J^k(-) \right) = \alpha_i \varepsilon^{ik} (P^j(+)-P^j(-)).$$

□

Let $C^k(\pm)$ be the limits of the CWY center of mass in the u coordinate and $C_f^k(\pm)$ be the limits of the CWY center of mass in the \bar{u} coordinate.

Theorem 5.5. *Suppose the news tensor decays as*

$$N_{AB}(u, x) = O(|u|^{-1-\varepsilon}) \text{ as } u \rightarrow \pm\infty,$$

then the total flux of C^k is supertranslation invariant. Namely,

$$\left(C_f^k(+)-C_f^k(-) \right) - \left(C^k(+)-C^k(-) \right) = \alpha_0 \left(P^k(+)-P^k(-) \right) + \alpha_k (E(+)-E(-)).$$

Proof. We write

$$C^k = \tilde{C}^k - \int_{S^2} c \nabla^A \tilde{X}^k \nabla_A m + 3 \int_{S^2} c \tilde{X}^k m + \Xi(\underline{c}, m), \quad (5.15)$$

where Ξ is an integral over S^2 that involves only \underline{c} and m . Since the last three integrals have limits at $u = \pm\infty$, the mass loss formula now implies

$$\begin{aligned} & C^k(+)-C^k(-) \\ &= \frac{1}{4} \int_{-\infty}^{+\infty} \int_{S^2} u |N|^2(u, x) \tilde{X}^k + \nabla^A \tilde{X}^k [C_{AB} \nabla_D N^{BD} - N_{AB} \nabla_D C^{BD}] (u, x) dS^2 du \\ & \quad - \int_{S^2} c(+)\nabla^A \tilde{X}^k \nabla_A m(+) + \int_{S^2} c(-)\nabla^A \tilde{X}^k \nabla_A m(-) \\ & \quad + 3 \int_{S^2} c(+)\tilde{X}^k m(+) - 3 \int_{S^2} c(-)\tilde{X}^k m(-) \\ & \quad + \Xi(\underline{c}(+), m(+)) - \Xi(\underline{c}(-), m(-)). \end{aligned}$$

By (5.13), $\underline{c}(\pm)$ is invariant under supertranslation. We apply (5.12) and (5.14) to get

$$\begin{aligned} & \left(C_f^k(+)-C_f^k(-) \right) - \left(C^k(+)-C^k(-) \right) \\ &= \left(\tilde{C}_f^k(+)-\tilde{C}_f^k(-) \right) - \left(\tilde{C}^k(+)-\tilde{C}^k(-) \right) \\ & \quad + 2 \int_{S^2} f_{\ell \geq 2} \nabla^A \tilde{X}^k \nabla_A (m(+)-m(-)) - 6 \int_{S^2} f_{\ell \geq 2} \tilde{X}^k (m(+)-m(-)) \\ &= -2 \int_{S^2} f_{\ell \leq 1} \nabla^A \tilde{X}^k \nabla_A (m(+)-m(-)) + 6 \int_{S^2} f_{\ell \leq 1} \tilde{X}^k (m(+)-m(-)). \end{aligned}$$

We obtain

$$\begin{aligned} \left(C_f^k(+)-C_f^k(-) \right) - \left(C^k(+)-C^k(-) \right) &= 2 \int_{S^2} (\alpha_0 \tilde{X}^k + \alpha_k) (m(+)-m(-)) \\ &= \alpha_0 \left(P^k(+)-P^k(-) \right) + \alpha_k (E(+)-E(-)). \end{aligned}$$

□

6. SPACETIME WITH ZERO NEWS

In this section, we consider a non-radiative spacetime in the sense that the news vanishes. This includes all model spacetimes such as Minkowski and Kerr. First, we show that the CWY angular momentum and center of mass are constant.

Lemma 6.1. *Suppose the news $N_{AB}(u, x) \equiv 0$ in a Bondi-Sachs coordinate system (u, x) , then the CWY angular momentum $J^k(u)$ and CWY center of mass $C^k(u)$ are constant, i.e. independent of the retarded time u .*

Proof. The assumption implies $\partial_u m(u, x) = 0$, $\partial_u C_{AB}(u, x) = 0$ and thus

$$m(u, x) \equiv \dot{m}(x), C_{AB}(u, x) \equiv \dot{C}_{AB}(x)$$

and both potentials c and \underline{c} are independent of u as well.

We recall the definition of CWY angular momentum

$$J^k(u) = \int_{S^2} Y^A \left(N_A - \frac{1}{4} C_{AB} \nabla_D C^{DB} - c \nabla_A m \right) \quad (6.1)$$

where $Y^A = \epsilon^{AB} \nabla_B \tilde{X}^k$. Since c and m are both independent of u , our previous calculation shows

$$\begin{aligned} & \partial_u \int_{S^2} Y^A \left(N_A - \frac{1}{4} C_{AB} \nabla_D C^{DB} \right) \\ &= \frac{1}{4} \int_{S^2} \left[Y^A (C_{AB} \nabla_D N^{BD} - N_{AB} \nabla_D C^{BD}) + \tilde{X}^k \epsilon^{AB} (C_A^D N_{DB}) \right], \end{aligned}$$

the conclusion follows.

On the other hand, the CWY center of mass C^k is given by

$$\begin{aligned} & \int_{S^2} \nabla^A \tilde{X}^k \left(N_A - \frac{1}{4} C_{AB} \nabla_D C^{DB} - \frac{1}{16} \nabla_A (C^{DE} C_{DE}) \right) - \nabla^A \tilde{X}^k (c + u) \nabla_A m \\ & + \int_{S^2} \left(3\tilde{X}^k c m + 2\nabla^A \tilde{X}^k \epsilon_{AB} (\nabla^B \underline{c}) m - \frac{1}{16} \tilde{X}^k \nabla_A (\Delta + 2) \underline{c} \nabla^A (\Delta + 2) \underline{c} \right) \end{aligned} \quad (6.2)$$

Since all m , c , and \underline{c} are independent of u ,

$$\begin{aligned} & \partial_u C^k \\ & = \partial_u \int_{S^2} \nabla^A \tilde{X}^k \left(N_A - \frac{1}{4} C_{AB} \nabla_D C^{DB} - \frac{1}{16} \nabla_A (C^{DE} C_{DE}) \right) - \int_{S^2} \nabla^A \tilde{X}^k \nabla_A m \end{aligned} \quad (6.3)$$

Our previous calculation shows that the first term on the right hand side is

$$\int_{S^2} \nabla^A \tilde{X}^k \left(\nabla_A m + \frac{1}{4} C_{AB} \nabla_D N^{BD} - \frac{1}{4} N_{AB} \nabla_D C^{BD} \right),$$

and the conclusion follows. \square

Finally, we show that in a spacetime with vanishing news tensor, the angular momentum and center of mass themselves, not just their total flux, are invariant under supertranslation.

We pin down the exact formula for the angular momentum aspect on a spacetime with vanishing news. In this case, we have

$$\partial_u N_A(u, x) = \nabla_A m(u, x) - \frac{1}{4} \nabla^B P_{BA}(u, x)$$

where

$$P_{BA}(u, x) = (\nabla_B \nabla^E C_{EA} - \nabla_A \nabla^E C_{EB})(u, x)$$

Therefore

$$\partial_u N_A(u, x) = \nabla_A \dot{m}(x) - \frac{1}{4} \nabla^B \dot{P}_{BA}(x)$$

is independent of u . Integrating gives

$$N_A(u, x) = N_A(u_0, x) + (u - u_0) (\nabla_A \dot{m} - \frac{1}{4} \nabla^B \dot{P}_{BA}) \quad (6.4)$$

for any u and fixed u_0 .

Suppose (\bar{u}, x) is another Bondi-Sachs coordinate system that is related to (u, x) by a supertranslation $u = \bar{u} + f$ for $f \in C^\infty(S^2)$.

Recall the mass aspect $\bar{m}(\bar{u}, x)$, the shear $\bar{C}_{AB}(\bar{u}, x)$, and the news $\bar{N}_{AB}(\bar{u}, x)$ in the (\bar{u}, x) coordinate system are related to the mass aspect $m(u, x)$, the shear $C_{AB}(u, x)$, and the news $N_{AB}(u, x)$ in the (u, x) coordinate system through:

$$\begin{aligned}
\bar{m}(\bar{u}, x) &= m(\bar{u} + f, x) + \frac{1}{2}(\nabla^B N_{BD})(\bar{u} + f, x)\nabla^D f \\
&\quad + \frac{1}{4}(\partial_u N_{BD})(\bar{u} + f, x)\nabla^B f\nabla^D f + \frac{1}{4}N_{BD}(\bar{u} + f, x)\nabla^B\nabla^D f \\
\bar{C}_{AB}(\bar{u}, x) &= C_{AB}(\bar{u} + f(x), x) - 2\nabla_A\nabla_B f + \Delta f\sigma_{AB} \\
\bar{N}_{AB}(\bar{u}, x) &= N_{AB}(\bar{u} + f(x), x)
\end{aligned} \tag{6.5}$$

In particular, $\bar{N}_{AB}(\bar{u}, x) \equiv 0$ and $\bar{J}^k(\bar{u})$ and $\bar{C}^k(\bar{u})$ are independent of \bar{u} . In addition, we have

$$\begin{aligned}
\bar{m}(\bar{u}, x) &= \mathring{m}(x) = \mathring{m}(x) \\
\bar{C}_{AB}(\bar{u}, x) &= \mathring{C}_{AB}(x) = \mathring{C}_{AB}(x) - F_{AB} \\
\mathring{\bar{c}} &= \mathring{c} - 2f_{\ell \geq 2} \\
\mathring{\bar{c}} &= \mathring{c}
\end{aligned} \tag{6.6}$$

where $F_{AB} = 2\nabla_A\nabla_B f - \Delta f\sigma_{AB}$.

Finally the angular momentum aspect transforms by

$$\begin{aligned}
\bar{N}_A(\bar{u}, x) &= N_A(\bar{u} + f, x) + 3m(\bar{u} + f, x)\nabla_A f - \frac{3}{4}P_{BA}(\bar{u} + f, x)\nabla^B f \\
&= N_A(\bar{u} + f, x) + 3\mathring{m}\nabla_A f - \frac{3}{4}\mathring{P}_{BA}\nabla^B f.
\end{aligned}$$

See [10, (C.123)]. Note that the convention of angular momentum aspect there is $-3N_A$.

Combining with (6.4) and setting $u = \bar{u} + f$, we obtain

$$\bar{N}_A(\bar{u}, x) = N_A(u_0, x) + (\bar{u} - u_0 + f)(\nabla_A \mathring{m} - \frac{1}{4}\nabla^B \mathring{P}_{BA}) + 3\mathring{m}\nabla_A f - \frac{3}{4}\mathring{P}_{BA}\nabla^B f \tag{6.7}$$

for any \bar{u} and fixed u_0 .

Now fixing $\bar{u} = \bar{u}_0$, we consider the angular momentums

$$\begin{aligned}
\bar{J} &= \bar{J}(\bar{u}_0) = \int_{S^2} Y^A \left(\bar{N}_A - \frac{1}{4}\bar{C}_{AB}\nabla_D\bar{C}^{DB} - \mathring{\bar{c}}\nabla_A\mathring{m} \right) (\bar{u}_0, x) \\
J &= J(u_0) = \int_{S^2} Y^A \left(N_A - \frac{1}{4}\mathring{C}_{AB}\nabla_D\mathring{C}^{DB} - \mathring{c}\nabla_A\mathring{m} \right) (u_0, x)
\end{aligned}$$

where $Y^A = \epsilon^{AB} \nabla_B \tilde{X}^k$ and the center of mass

$$\begin{aligned}
\bar{C} &= \bar{C}(\bar{u}_0) = \int_{S^2} \nabla^A \tilde{X}^k \left(\bar{N}_A - \frac{1}{4} \bar{C}_{AB} \nabla_D \bar{C}^{DB} - \frac{1}{16} \nabla_A (\bar{C}^{DE} \bar{C}_{DE}) \right) (\bar{u}_0, x) \\
&+ \int_{S^2} \left(3 \tilde{X}^k \dot{\bar{c}} \dot{\bar{m}} - \nabla^A \tilde{X}^k (\dot{\bar{c}} + \bar{u}_0) \nabla_A \dot{\bar{m}} \right) \\
&+ \int_{S^2} \left(2 \nabla^A \tilde{X}^k \epsilon_{AB} (\nabla^B \dot{\bar{c}}) \dot{\bar{m}} - \frac{1}{16} \tilde{X}^k \nabla_A (\Delta + 2) \dot{\bar{c}} \nabla^A (\Delta + 2) \dot{\bar{c}} \right) \\
C &= C(u_0) = \int_{S^2} \nabla^A \tilde{X}^k \left(N_A - \frac{1}{4} \dot{C}_{AB} \nabla_D \dot{C}^{DB} - \frac{1}{16} \nabla_A (\dot{C}^{DE} \dot{C}_{DE}) \right) (u_0, x) \\
&+ \int_{S^2} \left(3 \tilde{X}^k \dot{c} \dot{m} - \nabla^A \tilde{X}^k (\dot{c} + u_0) \nabla_A \dot{m} \right) \\
&+ \int_{S^2} \left(2 \nabla^A \tilde{X}^k \epsilon_{AB} (\nabla^B \dot{c}) \dot{m} - \frac{1}{16} \tilde{X}^k \nabla_A (\Delta + 2) \dot{c} \nabla^A (\Delta + 2) \dot{c} \right)
\end{aligned}$$

We prove the following theorem:

Theorem 6.2. *On a spacetime with vanishing news, the CWY angular momentum and center of mass satisfy*

$$\bar{J} - J = -2 \int_{S^2} \epsilon^{AB} \nabla_B \tilde{X}^k f_{\ell \leq 1} \nabla_A \dot{m} \quad (6.8)$$

$$\bar{C} - C = \int_{S^2} \left(6 f_{\ell \leq 1} \tilde{X}^k \dot{m} - 2 f_{\ell \leq 1} \nabla^A \tilde{X}^k \nabla_A \dot{m} \right) \quad (6.9)$$

Proof. Taking the difference of \bar{J} and J and applying (6.6), we obtain

$$\begin{aligned}
\bar{J} - J &= \int_{S^2} Y^A [\bar{N}_A(\bar{u}_0, x) - N_A(u_0, x)] \\
&+ \frac{1}{4} \int_{S^2} Y^A [\dot{C}_{AB} \nabla_D F^{BD} + F_{AB} \nabla_D \dot{C}^{BD} - F_{AB} \nabla_D F^{BD}] \\
&+ 2 \int_{S^2} Y_A f_{\ell \geq 2} \nabla_A \dot{m}
\end{aligned}$$

We observe that $\int_{S^2} Y^A (F_{AB} \nabla_D F^{BD}) = 0$ and compute

$$\begin{aligned}
&\int_{S^2} Y^A [\bar{N}_A(\bar{u}_0, x) - N_A(u_0, x)] \\
&= \int_{S^2} Y^A \left[(\bar{u} - u_0 + f) (\nabla_A \dot{m} - \frac{1}{4} \nabla^B \dot{P}_{BA}) + 3 \dot{m} \nabla_A f - \frac{3}{4} \dot{P}_{BA} \nabla^B f \right] \\
&= \int_{S^2} Y^A \left[f \nabla_A \dot{m} - \frac{1}{4} f \nabla^B \dot{P}_{BA} + 3 \dot{m} \nabla_A f - \frac{3}{4} \dot{P}_{BA} \nabla^B f \right] \\
&= \int_{S^2} Y^A \left[-2f \nabla_A \dot{m} - \frac{1}{4} f \nabla^B \dot{P}_{BA} - \frac{3}{4} \dot{P}_{BA} \nabla^B f \right],
\end{aligned}$$

where we use $\int_{S^2} Y^A \nabla^B \dot{P}_{BA} = 0$ and $\int_{S^2} Y^A \nabla_A \dot{m} = 0$. Therefore,

$$\begin{aligned} \bar{J} - J &= \frac{1}{4} \int_{S^2} Y^A \left[\dot{C}_{AB} \nabla_D F^{BD} + F_{AB} \nabla_D \dot{C}^{BD} - f \nabla^B \dot{P}_{BA} - 3 \dot{P}_{BA} \nabla^B f \right] \\ &\quad - 2 \int_{S^2} Y_A f_{\ell \leq 1} \nabla_A \dot{m}. \end{aligned}$$

By Theorem B.1, the first integral vanishes and the result follows.

Taking the difference of \bar{C} and C and applying (6.6), we obtain

$$\begin{aligned} \bar{C} - C &= \int_{S^2} \nabla^A \tilde{X}^k \left[\bar{N}_A(\bar{u}_0, x) - N_A(u_0, x) \right] \\ &\quad + \frac{1}{4} \int_{S^2} \nabla^A \tilde{X}^k \left[\dot{C}_{AB} \nabla_D F^{BD} + F_{AB} \nabla_D \dot{C}^{BD} - F_{AB} \nabla_D F^{DB} \right. \\ &\quad \quad \quad \left. + \frac{1}{2} \nabla_A (C_{BD} F^{BD}) - \frac{1}{4} \nabla_A (F_{BD} F^{BD}) \right] \\ &\quad + \int_{S^2} \left(-6 f_{\ell \geq 2} \tilde{X}^k \dot{m} + (2 f_{\ell \geq 2} - \bar{u}_0 + u_0) \nabla^A \tilde{X}^k \nabla_A \dot{m} \right) \end{aligned}$$

We observe that

$$\int_{S^2} \nabla^A \tilde{X}^k \left[-F_{AB} \nabla_D F^{DB} - \frac{1}{4} \nabla_A (F_{BD} F^{BD}) \right] = 0$$

and compute

$$\begin{aligned} &\int_{S^2} \nabla^A \tilde{X}^k \left[\bar{N}_A(\bar{u}_0, x) - N_A(u_0, x) \right] \\ &= \int_{S^2} \nabla^A \tilde{X}^k \left[(\bar{u}_0 - u_0 + f) (\nabla_A \dot{m} - \frac{1}{4} \nabla^B \dot{P}_{BA}) + 3 \dot{m} \nabla_A f - \frac{3}{4} \dot{P}_{BA} \nabla^B f \right] \\ &= \int_{S^2} \nabla^A \tilde{X}^k \left[(\bar{u}_0 - u_0 + f) \nabla_A \dot{m} - \frac{1}{4} f \nabla^B \dot{P}_{BA} + 3 \dot{m} \nabla_A f - \frac{3}{4} \dot{P}_{BA} \nabla^B f \right] \\ &= \int_{S^2} \nabla^A \tilde{X}^k \left[(\bar{u}_0 - u_0 - 2f) \nabla_A \dot{m} - \frac{1}{4} f \nabla^B \dot{P}_{BA} - \frac{3}{4} \dot{P}_{BA} \nabla^B f \right] + \int_{S^2} (6f \tilde{X}^k \dot{m}), \end{aligned}$$

where we use $\int_{S^2} \nabla^A \tilde{X}^k \nabla^B \dot{P}_{BA} = 0$.

Putting everything together, we arrive at

$$\begin{aligned} &\bar{C} - C \\ &= \frac{1}{4} \int_{S^2} \nabla^A \tilde{X}^k \left[\dot{C}_{AB} \nabla_D F^{BD} + F_{AB} \nabla_D \dot{C}^{BD} + \frac{1}{2} \nabla_A (\dot{C}_{BD} F^{BD}) - f \nabla^B \dot{P}_{BA} - 3 \dot{P}_{BA} \nabla^B f \right] \\ &\quad + \int_{S^2} \left(6 f_{\ell \leq 1} \tilde{X}^k \dot{m} - 2 f_{\ell \leq 1} \nabla^A \tilde{X}^k \nabla_A \dot{m} \right) \end{aligned}$$

By Theorem B.2, the first integral vanishes and the result follows. \square

7. CONSERVATION LAW OF ANGULAR MOMENTUM AND A DUALITY
PARADIGM FOR NULL INFINITY

7.1. Conservation law of angular momentum. In this subsection, we derive a conservation law of angular momentum at \mathcal{I}^+ à la Christodoulou [7].

Suppose $I = (-\infty, +\infty)$ and \mathcal{I}^+ is complete extending from spatial infinity ($u = -\infty$) to timelike infinity ($u = +\infty$). Integrating the formula in Proposition 3.1 from $-\infty$ to $+\infty$ and projecting onto the $\ell = 1$ modes, we obtain

$$\epsilon^{AE}\nabla_E N_A(+\infty)_{\ell=1} - \epsilon^{AE}\nabla_E N_A(-\infty)_{\ell=1} = G_{\ell=1}, \quad (7.1)$$

where

$$G = \int_{-\infty}^{+\infty} \frac{1}{8} \nabla^A \nabla_A (\epsilon^{PQ} C_P^E N_{EQ}) + \frac{1}{2} \epsilon^{AE} \nabla_E (C_{AB} \nabla_D N^{DB})$$

Equation (7.1) should be considered as a conservation law for angular momentum that complements the conservation law for linear momentum of Christodoulou [7, Equation (13)], which in our notation is

$$\widehat{m}(+\infty)_{\ell=0,1} - \widehat{m}(-\infty)_{\ell=0,1} = -F_{\ell=0,1},$$

where

$$F = \frac{1}{8} \int_{-\infty}^{+\infty} N_{AB} N^{AB} \quad (7.2)$$

and follows from (3.3).

The above discussion can be carried over under the framework of stability of Minkowski spacetime, provided that we take Rizzi's definition of angular momentum [20, 21]. Recall from [7, 8] that two symmetric traceless 2-tensors Σ and Ξ are defined by

$$\lim_{C_u^+, r \rightarrow \infty} r^2 \widehat{\chi} = \Sigma, \quad \lim_{C_u^+, r \rightarrow \infty} r \widehat{\underline{\chi}} = \Xi$$

with

$$\frac{\partial \Sigma}{\partial u} = -\frac{1}{2} \Xi. \quad (7.3)$$

See Definition A.5 for the curvature components and their limits at null infinity.

Rizzi's definition of angular momentum [20, (3)] is given by (omitting the constant $\frac{1}{8\pi}$)

$$L(\Omega_{(i)}) = \int_{S^2} \Omega_{(i)}^A (I_A - \Sigma_{AB} \nabla_C \Sigma^{CB}), \quad i = 1, 2, 3 \quad (7.4)$$

where he *assumes* that the curvature component β satisfies $\lim_{r \rightarrow \infty} r^3 \beta_A = -I_A$. Here $\Omega_{(i)}^A$ corresponds to $\epsilon^{AB} \nabla_B \tilde{X}^i$. In the appendix, we show that I_A

and Σ_{AB} correspond to N_A and $-\frac{1}{2}C_{AB}$ in Bondi-Sachs coordinate system. Hence Rizzi's definition coincides with (1.2).

Using Bianchi identities, Rizzi derived the evolution formula [20, (4)]

$$\begin{aligned} \frac{\partial L}{\partial u} &= \int_{S^2} \Omega^A \left[\Xi_{AB} \nabla_C \Sigma^{CB} + \frac{1}{2} (\Sigma_B^C \nabla^B \Xi_{CA} - \Sigma_{AB} \nabla_C \Xi^{CB}) \right] \\ &= \frac{1}{2} \int_{S^2} \Omega^A (\Xi_{AB} \nabla_C \Sigma^{CB} - \Sigma_{AB} \nabla_C \Xi^{CB}) + \nabla^A \Omega^B \Sigma_B^C \Xi_{CA}, \end{aligned} \quad (7.5)$$

where the second line is obtained by integrating by parts the term $\Omega^A \Sigma_B^C \nabla^B \Xi_{CA}$.

Remark 7.1. The definition we take has the opposite sign to [20, (3),(4)]. The discrepancy comes from the fact that Kerr spacetime has angular momentum $-ma$ under our definition.

According to the main theorem of [9],

$$\underline{B} = O(|u|^{-\frac{3}{2}}) \quad (7.6)$$

as $|u| \rightarrow \infty$, where $\underline{B} = \lim_{r \rightarrow \infty} r^2 \underline{\beta}$. (see also [7], the paragraph after equation (8) where \underline{B} is denoted by \underline{B} there)

Estimate (7.6) and equation (2) of [7]

$$\operatorname{div} \Xi = \underline{B} \quad (7.7)$$

imply that

$$\Xi = O(|u|^{-\frac{3}{2}}) \quad (7.8)$$

as $|u| \rightarrow \infty$ and

$$\Sigma \rightarrow \Sigma^\pm \quad (7.9)$$

as $u \rightarrow \pm\infty$.

By (7.8) and (7.9), $\int_{-\infty}^{\infty} \frac{\partial L}{\partial u} du$ is finite and furnishes the difference of the angular momenta at timelike infinity ($u \rightarrow \infty$) and spatial infinity ($u \rightarrow -\infty$).

We can write this in the spirit of [7]. In general, the peeling fails and β decays as $\beta = o(r^{-\frac{7}{2}})$. Christodoulou [8] observed that Bianchi equation nevertheless implies that

$$R = \lim_{C_u^+, r \rightarrow \infty} r^4 \underline{D} \beta$$

exists. Moreover, one has

$$R = \nabla P + * \nabla Q + 2 \Sigma \cdot \underline{B}, \quad (7.10)$$

where $(P, Q) = \lim_{r \rightarrow \infty} (r^3 \rho, r^3 \sigma)$ and $\nabla, *, \cdot$ are taken with respect to standard metric σ on S^2 .

In order to exhibit a physically reasonable initial data set that has a complete Cauchy development without peeling, Christodoulou made the crucial assumption

$$\lim_{u \rightarrow -\infty} uR = R^- \neq 0, \quad (7.11)$$

which we adopt here.

From (7.11) he derived [8, (5)]

$$\beta = B_* r^{-4} \log r + Br^{-4} + o(r^{-4})$$

uniformly in u with 1-forms B_* and B on S^2 satisfying [8, (6)]

$$\frac{\partial B_*}{\partial u} = 0 \quad (7.12)$$

$$\frac{\partial B}{\partial u} = \frac{1}{2}R. \quad (7.13)$$

Moreover, using Bianchi equation, he derived that

$$\lim_{C_u^+, r \rightarrow \infty} r^4 \alpha = A_* \neq 0$$

exists. A_* is a symmetric traceless 2-tensor that is independent of u and satisfies

$$\operatorname{div} A_* = -B_*. \quad (7.14)$$

Definition 7.2. For a function f on S^2 , we denote the projection of f on the sum of zeroth and first eigenspaces of Δ by $f_{[1]}$. Namely, $f_{[1]} = f_{\ell=0} + f_{\ell=1}$. For a 1-form $\omega_A = \nabla_A f + \epsilon_{AB} \nabla^B g$, we denote $\omega_{A[1]} = \nabla_A f_{\ell=1} + \epsilon_{AB} \nabla^B g_{\ell=1}$.

Since the spherical tangent vectors ∂_A have length $O(r)$, we have the correspondence

$$B_A = -I_A. \quad (7.15)$$

By (7.14), $B_{*[1]} = 0$ and we integrate (7.10) to get

$$(I_A(u_2, x) - I_A(u_1, x))_{[1]} = -\frac{1}{2} \int_{u_1}^{u_2} \nabla_A P_{\ell=1} + \epsilon_{AB} \nabla^B Q_{\ell=1} + (2\Sigma_{AB} \underline{B}^B)_{[1]} du.$$

By (7.6) and (7.9), the last term is integrable on $(-\infty, \infty)$. For the first two terms, we observe that equations (10, 11) of [8]

$$\frac{\partial P}{\partial u} = -\frac{1}{2} \operatorname{div} \underline{B} + \frac{1}{2} \Sigma \cdot \frac{\partial \Xi}{\partial u} \quad (7.16)$$

$$\frac{\partial Q}{\partial u} = -\frac{1}{2} \operatorname{curl} \underline{B} + \frac{1}{2} \Sigma \wedge \frac{\partial \Xi}{\partial u} \quad (7.17)$$

infer that $P_{\ell=1}(u, x) = a_i \tilde{X}^i + O(|u|^{-\frac{3}{2}})$ and $Q_{\ell=1}(u, x) = b_i \tilde{X}^i + O(|u|^{-\frac{3}{2}})$ for some constants a_i, b_i independent of u . Thanks to the main theorem of [9], $P - P_{\ell=0}, Q - Q_{\ell=0} = O(|u|^{-\frac{1}{2}})$, we have $a_i = 0, b_i = 0$. Thus $P_{\ell=1}, Q_{\ell=1}$ are also integrable on $(-\infty, \infty)$.

We conclude $\lim_{u \rightarrow \infty} I_{A[1]}(u, x) - \lim_{u \rightarrow -\infty} I_{A[1]}(u, x)$ exists and is given by

$$-\frac{1}{2} \int_{-\infty}^{\infty} (\nabla_A P_{\ell=1} + \epsilon_{AB} \nabla^B Q_{\ell=1} + (2\Sigma_{AB} \underline{B}^B)_{[1]}) du'.$$

By (7.3) and Rizzi's definition (7.4), we can interpret the following formula as a conservation law of angular momentum

$$\begin{aligned} & \lim_{u \rightarrow \infty} (I_A - \Sigma_{AB} \nabla_C \Sigma^{CB})_{[1]} - \lim_{u \rightarrow -\infty} (I_A - \Sigma_{AB} \nabla_C \Sigma^{CB})_{[1]} \\ &= \frac{1}{2} \int_{-\infty}^{\infty} -\nabla_A P_{\ell=1} - \epsilon_{AB} \nabla^B Q_{\ell=1} + (\Xi_{AB} \nabla_C \Sigma^{CB} - \Sigma_{AB} \nabla_C \Xi^{CB})_{[1]} du. \end{aligned} \quad (7.18)$$

From Proposition A.2 and Proposition A.6, it follows that the co-closed part of the above conservation law is equivalent to the total flux of the classical angular momentum in a Bondi-Sachs coordinate system.

7.2. A duality paradigm for null infinity. In this subsection, we describe a duality paradigm for null infinity which creates a pair of dual spacetimes with the same classical conserved quantities.

Corollary 7.3 (Corollary 1.7). *Given a set of null infinity data (m, N_A, C_{AB}, N_{AB}) defined on $[u_1, u_2] \times S^2$, there exists a dual set of null infinity data $(m^*, N_A^*, C_{AB}^*, N_{AB}^*)$ that has the same (classical) energy, linear momentum, angular momentum, and center of mass.*

Proof. Define $C_{AB}^* = \varepsilon_2(C_{AB})$ on $[u_1, u_2] \times S^2$. Then $N_{AB}^* = \partial_u C_{AB}^* = \varepsilon_2(N_{AB})$. Define $m^*(u, x)$ by the differential equation

$$\begin{cases} m^*(u_1, x) = m(u_1, x) \\ \partial_u m^* = \frac{1}{4} \nabla^A \nabla^B N_{AB}^* - N_{AB}^* N^{*AB} \end{cases}$$

and then define N_A^* by the differential equation

$$\begin{cases} N_A^*(u_1, x) = N_A(u_1, x) \\ \partial_u N_A^* = \nabla_A m^* - \frac{1}{4} \nabla^D (\nabla_D \nabla^E C_{EA}^* - \nabla_A \nabla^E C_{ED}^*) \\ \quad + \frac{1}{4} \nabla_A (C_{BE}^* N^{*BE}) - \frac{1}{4} \nabla_B (C^{*BD} N_{DA}^*) + \frac{1}{2} C_{AB}^* \nabla_D N^{*DB}. \end{cases}$$

For this subsection alone, we denote the classical conserved quantities of the infinity data (m, N_A, C_{AB}, N_{AB}) by E, P^k, J^k, C^k and denote the classical conserved quantities of the data $(m^*, N_A^*, C_{AB}^*, N_{AB}^*)$ by $E^*, P^{*k}, J^{*k}, C^{*k}$, we have

$$E^*(u_1) = E(u_1), P^{*k}(u_1) = P^k(u_1)$$

and since $C_A^*{}^D \nabla^B C_{BD}^* = C_A^D \nabla^B C_{BD}, C_{DE}^* C^{*DE} = C_{DE} C^{DE}$,

$$J^{*k}(u_1) = J^k(u_1), C^{*k}(u_1) = C^k(u_1).$$

It remains to show that the evolutions of the conserved quantities are identical. Recall that the potentials of C_{AB}^* and N_{AB}^* are given by $(-\underline{c}, c)$

and $(-\underline{n}, n)$. We observe that replacing $(c, \underline{c}, n, \underline{n})$ by $(-\underline{c}, c, -\underline{n}, n)$ does not change the following expressions

$$\begin{aligned}\partial_u E &= -\frac{1}{8} \int_{S^2} [n\Delta(\Delta+2)n + \underline{n}\Delta(\Delta+2)\underline{n}], \\ \partial_u P^k &= -\frac{1}{8} \int_{S^2} \tilde{X}^k [((\Delta+2)n)^2 + ((\Delta+2)\underline{n})^2 - 4\epsilon^{AB}\nabla_A n \nabla_B(\Delta+2)\underline{n}], \\ \partial_u J^k &= \frac{1}{8} \int_{S^2} \tilde{X}^k \epsilon^{AB} [\nabla_A c \nabla_B \Delta(\Delta+2)n + \nabla_A \underline{c} \nabla_B \Delta(\Delta+2)\underline{n}] \\ \partial_u C^k &= \frac{1}{8} \int_{S^2} \tilde{X}^k [((\Delta+2)n)^2 + ((\Delta+2)\underline{n})^2 - 4\epsilon^{AB}\nabla_A n \nabla_B(\Delta+2)\underline{n}] \\ &\quad + \frac{1}{16} \int_{S^2} [\tilde{X}^k (\Delta(\Delta+2)\underline{c}(\Delta+2)\underline{n} - \Delta(\Delta+2)\underline{n}(\Delta+2)\underline{c})] \\ &\quad + \frac{1}{16} \int_{S^2} [\tilde{X}^k (\Delta(\Delta+2)c(\Delta+2)n - \Delta(\Delta+2)n(\Delta+2)c)]\end{aligned}$$

This finishes the proof. \square

8. THE CASE OF QUADRUPOLE MOMENTS

In this section, we consider the case of generalized quadrupole moments. Namely, all $c, \underline{c}, n, \underline{n}$ are (-6) eigenfunctions (or $\ell = 2$ spherical harmonics). Therefore, $c = \sum c_{ij}(u)\tilde{X}^i\tilde{X}^j$, $\underline{c} = \sum \underline{c}_{ij}(u)\tilde{X}^i\tilde{X}^j$, $n = \sum n_{ij}(u)\tilde{X}^i\tilde{X}^j$, $\underline{n} = \sum \underline{n}_{ij}(u)\tilde{X}^i\tilde{X}^j$ with $\partial_u c_{ij} = n_{ij}$ and $\partial_u \underline{c}_{ij} = \underline{n}_{ij}$.

8.1. Classical conserved quantities. Next we compute the evolution of classical angular momentum and center of mass for quadrupole moments.

Lemma 8.1. *Suppose f_{ij} and g_{ij} are both symmetric, traceless 3×3 matrices. Then*

$$\int_{S^2} (f_{ij}\tilde{X}^i\tilde{X}^j)^2 = \frac{8\pi}{15} \sum_{ij} f_{ij}^2 \quad (8.1)$$

$$\int_{S^2} \tilde{X}^p \epsilon^{AB} \nabla_A (f_{ij}\tilde{X}^i\tilde{X}^j) \nabla_B (g_{kl}\tilde{X}^k\tilde{X}^l) = \frac{16\pi}{15} \sum_j f_{ij} g_{jk} \epsilon^{ikp}. \quad (8.2)$$

Proof. Both formulae follow from Lemma 5.3 of [6]

$$\int_{S^2} \tilde{X}^i \tilde{X}^j \tilde{X}^k \tilde{X}^l = \frac{4\pi}{15} (\delta_{ij}\delta_{kl} + \delta_{ik}\delta_{jl} + \delta_{il}\delta_{jk}).$$

\square

Combining the above lemma with Theorem 1.3 and Proposition 4.1, we conclude that

Proposition 8.2. *Suppose $c = \sum c_{ij}(u)\tilde{X}^i\tilde{X}^j$, $\underline{c} = \sum \underline{c}_{ij}(u)\tilde{X}^i\tilde{X}^j$, $n = \sum n_{ij}(u)\tilde{X}^i\tilde{X}^j$, $\underline{n} = \sum \underline{n}_{ij}(u)\tilde{X}^i\tilde{X}^j$, then*

$$\begin{aligned}\partial_u E &= -\frac{8\pi}{5} \left(\sum_{ij} n_{ij}^2 + \sum_{ij} \underline{n}_{ij}^2 \right) \\ \partial_u P^k &= -\frac{32\pi}{15} \sum n_{ij} \underline{n}_{jp} \epsilon^{ipk} \\ \partial_u \tilde{J}^k &= \frac{16\pi}{5} \sum (c_{ij} n_{jp} + \underline{c}_{ij} \underline{n}_{jp}) \epsilon^{ipk} \\ \partial_u \tilde{C}^k &= \frac{32u\pi}{15} \sum n_{ij} \underline{n}_{jp} \epsilon^{ipk}.\end{aligned}\tag{8.3}$$

8.2. CWY angular momentum and center of mass. Next we compute the evolution of the CWY angular momentum and center of mass for quadrupole moments. We need the following lemma.

Lemma 8.3. *Suppose the potentials of the news tensor are of mode $\ell = 2$. Namely,*

$$N_{AB} = \nabla_A \nabla_B n - \frac{1}{2} \Delta n \sigma_{AB} + \frac{1}{2} (\epsilon_{AC} \nabla_B \nabla^C \underline{n} + \epsilon_{BC} \nabla_A \nabla^C \underline{n})$$

where $n = \sum_{ij} n_{ij} \tilde{X}^i \tilde{X}^j$ and $\underline{n} = \sum_{ij} \underline{n}_{ij} \tilde{X}^i \tilde{X}^j$ satisfy $\sum_i n_{ii} = \sum_i \underline{n}_{ii} = 0$. Introduce two $\ell = 2$ spherical harmonics

$$\begin{aligned}Q &= \sum_{i,k,l} (n_{ik} n_{il} \tilde{X}^k \tilde{X}^l) - \frac{1}{3} \sum_{i,j} n_{ij}^2, \\ \underline{Q} &= \sum_{i,k,l} (\underline{n}_{ik} \underline{n}_{il} \tilde{X}^k \tilde{X}^l) - \frac{1}{3} \sum_{i,j} \underline{n}_{ij}^2.\end{aligned}$$

Then

(1) the $\ell = 2$ component of $N_{AB} N^{AB}$ is

$$-\frac{48}{7} Q - \frac{48}{7} \underline{Q},$$

(2) the odd mode component of $N_{AB} N^{AB}$ is

$$8\epsilon_m^{ik} n_{ij} \underline{n}_{kl} \tilde{X}^m (\delta^{jl} - \tilde{X}^j \tilde{X}^l).$$

Here ϵ_{ijk} is the Levi-Civita symbol in three dimensions.

Proof. Note that the even-mode components ($\ell = 0, 2, 4$) and odd-mode components ($\ell = 1, 3$) of $N_{AB} N^{AB}$ are given by

$$\nabla_A \nabla_B n \nabla^A \nabla^B n - \frac{1}{2} (\Delta n)^2 + \nabla_A \nabla_B \underline{n} \nabla^A \nabla^B \underline{n} - \frac{1}{2} (\Delta \underline{n})^2$$

and

$$\begin{aligned}
& (\nabla^A \nabla^B n - \frac{1}{2} \Delta n \sigma^{AB}) (\epsilon_{AC} \nabla_B \nabla^C \underline{n} + \epsilon_{BC} \nabla_A \nabla^C \underline{n}) \\
&= 2n_{ij} \nabla^A \tilde{X}^i \nabla^B \tilde{X}^j (\epsilon_{AC} \nabla_B \nabla^C \underline{n} + \epsilon_{BC} \nabla_A \nabla^C \underline{n}) \\
&= 2n_{ij} \nabla^A \tilde{X}^i \nabla^B \tilde{X}^j \cdot 2\underline{n}_{kl} (\epsilon_{AC} \nabla_B \tilde{X}^k \nabla^C \tilde{X}^l + \epsilon_{BC} \nabla_A \tilde{X}^k \nabla^C \tilde{X}^l) \\
&= 8\epsilon^ik_m n_{ij} \underline{n}_{kl} \tilde{X}^m (\delta^{jl} - \tilde{X}^j \tilde{X}^l)
\end{aligned}$$

respectively. In the last equality we use the identity $\epsilon_{AB} \nabla_A \tilde{X}^i \nabla_B \tilde{X}^j = \epsilon_{ijk} \tilde{X}^k$ and $\nabla_B \tilde{X}^i \nabla^B \tilde{X}^j = \delta^{ij} - \tilde{X}^i \tilde{X}^j$.

For (1), we compute

$$\nabla_A \nabla_B n \nabla^A \nabla^B n = 20n^2 - 8Q + \frac{4}{3} \sum_{ij} n_{ij}^2.$$

Since the space of $\ell = 4$ spherical harmonics is spanned by

$$\begin{aligned}
& \tilde{X}^i \tilde{X}^j \tilde{X}^k \tilde{X}^l + \frac{1}{35} (\delta^{ij} \delta^{kl} + \delta^{ik} \delta^{jl} + \delta^{il} \delta^{jk}) \\
& - \frac{1}{7} (\tilde{X}^i \tilde{X}^j \delta^{kl} + \tilde{X}^i \tilde{X}^k \delta^{jl} + \tilde{X}^i \tilde{X}^l \delta^{jk} + \tilde{X}^j \tilde{X}^k \delta^{il} + \tilde{X}^j \tilde{X}^l \delta^{ik} + \tilde{X}^k \tilde{X}^l \delta^{ij}),
\end{aligned} \tag{8.4}$$

the $\ell = 2$ component of n^2 is $\frac{4}{7}Q$. Putting these together, we obtain (1). \square

In the case of quadrupole moments, the CWY angular momentum and center of mass take the form:

$$J^k = \int_{S^2} \epsilon^{AB} \nabla_B \tilde{X}^k [N_A - \frac{1}{4} C_A^D \nabla^B C_{DB} - c \nabla_A m] \tag{8.5}$$

$$C^k = \int_{S^2} \nabla^A \tilde{X}^k [N_A - u \nabla_A m - \frac{1}{4} C_A^D \nabla^B C_{DB} - \frac{1}{16} \nabla_A (C_{DE} C^{DE}) - 2\underline{c} \epsilon_{AB} \nabla^B m], \tag{8.6}$$

Therefore,

$$\begin{aligned}
J^k &= \tilde{J}^k - \int_{S^2} \tilde{X}^k \epsilon^{AB} \nabla_A c \nabla_B \hat{m} \\
C^k &= \tilde{C}^k + 2 \int_{S^2} \tilde{X}^k \epsilon^{AB} \nabla_A \underline{c} \nabla_B \hat{m} + \frac{1}{4} \int_{S^2} \tilde{X}^k \epsilon^{AB} \nabla_A \underline{c} \nabla_B \Delta (\Delta + 2) c,
\end{aligned}$$

where we use (3.2) and (2.7).

The evolution formulae for J^k and C^k are thus

$$\begin{aligned}
\partial_u J^k &= \partial_u \tilde{J}^k - \int_{S^2} \tilde{X}^k \epsilon^{AB} \nabla_A n \nabla_B \hat{m} - \int_{S^2} \tilde{X}^k \epsilon^{AB} \nabla_A c \nabla_B \partial_u \hat{m} \\
\partial_u C^k &= \partial_u \tilde{C}^k + 2 \int_{S^2} \tilde{X}^k \epsilon^{AB} \nabla_A \underline{n} \nabla_B \hat{m} + 2 \int_{S^2} \tilde{X}^k \epsilon^{AB} \nabla_A \underline{c} \nabla_B \partial_u \hat{m} \\
&\quad + \frac{1}{4} \int_{S^2} \tilde{X}^k \epsilon^{AB} \nabla_A \underline{n} \nabla_B \Delta (\Delta + 2) c + \frac{1}{4} \int_{S^2} \tilde{X}^k \epsilon^{AB} \nabla_A \underline{c} \nabla_B \Delta (\Delta + 2) n
\end{aligned}$$

By Lemma 2.1, only the $\ell = 2$ mode components of \widehat{m} and $\partial_u \widehat{m}$ will survive in the above integrals.

Denote the $\ell = 2$ mode of \widehat{m} by $\widehat{m}_{\ell=2} = \widehat{m}_{kl} \widetilde{X}^k \widetilde{X}^l$. By Lemma 8.3, we get

$$\partial_u \widehat{m}_{kl} = \frac{6}{7} \left[\sum_i (n_{ik} n_{il} + \underline{n}_{ik} \underline{n}_{il}) - \frac{1}{3} \delta_{kl} \sum_{i,j} (n_{ij}^2 + \underline{n}_{ij}^2) \right].$$

By (8.2), we obtain the evolution equation of J^k and C^k :

Proposition 8.4. *Suppose $c = \sum c_{ij}(u) \widetilde{X}^i \widetilde{X}^j$, $\underline{c} = \sum \underline{c}_{ij}(u) \widetilde{X}^i \widetilde{X}^j$, $n = \sum n_{ij}(u) \widetilde{X}^i \widetilde{X}^j$, $\underline{n} = \sum \underline{n}_{ij}(u) \widetilde{X}^i \widetilde{X}^j$, then*

$$\begin{aligned} \partial_u E &= -\frac{8\pi}{5} \left(\sum_{ij} n_{ij}^2 + \sum_{ij} \underline{n}_{ij}^2 \right) \\ \partial_u P^k &= -\frac{32\pi}{15} \sum_{i,j,p} n_{ij} \underline{n}_{jp} \epsilon^{ipk} \\ \partial_u J^k &= \frac{16\pi}{15} \sum_{i,j,p} (3c_{ij} n_{jp} + 3\underline{c}_{ij} \underline{n}_{jp} - n_{ij} \widehat{m}_{jp} - c_{ij} \partial_u \widehat{m}_{jp}) \epsilon^{ipk} \\ \partial_u C^k &= \frac{16\pi}{15} \sum_{i,j,p} (2\underline{n}_{ij} \widehat{m}_{jp} + 2\underline{c}_{ij} \partial_u \widehat{m}_{jp} + 6\underline{n}_{ij} c_{jp} + 6\underline{c}_{ij} n_{jp}) \epsilon^{ipk} \\ &\quad + \frac{32u\pi}{15} \sum n_{ij} \underline{n}_{jp} \epsilon^{ipk}, \end{aligned} \tag{8.7}$$

where \widehat{m}_{kl} is given by

$$\partial_u \widehat{m}_{kl} = \frac{6}{7} \left[\sum_i (n_{ik} n_{il} + \underline{n}_{ik} \underline{n}_{il}) - \frac{1}{3} \delta_{kl} \sum_{i,j} (n_{ij}^2 + \underline{n}_{ij}^2) \right]$$

APPENDIX A. CHRISTODOULOU-KLAINERMAN CONNECTION COEFFICIENTS AND CURVATURE COMPONENTS IN BONDI-SACHS FORMALISM

We write the limit of connection coefficients and curvature components defined in [9, 7, 8] in terms of the Bondi-Sachs metric coefficients.

We choose the null vector fields $L = \frac{\partial}{\partial r}$ and $\underline{L} = \frac{2}{U} (\partial_u - W^D \partial_D - \frac{V}{2} \partial_r)$, which satisfy $\langle L, \underline{L} \rangle = -2$.

Definition A.1. The second fundamental forms and torsion are defined by

$$\begin{aligned} \chi_{AB} &= \langle D_A L, \partial_B \rangle = \frac{1}{2} \text{tr} \chi g_{AB} + \widehat{\chi}_{AB} \\ \underline{\chi}_{AB} &= \langle D_A \underline{L}, \partial_B \rangle = \frac{1}{2} \text{tr} \underline{\chi} g_{AB} + \widehat{\underline{\chi}}_{AB} \\ \zeta_A &= \frac{1}{2} \langle D_A L, \underline{L} \rangle \end{aligned}$$

Their limit as $r \rightarrow \infty$ are defined by

$$\begin{aligned}\Sigma &= \lim_{r \rightarrow \infty} \widehat{\chi} \\ \Xi &= \lim_{r \rightarrow \infty} r^{-1} \widehat{\underline{\chi}} \\ Z &= \lim_{r \rightarrow \infty} r \zeta.\end{aligned}$$

They are related to the metric coefficients in the corresponding Bondi-Sachs coordinate system as follows:

Proposition A.2.

$$\begin{aligned}\Sigma_{AB} &= -\frac{1}{2} C_{AB} \\ \Xi_{AB} &= N_{AB} \\ Z_A &= -\frac{1}{2} \nabla^B C_{AB}\end{aligned}$$

Proof. Starting with $g_{AB} = r^2 \sigma_{AB} + r C_{AB} + O(1)$, the determinant condition gives $\text{tr} \chi = \frac{2}{r}$ and we compute

$$\chi_{AB} = r \sigma_{AB} + \frac{1}{2} C_{AB} + O(r^{-1})$$

to get $\Sigma_{AB} = -\frac{1}{2} C_{AB}$. Direct computation gives

$$\underline{\chi}_{AB} = r(-\sigma_{AB} + \partial_u C_{AB}) + O(1)$$

and hence $\text{tr} \underline{\chi} = -\frac{2}{r} + O(r^{-2})$ and $\widehat{\underline{\chi}}_{AB} = r \partial_u C_{AB} + O(1)$. The limit of torsion follows from $\zeta_A = -\frac{1}{r} W_A^{(-2)} + O(r^{-2})$. \square

Definition A.3. The mass aspect and conjugate mass aspect function of Christodoulou-Klainerman are defined by

$$\mu = K + \frac{1}{4} \text{tr} \chi \text{tr} \underline{\chi} - \text{div} \zeta \quad \underline{\mu} = K + \frac{1}{4} \text{tr} \chi \text{tr} \underline{\chi} + \text{div} \zeta.$$

Here K denotes the Gauss curvature of the two-sphere $r = \text{const}$. Their limits are defined by

$$\begin{aligned}N &= \lim_{r \rightarrow \infty} r^3 \mu \\ \underline{N} &= \lim_{r \rightarrow \infty} r^3 \underline{\mu}.\end{aligned}$$

We express them in terms of the Bondi-Sachs metric coefficients as follows:

Proposition A.4.

$$N = 2m + \frac{1}{2} \nabla^A \nabla^B C_{AB}, \quad \underline{N} = 2m - \frac{1}{2} \nabla^A \nabla^B C_{AB}$$

Proof. We compute $K = \frac{1}{r^2} + \frac{1}{2r^3} \nabla^A \nabla^B C_{AB} + O(r^{-4})$ and $\frac{1}{4} \text{tr} \chi \text{tr} \underline{\chi} = -\frac{1}{r^2} + \frac{1}{r^3} (2m - \frac{1}{2} \nabla^A \nabla^B C_{AB})$ and the assertion follows. \square

We turn to curvature components. The convention of Riemann curvature tensor is

$$\begin{aligned} R(X, Y)Z &= (D_X D_Y - D_Y D_X - D_{[X, Y]})Z \\ R(X, Y, W, Z) &= \langle R(X, Y)Z, W \rangle. \end{aligned}$$

Definition A.5. Define the curvature components

$$\begin{aligned} \underline{\alpha}_{AB} &= R(\partial_A, \underline{L}, \partial_B, \underline{L}) \\ \underline{\beta}_A &= \frac{1}{2} R(\partial_A, \underline{L}, \underline{L}, L) \\ \rho &= \frac{1}{4} R(\underline{L}, L, \underline{L}, L) \\ \sigma \not\epsilon_{AB} &= \frac{1}{2} R(\partial_A, \partial_B, \underline{L}, L) \\ \beta_A &= \frac{1}{2} R(\partial_A, L, \underline{L}, L) \end{aligned}$$

Here $\not\epsilon_{AB} dx^A \wedge dx^B$ is the area form of the two-sphere with respect to g_{AB} . Their limits are defined by

$$\begin{aligned} \underline{A}_{AB} &= \lim_{r \rightarrow \infty} r^{-1} \underline{\alpha}_{AB} \\ \underline{B}_A &= \lim_{r \rightarrow \infty} r \underline{\beta}_A \\ P &= \lim_{r \rightarrow \infty} r^3 \rho \\ Q &= \lim_{r \rightarrow \infty} r^3 \sigma \\ B_A &= \lim_{r \rightarrow \infty} r^3 \beta_A \end{aligned}$$

Note that $(\underline{A}, \underline{B})$ were denoted by (A, B) in [7].

We express them in terms of the Bondi-Sachs metric coefficients as follows:

Proposition A.6.

$$\begin{aligned} \underline{A}_{AB} &= -2\partial_u N_{AB} \\ \underline{B}_A &= \nabla^B N_{AB} \\ P &= -2m - \frac{1}{4} C_{AB} N^{AB} \\ Q &= \epsilon^{AB} \left(-\frac{1}{4} C_A^D N_{DB} - \frac{1}{2} \nabla_A \nabla^D C_{DB} \right) \\ B_A &= -N_A \end{aligned}$$

Proof. The formula for \underline{A} is obtained from (6) of [7], $2\frac{\partial \Xi}{\partial u} = -\underline{A}$, which is the rescaled limit of the propagation equation $\widehat{D}\widehat{\chi} = -\underline{\alpha}$.

The formula for \underline{B} is obtained from (2) of [7], $\nabla^B \Xi_{AB} = \underline{B}_A$, which is the rescaled limit of the Codazzi equation $\text{div}\widehat{\chi} - \widehat{\chi} \cdot \zeta = \frac{1}{2} (\nabla \text{tr}\widehat{\chi} - \text{tr}\widehat{\chi}\zeta) + \underline{\beta}$.

The formula for P and Q are obtained from (3) of [7],

$$\epsilon^{AB}\nabla_A Z_B = Q - \frac{1}{2}\Sigma \wedge \Xi, \quad \nabla^A Z_A = \underline{N} + P - \frac{1}{2}\Sigma \cdot \Xi,$$

which is the rescaled limit of the Hodge system

$$c\cancel{v}r\zeta = \sigma - \frac{1}{2}\widehat{\chi} \wedge \widehat{\underline{\chi}}, \quad d\cancel{v}\zeta = \underline{\mu} + \rho - \frac{1}{2}\widehat{\chi} \cdot \widehat{\underline{\chi}}.$$

Finally, we consider the Codazzi equation

$$d\cancel{v}\widehat{\chi} + \widehat{\chi} \cdot \zeta = \frac{1}{2}(\nabla\text{tr}\chi + \text{tr}\chi\zeta) - \beta.$$

Its leading order at $O(r^{-2})$ leads to (1) of [7] and its subleading order at $O(r^{-3})$ leads to

$$\begin{aligned} & \left(-\frac{1}{4}\partial_A|C|^2 + \frac{1}{2}C^{BD}\nabla_D C_{AB} + \frac{1}{4}\nabla_A C_D^E C_E^D + \frac{1}{2}\nabla_D C^{DE} C_{AE} \right) + \frac{1}{4}C_{AB}\nabla_D C^{BD} \\ & = \zeta_A^{(-2)} - B_A. \end{aligned}$$

We simplify the second term in the parentheses by the identity $\nabla_{(D}C_{B)A} = \nabla_A C_{BD} + \nabla^E C_{AE}\sigma_{BD} - \nabla^E C_{E(D}C_{B)A}$ and the left-hand side becomes $\frac{1}{8}\partial_A|C|^2 + \frac{1}{4}C_{AB}\nabla_D C^{BD}$. Direct computation yields $\zeta_A^{(-2)} = -N_A + \frac{1}{8}\partial_A|C|^2 + \frac{1}{4}C_{AB}\nabla_D C^{BD}$ and the formula for B follows. \square

APPENDIX B. INTEGRATION BY PART FORMULA

In this section, we prove two integration formula that are used to compute angular momentum and center of mass in spacetime with vanishing news.

Theorem B.1. *Let $Y^A = \epsilon^{AB}\nabla_B \tilde{X}^k$, $k = 1, 2, 3$. Let $F_{AB} = 2\nabla_A \nabla_B f - \Delta f \sigma_{AB}$ and $P_{BA} = \nabla_B \nabla^D C_{DA} - \nabla_A \nabla^D C_{DB}$. Then*

$$\int_{S^2} Y^A \left(\frac{1}{4}C_{AB}\nabla_D F^{DB} + \frac{1}{4}F_{AB}\nabla_D C^{DB} - \frac{3}{4}P_{BA}\nabla^B f - \frac{1}{4}\nabla^B P_{BA}f \right) = 0.$$

Proof. We integrate by parts the last two terms to get

$$\begin{aligned} & \int_{S^2} -\frac{1}{2}Y^A(\nabla_B \nabla^D C_{AD} - \nabla_A \nabla^D C_{BD})\nabla^B f + \frac{1}{2}\nabla^B Y^A \nabla_B \nabla^D C_{AD} \cdot f. \\ & = \int_{S^2} \frac{1}{2}\nabla_B Y^A \nabla^D C_{AD} \nabla^B f + \frac{1}{2}Y^A \nabla^D C_{AD} \Delta f - \frac{1}{2}Y^A \nabla^D C_D^B \nabla_A \nabla_B f \\ & \quad + \int_{S^2} \frac{1}{2}Y^A \nabla^D C_{AD} f - \frac{1}{2}\nabla^B Y^A \nabla^D C_{AD} \nabla_B f \\ & = \int_{S^2} -\frac{1}{2}Y^A \nabla^D C_D^B (\nabla_A \nabla_B f - \frac{1}{2}\Delta f \sigma_{AB}) + \frac{1}{4}Y^A \nabla^D C_{AD} (\Delta + 2)f \end{aligned}$$

\square

Theorem B.2. *Let $F_{AB} = 2\nabla_A\nabla_B f - \Delta f\sigma_{AB}$ and $P_{BA} = \nabla_B\nabla^D C_{DA} - \nabla_A\nabla^D C_{DB}$. Then*

$$\int_{S^2} \nabla^A \tilde{X}^k \left(C_{AB}\nabla_D F^{BD} + F_{AB}\nabla_D C^{BD} + \frac{1}{2}\nabla_A(C_{BD}F^{BD}) - f\nabla^B P_{BA} - 3P_{BA}\nabla^B f \right) = 0.$$

Proof. We integrate by parts the last two terms to get

$$\begin{aligned} & \int_{S^2} \nabla^A \tilde{X}^k (-2P_{BA})\nabla^B f \\ &= \int_{S^2} -2\nabla^A \tilde{X}^k (\nabla_B\nabla^D C_{DA} - \nabla_A\nabla^D C_{DB})\nabla^B f \\ &= \int_{S^2} -2\tilde{X}^k\nabla^D C_{DA}\nabla^A f + 2\nabla^A \tilde{X}^k\nabla^D C_{DA}\Delta f + 4\tilde{X}^k\nabla^D C_{DB}\nabla^B f - 2\nabla^A \tilde{X}^k\nabla^D C_{DB}\nabla_A\nabla^B f \\ &= \int_{S^2} 2\tilde{X}^k\nabla^D C_{DA}\nabla^A f - \nabla^A \tilde{X}^k\nabla^D C_{DB}F_{AB} + \nabla^A \tilde{X}^k\nabla^D C_{DA}\Delta f \\ &= \int_{S^2} -2\nabla^D \tilde{X}^k C_{DA}\nabla^A f - 2\tilde{X}^k C_{DA}\nabla^D\nabla^A f - \nabla^A \tilde{X}^k\nabla_D C^{DB}F_{AB} + \nabla^A \tilde{X}^k\nabla^D C_{DA}\Delta f \\ &= \int_{S^2} 2\nabla^A \tilde{X}^k\nabla^D C_{DA}\nabla^A f + \frac{1}{2}\Delta\tilde{X}^k C_{DA}F^{DA} - \nabla^A \tilde{X}^k\nabla_D C^{DB}F_{AB} + \nabla^A \tilde{X}^k\nabla^D C_{DA}\Delta f \\ &= \int_{S^2} -\nabla^A \tilde{X}^k C_{DA}\nabla^D(\Delta + 2)f - \frac{1}{2}\nabla^A \tilde{X}^k\nabla_A(C_{DB}F^{DB}) - \nabla^A \tilde{X}^k\nabla_D C^{DB}F_{AB}. \end{aligned}$$

□

REFERENCES

- [1] A. Ashtekar and R. O. Hansen, *A unified treatment of null and spatial infinity in general relativity. I. Universal structure, asymptotic symmetries, and conserved quantities at spatial infinity*, J. Math. Phys. **19** (1978), no. 7, 1542–1566.
- [2] H. Bondi, M. G. J. van der Burg and A. W. K. Metzner, *Gravitational waves in general relativity. VII. Waves from axi-symmetric isolated systems*, Proc. Roy. Soc. London Ser. A **269** (1962), 21–52.
- [3] P.-N. Chen, M.-T. Wang, and S.-T. Yau, *Evaluating quasi-local energy and solving optimal embedding equation at null infinity*, Comm. Math. Phys. **308** (2011), no.3, 845–863.
- [4] P.-N. Chen, M.-T. Wang, and S.-T. Yau, *Conserved quantities in general relativity: from the quasi-local level to spatial infinity*, Comm. Math. Phys. **338** (2015), no.1, 31–80.
- [5] P.-N. Chen, M.-T. Wang and S.-T. Yau, *Quasilocal angular momentum and center of mass in general relativity*, Adv. Theor. Math. Phys. **20** (2016), no. 4, 671–682.
- [6] P.-N. Chen, M.-T. Wang and S.-T. Yau, *Evaluating small sphere limit of the Wang-Yau quasi-local energy*, Comm. Math. Phys. **357** (2018), no. 2, 731–774.
- [7] D. Christodoulou, *Nonlinear nature of gravitation and gravitational-wave experiments*, Phys. Rev. Lett. **67** (1991), no. 12, 1486–1489.
- [8] D. Christodoulou, *The global initial value problem in general relativity*, Proceedings of the 9th Marcel Grossmann Meeting on General Relativity, MG9, World Scientific (2002) 44–54.
- [9] D. Christodoulou and S. Klainerman, *The global nonlinear stability of the Minkowski space*, Princeton Mathematical Series, 41. Princeton University Press, Princeton, NJ, 1993.

- [10] P. T. Chruściel, J. Jezierski and J. Kijowski, *Hamiltonian field theory in the radiating regime*, Lecture Notes in Physics. Monographs, 70, Springer-Verlag, Berlin, 2002.
- [11] P. T. Chruściel, M. A. H. MacCallum and D. B. Singleton, *Gravitational waves in general relativity. XIV. Bondi expansions and the “polyhomogeneity” of \mathcal{I}* , Philos. Trans. Roy. Soc. London Ser. A **350** (1995), no. 1692, 113–141
- [12] T. Dray and M. Streubel, *Angular momentum at null infinity*, *Classical Quantum Gravity* **1** (1984), no. 1, 15–26.
- [13] È. È. Flanagan and D. A. Nichols, *Conserved charges of the extended Bondi-Metzner-Sachs algebra*, Phys. Rev. D **95**, 044002 (2017).
- [14] S. W. Hawking, M. J. Perry, and A. Strominger, *Superrotation charge and supertranslation hair on black holes.*, Journal of High Energy Physics, 2017(5):161, 2017.
- [15] J. Keller, Y.-K. Wang, and S.-T. Yau, *Evaluating quasi-local angular momentum and center-of-mass at null infinity*, arXiv: 1811.02383
- [16] S. Klainerman and J. Szeftel, *Effective results on uniformization and intrinsic GCM spheres in perturbations of Kerr*, arXiv:1912.12195
- [17] T. Mädler and J. Winicour, *Bondi-Sachs formalism*, Scholarpedia, 11 (12): 33528, 2016. arXiv:1609.01731
- [18] R. Penrose, *Quasilocal mass and angular momentum in general relativity*, Proc. Roy. Soc. London Ser. A **381** (1982), no. 1780, 53–63. MR0661716
- [19] R. Penrose, *Some unsolved problems in classical general relativity*, in *Seminar on Differential Geometry*, 631–668, Ann. of Math. Stud., 102, Princeton Univ. Press, Princeton, NJ.
- [20] A. Rizzi, *Angular momentum in general relativity: a new definition*, Phys. Rev. Lett. **81** (1998), no. 6, 1150–1153.
- [21] A. Rizzi, *Angular momentum in general relativity*, PhD thesis, Princeton University, June 1997.
- [22] R. K. Sachs, *Gravitational waves in general relativity. VIII. Waves in asymptotically flat space-time*, Proc. Roy. Soc. London Ser. A **270** (1962), 103–126.
- [23] J. A. Valiente Kroon, *A comment on the outgoing radiation condition for the gravitational field and the peeling theorem*, Gen. Relativity Gravitation **31** (1999), no. 8, 1219–1224. MR1702757
- [24] M.-T. Wang, *Total mass and limits of quasilocal mass at future null infinity*, arXiv:2003.07732
- [25] M.-T. Wang, and S.-T. Yau, *Quasi-local mass in general relativity*, Phys. Rev. Lett. **102** (2009), no. 2, no. 021101.
- [26] M.-T. Wang, and S.-T. Yau, *Isometric embeddings into the Minkowski space and new quasi-local mass*, Comm. Math. Phys. **288** (2009), no. 3, 919–942.