

TRUNCATED AFFINE SPRINGER FIBERS AND ARTHUR'S WEIGHTED ORBITAL INTEGRALS

ZONGBIN CHEN 

*Yau Mathematical Science Center, Tsinghua University, Jinchunyuan West Building,
Office 255, 100084, Haidian District, Beijing, China*
(zbchen@tsinghua.edu.cn)

(Received 6 June 2017; revised 18 September 2021; accepted 22 September 2021)

Abstract We explain an algorithm to calculate Arthur's weighted orbital integral in terms of the number of rational points on the fundamental domain of the associated affine Springer fiber. The strategy is to count the number of rational points of the truncated affine Springer fibers in two ways: by the Arthur–Kottwitz reduction and by the Harder–Narasimhan reduction. A comparison of results obtained from these two approaches gives recurrence relations between the number of rational points on the fundamental domains of the affine Springer fibers and Arthur's weighted orbital integrals. As an example, we calculate Arthur's weighted orbital integrals for the groups GL_2 and GL_3 .

Keywords: Weighted orbital integrals, Affine Springer fibers, Harder–Narasimhan reduction, Arthur–Kottwitz reduction.

2020 *Mathematics subject classification*: Primary 22E67; 14M15
Secondary 11F85

1. Introduction

Let \mathbf{F}_q be the finite field with q elements. Let $F = \mathbf{F}_q((\epsilon))$ be the field of Laurent series with coefficients in \mathbf{F}_q , $\mathcal{O} = \mathbf{F}_q[[\epsilon]]$ the ring of integers of F , and $\mathfrak{p} = \epsilon\mathbf{F}_q[[\epsilon]]$ the maximal ideal of \mathcal{O} . We fix an algebraic closure $\overline{\mathbf{F}}_q$ of \mathbf{F}_q and also a compatible separable algebraic closure \overline{F} of F . Let $\mathrm{val} : \overline{F}^\times \rightarrow \mathbf{Q}$ be the discrete valuation normalized by $\mathrm{val}(\epsilon) = 1$.

Let G be a connected split reductive algebraic group over \mathbf{F}_q , and assume that $\mathrm{char}(\mathbf{F}_q) > |W|$, W being the Weyl group of G . Let G_F be the base change of G to F . Let T be a maximal torus of G_F . We make the assumption that the splitting field of T is totally ramified over F . Let $S \subset T$ be the maximal F -split subtorus of T , and let $M_0 = Z_{G_F}(S)$ be the centralizer of S in G_F ; then M_0 is a Levi subgroup of G_F and T is elliptic in M_0 . Given an algebraic group, we use the Gothic letter to denote its Lie algebra.

Let $\gamma \in \mathfrak{t}(F)$ be a regular element, elliptic in $\mathfrak{m}_0(F)$. Let $\mathcal{L}(M_0)$ be the set of Levi subgroups of G containing M_0 . For $M \in \mathcal{L}(M_0)$, consider Arthur's weighted orbital integral

$$J_M(\gamma) = J_M(\gamma, \mathbb{1}_{\mathfrak{g}(\mathcal{O})}) = \int_{T(F) \backslash G(F)} \mathbb{1}_{\mathfrak{g}(\mathcal{O})}(\mathrm{Ad}(g)^{-1}\gamma) \nu_M(g) \frac{dg}{dt}, \quad (1.1)$$

where $\mathbb{1}_{\mathfrak{g}(\mathcal{O})}$ is the characteristic function of the lattice $\mathfrak{g}(\mathcal{O})$ in $\mathfrak{g}(F)$, $v_M(g)$ is Arthur's weight factor, and dg and dt are Haar measures on $G(F)$ and $T(F)$, respectively. One of our main results states that it can be expressed in terms of the number of rational points of the *fundamental domains* F_γ^L of the affine Springer fiber \mathcal{X}_γ^L , $L \in \mathcal{L}(M_0)$. The main idea is to count the number of rational points of the truncated affine Springer fibers in two different ways: by the *Arthur–Kottwitz reduction* and by the *Harder–Narasimhan reduction*.

Before entering into the details of our approach, we give examples of results that can be obtained in this way. The calculations for the group $G = \mathrm{GL}_2$ are easy; the results are summarized in Theorems 5.1 and 5.2. But for the group $G = \mathrm{GL}_3$, the calculations are already quite nontrivial. There are three cases to deal with: the element γ can be split, mixed, or elliptic. When γ is split, we can find a set of simple roots $\{\alpha_1, \alpha_2\}$ in the root system $\Phi(G, T)$ of G with respect to T such that

$$\mathrm{val}(\alpha_1(\gamma)) = \mathrm{val}((\alpha_1 + \alpha_2)(\gamma)) \leq \mathrm{val}(\alpha_2(\gamma)).$$

We call $(n_1, n_2) = (\mathrm{val}(\alpha_1(\gamma)), \mathrm{val}(\alpha_2(\gamma)))$ the *root valuation* of γ .

Theorem 1.1. *Let $G = \mathrm{GL}_3$ and T the maximal torus of diagonal matrices. Let $\gamma \in \mathfrak{t}(\mathcal{O})$ be a regular element with root valuation $(n_1, n_2) \in \mathbf{N}^2$, with $n_1 \leq n_2$. Up to an explicit volume factor, we have*

$$J_T(\gamma) \doteq \sum_{i=1}^{n_1} i(q^{2i-1} + q^{2i-2}) + \sum_{i=n_1+n_2}^{2n_1+n_2-1} (4n_1 + 2n_2 - 4i - 3)q^i + (n_1^2 + 2n_1n_2)q^{2n_1+n_2}.$$

For $\alpha \in \Phi(G, T)$, let M_α be the unique Levi subgroup containing T with root system $\{\pm\alpha\}$; then, up to an explicit volume factor,

$$J_{M_{\alpha_1}}(\gamma) = J_{M_{\alpha_1+\alpha_2}}(\gamma) \doteq (n_1 + n_2)q^{2n_1+n_2} - q^{n_1+n_2}(1 + q + \cdots + q^{n_1-1}) - q^{2n_1}(1 + q + \cdots + q^{n_2-1})$$

and

$$J_{M_{\alpha_2}}(\gamma) \doteq 2n_1q^{2n_1+n_2} - 2q^{n_1+n_2}(1 + q + \cdots + q^{n_1-1}).$$

When γ is mixed – that is, T is isomorphic to $F^\times \times \mathrm{Res}_{E_2/F} E_2^\times$, where $E_2 = \mathbf{F}_q((\epsilon^{\frac{1}{2}}))$ is the unique totally ramified extension of F of degree 2 – it can be conjugate to a matrix of the form

$$\gamma = \begin{bmatrix} a & & \\ & b & \\ & b\epsilon & \end{bmatrix}. \quad (1.2)$$

Let $m = \mathrm{val}(a)$ and $n = \mathrm{val}(b)$; then we have the following:

Theorem 1.2. Let $G = GL_3$ and let γ be a matrix in the form of equation (1.2). When $\text{val}(a) = m \leq n$, up to an explicit volume factor,

$$J_{M_0}(\gamma) \doteq 2mq^{2m+n} + \sum_{j=m+n+1}^{2m+n-1} 2(j-m-n)q^j - \sum_{j=0}^{2m-1} \left(\left\lfloor \frac{j}{2} \right\rfloor + 1 \right) q^j.$$

Similarly, when $\text{val}(a) = m > n$, up to an explicit volume factor,

$$J_{M_0}(\gamma) \doteq (2n+1)q^{3n+1} + \sum_{j=2n+1}^{3n} (2j-4n-1)q^j - \sum_{j=0}^{2n} \left(\left\lfloor \frac{j}{2} \right\rfloor + 1 \right) q^j,$$

where $\lfloor x \rfloor$ denotes the maximal integer less than or equal to x .

When γ is anisotropic, Arthur's weighted orbital integral is just the orbital integral, and the result was essentially obtained by Goresky, Kottwitz, and MacPherson [16]. See Theorems 8.1 and 8.2 for the counting result.

Now we explain our approach to calculating Arthur's weighted orbital integrals using the geometry of the affine Springer fibers. For simplicity, we restrict to $J_{M_0}(\gamma)$. The affine Springer fiber \mathcal{X}_γ is the closed subscheme of the affine Grassmannian $\mathcal{X} = G(F)/G(\mathcal{O})$ defined by the equation

$$\mathcal{X}_\gamma = \{g \in G(F)/G(\mathcal{O}) \mid \text{Ad}(g^{-1})\gamma \in \mathfrak{g}(\mathcal{O})\}.$$

They can be used to geometrize Arthur's weighted orbital integrals. The group $T(F)$ acts on \mathcal{X}_γ by left translation. For $\mu \in X_*(S)$, we write ϵ^μ for $\mu(\epsilon) \in S(F)$. The map $\mu \rightarrow \epsilon^\mu$ identifies $X_*(S)$ with a subgroup of $S(F) \subset T(F)$, which we denote by Λ . It acts freely on \mathcal{X}_γ , and the quotient $\Lambda \backslash \mathcal{X}_\gamma$ is a projective scheme of finite type over \mathbf{F}_q (see [17, §3]). A simple reformulation shows that

$$\int_{T(F) \backslash G(F)} \mathbb{1}_{\mathfrak{g}(\mathcal{O})}(\text{Ad}(g)^{-1}\gamma) v_{M_0}(g) \frac{dg}{dt} = c \cdot \sum_{[g] \in \Lambda \backslash \mathcal{X}_\gamma(\mathbf{F}_q)} v_{M_0}(g),$$

where $[g]$ denotes the point $gG(\mathcal{O}) \in \mathcal{X}$ and c is a volume factor.

But this expression does not facilitate the calculations of Arthur's weighted orbital integral. We have to proceed in an indirect way. Let $\xi \in \mathfrak{a}_{M_0}^G$ be a generic element (for the definition of $\mathfrak{a}_{M_0}^G$, see §1.1). Chaudouard and Laumon [8] introduce a variant of the weighted orbital integral

$$J_{M_0}^\xi(\gamma) = J_{M_0}^\xi(\gamma, \mathbb{1}_{\mathfrak{g}(\mathcal{O})}) = \int_{T(F) \backslash G(F)} \mathbb{1}_{\mathfrak{g}(\mathcal{O})}(\text{Ad}(g)^{-1}\gamma) w_{M_0}^\xi(g) \frac{dg}{dt}, \quad (1.3)$$

with a slightly different weight factor $w_{M_0}^\xi(g)$. The two weight factors are closely related to each other. When G is semisimple, Chaudouard and Laumon show that

$$J_{M_0}(\gamma) = \text{vol}(\mathfrak{a}_{M_0}/X_*(M_0)) \cdot J_{M_0}^\xi(\gamma).$$

The variant $J_{M_0}^\xi(\gamma)$ has a better geometric interpretation. In fact, we can introduce a notion of ξ -stability on the affine Springer fiber \mathcal{X}_γ and show that

$$J_{M_0}^\xi(\gamma) = \text{vol}_{dt} (T(F)^1)^{-1} \cdot |\mathcal{X}_\gamma^\xi(\mathbf{F}_q)|.$$

The advantage of this variant is clear: it is a plain count rather than a weighted count. Moreover, we can use the Harder–Narasimhan reduction to get $|\mathcal{X}_\gamma^\xi(\mathbf{F}_q)|$ recursively from $|\mathcal{X}_\gamma(\mathbf{F}_q)|$, if only the latter is finite. Unfortunately this is not the case, as can be seen from the fact that the free abelian group Λ acts freely on \mathcal{X}_γ .

Let Π be a positive (G, M_0) -orthogonal family. We can introduce a truncation $\mathcal{X}_\gamma(\Pi)$ to overcome the finiteness issue. When Π is sufficiently regular, we can reduce the calculation of the rational points on $\mathcal{X}_\gamma(\Pi)$ to that of the fundamental domains F_γ^L , by the *Arthur–Kottwitz reduction*. Recall that the fundamental domain F_γ is introduced in [11] to play the role of an irreducible component of \mathcal{X}_γ . (All the irreducible components of \mathcal{X}_γ are isomorphic, because $T(F)$ acts transitively on a dense open subscheme of it.) The Arthur–Kottwitz reduction is a construction that decomposes $\mathcal{X}_\gamma(\Pi)$ into locally closed subschemes, which are iterated affine fibrations over the fundamental domains F_γ^L , $L \in \mathcal{L}(M_0)$. The counting result is summarized in Corollary 3.7. In particular, it shows that $\mathcal{X}_\gamma(\Pi)$ depends quasi-polynomially on the truncation parameter.

On the other hand, the Harder–Narasimhan reduction does not behave well on $\mathcal{X}_\gamma(\Pi)$. In fact, near the boundary, the Harder–Narasimhan strata are generally not affine fibrations over truncations of $\mathcal{X}_\gamma^{L, \xi^L}$. To overcome this difficulty, we cut $\mathcal{X}_\gamma(\Pi)$ into two parts: the *tail* and the *main body*. Roughly speaking, the tail is the union of the ‘boundary irreducible components’ of $\mathcal{X}_\gamma(\Pi)$, and the main body is its complement. The Harder–Narasimhan reduction works well on the main body, and we can use it to count the number of rational points. The result is summarized in Theorem 4.8; it can be expressed in terms of $|\mathcal{X}_\gamma^{L, \xi^L}(\mathbf{F}_q)|$, $L \in \mathcal{L}(M_0)$. Counting points on the tail proceeds by the Arthur–Kottwitz reduction, and can be expressed in terms of $|F_\gamma^L(\mathbf{F}_q)|$ s. But we are not able to obtain an explicit expression; we get a recursion.

These two different approaches to counting rational points on $\mathcal{X}_\gamma(\Pi)$ give us a recursive equation that involves the $|F_\gamma^L(\mathbf{F}_q)|$ s and the $|\mathcal{X}_\gamma^{L, \xi^L}(\mathbf{F}_q)|$ s. Solving it, we can express the latter in terms of the former. The problem of calculating $J_{M_0}(\gamma)$ is thus reduced to counting points on F_γ .

The geometry of F_γ is simpler than that of \mathcal{X}_γ^ξ : Goresky, Kottwitz, and MacPherson [14] have conjectured that the cohomology of \mathcal{X}_γ is pure in the sense of Deligne. As we have shown in [11], this is equivalent to the cohomological purity of F_γ . In fact, it is even expected that F_γ admits a *Hessenberg paving*. (This notion was introduced by Goresky, Kottwitz, and MacPherson [16].) On the contrary, \mathcal{X}_γ^ξ is generally not cohomologically pure, as one can see in case $G = \text{SL}_2$ or from the appearance of a minus sign in the counting-points result of Theorems 6.6, 6.10, and 6.14. Although one can still look at the quotient $\mathcal{X}_\gamma^\xi/A_{M_0}$, where A_{M_0} is the maximal F -split torus of the center of M_0 , it is clear that the quotient no longer admits a torus action, and hence it has much less structure to explore than F_γ .

When the torus T splits, we make a conjecture on the Poincaré polynomial of F_γ [12], assuming the cohomological purity of F_γ . This gives a conjectural expression for $|F_\gamma(\mathbf{F}_q)|$. We reproduce it here for the convenience of the reader. Following Chaudouard and Laumon [7], under the purity assumption the cohomology of F_γ can be expressed in terms of its 1-skeleton under the T -action. Indeed, the T -equivariant cohomology $H_T^*(F_\gamma, \overline{\mathbf{Q}}_\ell)^1$ will then be a free $H_T^*(\text{pt}, \overline{\mathbf{Q}}_\ell)$ -algebra, and we have

$$H^*(F_\gamma, \overline{\mathbf{Q}}_\ell) = H_T^*(F_\gamma, \overline{\mathbf{Q}}_\ell) \otimes_{H_T^*(\text{pt}, \overline{\mathbf{Q}}_\ell)} \overline{\mathbf{Q}}_\ell. \quad (1.4)$$

The torus T acts on F_γ with finitely many fixed points, but the 1-dimensional T -orbits form a higher-dimensional variety which we denote by $F_\gamma^{T,1}$. The bigger torus $\tilde{T} = T \times \mathbb{G}_m$, where \mathbb{G}_m is the rotational torus, acts on $F_\gamma^{T,1}$ with finitely many 1-dimensional \tilde{T} -orbits; let $F_\gamma^{\tilde{T},1}$ be their union. Let $F_\gamma^{\tilde{T}}$ be the set of \tilde{T} -fixed points on F_γ , and let

$$H_{\tilde{T}}^*(F_\gamma, \overline{\mathbf{Q}}_\ell) := H_T^*(F_\gamma, \overline{\mathbf{Q}}_\ell) \otimes_{H_T^*(\text{pt}, \overline{\mathbf{Q}}_\ell)} H_{\tilde{T}}^*(\text{pt}, \overline{\mathbf{Q}}_\ell). \quad (1.5)$$

Then the localization theorem of Goresky, Kottwitz, and MacPherson [13] implies an exact sequence of equivariant cohomology

$$0 \rightarrow H_{\tilde{T}}^*(F_\gamma, \overline{\mathbf{Q}}_\ell) \rightarrow H_{\tilde{T}}^*(F_\gamma^{\tilde{T}}, \overline{\mathbf{Q}}_\ell) \rightarrow H_{\tilde{T}}^*(F_\gamma^{\tilde{T},1}, F_\gamma^{\tilde{T}}; \overline{\mathbf{Q}}_\ell). \quad (1.6)$$

Let Γ be the graph with vertices $F_\gamma^{\tilde{T}}$ and edges $F_\gamma^{\tilde{T},1}$. Two vertices are linked by an edge if and only if they lie on the closure of the corresponding 1-dimensional \tilde{T} -orbit. We call it the *moment graph* of F_γ with respect to the action of \tilde{T} . The foregoing result implies that the information about the cohomology of F_γ is encoded in Γ . A direct calculation of the cohomology via formulas (1.4), (1.5), and (1.6) turns out to be very hard, and we look for a combinatorial way to get around it.

Let \mathbf{o} be a total order among the vertices of the graph Γ ; it will serve as the paving order. We associate to it an *acyclic* oriented graph (Γ, \mathbf{o}) such that the source of each arrow is greater than its target with respect to \mathbf{o} . For $v \in \Gamma$, denote by $n_v^\mathbf{o}$ the number of arrows having source v .

Definition 1.1. The formal Betti number $b_{2i}^\mathbf{o}$ associated to the order \mathbf{o} is defined as

$$b_{2i}^\mathbf{o} = \# \{v \in \Gamma : n_v^\mathbf{o} = i\}.$$

We call

$$P^\mathbf{o}(t) = \sum_i b_{2i}^\mathbf{o} t^{2i}$$

the *formal Poincaré polynomial* associated to the order \mathbf{o} .

Definition 1.2. For $P_1(t), P_2(t) \in \mathbf{Z}[t]$, we say that $P_1(t) < P_2(t)$ if the leading coefficient of $P_2(t) - P_1(t)$ is positive.

¹Here we actually mean the *geometric* $H_{\overline{\mathbf{F}}_q}^*(F_{\gamma, \overline{\mathbf{F}}_q}, \overline{\mathbf{Q}}_\ell)$; to simplify the notation, we do not specify the base change to $\overline{\mathbf{F}}_q$. A similar convention applies for the other cohomology groups.

Conjecture 1.1. *Let $P(t)$ be the Poincaré polynomial of F_γ . Then*

$$P(t) = \min_{\mathfrak{o}} \{P^{\mathfrak{o}}(t)\},$$

where \mathfrak{o} runs through all the total orders among the vertices of Γ .

The conjecture can be thought of as a kind of *Morse inequality*; it has been verified in a lot of examples. For the group $G = \mathrm{GL}_2$ and $\gamma = \mathrm{diag}(\gamma_1, \gamma_2)$, with $\gamma_1, \gamma_2 \in F$ and $\mathrm{val}(\gamma_1 - \gamma_2) = n \in \mathbf{N}$, the moment graph of F_γ contains $n+1$ vertices, which are pairwise connected by an edge. It is clear that the conjecture holds in this case. In general, the moment graph of F_γ is easy to describe, and we have an algorithm to find an order \mathfrak{o} which conjecturally should attain the minimum. Although we are not able to prove both conjectures at the moment, they have been very helpful in constructing affine pavings of F_γ in concrete examples.

Under the purity assumption, Conjecture 1.1 implies a point-counting result for F_γ . Indeed, the formulas (1.4), (1.5), and (1.6) are $\mathrm{Gal}(\overline{\mathbf{F}}_q/\mathbf{F}_q)$ -equivariant, and the Frobenius endomorphism acts on $H_{\overline{T}}^{2i}(F_\gamma^{\overline{T}}, \overline{\mathbf{Q}}_\ell)$ by q^i (the odd-degree cohomologies vanish), and hence it acts on $H^*(F_\gamma, \overline{\mathbf{Q}}_\ell)$ in the same way and so

$$|F_\gamma(\mathbf{F}_q)| = \min_{\mathfrak{o}} \left\{ P^{\mathfrak{o}}(q^{1/2}) \right\}.$$

Together with the recurrence relation between $|\mathcal{X}_\gamma^\xi(\mathbf{F}_q)|$ and $|F_\gamma(\mathbf{F}_q)|$, it gives a conjectural complete answer to the calculation of Arthur's weighted orbital integrals in the split case.

1.1. Notation

We fix a split maximal torus A of G over \mathbf{F}_q . Without loss of generality, we suppose that $A \subset M_0$. Let $\Phi = \Phi(G, A)$ be the root system of G with respect to A and let W be the Weyl group of G with respect to A . For any subgroup H of G which is stable under the conjugation of A , we write $\Phi(H, A)$ for the roots appearing in $\mathrm{Lie}(H)$. We fix a Borel subgroup B_0 of G containing A . Let $\Delta = \{\alpha_1, \dots, \alpha_r\}$ be the set of simple roots with respect to B_0 and let $\{\varpi_i\}_{i=1}^r$ be the corresponding fundamental weights. For an element $\alpha \in \Delta$, we have a unique maximal parabolic subgroup P_α of G containing B_0 such that $\Phi(N_{P_\alpha}, A) \cap \Delta = \{\alpha\}$, where N_{P_α} is the unipotent radical of P_α . This gives a bijective correspondence between the simple roots in Δ and the maximal parabolic subgroups of G containing B_0 . Any semistandard maximal parabolic subgroup P of G is conjugate to certain P_α by an element $w \in W$; when the element $w\varpi_\alpha$ does not depend on the choice of w , we denote it by ϖ_P .

We use the (G, M) notation of Arthur. Let $\mathcal{F}(A)$ be the set of parabolic subgroups of G containing A and let $\mathcal{L}(A)$ be the set of Levi subgroups of G containing A . For every $M \in \mathcal{L}(A)$, we denote by $\mathcal{P}(M)$ the set of parabolic subgroups of G whose Levi factor is M , by $\mathcal{L}(M)$ the set of Levi subgroups of G containing M , and by $\mathcal{F}(M)$ the set of parabolic subgroups of G containing M . For $P \in \mathcal{P}(M)$, we denote by $P^- \in \mathcal{P}(M)$ the opposite of P with respect to M .

Let $X^*(M) = \text{Hom}(M, \mathbb{G}_m)$ and $X_*(M) = \text{Hom}(X^*(M), \mathbb{Z})$. Let $\mathfrak{a}_M^* = X^*(M) \otimes \mathbb{R}$ and $\mathfrak{a}_M = X_*(M) \otimes \mathbb{R}$. The restriction $X^*(M) \rightarrow X^*(A)$ induces an injection $\mathfrak{a}_M^* \hookrightarrow \mathfrak{a}_A^*$. Let $(\mathfrak{a}_A^M)^*$ be the subspace of \mathfrak{a}_A^* generated by $\Phi(M, A)$. We have the decomposition in direct sums

$$\mathfrak{a}_A^* = (\mathfrak{a}_A^M)^* \oplus \mathfrak{a}_M^*.$$

The canonical pairing $X_*(A) \times X^*(A) \rightarrow \mathbb{Z}$ can be extended bilinearly to $\mathfrak{a}_A \times \mathfrak{a}_A^* \rightarrow \mathbb{R}$. For $M \in \mathcal{L}(A)$, we can embed \mathfrak{a}_M in \mathfrak{a}_A as the orthogonal subspace to $(\mathfrak{a}_A^M)^*$. Let $\mathfrak{a}_A^M \subset \mathfrak{a}_A$ be the subspace orthogonal to \mathfrak{a}_M^* . We have the dual decomposition

$$\mathfrak{a}_A = \mathfrak{a}_M \oplus \mathfrak{a}_A^M.$$

Let π_M, π^M be the projections to the two factors. More generally, for $L, M \in \mathcal{F}(A), M \subset L$, we also have a decomposition

$$\mathfrak{a}_M^G = \mathfrak{a}_L^G \oplus \mathfrak{a}_M^L.$$

Let $\pi_{M,L}, \pi_M^L$ be the projections to the two factors. If the context is clear, we also simplify them to π_L, π^L .

We identify $X_*(A)$ with $A(F)/A(\mathcal{O})$ by sending χ to $\chi(\epsilon)$. With this identification, the canonical surjection $A(F) \rightarrow A(F)/A(\mathcal{O})$ can be viewed as

$$A(F) \rightarrow X_*(A). \quad (1.7)$$

We use Λ_G to denote the quotient of $X_*(A)$ by the coroot lattice of G (the subgroup of $X_*(A)$ generated by the coroots of A in G). It is independent of the choice of A ; this is the algebraic fundamental group introduced by Borovoi [6]. According to Kottwitz [18], we have a canonical homomorphism

$$\nu_G : G(F) \rightarrow \Lambda_G, \quad (1.8)$$

which is characterized by the following properties: it is trivial on the image of $G_{\text{sc}}(F)$ in $G(F)$ (G_{sc} is the simply connected cover of the derived group of G), and its restriction to $A(F)$ coincides with the composition of formula (1.7) with the projection of $X_*(A)$ to Λ_G . Since the morphism (1.8) is trivial on $G(\mathcal{O})$, it descends to a map

$$\nu_G : \mathcal{X} \rightarrow \Lambda_G,$$

whose fibers are the connected components of \mathcal{X} . For $\mu \in \Lambda_G$, we denote the connected component $\nu_G^{-1}(\mu)$ by \mathcal{X}^μ .

Finally, we suppose that $\gamma \in \mathfrak{t}(\mathcal{O})$ satisfies $\gamma \equiv 0 \pmod{\epsilon}$, to avoid unnecessary complications.

2. (Weighted) orbital integrals and the affine Springer fibers

We recall briefly the geometrization of the (weighted) orbital integrals using the affine Springer fibers. We fix a regular element $\gamma \in \mathfrak{t}(\mathcal{O})$ as in the introduction. Let $P_0 = M_0 N_0$ be the unique element in $\mathcal{P}(M_0)$ which contains B_0 .

2.1. Orbital integrals

We begin by fixing the Haar measures. Let dg be the Haar measure on $G(F)$ normalized by the condition $\text{vol}_{dg}(G(\mathcal{O})) = 1$. For the group $T(F)$, the definition is more involved, as there is no natural \mathcal{O} -structure on T . Let $F^{\text{ur}} = \overline{\mathbf{F}}_q((\epsilon))$; it is the completion of the maximal unramified extension of F . Let σ be the Frobenius automorphism of both $\overline{\mathbf{F}}_q/\mathbf{F}_q$ and F^{ur}/F . We fix a separable algebraic closure $\overline{F^{\text{ur}}}$ of F^{ur} and let $I_F = \text{Gal}(\overline{F^{\text{ur}}}/F^{\text{ur}})$ – this is the inertia subgroup of $\Gamma = \text{Gal}(\overline{F}/F)$. According to Kottwitz [19, §7.6], we have an exact sequence

$$1 \rightarrow T(F^{\text{ur}})_1 \rightarrow T(F^{\text{ur}}) \xrightarrow{w_T} X_*(T)_{I_F} \rightarrow 1, \quad (2.1)$$

which implies another exact sequence if we take the $\langle \sigma \rangle$ -invariants:

$$1 \rightarrow T(F)_1 \rightarrow T(F) \xrightarrow{w_T} (X_*(T)_{I_F})^{(\sigma)} \rightarrow 1, \quad (2.2)$$

with $T(F)_1 := T(F) \cap T(F^{\text{ur}})_1$. We fix the Haar measure dt on $T(F)$ by setting $\text{vol}_{dt}(T(F)_1) = 1$. The group Λ is discrete and cocompact in $T(F)$. The volume of the quotient $\Lambda \backslash T(F)$ is calculated in [14, §15.3]:

$$\text{vol}_{dt}(\Lambda \backslash T(F)) = \frac{|\text{coker}[X_*(S)_{\Gamma} \rightarrow X_*(T)_{\Gamma}]|}{|\text{ker}[X_*(S)_{\Gamma} \rightarrow X_*(T)_{\Gamma}]|}.$$

Consider the orbital integral

$$I_{\gamma}^G = \int_{T(F) \backslash G(F)} \mathbb{1}_{\mathfrak{g}(\mathcal{O})}(\text{Ad}(g^{-1})\gamma) \frac{dg}{dt}. \quad (2.3)$$

It can be interpreted as counting points on the affine Springer fiber:

Proposition 2.1 (Goresky, Kottwitz, MacPherson [14]).

$$I_{\gamma}^G = \frac{|\text{ker}[X_*(S)_{\Gamma} \rightarrow X_*(T)_{\Gamma}]|}{|\text{coker}[X_*(S)_{\Gamma} \rightarrow X_*(T)_{\Gamma}]|} \cdot |\Lambda \backslash (\mathcal{X}_{\gamma}(\mathbf{F}_q))|.$$

The $T(F)$ -action on \mathcal{X}_{γ} can be exploited to further simplify the computations. Let $\mathcal{X}_{\gamma}^{\text{reg}}$ be the open subscheme of \mathcal{X}_{γ} consisting of the points $[g] \in \mathcal{X}_{\gamma}$ such that the image of $\text{Ad}(g^{-1})\gamma$ under the reduction $\mathfrak{g}(\mathcal{O}) \rightarrow \mathfrak{g}$ is regular nilpotent.

Proposition 2.2 (Bezrukavnikov [5]). *The group $T(F)$ acts transitively on $\mathcal{X}_{\gamma}^{\text{reg}}$.*

Proposition 2.3 (Ngô [20, Proposition 3.10.1]). *The open subscheme $\mathcal{X}_{\gamma}^{\text{reg}}$ is dense in \mathcal{X}_{γ} .*

Consequently, all the irreducible components of \mathcal{X}_{γ} are isomorphic to each other, and they are parametrized by $\pi_0(T(F))$. In particular, all the connected components of \mathcal{X}_{γ} are isomorphic and can be translated to each other under the $T(F)$ -action. In calculating the orbital integral (2.3), we can thus restrict to the central connected component of \mathcal{X}_{γ} , which often simplifies calculations.

The calculation of I_{γ}^G can be reduced to that of $I_{\gamma}^{M_0}$; it dates back at least to Harish-Chandra that

$$I_{\gamma}^G = q^{\frac{1}{2}\text{val}(\det(\text{ad}(\gamma)|_{\mathfrak{g}_F/\mathfrak{m}_{0,F}}))} \cdot I_{\gamma}^{M_0}. \quad (2.4)$$

Geometrically, this is a reflection of the existence of an affine fibration $f_P : \mathcal{X}_\gamma \rightarrow \mathcal{X}_\gamma^{M_0}$ for each $P \in \mathcal{P}(M_0)$. Recall that for $Q = LN_Q \in \mathcal{F}(A)$, we have the retraction

$$f_Q : \mathcal{X} \rightarrow \mathcal{X}^L,$$

which sends $[g] = gK$ to $[h] := hL(\mathcal{O})$, where $g = nhk, n \in N_Q(F), h \in L(F), k \in K$, is the Iwasawa decomposition. We want to point out that the retraction f_Q is not a morphism between $\text{ind-}\mathbf{F}_q$ -schemes, but its restriction to the inverse image of each connected component of \mathcal{X}^L ,

$$f_Q : f_Q^{-1}(\mathcal{X}^{L,\nu}) \rightarrow \mathcal{X}^{L,\nu}, \quad \nu \in \Lambda_L,$$

is actually a morphism over \mathbf{F}_q between $\text{ind-}\mathbf{F}_q$ -schemes. Moreover, these retractions satisfy the obvious transitivity property.

Restricted to the affine Springer fibers, the retraction f_Q sends \mathcal{X}_γ to \mathcal{X}_γ^L . To see this, for $[g] \in \mathcal{X}_\gamma$, let $g = nhk$ be the Iwasawa decomposition as before. We can write $g = hn'k$, with $n' = h^{-1}nh \in N_Q(F)$. Now that $\text{Ad}(h^{-1})\gamma \in \mathfrak{l}(F)$, we have

$$\text{Ad}(n'^{-1})\text{Ad}(h^{-1})\gamma = \text{Ad}(h^{-1})\gamma + n''$$

for some $n'' \in \mathfrak{n}_Q(F)$. This implies that

$$\text{Ad}(h)^{-1}\gamma \in [\mathfrak{g}(\mathcal{O}) + \mathfrak{n}_{Q,F}] \cap \mathfrak{l}(F) = \mathfrak{l}(\mathcal{O}),$$

which means that $f_Q([g]) = [h] \in \mathcal{X}_\gamma^L$.

Proposition 2.4 (Kazhdan and Lusztig [17, §5, Proposition 1]). *For any $\nu \in \Lambda_L$, the retraction*

$$f_Q : \mathcal{X}_\gamma \cap f_Q^{-1}(\mathcal{X}_\gamma^{L,\nu}) \rightarrow \mathcal{X}_\gamma^{L,\nu}$$

is an iterated affine fibration over \mathbf{F}_q of relative dimension $\text{val}(\det(\text{ad}(\gamma) | \mathfrak{n}_Q(F)))$.

The reader can also consult [11, Proposition 3.2] for a proof.

2.2. Arthur's weighted orbital integral

2.2.1. The weight factor v_M . Set $M \in \mathcal{L}(M_0)$. Roughly speaking, the weight factor $v_M(g)$ is the volume of a polytope in \mathfrak{a}_M generated by the point $[g] \in \mathcal{X}$. Let $H_M : M(F) \rightarrow \mathfrak{a}_M$ be the unique map² satisfying

$$\chi(H_M(m)) = \text{val}(\chi(m)), \quad \forall \chi \in X^*(M), m \in M(F).$$

Notice that it is a group homomorphism. Moreover, it is invariant under the right K -action, so it induces a map from \mathcal{X}^M to \mathfrak{a}_M , still denoted by H_M . For $P = MN \in \mathcal{F}(A)$, let $H_P : \mathcal{X} \rightarrow \mathfrak{a}_M$ be the composition

$$H_P : \mathcal{X} \xrightarrow{f_P} \mathcal{X}^M \xrightarrow{H_M} \mathfrak{a}_M.$$

²Our definition differs from the conventional one by a minus sign. But as we will see, it simplifies computations.

As shown in [7, Lemma 6.1], the map H_M is constant on each connected component of \mathcal{X}^M , so it has a factorization $H_M : \mathcal{X}^M \xrightarrow{\nu_M} \Lambda_M \rightarrow \mathfrak{a}_M$. A simple calculation of the restriction of the map to $\mathcal{X}^A \subset \mathcal{X}^M$ shows that the map $\Lambda_M \rightarrow \mathfrak{a}_M$ is just the one induced from the natural inclusion $X_*(A) \hookrightarrow \mathfrak{a}_A = X_*(A) \otimes \mathbf{R}$. Hence H_P is also the composition

$$H_P : \mathcal{X} \xrightarrow{f_P} \mathcal{X}^M \xrightarrow{\nu_M} \Lambda_M \rightarrow \mathfrak{a}_M.$$

The map H_P has the following remarkable property. There is a notion of adjacency among the parabolic subgroups in $\mathcal{P}(M)$: two parabolic subgroups $P_1 = MN_1, P_2 = MN_2 \in \mathcal{P}(M)$ are said to be *adjacent* if both of them are contained in a parabolic subgroup $Q = LN_Q$ such that $L \supset M$ and $\text{rk}(L) = \text{rk}(M) + 1$. Given such an adjacent pair, we define an element $\beta_{P_1, P_2} \in \Lambda_M$ in the following way: consider the collection of elements in Λ_M obtained from coroots of A in $\mathfrak{n}_1 \cap \mathfrak{n}_2^-$. We define β_{P_1, P_2} to be the minimal element in this collection – that is, all the other elements are positive integral multiples of it. Note that $\beta_{P_2, P_1} = -\beta_{P_1, P_2}$, and if $M = A$, then β_{P_1, P_2} is the unique coroot which is positive for P_1 and negative for P_2 . We also denote its image in \mathfrak{a}_M by β_{P_1, P_2} if no confusion is caused.

Proposition 2.5 (Arthur [1, Lemma 3.6]). *Let $P_1, P_2 \in \mathcal{P}(M)$ be two adjacent parabolic subgroups. For any $x \in \mathcal{X}$, we have*

$$H_{P_1}(x) - H_{P_2}(x) = n(x, P_1, P_2) \cdot \beta_{P_1, P_2},$$

with $n(x, P_1, P_2) \in \mathbf{Z}_{\geq 0}$.

The reader can consult [11, Proposition 2.1] for a proof. For any point $x \in \mathcal{X}$, we write $\text{Ec}_M(x)$ for the convex hull in \mathfrak{a}_M of the $H_P(x), P \in \mathcal{P}(M)$. For any $Q \in \mathcal{F}(M)$, we denote by $\text{Ec}_M^Q(x)$ the face of $\text{Ec}_M(x)$ whose vertices are $H_P(x), P \in \mathcal{P}(M), P \subset Q$. When $M = A$, we omit the subscript A to simplify the notation.

To define the volume, we need to choose a Lebesgue measure on \mathfrak{a}_M^G . We fix a W -invariant inner product $\langle \cdot, \cdot \rangle$ on the vector space \mathfrak{a}_A^G . Notice that \mathfrak{a}_A^M and \mathfrak{a}_M are orthogonal to each other with respect to the inner product for any $M \in \mathcal{L}(A)$. We fix a Lebesgue measure on \mathfrak{a}_M^G normalized by the condition that the lattice generated by the orthonormal bases in \mathfrak{a}_M^G has covolume 1.

The weight factor $v_M(g)$ is the volume of the projection $\pi_M^G(\text{Ec}_M(g)) \subset \mathfrak{a}_M^G$. We have to pass to \mathfrak{a}_M^G because the polytope $\text{Ec}_M(g)$ will lie in a hyperplane of \mathfrak{a}_M if G has nontrivial connected center. The weight factor $v_M(g)$ has the following invariance properties: it is invariant under the right action of K – that is,

$$v_M(gk) = v_M(g), \quad \forall k \in K.$$

This is evident from the definition of $v_M(g)$. It is not so evident, but also true, that

$$v_M(mg) = v_M(g), \quad \forall m \in M(F).$$

Indeed, for any $P \in \mathcal{P}(M)$, we have $f_P(mg) = mf_P(g)$. As H_M is a group homomorphism, this implies

$$H_P(mg) = H_M(m) + H_P(g),$$

so $\text{Ec}_M(mg)$ is just the translation of $\text{Ec}_M(g)$ by $H_M(m)$. In particular, they have the same volume.

Similar to Proposition 2.1, we can interpret Arthur's weighted orbital integral as

$$\int_{T(F) \backslash G(F)} \mathbb{1}_{\mathfrak{g}(\mathcal{O})}(\text{Ad}(g)^{-1}\gamma) v_M(g) \frac{dg}{dt} = \frac{|\ker[X_*(S)_\Gamma \rightarrow X_*(T)_\Gamma]|}{|\text{coker}[X_*(S)_\Gamma \rightarrow X_*(T)_\Gamma]|} \sum_{[g] \in \Lambda \backslash (\mathcal{X}_\gamma(\mathbf{F}_q))} v_M(g).$$

That is, it is a weighted count of the rational points on the affine Springer fiber. Notice also that $J_G(\gamma) = I_\gamma^G$, as $v_G(g) = 1$ for all $g \in G(F)$.

2.2.2. A variant. In their work on the weighted fundamental lemma [8], Chaudouard and Laumon introduce a variant of the weighted orbital integral.

Assume that G is semisimple and let $\xi \in \mathfrak{a}_M$ be a generic element. For $g \in G(F)$, they introduce the weight factor

$$w_M^\xi(g) = |\{\lambda \in X_*(M) \mid \lambda + \xi \in \text{Ec}_M(g)\}|.$$

It is the number of integral points in the polytope $\text{Ec}_M(g) - \xi$. Similar to $v_M(g)$, the weight factor $w_M^\xi(g)$ is invariant under the right K -action and the left $M(F)$ -action. In particular, it descends to a function on \mathcal{X} . Consider the following weighted orbital integral:

$$J_M^\xi(\gamma) = \int_{T(F) \backslash G(F)} \mathbb{1}_{\mathfrak{g}(\mathcal{O})}(\text{Ad}(g)^{-1}\gamma) w_M^\xi(g) \frac{dg}{dt}.$$

Remark 2.1. For a general reductive algebraic group G , $\xi \in \mathfrak{a}_M^G$, as $G(F) = M(F) \cdot G_{\text{der}}(F)$, we can define the weight factor w_M^ξ uniquely by requiring it to be invariant under the left $M(F)$ -action and the right K -action, and that as a function on \mathcal{X} its restriction to $\mathcal{X}^{G_{\text{der}}}$ coincide with the given definition for G_{der} . In other words, for generic $\xi \in \mathfrak{a}_M^G$, we define

$$w_M^\xi(g) = |\{\lambda \in X_*(M_{G_{\text{der}}}) \mid \lambda + \xi \in \pi_M^G(\text{Ec}_M(g))\}|,$$

where $M_{G_{\text{der}}} = M \cap G_{\text{der}}$. Notice that the weight factor v_M satisfies these conditions as well, and this justifies our definition in the general case.

The variant $J_M^\xi(\gamma)$ has a better geometric interpretation.

Lemma 2.6. *Let $T(F)^1 = T(F) \cap \ker(H_{M_0})$. Then it is of finite volume and we have an exact sequence*

$$1 \rightarrow T(F)^1 \rightarrow T(F) \xrightarrow{H_{M_0}} X_*(M_0) \rightarrow 1.$$

Proof. The first assertion is due to the fact that T is anisotropic modulo the center of M_0 . For the second assertion, only the surjectivity is nontrivial. Recall that we have the exact sequence

$$1 \rightarrow T(F)_1 \rightarrow T(F) \xrightarrow{w_T} (X_*(T)_{I_F})^{(\sigma)} \rightarrow 1,$$

and that the map w_T is defined via a map $v_T : T(F^{\text{ur}}) \rightarrow \text{Hom}(X^*(T)^{I_F}, \mathbf{Z})$, similar to the definition of H_{M_0} . Hence the morphism H_{M_0} factors through w_T , and the surjectivity

results from those of w_T and the homomorphism

$$(X_*(T)_{I_F})^{(\sigma)} = X_*(T)_{I_F} \rightarrow \operatorname{Hom}(X^*(M_0), \mathbf{Z}) = X_*(M_0).$$

□

As a consequence, let $T(F)_M^1 = T(F) \cap \ker(H_M)$ and let $\Lambda^{H_M} = \Lambda \cap \ker(H_M)$. Then the quotient $\Lambda^{H_M} \backslash T(F)_M^1$ is of finite volume and we have an exact sequence

$$1 \rightarrow T(F)_M^1 \rightarrow T(F) \xrightarrow{H_M} X_*(M) \rightarrow 1.$$

Proposition 2.7. *We have the equality*

$$J_M^\xi(\gamma) = \operatorname{vol}_{dt} \left(\Lambda^{H_M} \backslash T(F)_M^1 \right)^{-1} \cdot \left| \Lambda^{H_M} \backslash \{[g] \in \mathcal{X}_\gamma(\mathbf{F}_q) \mid \xi \in \operatorname{Ec}_M(g)\} \right|.$$

In particular,

$$J_{M_0}^\xi(\gamma) = \operatorname{vol}_{dt} \left(T(F)^1 \right)^{-1} \cdot \left| \{[g] \in \mathcal{X}_\gamma(\mathbf{F}_q) \mid \xi \in \operatorname{Ec}_{M_0}(g)\} \right|.$$

Proof. Let $\mathbb{1}_{M,g}$ be the characteristic function of $\operatorname{Ec}_M(g)$. As

$$\operatorname{Ec}_M(tg) = \operatorname{Ec}_M(g) + H_M(t), \quad \forall t \in T(F), g \in G(F),$$

we have

$$\begin{aligned} \sum_{t \in T(F)_M^1 \backslash T(F)} \mathbb{1}_{M,tg}(\xi) &= |\{\lambda \in X_*(M) \mid \xi \in \operatorname{Ec}_M(g) + \lambda\}| \\ &= w_M^\xi(g). \end{aligned}$$

Now we can rewrite

$$\begin{aligned} J_M^\xi(\gamma) &= \int_{T(F) \backslash G(F)} \mathbb{1}_{\mathfrak{g}(\mathcal{O})}(\operatorname{Ad}(g)^{-1}\gamma) w_M^\xi(g) \frac{dg}{dt} \\ &= \int_{T(F) \backslash G(F)} \mathbb{1}_{\mathfrak{g}(\mathcal{O})}(\operatorname{Ad}(g)^{-1}\gamma) \sum_{t \in T(F)_M^1 \backslash T(F)} \mathbb{1}_{M,tg}(\xi) \frac{dg}{dt} \\ &= \int_{T(F)_M^1 \backslash G(F)} \mathbb{1}_{\mathfrak{g}(\mathcal{O})}(\operatorname{Ad}(g)^{-1}\gamma) \mathbb{1}_{M,g}(\xi) \frac{dg}{dt} \\ &= \operatorname{vol}_{dt} \left(\Lambda^{H_M} \backslash T(F)_M^1 \right)^{-1} \int_{\Lambda^{H_M} \backslash G(F)} \mathbb{1}_{\mathfrak{g}(\mathcal{O})}(\operatorname{Ad}(g)^{-1}\gamma) \mathbb{1}_{M,g}(\xi) dg \\ &= \operatorname{vol}_{dt} \left(\Lambda^{H_M} \backslash T(F)_M^1 \right)^{-1} \cdot \left| \Lambda^{H_M} \backslash \{[g] \in \mathcal{X}_\gamma(\mathbf{F}_q) \mid \xi \in \operatorname{Ec}_M(g)\} \right|. \end{aligned}$$

□

In particular, $J_M^\xi(\gamma)$ is a plain count of a subset of $\mathcal{X}_\gamma(\mathbf{F}_q)$. In §4.1, we will see that the condition $\xi \in \operatorname{Ec}_M(g)$ behaves as a stability condition. (We believe that it is in fact a stability condition in the sense of Mumford.) In particular, there is a Harder–Narasimhan-type decomposition of \mathcal{X}_γ associated with it.

Remark 2.2. It is time to explain why we have imposed the assumption that T is totally ramified over F . Without it, the Frobenius $\sigma \in \operatorname{Gal}(\mathbf{F}_q/\mathbf{F}_q)$ acts nontrivially on

$X_*(T)_{I_F}$, and the morphism $T(F) \rightarrow X_*(M_0)$ in Lemma 2.6 might fail to be surjective. (Indeed, it does fail for T an unramified maximal torus in GL_n .) As a consequence, the interpretation of $J_M^\xi(\gamma)$ as in Proposition 2.7 no longer holds.

For completeness, we compute the volume factors in Proposition 2.7. We have the exact sequence

$$1 \rightarrow \Lambda^{H_M} \rightarrow T(F)_M^1 \rightarrow \Lambda \backslash T(F) \xrightarrow{H_M} X_*(M)/H_M(\Lambda) \rightarrow 1.$$

Because Λ is of finite index in $X_*(M_0)$ and the morphism $X_*(M_0) \rightarrow X_*(M)$ is surjective, the quotient $X_*(M)/H_M(\Lambda)$ is finite, and so

$$\begin{aligned} \mathrm{vol}_{dt}(\Lambda^{H_M} \backslash T(F)_M^1) &= \mathrm{vol}_{dt}(\Lambda \backslash T(F)) \cdot |X_*(M)/H_M(\Lambda)|^{-1} \\ &= \frac{|\mathrm{coker}[X_*(S)_\Gamma \rightarrow X_*(T)_\Gamma]|}{|\mathrm{ker}[X_*(S)_\Gamma \rightarrow X_*(T)_\Gamma]| \cdot |X_*(M)/H_M(X_*(S))|}. \end{aligned} \quad (2.5)$$

2.2.3. Comparison of weighted orbital integrals. The weight factors v_M and w_M^ξ are closely related, so we can compare the associated weighted orbital integrals.

Theorem 2.8 (Chaudouard and Laumon [8]). *We have the equality*

$$J_M(\gamma) = \mathrm{vol}(\mathfrak{a}_M/X_*(M)) \cdot J_M^\xi(\gamma).$$

Remark 2.3. For a general reductive algebraic group G , with the definition of w_M^ξ as explained in Remark 2.1, the comparison theorem becomes

$$J_M(\gamma) = \mathrm{vol}\left(\mathfrak{a}_{M_{G_{\mathrm{der}}}}^{G_{\mathrm{der}}} / X_*(M_{G_{\mathrm{der}}})\right) \cdot J_M^\xi(\gamma),$$

as can be seen from the proof below.

Chaudouard and Laumon work over the ring of adèles, but their proof carries over to the local setting. We reproduce their proof here, but to simplify the exposition, we assume moreover that G is simply connected. The key is to rewrite the convex polytope $\mathrm{Ec}_M(g)$ as alternating differences of translations of cones. We need some notation. For $P = MN_P \in \mathcal{F}(A)$, take a Borel subgroup $B \in \mathcal{P}(A)$ contained in P . Let Δ_B be the simple roots of $\Phi(G, A)$ with respect to B , and let $\Delta_{B,P} = \Delta_B \cap \Phi(N_P, A)$ and $\Delta_{B,P}^\vee$ be the associated coroots. The restriction $X^*(A) \rightarrow X^*(A_M)$ induces a bijection from $\Delta_{B,P}$ to a subset of $X^*(A_M)$ denoted Δ_P . Similarly, the projection $\mathfrak{a}_A \rightarrow \mathfrak{a}_M$ induces a bijection from $\Delta_{B,P}^\vee$ to a subset Δ_P^\vee . Obviously, the definition of Δ_P and Δ_P^\vee is independent of the choice of B . Moreover, they form bases of \mathfrak{a}_M^* and \mathfrak{a}_M , respectively. Let $(\varpi_\alpha)_{\alpha \in \Delta_P}$ be the basis of \mathfrak{a}_M^* dual to Δ_P^\vee .

For a generic element $\lambda \in \mathfrak{a}_M^*$, let

$$\Delta_P^\lambda = \{\alpha \in \Delta_P \mid \langle \lambda, \alpha \rangle < 0\},$$

and let φ_P^λ be the characteristic function of the cone

$$\{a \in \mathfrak{a}_M \mid \varpi_\alpha(a) > 0, \forall \alpha \in \Delta_P^\lambda; \varpi_\alpha(a) \leq 0, \forall \alpha \in \Delta_P \setminus \Delta_P^\lambda\}.$$

According to Arthur [3], the characteristic function of the convex polytope $\text{Ec}_M(g)$ is equal to the function

$$a \in \mathfrak{a}_M \mapsto \sum_{P \in \mathcal{P}(M)} (-1)^{|\Delta_P^\lambda|} \varphi_P^\lambda(-H_P(g) + a).$$

The proof is best illustrated by [4, Figure 11.1, p. 63]. It relies on the combinatorial identity

$$\sum_{F \subset S} (-1)^{|F|} = \begin{cases} 1 & \text{if } S = \emptyset, \\ 0 & \text{otherwise,} \end{cases}$$

for any finite set S . Now we can rewrite

$$w_M^\xi(g) = \sum_{\chi \in X_*(M)} \sum_{P \in \mathcal{P}(M)} (-1)^{|\Delta_P^\lambda|} \varphi_P^\lambda(-H_P(g) + \chi + \xi), \quad (2.6)$$

$$v_M(g) = \int_{\mathfrak{a}_M} \sum_{P \in \mathcal{P}(M)} (-1)^{|\Delta_P^\lambda|} \varphi_P^\lambda(-H_P(g) + a) da. \quad (2.7)$$

We introduce an extra exponential factor to treat the infinite sum in equation (2.6):

$$S_P(\lambda) = \sum_{\chi \in X_*(M)} \varphi_P^\lambda(-H_P(g) + \chi + \xi) e^{\langle \lambda, \chi \rangle}.$$

The series converges absolutely for generic λ , and hence

$$w_M^\xi(g) = \lim_{\lambda \rightarrow 0} \sum_{P \in \mathcal{P}(M)} (-1)^{|\Delta_P^\lambda|} S_P(\lambda),$$

where the limit is taken for generic $\lambda \in \mathfrak{a}_M^*$.

We can calculate $S_P(\lambda)$ explicitly. Let $\xi = [\xi]_P + \{\xi\}_P$, with $[\xi]_P \in X_*(M)$ and $\{\xi\}_P = \sum_{\alpha \in \Delta_P} r_\alpha \alpha^\vee$ for some $0 < r_\alpha < 1$. After a simple change of variables, we get

$$\begin{aligned} S_P(\lambda) &= e^{\langle \lambda, H_P(g) - [\xi]_P \rangle} \sum_{\chi \in X_*(M)} \varphi_P^\lambda(\chi + \{\xi\}_P) e^{\langle \lambda, \chi \rangle} \\ &= e^{\langle \lambda, H_P(g) - [\xi]_P \rangle} \sum_{(m_\alpha)_{\alpha \in \Delta_P}} e^{\langle \lambda, \sum_{\alpha} m_\alpha \alpha^\vee \rangle}, \end{aligned}$$

where $(m_\alpha)_{\alpha \in \Delta_P}$ runs over the integers satisfying $m_\alpha \geq 0$ for $\alpha \in \Delta_P^\lambda$ and $m_\alpha \leq -1$ for $\alpha \in \Delta_P \setminus \Delta_P^\lambda$. The geometric series can be calculated to be

$$S_P(\lambda) = (-1)^{|\Delta_P^\lambda|} e^{\langle \lambda, H_P(g) - [\xi]_P \rangle} \prod_{\alpha \in \Delta_P} \frac{1}{e^{\langle \lambda, \alpha^\vee \rangle} - 1}.$$

Let $c_P(\lambda) = \prod_{\alpha \in \Delta_P} (e^{\langle \lambda, \alpha^\vee \rangle} - 1)$. Taking everything together, we get

$$w_M^\xi(g) = \lim_{\lambda \rightarrow 0} \sum_{P \in \mathcal{P}(M)} c_P(\lambda)^{-1} e^{\langle \lambda, H_P(g) - [\xi]_P \rangle}. \quad (2.8)$$

Similarly, we can rewrite equation (2.7) as

$$\begin{aligned} v_M(g) &= \lim_{\lambda \rightarrow 0} \int_{\mathfrak{a}_M} \sum_{P \in \mathcal{P}(M)} (-1)^{|\Delta_P^\lambda|} \varphi_P^\lambda(-H_P(g) + a) e^{\langle \lambda, a \rangle} da \\ &= \lim_{\lambda \rightarrow 0} \sum_{P \in \mathcal{P}(M)} (-1)^{|\Delta_P^\lambda|} \int_{\mathfrak{a}_M} \varphi_P^\lambda(-H_P(g) + a) e^{\langle \lambda, a \rangle} da \\ &= \lim_{\lambda \rightarrow 0} \sum_{P \in \mathcal{P}(M)} e^{\langle \lambda, H_P(g) \rangle} \cdot \text{vol}(\mathfrak{a}_M / X_*(M)) \prod_{\alpha \in \Delta_P} \langle \lambda, \alpha^\vee \rangle^{-1}. \end{aligned}$$

Letting $d_P(\lambda) = \text{vol}(\mathfrak{a}_M / X_*(M))^{-1} \prod_{\alpha \in \Delta_P} \langle \lambda, \alpha^\vee \rangle$, we get

$$v_M(g) = \lim_{\lambda \rightarrow 0} \sum_{P \in \mathcal{P}(M)} d_P(\lambda)^{-1} \cdot e^{\langle \lambda, H_P(g) \rangle}. \quad (2.9)$$

To deal with limits of the form in equations (2.8) and (2.9) systematically, we need Arthur's notion of a (G, M) -family [2]. It is a family of smooth functions $(r_P(\lambda))_{P \in \mathcal{P}(M)}$ on \mathfrak{a}_M^* which satisfy, for any adjacent parabolic subgroups (P, P') , the property that $r_P(\lambda) = r_{P'}(\lambda)$ for any λ on the hyperplane defined by the unique coroot in $\Delta_P^\vee \cap (-\Delta_{P'}^\vee)$. For any such family, we define

$$r_M(\lambda) = \sum_{P \in \mathcal{P}(M)} d_P(\lambda)^{-1} r_P(\lambda),$$

for generic $\lambda \in \mathfrak{a}_M^*$. Arthur has shown in [2] that the function extends smoothly over all \mathfrak{a}_M^* . Let

$$r_M = \lim_{\lambda \rightarrow 0} r_M(\lambda).$$

It generalizes equation (2.9). Indeed, the functions

$$v_P(\lambda, g) = e^{\langle \lambda, H_P(g) \rangle}, \quad P \in \mathcal{P}(M),$$

form a (G, M) -family, and the resulting $v_M(g)$ is exactly Arthur's weight factor. From this point of view, we call r_M the *volume* of the (G, M) -family $(r_P(\lambda))_{P \in \mathcal{P}(M)}$.

Notice that the summands in equations (2.8) and (2.9) differ by a factor

$$w_P(\lambda, \xi) = \frac{d_P(\lambda)}{c_P(\lambda)} e^{-\langle \lambda, [\xi]_P \rangle},$$

and that they form a (G, M) -family. Letting $w_P(\lambda, g, \xi) = v_P(\lambda, g) w_P(\lambda, \xi)$, $P \in \mathcal{P}(M)$, they form a (G, M) -family and equation (2.8) can be rewritten as

$$w_M^\xi(g) = w_M(g, \xi). \quad (2.10)$$

In other words, we have expressed the lattice point-counting weight factor $w_M^\xi(g)$ as the volume of the product of two (G, M) -families.

We need a result of Arthur on the volume of the product of two (G, M) -families. Let $\{r_P(\lambda)\}_{P \in \mathcal{P}(M)}$ and $\{s_P(\lambda)\}_{P \in \mathcal{P}(M)}$ be two (G, M) -families. For $Q = LN_Q \in \mathcal{F}(M)$, let

$$r_R^Q(\lambda) = r_{RN_Q}(\lambda), \quad \forall R \in \mathcal{P}^L(M).$$

It is easy to see that $r_R^Q(\lambda)$, $R \in \mathcal{P}^L(M)$, form an (L, M) -family. The function $r_M^Q(\lambda)$ and the volume r_M^Q are defined in a similar way. From the (G, M) -family $\{s_P(\lambda)\}_{P \in \mathcal{P}(M)}$, Arthur has defined a smooth function $s'_Q(\lambda)$ on \mathfrak{a}_Q^* . The definition is quite involved, and we refer the reader to [2, §6]. Let $s'_Q = s'_Q(0)$.

Lemma 2.9 (Arthur [2, Lemma 6.3 and Corollary 6.4]). *Let $\{r_P(\lambda)\}_{P \in \mathcal{P}(M)}$ and $\{s_P(\lambda)\}_{P \in \mathcal{P}(M)}$ be two (G, M) -families, and let $r \cdot s$ be their product. Then for any $\lambda \in \mathfrak{a}_M^*$, we have*

$$(r \cdot s)_M(\lambda) = \sum_{Q \in \mathcal{F}(M)} r_M^Q(\lambda) s'_Q(\lambda).$$

In particular,

$$s_M(\lambda) = \sum_{P \in \mathcal{P}(M)} s'_P(\lambda).$$

In our situation, this implies

$$w_M^\xi(g) = (v(g) \cdot w(\xi))(0) = \sum_{Q \in \mathcal{F}(M)} v_M^Q(g) w'_Q(\xi) \quad (2.11)$$

and

$$w_M(\xi) = \sum_{P \in \mathcal{P}(M)} w'_P(\xi). \quad (2.12)$$

Similar results hold for Levi subgroups L containing M :

$$w_L^{\xi_L}(g) = \sum_{R \in \mathcal{F}(L)} v_L^R(g) w'_R(\xi_L), \quad (2.13)$$

$$w_L(\xi_L) = \sum_{Q \in \mathcal{P}(L)} w'_Q(\xi_L), \quad (2.14)$$

with $w'_R(\xi_L)$ deduced from the (G, L) -family

$$w_Q(\lambda, \xi_L) = \frac{d_Q(\lambda)}{c_Q(\lambda)} e^{-\langle \lambda, [\xi_L]_Q \rangle}, \quad \forall Q \in \mathcal{P}(L).$$

Setting $g = e \in G$ in equation (2.13), and noting that

$$v_L^R(e) = \begin{cases} 1 & \text{if } R \in \mathcal{P}(L), \\ 0 & \text{otherwise,} \end{cases}$$

we get

$$w_L^{\xi_L}(e) = \sum_{Q \in \mathcal{P}(L)} w'_Q(\xi_L) = w_L(\xi_L), \quad (2.15)$$

where the second equality is just equation (2.14).

Lemma 2.10.

$$\sum_{Q \in \mathcal{P}(L)} w'_Q(\xi) = \frac{\text{vol}(\mathfrak{a}_L/X_*(L))}{\text{vol}(\mathfrak{a}_M/X_*(M))} \cdot w_L^{\xi_L}(e) = \begin{cases} \text{vol}(\mathfrak{a}_M/X_*(M))^{-1} & \text{if } L = G, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Recall that given a (G, M) -family $\{s_P(\lambda)\}_{P \in \mathcal{P}(M)}$, we can define a (G, L) -family by setting

$$s_Q(\lambda) = s_P(\lambda), \quad \forall \lambda \in \mathfrak{a}_L^* \subset \mathfrak{a}_M^*,$$

for any $P \in \mathcal{P}(M)$, $P \subset Q$. Moreover, the function $s'_Q(\lambda)$ deduced from the (G, M) -family $\{s_P(\lambda)\}_{P \in \mathcal{P}(M)}$ is the same as that from the (G, L) -family $\{s_Q(\lambda)\}_{Q \in \mathcal{P}(L)}$, by [2, formula 6.3]. In this way, we get the (G, L) -family $\{w_Q(\lambda, \xi)\}_{Q \in \mathcal{P}(L)}$ and the equality

$$\sum_{Q \in \mathcal{P}(L)} w'_Q(\xi) = w_L(\xi) = \lim_{\lambda \rightarrow 0} \sum_{Q \in \mathcal{P}(L)} d_Q(\lambda)^{-1} \cdot \frac{d_P(\lambda)}{c_P(\lambda)} \cdot e^{-\langle \lambda, [\xi]_P \rangle},$$

by the second assertion of Lemma 2.9, where for each $Q \in \mathcal{P}(L)$ we take $P \in \mathcal{P}(M)$, $P \subset Q$, and the limit is taken for $\lambda \in \mathfrak{a}_L^*$ generic. Now that

$$\frac{d_P(\lambda)}{c_P(\lambda)} = \frac{\text{vol}(\mathfrak{a}_L/X_*(L))}{\text{vol}(\mathfrak{a}_M/X_*(M))} \cdot \frac{d_Q(\lambda)}{c_Q(\lambda)}$$

and $\langle \lambda, [\xi]_P \rangle = \langle \lambda, [\xi_L]_Q \rangle$ for any $\lambda \in \mathfrak{a}_L^*$, we get

$$\sum_{Q \in \mathcal{P}(L)} w'_Q(\xi) = \frac{\text{vol}(\mathfrak{a}_L/X_*(L))}{\text{vol}(\mathfrak{a}_M/X_*(M))} \cdot w_L(\xi_L) = \frac{\text{vol}(\mathfrak{a}_L/X_*(L))}{\text{vol}(\mathfrak{a}_M/X_*(M))} \cdot w_L^{\xi_L}(e),$$

where the last equality follows from equation (2.15). \square

By equation (2.11), we can rewrite $J_M^\xi(\gamma)$ as

$$\begin{aligned} J_M^\xi(\gamma) &= \int_{T(F) \backslash G(F)} \mathbb{1}_{\mathfrak{g}(\mathcal{O})}(\text{Ad}(g)^{-1}\gamma) w_M^\xi(g) \frac{dg}{dt} \\ &= \int_{T(F) \backslash G(F)} \mathbb{1}_{\mathfrak{g}(\mathcal{O})}(\text{Ad}(g)^{-1}\gamma) \left[\sum_{Q \in \mathcal{F}(M)} v_M^Q(g) w'_Q(\xi) \right] \frac{dg}{dt}. \end{aligned}$$

As v_M^Q is also left $T(F)$ -invariant, we can define

$$J_M^Q(\gamma) = \int_{T(F) \backslash G(F)} \mathbb{1}_{\mathfrak{g}(\mathcal{O})}(\text{Ad}(g)^{-1}\gamma) v_M^Q(g) \frac{dg}{dt}.$$

Let $Q = LN_Q$ be the standard Levi decomposition. Let dl be the Haar measure on $L(F)$ normalized by $\text{vol}_{dl}(L(\mathcal{O})) = 1$ and let dn be the Haar measure on $N_Q(F)$ normalized by

$\text{vol}_{dn}(N_Q(\mathcal{O})) = 1$. Using Iwasawa decomposition, we can rewrite $J_M^Q(\gamma)$ as

$$\begin{aligned} J_M^Q(\gamma) &= \int_{T(F) \backslash L(F)} \int_{N_Q(F)} \int_K \mathbb{1}_{\mathfrak{g}(\mathcal{O})}(\text{Ad}(nlk)^{-1}\gamma) v_M^Q(nlk) dk \cdot dn \cdot \frac{dl}{dt} \\ &= \int_{T(F) \backslash L(F)} \int_{N_Q(F)} \int_K \mathbb{1}_{\mathfrak{g}(\mathcal{O})}(\text{Ad}(nl)^{-1}\gamma) v_M^Q(l) dk \cdot dn \cdot \frac{dl}{dt} \\ &= \int_{T(F) \backslash L(F)} \left[\int_{N_Q(F)} \int_K \mathbb{1}_{\mathfrak{g}(\mathcal{O})}(\text{Ad}(nl)^{-1}\gamma) dk \cdot dn \right] v_M^Q(l) \frac{dl}{dt}, \end{aligned} \quad (2.16)$$

where in the second and third lines we have used the equalities $v_M^Q(nlk) = v_M^Q(l)$ and $v_M^Q(l) = v_M^L(l)$ respectively, which follow directly from definitions. Notice that

$$\begin{aligned} \int_{N_Q(F)} \int_K \mathbb{1}_{\mathfrak{g}(\mathcal{O})}(\text{Ad}(nl)^{-1}\gamma) dk \cdot dn &= |\{[nl] \in N_Q(F)lK/K \mid \text{Ad}(nl)^{-1}\gamma \in \mathfrak{g}(\mathcal{O})\}| \\ &= \left| \left(f_Q^{-1}([l]) \cap \mathcal{X}_\gamma \right) (\mathbf{F}_q) \right| \\ &= q^{\text{val}(\det(\text{ad}\gamma|_{\mathfrak{n}_Q, F}))} \cdot \mathbb{1}_{\mathfrak{l}(\mathcal{O})}(\text{Ad}(l)^{-1}\gamma), \end{aligned}$$

where the last equality follows from Proposition 2.4. Continuing the calculation of equation (2.16), we get

$$\begin{aligned} J_M^Q(\gamma) &= q^{\text{val}(\det(\text{ad}\gamma|_{\mathfrak{n}_Q, F}))} \cdot \int_{T(F) \backslash L(F)} \mathbb{1}_{\mathfrak{l}(\mathcal{O})}(\text{Ad}(l)^{-1}\gamma) v_M^L(l) \frac{dl}{dt} \\ &= q^{\text{val}(\det(\text{ad}\gamma|_{\mathfrak{n}_Q, F}))} \cdot J_M^L(\gamma). \end{aligned}$$

Combining all the foregoing calculations, we get

$$\begin{aligned} J_M^\xi(\gamma) &= \sum_{L \in \mathcal{L}(M)} \sum_{Q \in \mathcal{P}(L)} J_M^Q(\gamma) \cdot w'_Q(\xi) = \sum_{L \in \mathcal{L}(M)} \sum_{Q \in \mathcal{P}(L)} q^{\text{val}(\det(\text{ad}\gamma|_{\mathfrak{n}_Q, F}))} J_M^L(\gamma) \cdot w'_Q(\xi) \\ &= \sum_{L \in \mathcal{L}(M)} q^{\frac{1}{2}\text{val}(\det(\text{ad}\gamma|_{\mathfrak{g}_F/\mathfrak{l}_F}))} J_M^L(\gamma) \sum_{Q \in \mathcal{P}(L)} w'_Q(\xi) \\ &= \text{vol}(\mathfrak{a}_M/X_*(M))^{-1} \cdot J_M(\gamma), \end{aligned}$$

where the last equality follows from Lemma 2.10. This finishes the proof of Theorem 2.8.

3. Counting points by Arthur–Kottwitz reduction

From now on, we will assume that G_{der} is simply connected. The general case can be reduced to this one by focusing on each connected component. This extra assumption gives some technical convenience – for example, $M_{0, \text{der}}$ will be simply connected, Λ_{M_0} will be torsion-free, and we get an inclusion $\Lambda_{M_0} \hookrightarrow \mathfrak{a}_{M_0}$. Moreover, we have $X_*(M_0) = \Lambda_{M_0}$, according to [8, Lemma 11.6.1].

Fix $M \in \mathcal{L}(M_0)$ and let Π be a sufficiently regular positive (G, M) -orthogonal family. We count the number of points on $\Lambda^{H_M} \backslash \mathcal{X}_\gamma^{\nu_0}(\Pi)$, $\nu_0 \in \Lambda_G$. Generalizing our previous work [11], we show that it can be reduced to counting points on the *intermediate fundamental domains* $F_\gamma^{L, M}$, $L \in \mathcal{L}(M)$, and the counting result depends *quasi-polynomially* on the

truncation parameter. Moreover, counting points on $\Lambda^{H_M} \backslash F_\gamma^{L,M}$ can be further reduced to counting points on the fundamental domains $F_\gamma^{M'}$ for some $M' \in \mathcal{L}(M_0)$ ‘transversal’ to M .

3.1. Truncations on the affine Grassmannian

Recall the following definition of Arthur [1], which is a formalization of the orthogonal properties in Proposition 2.5:

Definition 3.1. A family $\Pi = (\lambda_P)_{P \in \mathcal{P}(M)}$ of elements in \mathfrak{a}_M^G is called a *positive* (G, M) -orthogonal family if it satisfies

$$\lambda_{P_1} - \lambda_{P_2} = n_{P_1, P_2} \cdot \pi_M^G(\beta_{P_1, P_2}), \quad \text{with } n_{P_1, P_2} \in \mathbf{R}_{\geq 0},$$

for any two adjacent parabolic subgroups $P_1, P_2 \in \mathcal{P}(M)$.

Given such a positive (G, M) -orthogonal family, we will denote again by Π the convex hull of the λ_P s. For $Q = LN_Q \in \mathcal{F}(M)$, parallel to $\text{Ec}_M^Q(x)$, we denote by Π^Q the face of Π whose vertices are λ_P , $P \in \mathcal{P}(M), P \subset Q$. With the projection π_M^L , it can be seen as a positive (L, M) -orthogonal family. This sets up a bijection between the set $\mathcal{F}(M)$ and the set of the faces of Π . Moreover, we denote by λ_Q or $\lambda_Q(\Pi)$ the element $\pi_{M, L}(\lambda_{P'})$ for any $P' \in \mathcal{P}(M), P' \subset Q$. One can show that $(\lambda_Q(\Pi))_{Q \in \mathcal{P}(L)}$ forms a positive (G, L) -orthogonal family. Later on, we also use the notation $(\lambda_{\bar{w}})_{\bar{w} \in W/W_M}$ for $(\lambda_{\bar{w} \cdot P})_{\bar{w} \in W/W_M}$, and we use the notation $\lambda_w(\Pi)$ or $\lambda_{w \cdot P}(\Pi)$ to indicate the vertex of Π indexed by $w \cdot P$.

Following Chaudouard and Laumon [7], we define the *truncated affine Grassmannian* $\mathcal{X}(\Pi)$ to be

$$\mathcal{X}(\Pi) = \{x \in \mathcal{X} \mid \pi_M^G(\text{Ec}_M(x)) \subset \Pi\}.$$

We want to point out that its connected components are also parametrized by Λ_G , but they are not isomorphic in general. However, there is periodicity in the connected components: let G^{ad} be the adjoint group of G and let $c_G : \Lambda_G \rightarrow \Lambda_{G^{\text{ad}}}$ be the projection induced by the natural projection $T \rightarrow T/Z_G$. For $\nu, \nu' \in \Lambda_G$, we have

$$\mathcal{X}^\nu(\Pi) = \mathcal{X}^{\nu'}(\Pi), \quad \text{if } c_G(\nu) = c_G(\nu'),$$

because they can be translated to each other by elements in $Z_G(F)$.

For a regular element $\gamma \in \mathfrak{t}(\mathcal{O})$, we can truncate the affine Springer fiber \mathcal{X}_γ similarly by defining

$$\mathcal{X}_\gamma(\Pi) = \mathcal{X}_\gamma \cap \mathcal{X}(\Pi),$$

and the same observation on the connected components of $\mathcal{X}(\Pi)$ holds also for $\mathcal{X}_\gamma(\Pi)$.

3.2. The intermediate fundamental domain

We generalize our construction of the fundamental domain F_γ in [11].³

³In [11], we confused Λ , Λ_{M_0} , and $\pi_0(T(F))$. With our current notation, there are morphisms $\Lambda \rightarrow \Lambda_{M_0}$ and $\Lambda \rightarrow \pi_0(T(F))$. Generally, they are not isomorphic. In particular, F_γ is

Let $P_1 = MN_1$ and $P_2 = MN_2 \in \mathcal{P}(M)$ be two adjacent parabolic subgroups. Let m_α be the unique positive integer such that the image of α^\vee in Λ_M is equal to $m_\alpha \cdot \beta_{P_1, P_2}$, and let

$$n(\gamma, P_1, P_2) = \sum_{\alpha \in \Phi(N_1, T_{\overline{F}}) \cap \Phi(N_2^-, T_{\overline{F}})} \text{val}(\alpha(\gamma)) \cdot m_\alpha.$$

It can be verified that $n(\gamma, P_1, P_2)$ is an integer.

Proposition 3.1 (Goresky-Kottwitz-MacPherson, [15]). *Set $x \in \mathcal{X}_\gamma$.*

(1) *For any two adjacent parabolic subgroups $P_1, P_2 \in \mathcal{P}(M)$, we have*

$$n(x, P_1, P_2) \leq n(\gamma, P_1, P_2).$$

(2) *The point x is regular in \mathcal{X}_γ if and only if the following two conditions hold:*

(a) *The point $f_P(x)$ is regular in \mathcal{X}_γ^M for all $P \in \mathcal{P}(M)$.*

(b) *For any two adjacent parabolic subgroups P_1, P_2 in $\mathcal{P}(M)$, one has*

$$n(x, P_1, P_2) = n(\gamma, P_1, P_2).$$

Notice that although Goresky, Kottwitz, and MacPherson work over the field $F = \mathbf{C}((\epsilon))$, their proof works for any field $F = k((\epsilon))$ with $\text{char}(k) > |W|$. Their result motivates our definition:

Definition 3.2. Take a regular point $x_0 \in \mathcal{X}_\gamma^{\text{reg}}$. Let

$$F_\gamma^{G, M} = \{x \in \mathcal{X}_\gamma \mid \text{Ec}_M(x) \subset \text{Ec}_M(x_0), \nu_G(x) = \nu_G(x_0)\}.$$

We call it an *intermediate fundamental domain* of \mathcal{X}_γ with respect to M .

We should have used the notation $F_{\gamma, x_0}^{G, M}$ to indicate the dependence on x_0 , but they are isomorphic to each other for any choice of the regular point x_0 . Indeed, for any two regular points x_1, x_2 , we can find $t \in T(F)$ such that $x_1 = t \cdot x_2$. Now that $\text{Ec}_M(tx) = \text{Ec}_M(x) + H_M(t)$, $\forall x \in \mathcal{X}$, the intermediate fundamental domain given by x_1 is just the translation by t of that given by x_2 . Notice that for $M = M_0$, we recover the fundamental domain F_γ . For simplicity, we assume that $\nu_G(x_0) = 0$.

Unlike the fundamental domain, the intermediate $F_\gamma^{G, M}$ is no longer of finite type for $M \supsetneq M_0$. Nonetheless, we have the following:

not a fundamental domain for the Λ -action – that is, $\mathcal{X}_\gamma \neq \bigcup_{\lambda \in \Lambda} \lambda \cdot F_\gamma$. Moreover, the group $\pi_0(T(F))$ may have a complicated torsion subgroup, which implies that F_γ may have complicated irreducible components as well, contrary to our expectation there. Actually, there should be a bijection between $\pi_0(F_\gamma)$ and $\pi_0(F_\gamma^{M_0})$, and both are isomorphic to $\pi_0(T(F))_{\text{tor}}$. Nevertheless, other results of [11] hold if we assume that G_{der} is simply connected, and the general case can be reduced to that one. This extra assumption is to make sure that for any Levi subgroup $M \in \mathcal{L}(M_0)$ we have Λ_M being torsion-free and we get an inclusion $\Lambda_M \hookrightarrow \mathfrak{a}_M$; they hold, as M_{der} is simply connected. Moreover, we have $X_*(M) = \Lambda_M$, according to [8, Lemma 11.6.1].

Proposition 3.2. *The free discrete abelian group Λ^{H_M} acts freely on $F_\gamma^{G,M}$, and the quotient $\Lambda^{H_M} \backslash F_\gamma^{G,M}$ is of finite type.*

Proof. Recall that $\Lambda^{H_M} = \Lambda \cap \ker(H_M)$ by definition, and hence it preserves $F_\gamma^{G,M}$ because left translation by $m \in \ker(H_M)$ does not change the polytope $\text{Ec}_M(x)$, due to the property

$$H_P(mx) = H_M(m) + H_P(x), \quad \forall m \in M(F), x \in \mathcal{X}, P \in \mathcal{P}(M).$$

For the finiteness issue, let $\Lambda_{M_0}^{H_M} \subset \Lambda_{M_0}$ be the kernel of the natural projection $\Lambda_{M_0} \rightarrow \Lambda_M$. By definition, we have

$$\pi_{M_0, M}^{-1}(\text{Ec}_M(x_0)) = \bigcup_{\nu \in \Lambda_{M_0}^{H_M}} (\nu + \text{Ec}_{M_0}(x_0)),$$

which implies that

$$F_\gamma^{G,M} = \bigcup_{\nu \in \Lambda_{M_0}^{H_M}} \mathcal{X}_\gamma^0(\nu + \text{Ec}_{M_0}(x_0)).$$

Now that $\Lambda \cong X_*(S)$ and $X_*(S) \hookrightarrow \Lambda_{M_0}$ is of finite index, the quotient $\Lambda^{H_M} \backslash \Lambda_{M_0}^{H_M}$ is of finite cardinality. Hence the quotient $\Lambda^{H_M} \backslash F_\gamma^{G,M}$ is dominated by the union of finitely many translations of F_γ under the natural projection $F_\gamma^{G,M} \rightarrow \Lambda^{H_M} \backslash F_\gamma^{G,M}$. As F_γ is of finite type, so is the quotient $\Lambda^{H_M} \backslash F_\gamma^{G,M}$. \square

A similar proof applies to the following:

Proposition 3.3. *Let Π be a regular positive (G, M) -orthogonal family. For any $\nu \in \Lambda_G$, the free discrete abelian group Λ^{H_M} acts freely on $\mathcal{X}_\gamma^\nu(\Pi)$, and the quotient $\Lambda^{H_M} \backslash \mathcal{X}_\gamma^\nu(\Pi)$ is of finite type. In particular,*

$$|(\Lambda^{H_M} \backslash \mathcal{X}_\gamma^\nu(\Pi))(\mathbf{F}_q)| < \infty.$$

Remark 3.1. In the definition of the (weighted) orbital integral, we are concerned more about analogues of $\Lambda^{H_M} \backslash (\mathcal{X}_\gamma^\nu(\Pi)(\mathbf{F}_q))$, but notice that there is bijection between

$$(\Lambda^{H_M} \backslash \mathcal{X}_\gamma^\nu(\Pi))(\mathbf{F}_q) \quad \text{and} \quad \Lambda^{H_M} \backslash (\mathcal{X}_\gamma^\nu(\Pi)(\mathbf{F}_q)),$$

because Λ^{H_M} acts freely on $\mathcal{X}_\gamma^\nu(\Pi)$ and the Galois group $\text{Gal}(\overline{\mathbf{F}}_q/\mathbf{F}_q)$ acts trivially on Λ^{H_M} . We will decompose the scheme $\Lambda^{H_M} \backslash \mathcal{X}_\gamma^\nu(\Pi)$ in different ways, and the bijection given here implies that we can deduce equality of rational points over \mathbf{F}_q from the decomposition of schemes.

In the following, we simplify the notation $\text{Ec}_M(x_0)$ to $\Sigma_\gamma^{G,M}$. For $\nu \in \Lambda_G$, let

$$F_\gamma^{G,M,\nu} := \mathcal{X}_\gamma^\nu(\Sigma_\gamma^{G,M}).$$

As we have explained before, it depends only on the class $c_G(\nu) \in \Lambda_{G^{\text{ad}}}$. For $M = M_0$, we simplify Σ_γ^{G,M_0} to Σ_γ and $F_\gamma^{G,M_0,\nu}$ to F_γ^ν .




Figure 1. Partition of \mathfrak{a}_A^G for GL_3 .

3.3. The Arthur–Kottwitz reduction

Recall that we can reduce the geometry of \mathcal{X}_γ to that of its fundamental domain by the Arthur–Kottwitz reduction [11]. The construction can be generalized to our current setting.

Let Q_0 be the unique parabolic subgroup in $\mathcal{P}(M)$ which contains P_0 . Let $\varsigma \in \mathfrak{a}_M^G$ be such that $\alpha(\varsigma)$ is positive but almost equal to 0 for any $\alpha \in \Delta_{Q_0}$. Let $D_M = (\lambda_P)_{P \in \mathcal{P}(M)}$ be the (G, M) -orthogonal family given by

$$\lambda_P = H_P(x_0) + w \cdot \varsigma, \quad (3.1)$$

where $w \in W$ is any element satisfying $P = w \cdot Q_0$. For $Q = LN_Q \in \mathcal{F}(M)$, define $R_Q^{G, M}$ to be the subset of \mathfrak{a}_M^G satisfying the conditions

$$\begin{aligned} \pi_M^L(a) &\subset D_M^Q, \\ \alpha(\pi_{M, L}(a)) &\geq \alpha(\pi_{M, L}(\lambda_Q)), \quad \forall \alpha \in \Delta_Q. \end{aligned}$$

This gives us a partition which dates back at least to Arthur [3]:

$$\mathfrak{a}_M^G = \bigcup_{Q \in \mathcal{F}(M)} R_Q^{G, M}. \quad (3.2)$$

It induces a disjoint partition of Λ_M via the map $\Lambda_M \rightarrow \mathfrak{a}_M^G$, as we have perturbed $(H_P(x_0))_{P \in \mathcal{P}(M)}$ with ς . Figure 1 gives an illustration of the partition for the group GL_3 and $M_0 = T = A$.

Similar to [11, Lemma 3.1], we have the following result due to Proposition 3.1:

Lemma 3.4. *For any $x \in \mathcal{X}_\gamma$, there exists a unique $Q \in \mathcal{F}(M)$ such that*

$$\pi_M^G(Ec_M^Q(x)) \subset R_Q^{G, M}.$$

The referee has suggested an equivalent form of the lemma, which is much easier to understand and to prove: let $\mathfrak{a}_M^G = \bigcup_{Q \in \mathcal{F}(M)} R'_Q$ be the partition attached to the

positive (G, M) -orthogonal family $(H_P(x_0) - H_P(x) + w \cdot \varsigma)_{P \in \mathcal{P}(M)}$, then the statement is equivalent to the existence of a unique $Q \in \mathcal{F}(M)$ such that $0 \in R'_Q$. Here the positiveness of the (G, M) -orthogonal family is due to Proposition 3.1. Let

$$S_Q^{G,M} := \left\{ x \in \mathcal{X}_\gamma \mid \pi_M^G \left(\text{Ec}_M^Q(x) \right) \subset R_Q^{G,M} \right\}.$$

We thus get a disjoint partition

$$\mathcal{X}_\gamma = \mathcal{X}_\gamma(D_M) \cup \bigcup_{\substack{Q \in \mathcal{F}(M) \\ Q \neq G}} S_Q^{G,M}. \quad (3.3)$$

For each parabolic subgroup $Q = LN_Q \in \mathcal{F}(M)$, consider the restriction of the retraction $f_Q : \mathcal{X} \rightarrow \mathcal{X}^L$ to $S_Q^{G,M}$; its image is $S_Q^{G,M} \cap \mathcal{X}_\gamma^L$. Recall that the connected components of \mathcal{X}_γ^L are fibers of the map $\nu_L : \mathcal{X}_\gamma^L \rightarrow \Lambda_L$. For $\nu \in \Lambda_L$, let $\mathcal{X}_\gamma^{L,\nu}$ be its fiber at ν . Letting

$$S_Q^{G,M,\nu} = S_Q^{G,M} \cap f_Q^{-1}(\mathcal{X}_\gamma^{L,\nu}),$$

we have

$$S_Q^{G,M,\nu} \cap \mathcal{X}_\gamma^{L,\nu} = \mathcal{X}_\gamma^{L,\nu} \left(D_M^Q \right).$$

Proposition 3.5. *The strata $S_Q^{G,M,\nu}$ are locally closed subschemes of \mathcal{X}_γ , and the retraction $f_Q : S_Q^{G,M,\nu} \rightarrow \mathcal{X}_\gamma^{L,\nu} \left(D_M^Q \right)$ is an iterated affine fibration over \mathbf{F}_q of dimension*

$$\text{val}(\det(\text{ad}(\gamma \mid \mathfrak{n}_{Q,F}))).$$

Indeed, by the bound on $\text{Ec}_M(x)$ given by Proposition 3.1, we get

$$S_Q^{G,M,\nu} = \mathcal{X}_\gamma \cap f_Q^{-1} \left(\mathcal{X}_\gamma^{L,\nu} \left(D_M^Q \right) \right).$$

It is an iterated affine fibration over $\mathcal{X}_\gamma^{L,\nu} \left(D_M^Q \right)$ by Proposition 2.4.

The decomposition (3.3) can thus be refined to

$$\mathcal{X}_\gamma = \mathcal{X}_\gamma(D_M) \cup \bigcup_{\substack{Q = LN_Q \in \mathcal{F}(M) \\ Q \neq G}} \bigcup_{\nu \in \Lambda_L \cap \pi_L(R_Q^{G,M})} S_Q^{G,M,\nu}, \quad (3.4)$$

where we have loosely used $\Lambda_L \cap \pi_L \left(R_Q^{G,M} \right)$ to mean elements in Λ_L whose projection to \mathfrak{a}_L^G lies in $\pi_L \left(R_Q^{G,M} \right)$. Similar notations will be used later on. The decomposition (3.4) will also be called the *Arthur–Kottwitz reduction*. Notice that the stratum $S_Q^{G,M,\nu}$ is an iterated affine fibration over $\mathcal{X}_\gamma^{L,\nu} \left(D_M^Q \right) = F_\gamma^{L,M,\nu}$, and the latter is related to $F_\gamma^{L,M}$ again by the Arthur–Kottwitz reduction, similar to what is explained in [11, Lemma 3.4].

As in [11], the existence of Arthur–Kottwitz reduction implies the following:

Corollary 3.6. *For any $\gamma \in \mathfrak{t}(\mathcal{O})$, suppose that $F_\gamma^{L,M}$ is cohomologically pure for any proper Levi subgroup $L \in \mathcal{L}(M)$. Then \mathcal{X}_γ is cohomologically pure if and only if $F_\gamma^{G,M}$ is.*

We can restrict the Arthur–Kottwitz reduction to the truncated affine Springer fibers. A positive (G, M) -orthogonal family $\Pi = (\mu_P)_{P \in \mathcal{P}(M)}$ is said to be *regular* with respect to D_M if $\mu_P \in R_P^{G, M}$, $\forall P \in \mathcal{P}(M)$. In this case, each $S_Q^{G, M, \nu}$ is either contained in $\mathcal{X}_\gamma^\nu(\Pi)$ or disjoint from it. So we have

$$\mathcal{X}_\gamma(\Pi) = \mathcal{X}_\gamma(D_M) \cup \bigcup_{\substack{Q=LN_Q \in \mathcal{F}(M) \\ Q \neq G}} \bigcup_{\substack{\nu \in \Lambda_L \cap \pi_L(R_Q^{G, M}) \\ \cap \pi_L(\Pi)}} S_Q^{G, M, \nu}. \quad (3.5)$$

The reduction can be further restricted to each connected component of $\mathcal{X}_\gamma(\Pi)$:

$$\mathcal{X}_\gamma^{\nu_0}(\Pi) = \mathcal{X}_\gamma^{\nu_0}(D_M) \cup \bigcup_{\substack{Q=LN_Q \in \mathcal{F}(M) \\ Q \neq G}} \bigcup_{\substack{\nu \in \Lambda_L^{\nu_0} \cap \pi_L(R_Q^{G, M}) \\ \cap \pi_L(\Pi)}} S_Q^{G, M, \nu}. \quad (3.6)$$

As we have explained, left translation by elements in $\ker(H_M)$ does not change the polytope $\text{Ec}_M(x)$, and hence the group Λ^{H_M} acts on each item of equation (3.6). Now that we have finiteness results – Propositions 3.2 and 3.3, combined with Proposition 3.5 and the periodicity of $F_\gamma^{L, M, \nu}$ in $\nu \in \Lambda_L$ – equation (3.6) implies equality of counting points:

Corollary 3.7. *We have the equality*

$$\begin{aligned} |(\Lambda^{H_M} \setminus \mathcal{X}_\gamma^{\nu_0}(\Pi))(\mathbf{F}_q)| &= |(\Lambda^{H_M} \setminus F_\gamma^{G, M, \nu_0})(\mathbf{F}_q)| + \sum_{\substack{Q=LN_Q \in \mathcal{F}(M) \\ Q \neq G}} \sum_{\mu \in \Lambda_{L, \text{ad}}} q^{\text{val}(\det(\text{ad}_\gamma | \mathfrak{n}_{Q, F}))} \\ &\quad \cdot |(\Lambda^{H_M} \setminus F_\gamma^{L, M, \mu})(\mathbf{F}_q)| \cdot \left| \Lambda_L^{\nu_0} \cap \pi_L(R_Q^{G, M}) \cap \pi_L(\Pi) \cap c_L^{-1}(\mu) \right|. \end{aligned}$$

Notice that the term $\left| \Lambda_L^{\nu_0} \cap \pi_L(R_Q^{G, M}) \cap \pi_L(\Pi) \cap c_L^{-1}(\mu) \right|$ counts the number of lattice points in a polytope. Well-known techniques from toric geometry tells us that the counting result depends *quasi-polynomially* on the size of the polytope.

Remark 3.2. As the foregoing constructions rely ultimately on the bound of $\text{Ec}_M(x)$ given by Proposition 3.1, they continue to work if we replace $\Sigma_\gamma^{G, M}$ from the beginning by any integral positive (G, M) -orthogonal family Σ which satisfies

$$\lambda_{P_1}(\Sigma) - \lambda_{P_2}(\Sigma) = n_{P_1, P_2} \cdot \beta_{P_1, P_2}, \quad \text{with } n_{P_1, P_2} \geq n(\gamma, P_1, P_2), \quad (3.7)$$

for any two adjacent parabolic subgroups $P_1, P_2 \in \mathcal{P}(M)$. The resulting decomposition will also be called the Arthur–Kottwitz reduction.

3.4. Counting points on the intermediate fundamental domains

Although the intermediate fundamental domains $F_\gamma^{G, M}$ look like something new, it turns out that counting points of $\Lambda^{H_M} \setminus F_\gamma^{G, M}$ can be reduced to counting points of the fundamental domains.

As explained in the proof of Proposition 3.2, we have

$$\pi_{M_0, M}^{-1}(\Sigma_\gamma^{G, M}) = \bigcup_{\nu \in \Lambda_{M_0}^{H_M}} (\nu + \Sigma_\gamma)$$

and

$$F_\gamma^{G, M} = \bigcup_{\nu \in \Lambda_{M_0}^{H_M}} \mathcal{X}_\gamma^0(\nu + \Sigma_\gamma).$$

Let $P_0^M = P_0 \cap M$, and let $\beta_1^\vee, \dots, \beta_{r'}^\vee \in \Lambda_{M_0}^{H_M}$ be a basis of $\mathfrak{a}_{M_0}^M$ which is positive with respect to P_0^M . For $\mu_1, \mu_2 \in \Lambda_{M_0}^{H_M}$, we say that $\mu_1 \preccurlyeq_{P_0^M} \mu_2$ if $\mu_2 - \mu_1$ is a linear combination of β_i^\vee s with positive coefficients. This defines a partial order $\preccurlyeq_{P_0^M}$ on $\Lambda_{M_0}^{H_M}$. For $\mu \in \Lambda_{M_0}^{H_M}$, define

$$\Lambda_{M_0, \preccurlyeq \mu}^{H_M} := \left\{ \nu \in \Lambda_{M_0}^{H_M} \mid \nu \preccurlyeq_{P_0^M} \mu \right\} \quad \text{and} \quad \Pi_{\gamma, \preccurlyeq \mu}^{G, M} := \bigcup_{\nu \in \Lambda_{M_0, \preccurlyeq \mu}^{H_M}} (\nu + \Sigma_\gamma).$$

Then $\Pi_{\gamma, \preccurlyeq \mu}^{G, M}$ is a semi-infinite polytope in $\mathfrak{a}_{M_0}^G$ and $\Lambda_{M_0, \preccurlyeq \mu}^{H_M}$ is the integral points in it. Similar definitions hold for $\Lambda_{M_0, \prec \mu}^{H_M}$ and $\Pi_{\gamma, \prec \mu}^{G, M}$. But notice that $\Pi_{\gamma, \prec \mu}^{G, M}$ is not a semi-infinite polytope: it is the union of finitely many semi-infinite polytopes of the form $\Pi_{\gamma, \preccurlyeq \mu'}^{G, M}$, $\mu' \in \Lambda_{M_0}^{H_M}$. Define

$$F_{\gamma, \preccurlyeq \mu}^{G, M} := \mathcal{X}_\gamma^0(\Pi_{\gamma, \preccurlyeq \mu}^{G, M}) = \bigcup_{\nu \in \Lambda_{M_0, \preccurlyeq \mu}^{H_M}} \mathcal{X}_\gamma^0(\nu + \Sigma_\gamma)$$

and similarly

$$F_{\gamma, \prec \mu}^{G, M} := \bigcup_{\nu \in \Lambda_{M_0, \prec \mu}^{H_M}} \mathcal{X}_\gamma^0(\nu + \Sigma_\gamma).$$

It is the union of finitely many $F_{\gamma, \preccurlyeq \mu'}^{G, M}$, $\mu' \in \Lambda_{M_0}^{H_M}$. Define

$$F_{\gamma, \mu}^{G, M} := F_{\gamma, \preccurlyeq \mu}^{G, M} \setminus F_{\gamma, \prec \mu}^{G, M}.$$

Being a difference of closed subschemes, $F_{\gamma, \mu}^{G, M}$ is locally closed in \mathcal{X}_γ^0 . As $F_{\gamma, \preccurlyeq \mu}^{G, M}$ is semi-infinite unions of translations of the fundamental domains, they are all isomorphic, and similarly for $F_{\gamma, \prec \mu}^{G, M}$. Hence $F_{\gamma, \mu}^{G, M}$ are all isomorphic. Moreover,

$$F_{\gamma, \preccurlyeq \mu}^{G, M} = \bigsqcup_{\nu \in \Lambda_{M_0, \preccurlyeq \mu}^{H_M}} F_{\gamma, \nu}^{G, M}$$

by induction, and

$$F_\gamma^{G, M} = \lim_{\mu \rightarrow \infty} F_{\gamma, \preccurlyeq \mu}^{G, M} = \bigsqcup_{\nu \in \Lambda_{M_0}^{H_M}} F_{\gamma, \nu}^{G, M}.$$

From all these we conclude the following:

Proposition 3.8. $F_{\gamma,\mu}^{G,M}$ is isomorphic to each other for all $\mu \in \Lambda_{M_0}^{H_M}$. In particular,

$$\left| (\Lambda^{H_M} \setminus F_{\gamma}^{G,M}) (\mathbf{F}_q) \right| = \left| \Lambda^{H_M} \setminus \Lambda_{M_0}^{H_M} \right| \cdot \left| F_{\gamma,\mu}^{G,M} (\mathbf{F}_q) \right|.$$

Counting points on $F_{\gamma,\mu}^{G,M}$ can be reduced to the fundamental domains via a process similar to the Arthur–Kottwitz reduction. To begin with, notice that $\Pi_{\gamma,\mu}^{G,M}$ is bounded only in directions that are positive with respect to P_0^M . Indeed, its vertices are indexed by $P \in \mathcal{P}(M_0)$ satisfying $P \cap M = P_0^M$, and its faces by $Q \in \mathcal{F}(M_0)$ such that $Q \cap M \supset P_0^M$. Then we define a semi-infinite polytope $\Pi_{\gamma,\mu^-}^{G,M}$ which is a translation of $\Pi_{\gamma,\mu}^{G,M}$: let Δ^M be the set of simple roots in $\Phi(M, A)$ with respect to $B_0 \cap M$, and let $\Delta_{P_0^M}^M = \Delta^M \cap \Phi(N_{P_0^M}, A)$, with $N_{P_0^M}$ the unipotent radical of P_0^M . For $\alpha \in \Delta_{P_0^M}^M$, let $\{\omega_\alpha^\vee\}$ be the corresponding fundamental coweights. Let

$$\mu^- = \mu - \pi_2 \left(\sum_{\alpha \in \Delta_{P_0^M}^M} \omega_\alpha^\vee \right), \quad (3.8)$$

where π_2 is the projection to the second factor in the orthogonal decomposition $\mathfrak{a}_A^G = \mathfrak{a}_A^{M_0} \oplus \mathfrak{a}_{M_0}^M \oplus \mathfrak{a}_M^G$. Then $\Pi_{\gamma,\mu^-}^{G,M}$ is a translation of $\Pi_{\gamma,\mu}^{G,M}$ by the same vector. Now let $\varsigma \in \mathfrak{a}_{M_0}^G$ be a generic element such that $\alpha(\varsigma)$ is positive but almost equal to 0 for any $\alpha \in \Delta_{P_0}^G$. We perturb the semi-infinite polytope $\Pi_{\gamma,\mu^-}^{G,M}$ to a similar one Π' , with vertices

$$\lambda_P(\Pi') = \lambda_P \left(\Pi_{\gamma,\mu^-}^{G,M} \right) + w \cdot \varsigma, \quad \forall P \in \mathcal{P}(M_0), \quad P \cap M = P_0^M,$$

where $w \in W$ is any element satisfying $P = w \cdot P_0$. Both $\Pi_{\gamma,\mu}^{G,M}$ and Π' can be seen as limits of positive (G, M_0) -orthogonal families containing Σ_γ , hence we can apply an analogue of the Arthur–Kottwitz reduction to get a decomposition of the complement $\mathcal{X}_\gamma^0 \left(\Pi_{\gamma,\mu}^{G,M} \right) \setminus \mathcal{X}_\gamma^0(\Pi')$. For $Q = LN_Q \in \mathcal{F}(M_0)$ satisfying $Q \cap M \supset P_0^M$, define $R_{\Pi',Q}$ to be the subset of $\mathfrak{a}_{M_0}^G$ satisfying conditions

$$\begin{aligned} \pi_{M_0}^L(a) &\subset \Pi'^Q, \\ \alpha(\pi_{M_0,L}(a)) &\geq \alpha(\pi_{M_0,L}(\lambda_Q(\Pi'))), \quad \forall \alpha \in \Delta_Q. \end{aligned}$$

This gives us a partition

$$\mathfrak{a}_{M_0}^G = \Pi' \cup \bigcup_{\substack{Q \in \mathcal{F}(M_0), Q \neq G \\ Q \cap M \supset P_0^M}} R_{\Pi',Q}. \quad (3.9)$$

It induces a disjoint partition of Λ_{M_0} . For $G = \mathrm{GL}_3$, γ split, and $M = M_{\alpha_{12}}$, we get Figure 2.

Running the same construction as in §3.3, for $Q = LN_Q \in \mathcal{F}(M_0)$, $Q \neq G$ and $Q \cap M \supset P_0^M$, define

$$S_{\Pi',Q} := \left\{ x \in \mathcal{X}_\gamma \mid \pi_{M_0}^G \left(\mathrm{Ec}_{M_0}^Q(x) \right) \subset R_{\Pi',Q} \right\},$$




Figure 2. Arthur–Kottwitz reduction for $\Pi_{\gamma, \leq \mu}^{G, M}$.

and let

$$S_{\Pi', Q}^{\nu} = S_{\Pi', Q} \cap f_Q^{-1}(\mathcal{X}_{\gamma}^{L, \nu}), \quad \forall \nu \in \Lambda_L.$$

We get a disjoint partition

$$\mathcal{X}_{\gamma}^0(\Pi_{\gamma, \leq \mu}^{G, M}) = \mathcal{X}_{\gamma}^0(\Pi') \cup \bigcup_{\substack{Q \in \mathcal{F}(M_0), Q \neq G \\ Q \cap M \supset P_0^M}} \bigcup_{\substack{\nu \in \Lambda_L^0 \cap \pi_L(R_{\Pi', Q}) \\ \cap \pi_L(\Pi_{\gamma, \leq \mu}^{G, M})}} S_{\Pi', Q}^{\nu}. \quad (3.10)$$

The strata $S_{\Pi', Q}^{\nu}$ are locally closed subschemes of \mathcal{X}_{γ} , and the retraction $f_Q : S_{\Pi', Q}^{\nu} \rightarrow \mathcal{X}_{\gamma}^{L, \nu}(\Pi'Q)$ is an iterated affine fibration over \mathbf{F}_q of dimension $\text{val}(\det(\text{ad}(\gamma | \mathfrak{n}_{Q, F})))$.

Proposition 3.9. *We have the equality*

$$F_{\gamma, \mu}^{G, M} = \bigcup_{\substack{Q \in \mathcal{F}(M_0), Q \neq G \\ Q \cap M = P_0^M}} \bigcup_{\substack{\nu \in \Lambda_L^0 \cap \pi_L(R_{\Pi', Q}) \\ \cap \pi_L(\Pi_{\gamma, \leq \mu}^{G, M})}} S_{\Pi', Q}^{\nu}.$$

Moreover, the index set $\Lambda_L^0 \cap \pi_L(R_{\Pi', Q}) \cap \pi_L(\Pi_{\gamma, \leq \mu}^{G, M})$ consists of at most one element, and is nonempty if and only if Q is not contained in any $Q' \in \mathcal{F}(M)$.

Proof. For the first assertion, by construction, the points $x \in F_{\gamma, \mu}^{G, M}$ are characterized by the property

$$\text{Ec}_{M_0}^Q(x) \subset (\mu + \Sigma_{\gamma})^Q$$

for some $Q \in \mathcal{F}(M_0)$, $Q \neq G$, $Q \cap M = P_0^M$. Since this is also the property characterizing points on the right-hand side of the equality, we get the equality as claimed. The second assertion follows from the observation that Π' is a slight expansion of $\Pi_{\gamma, \leq \mu^-}^{G, M}$, and hence

the regions $R_{\Pi', Q}$, Q contained in some maximal parabolic subgroup in $\mathcal{F}(M)$, contain no elements in $\Lambda_L^0 \cap \pi_L \left(\Pi_{\gamma, \preccurlyeq \mu}^{G, M} \right)$. \square

If the index set $\Lambda_L^0 \cap \pi_L (R_{\Pi', Q}) \cap \pi_L \left(\Pi_{\gamma, \preccurlyeq \mu}^{G, M} \right)$ is nonempty, we denote by ν_Q the unique element in it. Let $\mu_Q \in \Lambda_L^0$ be the unique element such that $\alpha(\mu_Q) = 1$ for all $\alpha \in \Delta_Q$; then we have

$$\mathcal{X}_{\gamma}^{L, \nu_Q} \left(\Pi' Q \right) \cong F_{\gamma}^{L, \mu_Q}.$$

Combining with the fact that

$$f_Q : S_{\Pi', Q}^{\nu_Q} \rightarrow \mathcal{X}_{\gamma}^{L, \nu_Q} \left(\Pi' Q \right)$$

is an iterated affine fibration, we get the following:

Corollary 3.10. *For $M \in \mathcal{L}(M_0)$, $M \neq M_0$, we have the equality*

$$\left| F_{\gamma, \mu}^{G, M}(\mathbf{F}_q) \right| = \sum_{\substack{Q = LN_Q \in \mathcal{F}(M_0), Q \neq G \\ \text{satisfying } (*)}} q^{\frac{1}{2} \text{val}(\det(\text{ad} \gamma | \mathfrak{g}_F / \mathfrak{l}_F))} \cdot \left| F_{\gamma}^{L, \mu_Q}(\mathbf{F}_q) \right|,$$

where $(*)$ refers to the condition that $Q \cap M = P_0^M$ and $Q \not\subseteq Q', \forall Q' \in \mathcal{F}(M)$.

Notice that the equation does not involve the fundamental domain F_{γ} . Together with Corollary 3.7 and Proposition 3.8, we get an expression of $\left| (\Lambda^{H_M} \setminus \mathcal{X}_{\gamma}^{\nu_0}(\Pi))(\mathbf{F}_q) \right|$ in terms of $\left| F_{\gamma}^{L, \mu_Q}(\mathbf{F}_q) \right|$, $L \in \mathcal{L}(M_0)$, $L \neq G$. Recalling that counting points on F_{γ}^{L, μ_Q} can be reduced to counting points on $F_{\gamma}^{L'}$, $L' \in \mathcal{L}(M_0)$, $L' \subset L$, by the Arthur–Kottwitz reduction we get an expression of $\left| (\Lambda^{H_M} \setminus \mathcal{X}_{\gamma}^{\nu_0}(\Pi))(\mathbf{F}_q) \right|$ in terms of $\left| F_{\gamma}^{L'}(\mathbf{F}_q) \right|$, $L' \in \mathcal{L}(M_0)$.

4. Counting points by Harder–Narasimhan reduction

The number of points $\left| (\Lambda^{H_M} \setminus \mathcal{X}_{\gamma}^{\nu_0}(\Pi))(\mathbf{F}_q) \right|$, $\nu_0 \in \Lambda_G$, can also be counted by the Harder–Narasimhan reduction. Comparison with results from the last section gives us a recursive relation between Arthur’s weighted orbital integrals and the number of rational points on the fundamental domains.

4.1. Harder–Narasimhan reduction on the affine Springer fibers

We have introduced a notion of ξ -stability on the affine Grassmannian and constructed the associated Harder–Narasimhan reduction in [10]. In this section, we generalize it to a broader setup. The following lemma is an analogue of [8, Proposition 5.6.1]. Let S be an affine \mathbf{F}_q -scheme and set $x \in \mathcal{X}(S)$. For every point $s \in S$, let $x_s \in \mathcal{X}(k(s))$ be the base change of x to the residue field $k(s)$ of S at s . Let \mathcal{C}_x be the map on S which sends every point $s \in S$ to the convex polytope $\text{Ec}(x_s)$.

Lemma 4.1. *Suppose that S is noetherian. The map \mathcal{C}_x from S to the set of convex polytopes in \mathfrak{a}_A^G ordered by inclusion is lower semicontinuous. In other words, for any*

convex polytope Ξ , the set

$$\{s \in S \mid \mathcal{C}_x(s) \supset \Xi\}$$

is open.

Proof. To begin with, we show that \mathcal{C}_x is constructible and takes only finitely many values. Passing to the irreducible components of S , we can suppose that S is irreducible. Let η be the generic point of S , and let $g_\eta \in G(k(\eta)((\epsilon)))$ be a representative of x_η . For $B = AN \in \mathcal{P}(A)$, we have the Iwasawa decomposition

$$g_\eta = n_\eta a_\eta k_\eta,$$

where $n_\eta \in N(k(\eta)((\epsilon)))$, $a_\eta \in A(k(\eta)((\epsilon)))$, and $k_\eta \in G(k(\eta)[[\epsilon]])$. Because η is the generic point and the map $\nu_A : \mathcal{X}^A \rightarrow X_*(A)$ is essentially the valuation map, there exists an open subscheme U of S such that $H_B(x_s) = \nu_A(a_\eta)$ for any $x \in U$. As $\text{Ec}(x_s)$ is the convex hull of $H_B(x_s)$, $B \in \mathcal{P}(A)$, the map \mathcal{C}_x takes the constant value $\text{Ec}(x_\eta)$ on the intersection of all such open subschemes U . This proves the constructibility of \mathcal{C}_x . By noetherian induction, the map \mathcal{C}_x takes only finitely many values.

To finish the proof, we only need to show that the map \mathcal{C}_x decreases under specialization. In other words, let S be the spectrum of a discrete valuation ring and let s be its special point and η its generic point. Then

$$\text{Ec}(x_s) \subset \text{Ec}(x_\eta).$$

This is equivalent to the assertion that

$$f_B(x_s) \prec_B f_B(x_\eta), \quad \forall B \in \mathcal{P}(A), \quad (4.1)$$

where \prec_B is the order on $X_*(A)$ such that $\mu_1 \prec_B \mu_2$ if and only if $\mu_2 - \mu_1$ is a positive linear combination of positive coroots with respect to B .

Let $\mu = f_B(x_\eta) \in X_*(A)$. By definition, we have

$$x_\eta \in U_B((\epsilon))\epsilon^\mu G[[\epsilon]]/G[[\epsilon]],$$

where U_B is the unipotent radical of B . So

$$x_s \in \overline{x_\eta} \subset \overline{U_B((\epsilon))\epsilon^\mu G[[\epsilon]]/G[[\epsilon]]} = \bigcup_{\substack{\lambda \in X_*(A) \\ \lambda \prec_B \mu}} U_B((\epsilon))\epsilon^\lambda G[[\epsilon]]/G[[\epsilon]],$$

which implies the relation (4.1). \square

Definition 4.1. Let $\xi \in \mathfrak{a}_M^G$ be a generic element. A point $x \in \mathcal{X}$ is said to be ξ -stable if the polytope $\pi^G(\text{Ec}_M(x))$ contains ξ .

As $\text{Ec}_M(x) = \pi_M(\text{Ec}(x))$, the subset

$$\mathcal{X}^\xi = \{x \in \mathcal{X} \mid \xi \in \pi^G(\text{Ec}_M(x))\}$$

is an open sub-ind- \mathbf{F}_q -scheme of \mathcal{X} by Lemma 4.1. This being shown, all the other constructions of [10] generalize.

Remark 4.1. When $M = A$, we recover the ξ -stability of [10]. In that work, we prove that the notion of ξ -stability coincides with the notion of stability for a twisted action of A on \mathcal{X} . We believe that this holds also in the current setting, with the torus A_M playing the role of A . If this holds, we can conclude that the quotient \mathcal{X}^ξ/A_M exists as an ind- \mathbf{F}_q -scheme.

Harder–Narasimhan reduction works as well in this setting. For $Q = LN_Q \in \mathcal{F}(M)$, let $\Phi_Q(G, L)$ be the image of $\Phi(N_Q, A)$ in $(\mathfrak{a}_L^G)^*$. For any point $a \in \mathfrak{a}_L^G$, we define a cone in \mathfrak{a}_L^G ,

$$D_Q(a) = \{y \in \mathfrak{a}_L^G \mid \alpha(y - a) \geq 0, \forall \alpha \in \Phi_Q(G, L)\}.$$

Definition 4.2. For any geometric point $x \in \mathcal{X}$, we define a semicylinder $C_Q(x)$ in \mathfrak{a}_M^G by

$$C_Q(x) = \pi_M^{L, -1}(\text{Ec}_M^L(f_Q(x))) \cap \pi_{M, L}^{-1}(D_Q(H_Q(x))).$$

By definition, we get a partition

$$\mathfrak{a}_M^G = \pi^G(\text{Ec}_M(x)) \cup \bigcup_{\substack{Q \in \mathcal{F}(M) \\ Q \neq G}} C_Q(x),$$

for which the interior of any two parts does not intersect. The picture is similar to Figure 1. Hence for any $x \notin \mathcal{X}^\xi$, there exists a unique parabolic subgroup $Q \in \mathcal{F}(M)$ such that $\xi \in C_Q(x)$, as ξ is generic. In this case, $f_Q(x) \in \mathcal{X}^L$ is ξ^L -stable, where $\xi^L = \pi_M^L(\xi) \in \mathfrak{a}_M^L$. Let

$$X_Q = \{x \in \mathcal{X} \mid \xi \in C_Q(x)\}.$$

We have the decomposition of the affine Grassmannian

$$\mathcal{X} = \mathcal{X}^\xi \sqcup \bigsqcup_{\substack{Q \in \mathcal{F}(M) \\ Q \neq G}} X_Q. \quad (4.2)$$

For $Q \in \mathcal{P}(L)$, let Q^- be the parabolic subgroup opposite to Q with respect to L . Let $\Lambda_{L, Q}^\xi = (\pi_L^G)^{-1}(D_{Q^-}(\xi_L)) \cap \Lambda_L$, we have the disjoint partition

$$\Lambda_L = \bigsqcup_{Q \in \mathcal{P}(L)} \Lambda_{L, Q}^\xi.$$

For $\lambda \in \Lambda_L$, let $\mathcal{X}^{L, \lambda, \xi^L} = \mathcal{X}^{L, \xi^L} \cap \mathcal{X}^{L, \lambda}$. The stratum X_Q can be further decomposed into $N_Q((\epsilon))$ -orbits

$$X_Q = \bigsqcup_{\lambda \in \Lambda_{L, Q}^\xi} N_Q((\epsilon)) \mathcal{X}^{L, \lambda, \xi^L}.$$

Each orbit is locally closed in \mathcal{X} , and they are infinite-dimensional homogeneous affine fibrations on $\mathcal{X}^{L, \lambda, \xi^L}$ under the retraction f_Q . The foregoing discussions can be summarized as follows:

Theorem 4.2. *The affine Grassmannian can be decomposed as*

$$\mathcal{X} = \mathcal{X}^\xi \sqcup \bigsqcup_{\substack{Q=LN_Q \in \mathcal{F}(M) \\ Q \neq G}} \bigsqcup_{\lambda \in \Lambda_{L,Q}^\xi} N_Q((\epsilon)) \mathcal{X}^{L,\lambda,\xi^L}.$$

Each stratum $N_Q((\epsilon)) \mathcal{X}^{L,\lambda,\xi^L}$ is an infinite-dimensional homogeneous affine fibration over $\mathcal{X}^{L,\lambda,\xi^L}$.

Now that $\gamma \in \mathfrak{m}(F)$, we can restrict these constructions to \mathcal{X}_γ . Let $\mathcal{X}_\gamma^\xi = \mathcal{X}_\gamma \cap \mathcal{X}^\xi$; it is an open subscheme of \mathcal{X}_γ . As $T(F) \xrightarrow{H_M} X_*(M)$ is surjective, the connected components of \mathcal{X}_γ^ξ can be translated to each other by elements of $T(F)$. Moreover, for different choices of generic element $\xi, \xi' \in \mathfrak{a}_M^G$, the corresponding $\mathcal{X}_\gamma^\xi, \mathcal{X}_\gamma^{\xi'}$ can be translated to each other by elements of $T(F)$. Hence \mathcal{X}_γ^ξ does not depend on the choice of ξ .

The Harder–Narasimhan reduction restricts to

$$\mathcal{X}_\gamma = \mathcal{X}_\gamma^\xi \sqcup \bigsqcup_{\substack{Q=LN_Q \in \mathcal{F}(M) \\ Q \neq G}} \bigsqcup_{\lambda \in \Lambda_{L,Q}^\xi} \left(\mathcal{X}_\gamma \cap N_Q((\epsilon)) \mathcal{X}_\gamma^{L,\lambda,\xi^L} \right). \quad (4.3)$$

By Proposition 2.4, the retraction

$$f_Q : \mathcal{X}_\gamma \cap N_Q((\epsilon)) \mathcal{X}_\gamma^{L,\lambda,\xi^L} \rightarrow \mathcal{X}_\gamma^{L,\lambda,\xi^L}$$

is an iterated affine fibration over \mathbf{F}_q of relative dimension $\text{val}(\det(\text{ad}(\gamma) | \mathfrak{n}_Q(F)))$.

Coming back to the weighted orbital integrals, with the definition for general reductive algebraic groups as explained in Remark 2.1, Proposition 2.7 can be reformulated as follows:

Proposition 4.3. *Let $\xi \in \mathfrak{a}_M^G$ be a generic element. Then*

$$J_M^\xi(\gamma) = \text{vol}_{dt}(\Lambda^{H_M} \backslash T(F)_M^1)^{-1} \cdot |\Lambda^{H_M} \backslash ((\mathcal{X}^{G_{\text{der}}} \cap \mathcal{X}_\gamma^\xi)(\mathbf{F}_q))|.$$

In particular, let $\xi_0 \in \mathfrak{a}_{M_0}^G$ be a generic element. Then

$$J_{M_0}^{\xi_0}(\gamma) = \text{vol}_{dt}(T(F)^1)^{-1} \cdot |(\mathcal{X}^{G_{\text{der}}} \cap \mathcal{X}_\gamma^\xi)(\mathbf{F}_q)|.$$

Proof. When G is semisimple, the proposition is a reformulation of Proposition 2.7. The complexity arises when G has nontrivial connected center.

As T is totally ramified over F , with the exact sequence (2.2) we see that the morphism $T(F) \xrightarrow{\nu_G} \Lambda_G$ is surjective, hence $G(F) = T(F)G_{\text{der}}(F)$, and so

$$\begin{aligned} J_M^\xi(\gamma) &= \int_{T(F) \backslash G(F)} \mathbb{1}_{\mathfrak{g}(\mathcal{O})}(\text{Ad}(g)^{-1}\gamma) w_M^\xi(g) \frac{dg}{dt} \\ &= \int_{T_{G_{\text{der}}}(F) \backslash G_{\text{der}}(F)} \mathbb{1}_{\mathfrak{g}(\mathcal{O})}(\text{Ad}(g)^{-1}\gamma) w_{M_{G_{\text{der}}}}^\xi(g) \frac{dg}{dt}, \end{aligned}$$

with $T_{G_{\text{der}}} = T \cap G_{\text{der}}$ and $M_{G_{\text{der}}} = M \cap G_{\text{der}}$. Following calculations in Proposition 2.7, we get a result similar to what we claim, with Λ^{H_M} replaced by $\Lambda \cap G_{\text{der}}(F) \cap \ker(H_{M_{G_{\text{der}}}})$

and $T(F)_M^1$ replaced by $T_{G_{\text{der}}}(F)_{M_{G_{\text{der}}}}^1$. Noting that $\ker(H_M) = M_{\text{der}}(F) \cdot M(\mathcal{O})$ by [7, Lemma 6.1], we have

$$\Lambda \cap G_{\text{der}}(F) \cap \ker(H_{M_{G_{\text{der}}}}) = \Lambda \cap \ker(H_M) = \Lambda^{H_M}$$

and

$$T_{G_{\text{der}}}(F)_{M_{G_{\text{der}}}}^1 = T(F) \cap G_{\text{der}}(F) \cap \ker(H_{M_{G_{\text{der}}}}) = T(F) \cap \ker(H_M) = T(F)_M^1,$$

and the proposition is proved. \square

The volume factors have been calculated in equation (2.5).

4.2. Harder–Narasimhan reduction for the truncated affine Springer fibers

In contrast to the Arthur–Kottwitz reduction, the Harder–Narasimhan reduction does not work well on the truncated affine Springer fiber $\mathcal{X}_\gamma(\Pi)$. We need to cut it into two parts, the *tail* and the *main body*. The Harder–Narasimhan reduction works well on the main body, and we can handle the tail with the Arthur–Kottwitz reduction.

For $Q \in \mathcal{F}(M)$, $Q \neq G$, we define the positive (G, M) -orthogonal family $E_Q(\Pi)$, which as a polytope is the union of the translations $\pi^G(\Sigma_\gamma^{G, M} + \lambda)$, $\lambda \in \Lambda_M$, such that $\pi^G(\Sigma_\gamma^{G, M} + \lambda)^Q \subset \Pi^Q$. Let

$${}^t\mathcal{X}_\gamma(\Pi) = \bigcup_{\substack{Q \in \mathcal{F}(M) \\ Q \neq G}} \mathcal{X}_\gamma(E_Q(\Pi)), \quad {}^m\mathcal{X}_\gamma(\Pi) = \mathcal{X}_\gamma(\Pi) \setminus {}^t\mathcal{X}_\gamma(\Pi).$$

We call them the *tail* and the *main body* of $\mathcal{X}_\gamma(\Pi)$, and they are respectively closed and open subschemes of $\mathcal{X}_\gamma(\Pi)$. Figure 3 gives an example of $E_Q(\Pi)$ for the group $G = \text{GL}_3$ when $M = A$.


Before proceeding, we make precise the condition of Π being sufficiently regular. We would like it to satisfy the following conditions:

- (1) Π is Σ_γ -regular.
- (2) For all $P, Q \in \mathcal{F}(M)$, $E_P(\Pi) \cap E_Q(\Pi) = E_{P \cap Q}(\Pi)$.
- (3) The complement $\Pi \setminus \bigcup_{\substack{Q \in \mathcal{F}(M) \\ Q \neq G}} E_Q(\Pi)$ is a polytope associated to a positive (G, M) -orthogonal family; let Π_0 be a slight shrinking of it (the definition is similar to equation (3.1), with plus sign replaced by a minus). We require that Π_0 be sufficiently large: for all $Q = LN_Q \in \mathcal{F}(M)$, the face Π_0^Q contains the translations of Σ_γ^Q in \mathfrak{a}_M^L which have ξ^L as one of their vertices.

Remark 4.2. As Π_0 is convex, condition (3) implies that for any $\nu \in \Lambda_{L, Q}^\xi \cap \pi_L(\Pi_0)$, the intersection $\Pi_0 \cap \pi_{M_0, L}^{-1}(\nu)$ contains translations of Σ_γ^Q in the hyperplane $\pi_{M_0, L}^{-1}(\nu)$ which have ξ^L as one of their vertices. By the definition of ξ -stability, this implies

$$\mathcal{X}_\gamma^{L, \nu, \xi^L} \subset \mathcal{X}_\gamma^{L, \nu} \left(\Pi_0 \cap \pi_{M_0, L}^{-1}(\nu) \right), \quad \forall \nu \in \Lambda_{L, Q}^\xi \cap \pi_L(\Pi_0).$$

Actually, this is the reason to impose condition (3).

Figure 3. $E_Q(\Pi)$ for GL_3 when $M_0 = A$.

4.2.1. The main body. By definition, a Harder–Narasimhan stratum

$$N_Q((\epsilon)) \mathcal{X}_\gamma^{L, \nu, \xi^L} \cap \mathcal{X}_\gamma, \quad \nu \in \Lambda_{L, Q}^\xi,$$

intersects nontrivially with $\mathcal{X}_\gamma(\Pi)$ if and only if $\nu \in \Lambda_{L, Q}^\xi \cap \pi_L(\Pi)$. So, after restriction, the Harder–Narasimhan reduction becomes

$$\mathcal{X}_\gamma(\Pi) = \mathcal{X}_\gamma^\xi \sqcup \bigsqcup_{\substack{Q=LN_Q \in \mathcal{F}(M) \\ Q \neq G}} \bigsqcup_{\lambda \in \Lambda_{L, Q}^\xi \cap \pi_L(\Pi)} \left(\mathcal{X}_\gamma(\Pi) \cap N_Q((\epsilon)) \mathcal{X}_\gamma^{L, \lambda, \xi^L} \right).$$

The problem is that the retraction

$$f_Q : \mathcal{X}_\gamma(\Pi) \cap N_Q((\epsilon)) \mathcal{X}_\gamma^{L, \lambda, \xi^L} \rightarrow \mathcal{X}_\gamma^{L, \lambda, \xi^L}$$

is not necessarily an iterated affine fibration. This problem disappears on the main body ${}^m \mathcal{X}_\gamma(\Pi)$. We begin by analyzing the polytope $\mathrm{Ec}(x)$, $x \in \mathcal{X}_\gamma(\Pi)$.

Lemma 4.4. *For $x \in \mathcal{X}_\gamma(\Pi)$, suppose that*

$$\pi^G(\mathrm{Ec}_M(x)) \subset \bigcup_{\substack{Q \in \mathcal{F}(M) \\ Q \neq G}} E_Q(\Pi).$$

Then $\pi^G(\mathrm{Ec}_M(x)) \subset E_Q(\Pi)$ for some $Q \in \mathcal{F}(M)$, $Q \neq G$.

Proof. By Proposition 3.1, it is enough to prove the lemma for $x \in \mathcal{X}_\gamma^{\mathrm{reg}}$. In this case, the polytope $\pi^G(\mathrm{Ec}_M(x))$ is a translation of $\Sigma_\gamma^{G, M}$. As $\bigcup_{\substack{Q \in \mathcal{F}(M) \\ Q \neq G}} E_Q(\Pi)$ is the union of translations of $\Sigma_\gamma^{G, M}$ along the facets of Π , there must be a maximal parabolic subgroup

$Q \in \mathcal{F}(M)_{\max}$ such that $\pi^G(\text{Ec}_M^Q(x)) \subset \Pi^Q$. By definition, this means that $\pi^G(\text{Ec}_M(x)) \subset E_Q(\Pi)$. \square

Lemma 4.5. *Let $Q = LN \in \mathcal{F}(M)$, $\nu \in \Lambda_{L,Q}^\xi$. Suppose that*

$${}^m\mathcal{X}_\gamma(\Pi) \cap N((\epsilon))\mathcal{X}_\gamma^{L,\nu} \neq \emptyset.$$

Then $\nu \in \Lambda_{L,Q}^\xi \cap \pi_L(\Pi_0)$.

Proof. We only need to show that $\nu \in \pi_L(\Pi_0)$. Set $x \in {}^m\mathcal{X}_\gamma(\Pi) \cap N((\epsilon))\mathcal{X}_\gamma^{L,\nu}$. As $x \in {}^m\mathcal{X}_\gamma(\Pi)$, we have

$$\pi^G(\text{Ec}_M(x)) \not\subset E_{Q'}(\Pi), \quad \forall Q' \in \mathcal{F}(M), Q' \neq G.$$

By Lemma 4.4, this is equivalent to

$$\pi^G(\text{Ec}_M(x)) \not\subset \bigcup_{\substack{Q' \in \mathcal{F}(M) \\ Q' \neq G}} E_{Q'}(\Pi). \quad (4.4)$$

Suppose that $\nu \notin \pi_L(\Pi_0)$. Then $\nu \in \pi_L(E_{Q_0}(\Pi))$ for some $Q_0 \in \mathcal{F}(M), Q_0 \supset L$. As $\nu \in \Lambda_{L,Q}^\xi$, the parabolic subgroup Q_0 needs to satisfy $Q_0 \supset Q^-$. Now that $x \in N((\epsilon))\mathcal{X}_\gamma^{L,\nu}$ and $\text{Ec}_L(x)$ is a positive (G, L) -orthogonal family, we have

$$\alpha(H_{Q'}(x) - \nu) \geq 0, \quad \forall \alpha \in \Delta_{Q'}, Q' \in \mathcal{P}(L).$$

As $Q_0 \supset Q^-$, this implies that

$$H_{Q'}(x) \subset \pi_L(E_{Q_0}(\Pi)), \quad \forall Q' \in \mathcal{P}(L).$$

Hence $\pi_L(\text{Ec}_M(x)) \subset \pi_L(E_{Q_0}(\Pi))$, so

$$\text{Ec}_M(x) \subset \pi_L^{-1}(\pi_L(E_{Q_0}(\Pi))) \subset \bigcup_{\substack{Q' \in \mathcal{F}(M), Q' \neq G \\ Q' \cap Q_0 \neq \emptyset}} E_{Q'}(\Pi).$$

This is in contradiction to the relation (4.4), hence ν must lie in $\pi_L(\Pi_0)$. \square

Restricting the Harder–Narasimhan reduction (4.3) to the main part ${}^m\mathcal{X}_\gamma(\Pi)$, we have

$${}^m\mathcal{X}_\gamma(\Pi) = \mathcal{X}_\gamma^\xi \sqcup \bigsqcup_{\substack{Q = LN_Q \in \mathcal{F}(M) \\ Q \neq G}} \bigsqcup_{\lambda \in \Lambda_{L,Q}^\xi, \lambda \in \pi_L(\Pi_0)} \left({}^m\mathcal{X}_\gamma(\Pi) \cap N_Q((\epsilon))\mathcal{X}_\gamma^{L,\lambda,\xi^L} \right).$$

The retraction f_Q behaves much better on the stratum ${}^m\mathcal{X}_\gamma(\Pi) \cap N_Q((\epsilon))\mathcal{X}_\gamma^{L,\lambda,\xi^L}$:

Proposition 4.6. *Let $Q = LN_Q \in \mathcal{F}(M)$, $\nu \in \Lambda_{L,Q}^\xi \cap \pi_L(\Pi_0)$. We have*

$${}^m\mathcal{X}_\gamma(\Pi) \cap N_Q((\epsilon))\mathcal{X}_\gamma^{L,\nu,\xi^L} = \mathcal{X}_\gamma \cap N_Q((\epsilon))\mathcal{X}_\gamma^{L,\nu,\xi^L}.$$

Hence the retraction

$$f_Q : {}^m\mathcal{X}_\gamma(\Pi) \cap N_Q((\epsilon))\mathcal{X}_\gamma^{L,\nu,\xi^L} \rightarrow \mathcal{X}_\gamma^{L,\nu,\xi^L}$$

is an iterated affine fibration over \mathbf{F}_q .

Proof. Notice that the second assertion is the corollary of the first one, as follows from Proposition 2.4. It is thus enough to show the first one. In particular, it is enough to show

$$\mathcal{X}_\gamma \cap N_Q((\epsilon)) \mathcal{X}_\gamma^{L, \nu, \xi^L} \subset {}^m \mathcal{X}_\gamma(\Pi) \cap N_Q((\epsilon)) \mathcal{X}_\gamma^{L, \nu, \xi^L},$$

as the inclusion in the other direction is obvious.

Let $x \in \mathcal{X}_\gamma \cap N_Q((\epsilon)) \mathcal{X}_\gamma^{L, \nu, \xi^L}$. We claim that $\text{Ec}_M(x) \subset \Pi$. According to Remark 4.2, condition (3) of Π being sufficiently regular implies

$$\text{Ec}_M^L(f_Q(x)) \subset \Pi_0 \cap \pi_L^{-1}(\nu), \quad (4.5)$$

because $f_Q(x) \in \mathcal{X}_\gamma^{L, \nu, \xi^L}$. This implies that $\text{Ec}_M(x) \subset \Pi$ by Proposition 3.1, because of the inclusion

$$\bigcup_{\substack{\lambda \in \Lambda_M \\ \text{satisfying } (*)}} (\lambda + \Sigma_\gamma^{G, M}) \subset \Pi,$$

where the condition $(*)$ refers to

$$(\lambda + \Sigma_\gamma^{G, M}) \cap \Pi_0 \neq \emptyset.$$

The inclusion (4.5) also implies that

$$\text{Ec}_M(x) \not\subset \bigcup_{\substack{Q \in \mathcal{F}(M) \\ Q \neq G}} E_Q(\Pi).$$

So $x \in {}^m \mathcal{X}_\gamma(\Pi)$, and the proof is concluded. \square

We summarize the foregoing discussions in a proposition.

Proposition 4.7. *The main body has a decomposition*

$${}^m \mathcal{X}_\gamma(\Pi) = \mathcal{X}_\gamma^\xi \sqcup \bigsqcup_{\substack{Q=LN_Q \in \mathcal{F}(M) \\ Q \neq G}} \bigsqcup_{\lambda \in \Lambda_{L, Q}^\xi \cap \pi_L(\Pi_0)} \left({}^m \mathcal{X}_\gamma(\Pi) \cap N_Q((\epsilon)) \mathcal{X}_\gamma^{L, \lambda, \xi^L} \right),$$

and the retraction f_Q on each stratum

$$f_Q : {}^m \mathcal{X}_\gamma(\Pi) \cap N_Q((\epsilon)) \mathcal{X}_\gamma^{L, \nu, \xi^L} \rightarrow \mathcal{X}_\gamma^{L, \nu, \xi^L}$$

is an iterated affine fibration over \mathbf{F}_q of dimension $\text{val}(\det(\text{ad}(\gamma) | \mathfrak{n}_{Q, F}))$.

Of course, we can restrict the decomposition to each connected component ${}^m \mathcal{X}_\gamma^{\nu_0}(\Pi)$, $\nu_0 \in \Lambda_G$. Let $\Lambda_{L, Q}^{\nu_0, \xi} = \Lambda_{L, Q}^\xi \cap \Lambda_L^{\nu_0}$. The decomposition implies

$$\begin{aligned} & \left| \left(\Lambda^{H_M} \setminus {}^m \mathcal{X}_\gamma^{\nu_0}(\Pi) \right) (\mathbf{F}_q) \right| \\ &= \left| \left(\Lambda^{H_M} \setminus \mathcal{X}_\gamma^{\nu_0, \xi} \right) (\mathbf{F}_q) \right| + \sum_{\substack{Q=LN_Q \in \mathcal{F}(M) \\ Q \neq G}} \sum_{\lambda \in \Lambda_{L, Q}^{\nu_0, \xi} \cap \pi_L(\Pi_0)} q^{\frac{1}{2} \text{val}(\det(\text{ad}(\gamma) | \mathfrak{g}_F / \mathfrak{l}_F))} \\ & \quad \cdot \left| \left(\Lambda^{H_M} \setminus \mathcal{X}_\gamma^{L, \lambda, \xi^L} \right) (\mathbf{F}_q) \right| \end{aligned}$$

$$\begin{aligned}
&= |(\Lambda^{H_M} \setminus \mathcal{X}_\gamma^{0,\xi})(\mathbf{F}_q)| + \sum_{\substack{Q=LN_Q \in \mathcal{F}(M) \\ Q \neq G}} q^{\frac{1}{2} \text{val}(\det(\text{ad} \gamma|_{\mathfrak{g}_F/\mathfrak{l}_F}))} \cdot |(\Lambda^{H_M} \setminus \mathcal{X}_\gamma^{L,0,\xi^L})(\mathbf{F}_q)| \\
&\quad \cdot |\Lambda_{L,Q}^{\nu_0,\xi} \cap \pi_L(\Pi_0)| \\
&= |(\Lambda^{H_M} \setminus \mathcal{X}_\gamma^{0,\xi})(\mathbf{F}_q)| + \sum_{\substack{L \in \mathcal{L}(M) \\ L \neq G}} q^{\frac{1}{2} \text{val}(\det(\text{ad} \gamma|_{\mathfrak{g}_F/\mathfrak{l}_F}))} \cdot |(\Lambda^{H_M} \setminus \mathcal{X}_\gamma^{L,0,\xi^L})(\mathbf{F}_q)| \\
&\quad \cdot \sum_{Q \in \mathcal{P}(L)} |\Lambda_{L,Q}^{\nu_0,\xi} \cap \pi_L(\Pi_0)| \\
&= |(\Lambda^{H_M} \setminus \mathcal{X}_\gamma^{0,\xi})(\mathbf{F}_q)| + \sum_{\substack{L \in \mathcal{L}(M) \\ L \neq G}} q^{\frac{1}{2} \text{val}(\det(\text{ad} \gamma|_{\mathfrak{g}_F/\mathfrak{l}_F}))} \cdot |(\Lambda^{H_M} \setminus \mathcal{X}_\gamma^{L,0,\xi^L})(\mathbf{F}_q)| \\
&\quad \cdot |\Lambda_L^{\nu_0} \cap \pi_L(\Pi_0)|.
\end{aligned}$$

Here for the second equality we have used the fact that all the connected components of $\mathcal{X}_\gamma^{L,\xi^L}$ are isomorphic. Moreover, the last term in the equation counts the number of lattice points in a polytope; it can be calculated effectively with methods from toric geometry. The following theorem summarizes:

Theorem 4.8. *For any $\nu_0 \in \Lambda_G$, the number of rational points on the main body is*

$$\begin{aligned}
|(\Lambda^{H_M} \setminus {}^m \mathcal{X}_\gamma^{\nu_0}(\Pi))(\mathbf{F}_q)| &= |(\Lambda^{H_M} \setminus \mathcal{X}_\gamma^{0,\xi})(\mathbf{F}_q)| + \sum_{\substack{L \in \mathcal{L}(M) \\ L \neq G}} q^{\frac{1}{2} \text{val}(\det(\text{ad} \gamma|_{\mathfrak{g}_F/\mathfrak{l}_F}))} \\
&\quad \cdot |(\Lambda^{H_M} \setminus \mathcal{X}_\gamma^{L,0,\xi^L})(\mathbf{F}_q)| \cdot |\Lambda_L^{\nu_0} \cap \pi_L(\Pi_0)|.
\end{aligned}$$

4.2.2. The tail. As the polytope Π satisfies

$$E_P(\Pi) \cap E_Q(\Pi) = E_{P \cap Q}(\Pi), \quad \forall P, Q \in \mathcal{F}(M),$$

by the inclusion-exclusion principle we have

$$|(\Lambda^{H_M} \setminus {}^t \mathcal{X}_\gamma^{\nu_0}(\Pi))(\mathbf{F}_q)| = \sum_{\substack{Q \in \mathcal{F}(M) \\ Q \neq G}} (-1)^{\text{rk}(G) - \text{rk}(Q) - 1} |(\Lambda^{H_M} \setminus \mathcal{X}_\gamma^{\nu_0}(E_Q(\Pi)))(\mathbf{F}_q)|, \quad (4.6)$$

where the notation rk means the semisimple rank. Although the polytope $E_Q(\Pi)$ is not $\Sigma_\gamma^{G,M}$ -regular, we can use the general Arthur–Kottwitz reduction, as explained in Remark 3.2, repeatedly to decompose $\mathcal{X}_\gamma(E_Q(\Pi))$ into locally closed subschemes which are iterated affine fibrations over $F_\gamma^{L,M}$, $L \in \mathcal{L}(M)$. This gives a formula for $|(\Lambda^{H_M} \setminus \mathcal{X}_\gamma^{\nu_0}(E_Q(\Pi)))(\mathbf{F}_q)|$ in terms of the $|(\Lambda^{H_M} \setminus F_\gamma^{L,M})(\mathbf{F}_q)|$ s, which can be further reduced to counting points on the fundamental domains by Proposition 3.8 and Corollary 3.10. This process applies to a large family of truncated affine Springer fibers.

We introduce a family of operators on the set of all positive (G, M) -orthogonal families. Recall that $Q_0 = MU_0$ is the unique parabolic subgroup in $\mathcal{P}(M)$ which contains P_0 . For $L \in \mathcal{L}(M)$, let $Q_0^L = Q_0 \cap L$. For a positive (L, M) -orthogonal family, we say that

two faces of it are *conjugate* if their associated parabolic subgroups are conjugate to each other by the Weyl group W_L/W_M . In particular, the edges of the polytope are parametrized by minimal elements in $\mathcal{F}^L(M) \setminus \{M\}$. An edge is said to be of *type* $\alpha \in \Delta_{A_M}^{Q_0^L} := \Phi(L, A) \cap \Phi(U_0, A) \cap \Delta$ if it is conjugate to the edge having vertices $\lambda_{Q_0^L}, \lambda_{s_\alpha Q_0^L}$, where s_α is the simple reflection associated to α . Let $A_{M,\alpha}^{G,L}$ be the operator on the set of positive (G, M) -orthogonal families defined as follows: as a polytope, it increases by 1 the length of all the edges whose images in \mathfrak{a}_M^L under the projection π_M^L are of type α , and keeps the lengths of all the others invariant. To check that it actually sends a positive (G, M) -orthogonal family to another one, it suffices to verify for the faces of dimension 2, but this is clear. Then we set the vertex

$$\lambda_{Q_0} \left(A_{M,\alpha}^{G,L}(\Pi) \right) = \lambda_{Q_0}(\Pi) + \frac{1}{2} \pi_M(\varpi_\alpha^\vee),$$

to make $A_{M,\alpha}^{G,L}(\Pi)$ symmetric with respect to Π . Here ϖ_α^\vee is the fundamental coweight corresponding to α . By definition, we see that the operators $A_{M,\alpha}^{G,L}$ commute with each other. When $G = L$, we simplify the notation $A_{M,\alpha}^{G,G}$ to $A_{M,\alpha}^G$.

Given a tuple of nonnegative integers $\underline{n} = (n_\alpha)$, $\alpha \in \Delta_{A_M}^{Q_0^L}$, let

$$\Sigma_\gamma^n = \prod_{\alpha \in \Delta_{A_M}^{Q_0^L}} \left(A_{M,\alpha}^{G,L} \right)^{n_\alpha} (\Sigma_\gamma). \quad (4.7)$$

It is easy to see that the polytopes $E_Q(\Pi)$ can be made by iterating this process. For $\alpha \in \Delta_{A_M}^{Q_0^L}$, let 1_α be the tuple taking value 1 at α and 0 elsewhere. By Remark 3.2, the Arthur–Kottwitz reduction works for the complement $\mathcal{X}_\gamma \left(\Sigma_\gamma^{n+1_\alpha} \right) \setminus \mathcal{X}_\gamma \left(\Sigma_\gamma^n \right)$. The process is completely the same as explained in §3.3, so we do not repeat it here. The resulting strata are iterated affine fibrations over truncated affine Springer fibers of the form $\mathcal{X}_\gamma^{L,\nu'} \left(\Sigma_\gamma^{L,(\underline{n}')} \right)$, $L \in \mathcal{L}(M)$, $\nu' \in \Lambda_L$. Iterating this process, $\mathcal{X}_\gamma \left(\Sigma_\gamma^n \right)$ can be decomposed as a disjoint union of locally closed subschemes, which are iterated affine fibrations over $F_\gamma^{L,M,\mu}$, $L \in \mathcal{L}(M)$, $\mu \in \Lambda_L$. In particular, counting points on $\Lambda^{H_M} \setminus \mathcal{X}_\gamma \left(\Sigma_\gamma^n \right)$ can be reduced to counting points on $\Lambda^{H_M} \setminus F_\gamma^{L,M,\mu}$, which can be further reduced to counting points on the fundamental domains $F_\gamma^{L'}$, $L' \in \mathcal{L}(M_0)$, as we have explained in §3.2. This process applies to counting points on $\Lambda^{H_M} \setminus \mathcal{X}_\gamma^{\nu_0} (E_Q(\Pi))$. By equation (4.6), it gives an expression of $\left| (\Lambda^{H_M} \setminus {}^t \mathcal{X}_\gamma^{\nu_0}(\Pi)) (\mathbf{F}_q) \right|$ in terms of $\left| F_\gamma^L(\mathbf{F}_q) \right|$, $L \in \mathcal{L}(M_0)$.

4.3. Application to Arthur's weighted orbital integral

By Theorem 2.8 and Proposition 4.3, Arthur's weighted orbital integral $J_M(\gamma)$ calculates essentially $\left| (\Lambda^{H_M} \setminus \mathcal{X}_\gamma^{0,\xi}) (\mathbf{F}_q) \right|$, as $\mathcal{X}^{G_{\text{der}}} \cap \mathcal{X}_\gamma^\xi$ is the union of $|\Lambda_{G_{\text{der}}}|$ -copies of $\mathcal{X}_\gamma^{0,\xi}$. The two approaches in §3 and §4 to calculating $\left| (\Lambda^{H_M} \setminus \mathcal{X}_\gamma^0(\Pi)) (\mathbf{F}_q) \right|$ give us a recurrence relation involving $\left| (\Lambda^{H_M} \setminus F_\gamma^{L,M,\mu}) (\mathbf{F}_q) \right|$ and $\left| (\Lambda^{H_M} \setminus \mathcal{X}_\gamma^{L,0,\xi^L}) (\mathbf{F}_q) \right|$, for $L \in \mathcal{L}(M)$, $\mu \in \Lambda_{L_{\text{ad}}}$. If we are able to solve this recurrence relation, we will get an expression for

$|(\Lambda^{H_M} \backslash \mathcal{X}_\gamma^{0,\xi})(\mathbf{F}_q)|$ in terms of $|(\Lambda^{H_M} \backslash F_\gamma^{L,M,\mu})(\mathbf{F}_q)|$ s, which can be further reduced to counting points on fundamental domains, as explained in §3.2.

5. Calculations for the group GL_2

Let $G = \mathrm{GL}_2$ and let $\gamma \in \mathfrak{gl}_2(F)$ be a regular semisimple integral element. Assume that $\mathrm{char}(k) > 2$ and the splitting field of γ is totally ramified over F . The torus T is isomorphic either to $F^\times \times F^\times$ or to $\mathrm{Res}_{E/F} E^\times$, where E is a separable totally ramified field extension over F of degree 2. We call elements γ in these cases *split* and *anisotropic*, respectively.

5.1. Split elements

We can take T to be the maximal torus of G of the diagonal matrices and $\gamma \in \mathfrak{t}(\mathcal{O})$ a regular element. Let

$$n = \mathrm{val}(\alpha_{12}(\gamma)),$$

which we call the root valuation of γ . The dimension of the affine Springer fiber \mathcal{X}_γ is known to be

$$\dim(\mathcal{X}_\gamma) = n.$$

In the remainder of this section, we assume that $n \geq 1$, as the case $n = 0$ reduces to the group GL_1 . Recall that we have calculated F_γ in [11]. Let $X_*(T) \cong \mathbf{Z}^2$ be the usual identification, set $(n, 0) \in \mathbf{Z}^2$, and let

$$\mathrm{Sch}(n, 0) = \overline{K \begin{pmatrix} \epsilon^n & \\ & 1 \end{pmatrix} K / K}.$$

We have $F_\gamma \cong \mathrm{Sch}(n, 0)$, and its number of rational points is

$$|F_\gamma(\mathbf{F}_q)| = \sum_{i=0}^n q^i$$

by the Bruhat–Tits decomposition of $\mathrm{Sch}(n, 0)$. As $\Lambda_{\mathrm{PGL}_2} = \mathbf{Z}/2$, F_γ has only one variant F_γ^1 ; we can calculate its number of rational points to be

$$F_\gamma^1(\mathbf{F}_q) = \sum_{i=0}^{n-1} q^i.$$

Set $a \in \mathbf{N}$ and let Π be the positive (G, T) -orthogonal family defined by

$$\lambda_w(\Pi) = \lambda_w(\Sigma_\gamma) + w(a\alpha_{12}^\vee), \quad \forall w \in W.$$

Assume that $a \gg 0$. Then Π is sufficiently regular in the sense of §4.2. We can easily calculate

$$Q_\gamma^0(a) := |\mathcal{X}_\gamma^0(\Pi)(\mathbf{F}_q)| = \sum_{i=0}^n q^i + 2q^n a,$$

by the Arthur–Kottwitz reduction. We see that $Q_\gamma^0(a)$ is polynomial in a . By Theorem 4.8, we have

$$|{}^m\mathcal{X}_\gamma^0(\Pi)(\mathbf{F}_q)| = |\mathcal{X}_\gamma^{0,\xi}(\mathbf{F}_q)| + [2a - (n+1)]q^n.$$

The tail is the disjoint union of two fundamental domains, so its number of rational points is

$$|{}^t\mathcal{X}_\gamma^0(\Pi)(\mathbf{F}_q)| = 2 \sum_{i=0}^n q^i.$$

Because

$$|\mathcal{X}_\gamma^0(\Pi)(\mathbf{F}_q)| = |{}^m\mathcal{X}_\gamma^0(\Pi)(\mathbf{F}_q)| + |{}^t\mathcal{X}_\gamma^0(\Pi)(\mathbf{F}_q)|,$$

we get the equation

$$\sum_{i=0}^n q^i + 2aq^n = |\mathcal{X}_\gamma^{0,\xi}(\mathbf{F}_q)| + [2a - (n+1)]q^n + 2 \sum_{i=0}^n q^i.$$

Solving it, we get

$$|\mathcal{X}_\gamma^{0,\xi}(\mathbf{F}_q)| = nq^n - \sum_{i=0}^{n-1} q^i. \quad (5.1)$$

Now that $T(F)^1 = T(\mathcal{O}) = T(F)_1$ has volume 1, by Proposition 4.3 we have

$$J_T^\xi(\gamma) = |\mathcal{X}_\gamma^{0,\xi}(\mathbf{F}_q)| = nq^n - \sum_{i=0}^{n-1} q^i.$$

On the other hand, we can use equation (2.4) to easily calculate the orbital integral

$$I_\gamma^G = q^n.$$

Combined with Theorem 2.8, the foregoing calculations can be summarized as follows:

Theorem 5.1. *Let $\gamma \in \mathfrak{gl}_2(F)$ be a regular semisimple integral element of root valuation n . It has orbital integral $I_\gamma^G = q^n$. The number of rational points on $\mathcal{X}_\gamma^0(\Pi)$ is*

$$|\mathcal{X}_\gamma^0(\Pi)(\mathbf{F}_q)| = \sum_{i=0}^n q^i + 2aq^n,$$

and Arthur's weighted orbital integral $J_T(\gamma)$ equals

$$J_T(\gamma) = \text{vol}\left(\mathfrak{a}_{T_{SL_2}}^{SL_2} / X_*(T_{SL_2})\right) \cdot \left[nq^n - \sum_{i=0}^{n-1} q^i \right].$$

5.2. Anisotropic elements

In this case, $E = \mathbf{F}_q((\epsilon^{\frac{1}{2}}))$. Suppose that $\gamma = a + b\epsilon^{\frac{1}{2}}$ under the isomorphism $Z_{G(F)}(\gamma) \cong \text{Res}_{E/F} E^\times$, with $a, b \in \mathcal{O}$. Under the basis $\{\epsilon^{\frac{1}{2}}, 1\}$ of E over $\mathbf{F}_q((\epsilon))$, the element γ is of

the form

$$\gamma = \begin{bmatrix} a & b \\ b\epsilon & a \end{bmatrix}.$$

It is clear that the affine Springer fibers \mathcal{X}_γ and $\mathcal{X}_{-a+\gamma}$ are isomorphic, so we can assume that $a = 0$. Let $b = b_0\epsilon^n$, $b_0 \in \mathcal{O}^\times$. We can write

$$\gamma = \begin{bmatrix} & b_0\epsilon^n \\ b_0\epsilon^{n+1} & \end{bmatrix}. \quad (5.2)$$

Put in this form, it has been shown by Goresky, Kottwitz, and MacPherson [16] that \mathcal{X}_γ admits an affine paving which is induced by the standard Bruhat–Tits decomposition of the affine Grassmannian. More precisely, let I be the standard Iwahori subgroup – that is, it is the preimage of B_0 under the reduction $G(\mathcal{O}) \rightarrow G$. Then

$$\mathcal{X}_\gamma = \bigsqcup_{(a_1, a_2) \in \mathbf{Z}^2} \mathcal{X}_\gamma \cap I \begin{pmatrix} \epsilon^{a_1} & \\ & \epsilon^{a_2} \end{pmatrix} K/K,$$

and each intersection, denoted $S_{\mathbf{a}}$, is isomorphic to a standard affine space. We calculate that $S_{\mathbf{a}}$ is not empty if and only if

$$-(n+1) \leq a_1 - a_2 \leq n,$$

and that

$$\dim(S_{\mathbf{a}}) = \begin{cases} a_1 - a_2 & \text{if } a_1 \geq a_2, \\ a_2 - a_1 - 1 & \text{if } a_1 < a_2. \end{cases}$$

Notice that this is also the dimension of $I \begin{pmatrix} \epsilon^{a_1} & \\ & \epsilon^{a_2} \end{pmatrix} K/K$, so they must be the same.

Summarizing the foregoing calculations, and noting that $T(F)^1 = T(\mathcal{O}) = T(F)_1$ has volume 1, we get the following:

Theorem 5.2. *Let γ be matrix (5.2). For $(a_1, a_2) \in \mathbf{Z}^2$, we have*

$$\mathcal{X}_\gamma \cap I \begin{pmatrix} \epsilon^{a_1} & \\ & \epsilon^{a_2} \end{pmatrix} K/K = \begin{cases} I \begin{pmatrix} \epsilon^{a_1} & \\ & \epsilon^{a_2} \end{pmatrix} K/K & \text{if } -(n+1) \leq a_1 - a_2 \leq n, \\ \emptyset & \text{if not.} \end{cases}$$

As a corollary, we have

$$J_G(\gamma) = I_\gamma^G = |F_\gamma(\mathbf{F}_q)| = \sum_{i=0}^n q^i.$$

6. Calculations for GL_3 -split case

Let $G = \mathrm{GL}_3$ and let $\gamma \in \mathfrak{gl}_3(F)$ be a regular semisimple integral element. Assume that $\mathrm{char}(k) > 3$ and the splitting field of γ is totally ramified over F . The torus T is isomorphic to either $F^\times \times F^\times \times F^\times$ or $F^\times \times \mathrm{Res}_{E_2/F} E_2^\times$ or $\mathrm{Res}_{E_3/F} E_3^\times$, where E_2, E_3 are separable

totally ramified field extensions over F of degree 2 and 3, respectively. We call elements γ in these cases *split*, *mixed*, and *anisotropic* respectively. Notice that in all these cases, $T(F)^1 = T(\mathcal{O}) = T(F)_1$ has volume 1; hence by Proposition 4.3 we have

$$J_M^\xi(\gamma) = |\mathcal{X}_\gamma^{0,\xi}(\mathbf{F}_q)|,$$

and so

$$J_M(\gamma) = \text{vol}\left(\mathfrak{a}_{M_{\text{SL}_3}}^{\text{SL}_3} / X_*(M_{\text{SL}_3})\right) \cdot |\mathcal{X}_\gamma^{0,\xi}(\mathbf{F}_q)|$$

by Theorem 2.8 and Remark 2.3.

In this section, we restrict ourselves to the split case. After conjugation, we take T to be the maximal torus of G of the diagonal matrices. Then $M_0 = T$ and the other proper Levi subgroups in $\mathcal{L}(T)$ can be parametrized as follows: for a nonempty subset $I \subsetneq \{1, 2, 3\}$, let P_I be the parabolic subgroup of G which stabilizes the flag

$$\bigoplus_{i=1}^3 \mathbf{F}_q e_i \supsetneq \bigoplus_{i \notin I} \mathbf{F}_q e_i \supsetneq \emptyset.$$

Let $P_I = M_I N_I$ be the standard Levi factorization. We have $M_I \cong \text{GL}_2 \times \text{GL}_1$. As $M_I = M_{I^c}$, with I^c the complement of I , it is enough to calculate $J_T(\gamma)$ and $J_{M_{\{i\}}}(\gamma)$, $i = 1, 2, 3$.

Let $\gamma \in \mathfrak{t}(\mathcal{O})$ be a regular element. As we show in the appendix of [9], up to conjugation by the Weyl group we can suppose that

$$\text{val}(\alpha_{12}(\gamma)) \leq \text{val}(\alpha_{23}(\gamma)), \quad \text{val}(\alpha_{13}(\gamma)) = \text{val}(\alpha_{12}(\gamma)).$$

In this case, γ is said to be *in minimal form*, and we call

$$(n_1, n_2) = (\text{val}(\alpha_{12}(\gamma)), \text{val}(\alpha_{23}(\gamma)))$$

the *root valuation* of γ . The dimension of the affine Springer fiber \mathcal{X}_γ is known to be

$$\dim(\mathcal{X}_\gamma) = 2n_1 + n_2.$$

In the remainder of the section, we assume that $n_1 \geq 1$, as the case $n_1 = 0$ reduces to the group GL_2 . Recall that we have calculated the Poincaré polynomial of F_γ in [11].

Proposition 6.1. *The fundamental domain F_γ admits an affine paving. Its Poincaré polynomial, which depends only on the root valuation (n_1, n_2) , is*

$$\begin{aligned} P_{(n_1, n_2)}(t) &= \sum_{i=1}^{n_1} i(t^{4i-2} + t^{4i-4}) + \sum_{i=2n_1}^{n_1+n_2-1} (2n_1+1)t^{2i} \\ &\quad + \sum_{i=n_1+n_2}^{2n_1+n_2-1} 4(2n_1+n_2-i)t^{2i} + t^{4n_1+2n_2}. \end{aligned}$$

In particular, $|F_\gamma(\mathbf{F}_q)| = P_{(n_1, n_2)}(q^{1/2})$.

6.1. Calculation of $J_T(\gamma)$

Set $(a_1, a_2) \in \mathbf{N}^2$, and let Π be the positive (G, T) -orthogonal family defined by

$$\lambda_w(\Pi) = \lambda_w(\Sigma_\gamma) + w \sum_{i=1}^2 a_i \alpha_i^\vee, \quad \forall w \in W.$$

Assume that Π is sufficiently regular in the sense of §4.2, which means that $a_1, a_2 \gg 0$ and

$$2a_1 - a_2 > 0, \quad 2a_2 - a_1 > 0.$$

We will calculate

$$Q_\gamma^0(a_1, a_2) := |\mathcal{X}_\gamma^0(\Pi)(\mathbf{F}_q)|$$

following the two approaches that we have explained, and draw conclusions on Arthur's weighted orbital integral.

6.1.1. Counting points by Arthur–Kottwitz reduction. We will work out each term in Corollary 3.7. Look at the summands indexed by the Borel subgroups. Each stratum contributes $q^{2n_1+n_2}$, so it remains to count the number of lattice points

$$\sum_{B \in \mathcal{P}(T)} |\Lambda_T^0 \cap R_B \cap \Pi| = 6 |\Lambda_T^0 \cap R_{B_0} \cap \Pi|,$$

where the equality is due to the symmetry of Π with respect to Σ_γ . We identify

$$\Lambda_T^0 \cong \{(m_1, m_2, m_3) \in \mathbf{Z}^3 \mid m_1 + m_2 + m_3 = 0\}$$

in the usual way. Let $\bar{\mathfrak{a}}_{B_0}^G = \{a \in \mathfrak{a}_T^G \mid \alpha_1(a) \geq 0, \alpha_2(a) \geq 0\}$, and let

$$R_0 = \{a \in \bar{\mathfrak{a}}_{B_0}^G \mid \varpi_1(a) \leq a_1 - 1, \varpi_2(a) \leq a_2 - 1\}.$$

Up to a suitable translation, we have

$$|\Lambda_T^0 \cap R_{B_0} \cap \Pi| = |\Lambda_T^0 \cap R_0|.$$

We can express it as the difference of two lattice point-counting problems. Let

$$\begin{aligned} R_1 &= \{a \in \bar{\mathfrak{a}}_{B_0}^G \mid \varpi_1(a) \leq a_1 - 1, \varpi_2(a) \leq 2(a_1 - 1)\}, \\ R_2 &= \{a \in \bar{\mathfrak{a}}_{B_0}^G \mid \varpi_1(a) \leq a_1 - 1, \varpi_2(a) \geq a_2\}. \end{aligned}$$

Then we have

$$|\Lambda_T^0 \cap R_0| = |\Lambda_T^0 \cap R_1| - |\Lambda_T^0 \cap R_2|.$$

We count $|\Lambda_T^0 \cap R_1|$ as follows:

$$\begin{aligned} |\Lambda_T^0 \cap R_1| &= \sum_{n=0}^{+\infty} |R_1 \cap \{\mu \in \Lambda_T^0 \mid \varpi_1(\mu) = n\}| \\ &= \sum_{i=1}^{\lfloor \frac{a_1}{2} \rfloor} [(3i-2) + (3i-1)] + \frac{1 - (-1)^{a_1}}{2} \left(1 + \frac{3(a_1-1)}{2}\right) \\ &= 3 \left\lfloor \frac{a_1}{2} \right\rfloor^2 + \frac{1}{4} (1 - (-1)^{a_1}) (3a_1 - 1), \end{aligned}$$

where $\lfloor x \rfloor$ means the largest integer that is less than or equal to x . Similarly, we have

$$\begin{aligned} |\Lambda_T^0 \cap R_2| &= \sum_{n=2(a_1-1)}^{-\infty} |R_2 \cap \{\mu \in \Lambda_T^0 \mid \varpi_2(\mu) = n\}| \\ &= \sum_{i=1}^{\lfloor \frac{2a_1-a_2-1}{2} \rfloor} (i+i) + \frac{1 + (-1)^{2a_1-a_2}}{2} \cdot \frac{2a_1-a_2}{2} \\ &= \left\lfloor \frac{2a_1-a_2-1}{2} \right\rfloor \left(\left\lfloor \frac{2a_1-a_2-1}{2} \right\rfloor + 1 \right) + \frac{1}{4} (1 + (-1)^{2a_1-a_2}) (2a_1 - a_2). \end{aligned}$$

In summary, the summands in Corollary 3.7 indexed by the Borel subgroups contribute

$$\begin{aligned} 6q^{2n_1+n_2} &\left[3 \left\lfloor \frac{a_1}{2} \right\rfloor^2 - \left\lfloor \frac{2a_1-a_2-1}{2} \right\rfloor \left(\left\lfloor \frac{2a_1-a_2-1}{2} \right\rfloor + 1 \right) \right. \\ &\quad \left. + \frac{1}{4} (1 - (-1)^{a_1}) (3a_1 - 1) - \frac{1}{4} (1 + (-1)^{2a_1-a_2}) (2a_1 - a_2) \right]. \quad (6.1) \end{aligned}$$

Now we calculate the contributions of the summands indexed by the maximal parabolic subgroups. They are parametrized at the beginning of the section by nonempty subsets $I \subsetneq \{1, 2, 3\}$. For $\mu \in \Lambda_{M_I^{\text{ad}}} \cong \mathbf{Z}/2$, let $q_I^\mu = |F_\gamma^{M_I, \nu}(\mathbf{F}_q)|$, for any $\nu \in \Lambda_{M_I}$ which projects to $\mu \in \Lambda_{M_I^{\text{ad}}}$. Let α_I be the unique element in $\Phi_{B_0 \cap M_I}(M_I, T)$. A simple calculation with the affine Springer fibers for the group GL_2 shows that

$$q_I^{(0)} = \sum_{i=0}^{\text{val}(\alpha_I(\gamma))} q^i, \quad q_I^{(1)} = \sum_{i=0}^{\text{val}(\alpha_I(\gamma))-1} q^i.$$

For $I = \{i\}$, $i = 1, 2, 3$, it is easy to see that

$$\begin{aligned} |\Lambda_{M_I}^0 \cap \pi_{M_I}(R_{P_I}) \cap \pi_{M_I}(\Pi) \cap c_M^{-1}(0)| &= \left\lfloor \frac{a_1}{2} \right\rfloor, \\ |\Lambda_{M_I}^0 \cap \pi_{M_I}(R_{P_I}) \cap \pi_{M_I}(\Pi) \cap c_M^{-1}(1)| &= \left\lfloor \frac{a_1+1}{2} \right\rfloor. \end{aligned}$$

The summands indexed by P_I in Corollary 3.7 with $|I| = 1$ contribute in total

$$\left\lfloor \frac{a_1}{2} \right\rfloor \left(q^{2n_1} \sum_{i=0}^{n_2} q^i + 2q^{n_1+n_2} \sum_{i=0}^{n_1} q^i \right) + \left\lfloor \frac{a_1+1}{2} \right\rfloor \left(q^{2n_1} \sum_{i=0}^{n_2-1} q^i + 2q^{n_1+n_2} \sum_{i=0}^{n_1-1} q^i \right). \quad (6.2)$$

Similarly, the summands indexed by P_I with $|I| = 2$ contribute in total

$$\left\lfloor \frac{a_2}{2} \right\rfloor \left(q^{2n_1} \sum_{i=0}^{n_2} q^i + 2q^{n_1+n_2} \sum_{i=0}^{n_1} q^i \right) + \left\lfloor \frac{a_2+1}{2} \right\rfloor \left(q^{2n_1} \sum_{i=0}^{n_2-1} q^i + 2q^{n_1+n_2} \sum_{i=0}^{n_1-1} q^i \right). \quad (6.3)$$

Summing up the contributions from formulas (6.1), (6.2), and (6.3), we obtain the following:

Proposition 6.2. *We have*

$$\begin{aligned} Q_\gamma^0(a_1, a_2) = & \sum_{i=1}^{n_1} i (q^{2i-1} + q^{2i-2}) + \sum_{i=2n_1}^{n_1+n_2-1} (2n_1+1)q^i \\ & + \sum_{i=n_1+n_2}^{2n_1+n_2-1} 4(2n_1+n_2-i)q^i + q^{2n_1+n_2} \\ & + 6q^{2n_1+n_2} \left[3 \left\lfloor \frac{a_1}{2} \right\rfloor^2 - \left\lfloor \frac{2a_1-a_2-1}{2} \right\rfloor \left(\left\lfloor \frac{2a_1-a_2-1}{2} \right\rfloor + 1 \right) \right. \\ & \left. + \frac{1}{4}(1 - (-1)^{a_1})(3a_1-1) - \frac{1}{4}(1 + (-1)^{2a_1-a_2})(2a_1-a_2) \right] \\ & + \left(\left\lfloor \frac{a_1}{2} \right\rfloor + \left\lfloor \frac{a_2}{2} \right\rfloor \right) \left(q^{2n_1} \sum_{i=0}^{n_2} q^i + 2q^{n_1+n_2} \sum_{i=0}^{n_1} q^i \right) \\ & + \left(\left\lfloor \frac{a_1+1}{2} \right\rfloor + \left\lfloor \frac{a_2+1}{2} \right\rfloor \right) \left(q^{2n_1} \sum_{i=0}^{n_2-1} q^i + 2q^{n_1+n_2} \sum_{i=0}^{n_1-1} q^i \right). \end{aligned}$$

In particular, it depends quasi-polynomially on (a_1, a_2) .

6.1.2. Counting points by Harder–Narasimhan reduction. We begin by counting points on the main body; we need to work out each term in Theorem 4.8. For $L = T$, it is easy to see that $|\mathcal{X}_\gamma^{T,0,\xi^T}(\mathbf{F}_q)| = 1$, and we need to count the number of lattice points in $\Lambda_T^0 \cap \Pi_0$. Notice that for this we can shrink Π_0 to the convex hull of $\Lambda_T^0 \cap \Pi_0$. We conserve the notation Π_0 for the shrunken polytope. In [11, §6], we calculate $\text{Ec}(x_0)$ for a particular choice of regular point $x_0 \in \mathcal{X}_\gamma^{\text{reg}}$; we can adapt the result to our current setting. Let $(\sigma_1\sigma_2\sigma_3)$ be the permutation sending (123) to $(\sigma_1\sigma_2\sigma_3)$. The vertices of Σ_γ are

$$\begin{aligned} \lambda_{123}(\Sigma_\gamma) &= (0, 0, 0), & \lambda_{321}(\Sigma_\gamma) &= (-2n_1, n_1 - n_2, n_1 + n_2), \\ \lambda_{213}(\Sigma_\gamma) &= (-n_1, n_1, 0), & \lambda_{312}(\Sigma_\gamma) &= (-n_1, -n_2, n_1 + n_2), \\ \lambda_{132}(\Sigma_\gamma) &= (0, -n_2, n_2), & \lambda_{231}(\Sigma_\gamma) &= (-2n_1, n_1, n_1). \end{aligned}$$




Figure 4. Completing the hexagon to a triangle.

The vertices of Π_0 can be calculated to be

$$\begin{aligned}\lambda_{123}(\Pi_0) &= (a_1 - 2n_1 - 1, a_2 - a_1 + n_1 - n_2, -a_2 + n_1 + n_2 + 1), \\ \lambda_{321}(\Pi_0) &= (-a_2 + 1, a_2 - a_1, a_1 - 1), \\ \lambda_{213}(\Pi_0) &= (a_2 - a_1 - n_1, a_1 - n_2 - 1, -a_2 + n_1 + n_2 + 1), \\ \lambda_{312}(\Pi_0) &= (a_2 - a_1 - n_1, -a_2 + n_1 + 1, a_1 - 1), \\ \lambda_{132}(\Pi_0) &= (a_1 - 2n_1 - 1, -a_2 + n_1 + 1, a_2 - a_1 + n_1), \\ \lambda_{231}(\Pi_0) &= (-a_2 + 1, a_1 - n_2 - 1, a_2 - a_1 + n_2).\end{aligned}$$

We will count the lattice points in Π_0 indirectly. We complete the hexagon Π_0 to a triangle T_0 , whose vertices are

$$\begin{aligned}\lambda_{123}(T_0) &= \lambda_{132}(T_0) = (2a_2 - 2n_1 - n_2 - 2, -a_2 + n_1 + 1, -a_2 + n_1 + n_2 + 1), \\ \lambda_{321}(T_0) &= \lambda_{312}(T_0) = (-a_2 + 1, -a_2 + n_1 + 1, 2a_2 - n_1 - 2), \\ \lambda_{213}(T_0) &= \lambda_{231}(T_0) = (-a_2 + 1, 2a_2 - 2 - n_1 - n_2, -a_2 + n_1 + n_2 + 1).\end{aligned}$$

Let $T_1 \cup T_2 \cup T_3$ be the complement of Π_0 in T_0 , as shown in Figure 4. Notice that the T_i s do not contain their common boundary with Π_0 , so

$$|\Lambda_T^0 \cap \Pi_0| = |\Lambda_T^0 \cap T_0| - \sum_{i=1}^3 |\Lambda_T^0 \cap T_i|.$$

The right-hand side is much easier to calculate.

The length of the edges of T_0 is $3a_2 - 3 - 2n_1 - n_2$, so

$$\begin{aligned}|\Lambda_T^0 \cap T_0| &= \sum_{i=1}^{3a_2 - 3 - 2n_1 - n_2 + 1} i \\ &= \frac{1}{2}(3a_2 - 2 - 2n_1 - n_2)(3a_2 - 1 - 2n_1 - n_2).\end{aligned}$$

The length of the edges of T_1 is $2a_2 - a_1 - n_1 - 1$. As we do not count the lattice points on the common boundary of T_1 and Π_0 , we have

$$\begin{aligned} |\Lambda_T^0 \cap T_1| &= \sum_{i=1}^{2a_2 - a_1 - n_1 - 1} i \\ &= \frac{1}{2}(2a_2 - a_1 - n_1 - 1)(2a_2 - a_1 - n_1). \end{aligned}$$

Similarly, the length of the edges of T_2 is $2a_2 - a_1 - n_2 - 1$ and we have

$$|\Lambda_T^0 \cap T_2| = \frac{1}{2}(2a_2 - a_1 - n_2 - 1)(2a_2 - a_1 - n_2).$$

The triangle T_3 is of the same size as T_1 , so

$$|\Lambda_T^0 \cap T_3| = \frac{1}{2}(2a_2 - a_1 - n_1 - 1)(2a_2 - a_1 - n_1).$$

Finally,

$$\begin{aligned} |\Lambda_T^0 \cap \Pi_0| &= |\Lambda_T^0 \cap T_0| - \sum_{i=1}^3 |\Lambda_T^0 \cap T_i| \\ &= \frac{1}{2}(3a_2 - 2 - 2n_1 - n_2)(3a_2 - 1 - 2n_1 - n_2) \\ &\quad - (2a_2 - a_1 - n_1 - 1)(2a_2 - a_1 - n_1) \\ &\quad - \frac{1}{2}(2a_2 - a_1 - n_2 - 1)(2a_2 - a_1 - n_2). \end{aligned} \tag{6.4}$$

We go on to calculate $|\Lambda_L^0 \cap \pi_L(\Pi_0)|$ for the other Levi subgroups $L \in \mathcal{L}(T)$. Let d_L be the distance between the facets Π_0^Q and $\Pi_0^{Q^-}$, where $\mathcal{P}(L) = \{Q, Q^-\}$. It is easy to see that

$$|\Lambda_L^0 \cap \pi_L(\Pi_0)| = d_L + 1.$$

The set $\mathcal{L}(T) \setminus \{T, G\}$ consists of three elements, Levi factors $M_{\{i\}}$ of the parabolic subgroups $P_{\{i\}}$, $i = 1, 2, 3$. Using the explicit expression of the vertices of Π_0 , we can calculate

$$|\Lambda_{M_{\{1\}}}^0 \cap \pi_{M_{\{1\}}}(\Pi_0)| = d_{M_{\{1\}}} + 1 = a_1 + a_2 - 2n_1 - 1, \tag{6.5}$$

$$|\Lambda_{M_{\{2\}}}^0 \cap \pi_{M_{\{2\}}}(\Pi_0)| = d_{M_{\{2\}}} + 1 = a_1 + a_2 - n_1 - n_2 - 1, \tag{6.6}$$

$$|\Lambda_{M_{\{3\}}}^0 \cap \pi_{M_{\{3\}}}(\Pi_0)| = d_{M_{\{3\}}} + 1 = a_1 + a_2 - n_1 - n_2 - 1. \tag{6.7}$$

Now that $|\mathcal{X}_\gamma^{L,0,\xi^L}(\mathbf{F}_q)|$ has been calculated in equation (5.1), we can insert equations (6.4)–(6.7) into the equation in Theorem 4.8 to get the following:

Proposition 6.3. *The number of rational points on the main body is*

$$\begin{aligned} |\mathcal{X}_\gamma^0(\Pi)(\mathbf{F}_q)| &= |\mathcal{X}_\gamma^{0,\xi}(\mathbf{F}_q)| + q^{2n_1+n_2} \left[\frac{1}{2}(3a_2-2-2n_1-n_2)(3a_2-1-2n_1-n_2) \right. \\ &\quad - (2a_2-a_1-n_1-1)(2a_2-a_1-n_1) \\ &\quad \left. - \frac{1}{2}(2a_2-a_1-n_2-1)(2a_2-a_1-n_2) \right] \\ &\quad + q^{2n_1}(a_1+a_2-2n_1-1) \left(n_2 q^{n_2} - \sum_{i=0}^{n_2-1} q^i \right) \\ &\quad + 2q^{n_1+n_2}(a_1+a_2-n_1-n_2-1) \left(n_1 q^{n_1} - \sum_{i=0}^{n_1-1} q^i \right). \end{aligned}$$

Now we proceed to counting points on the tail. To begin with, we write down the vertices of Π :

$$\begin{aligned} \lambda_{123}(\Pi) &= (a_1, a_2 - a_1, -a_2), \\ \lambda_{321}(\Pi) &= (-a_2 - 2n_1, a_2 - a_1 + n_1 - n_2, a_1 + n_1 + n_2), \\ \lambda_{213}(\Pi) &= (a_2 - a_1 - n_1, a_1 + n_1, -a_2), \\ \lambda_{312}(\Pi) &= (a_2 - a_1 - n_1, -a_2 - n_2, a_1 + n_1 + n_2), \\ \lambda_{132}(\Pi) &= (a_1, -a_2 - n_2, a_2 - a_1 + n_2), \\ \lambda_{231}(\Pi) &= (-a_2 - 2n_1, a_1 + n_1, a_2 - a_1 + n_1). \end{aligned}$$

For nonempty subsets $I \subsetneq \{1, 2, 3\}$, we simplify the notation $E_{P_I}(\Pi)$ to $E_I(\Pi)$. Using the coordinates of vertices of Π , we can calculate the lengths of the edges of Π and find the following expression for $E_I(\Pi)$: when $|I| = 1$, we have

$$E_I(\Pi) = (A_{\alpha_I}^{G, M_I})^{2a_2 - a_1}(\Sigma_\gamma). \quad (6.8)$$

When $|I| = 2$, we have

$$E_I(\Pi) = (A_{\alpha_I}^{G, M_I})^{2a_1 - a_2}(\Sigma_\gamma). \quad (6.9)$$

As explained before, we can use the Arthur–Kottwitz reduction inductively to count the number of rational points on $\mathcal{X}_\gamma^0(E_I(\Pi))$. We give the details for $I = \{3\}$; the others can be calculated in the same way.

Applying Arthur–Kottwitz reduction to pass from $(A_{\alpha_{\{3\}}}^{G, M_{\{3\}}})^a(\Sigma_\gamma)$ to $(A_{\alpha_{\{3\}}}^{G, M_{\{3\}}})^{a+1}(\Sigma_\gamma)$, the picture is similar to Figure 2. We obtain

$$\begin{aligned}
\left| \mathcal{X}_\gamma^0 \left(\left(A_{\alpha_{\{3\}}}^{G, M_{\{3\}}} \right)^{a+1} (\Sigma_\gamma) \right) (\mathbf{F}_q) \right| &= \left| \mathcal{X}_\gamma^0 \left(\left(A_{\alpha_{\{3\}}}^{G, M_{\{3\}}} \right)^a (\Sigma_\gamma) \right) (\mathbf{F}_q) \right| + q^{2n_1} \left| F_\gamma^{M_{\{1\}}, 1} (\mathbf{F}_q) \right| \\
&\quad + q^{n_1+n_2} \left| F_\gamma^{M_{\{13\}}, 1} (\mathbf{F}_q) \right| + q^{2n_1+n_2} \\
&= \left| \mathcal{X}_\gamma^0 \left(\left(A_{\alpha_{\{3\}}}^{G, M_{\{3\}}} \right)^a (\Sigma_\gamma) \right) (\mathbf{F}_q) \right| + q^{2n_1} \sum_{i=0}^{n_2-1} q^i \\
&\quad + q^{n_1+n_2} \sum_{i=0}^{n_1-1} q^i + q^{2n_1+n_2}.
\end{aligned}$$

From this relation and equation (6.8), we deduce that

$$\begin{aligned}
\left| \mathcal{X}_\gamma^0 (E_{\{3\}}(\Pi)) (\mathbf{F}_q) \right| &= |F_\gamma (\mathbf{F}_q)| + (2a_2 - a_1) \left(q^{2n_1} \sum_{i=0}^{n_2-1} q^i \right. \\
&\quad \left. + q^{n_1+n_2} \sum_{i=0}^{n_1-1} q^i + q^{2n_1+n_2} \right). \tag{6.10}
\end{aligned}$$

Similarly, we have

$$\left| \mathcal{X}_\gamma^0 (E_{\{1\}}(\Pi)) (\mathbf{F}_q) \right| = |F_\gamma (\mathbf{F}_q)| + (2a_2 - a_1) \left(2q^{n_1+n_2} \sum_{i=0}^{n_1-1} q^i + q^{2n_1+n_2} \right), \tag{6.11}$$

$$\begin{aligned}
\left| \mathcal{X}_\gamma^0 (E_{\{2\}}(\Pi)) (\mathbf{F}_q) \right| &= |F_\gamma (\mathbf{F}_q)| + (2a_2 - a_1) \left(q^{2n_1} \sum_{i=0}^{n_2-1} q^i \right. \\
&\quad \left. + q^{n_1+n_2} \sum_{i=0}^{n_1-1} q^i + q^{2n_1+n_2} \right), \tag{6.12}
\end{aligned}$$

$$\begin{aligned}
\left| \mathcal{X}_\gamma^0 (E_{\{12\}}(\Pi)) (\mathbf{F}_q) \right| &= |F_\gamma (\mathbf{F}_q)| + (2a_1 - a_2) \left(q^{2n_1} \sum_{i=0}^{n_2-1} q^i \right. \\
&\quad \left. + q^{n_1+n_2} \sum_{i=0}^{n_1-1} q^i + q^{2n_1+n_2} \right), \tag{6.13}
\end{aligned}$$

$$\left| \mathcal{X}_\gamma^0 (E_{\{23\}}(\Pi)) (\mathbf{F}_q) \right| = |F_\gamma (\mathbf{F}_q)| + (2a_1 - a_2) \left(2q^{n_1+n_2} \sum_{i=0}^{n_1-1} q^i + q^{2n_1+n_2} \right), \tag{6.14}$$

$$\begin{aligned}
\left| \mathcal{X}_\gamma^0 (E_{\{13\}}(\Pi)) (\mathbf{F}_q) \right| &= |F_\gamma (\mathbf{F}_q)| + (2a_1 - a_2) \left(q^{2n_1} \sum_{i=0}^{n_2-1} q^i \right. \\
&\quad \left. + q^{n_1+n_2} \sum_{i=0}^{n_1-1} q^i + q^{2n_1+n_2} \right). \tag{6.15}
\end{aligned}$$

Inserting equations (6.10)–(6.15) into equation (4.6), we get the following:

Proposition 6.4. *The number of rational points on the tail equals*

$$|{}^t\mathcal{X}_\gamma^0(\Pi)(\mathbf{F}_q)| = (a_1 + a_2) \left(2q^{2n_1} \sum_{i=0}^{n_2-1} q^i + 4q^{n_1+n_2} \sum_{i=0}^{n_1-1} q^i + 3q^{2n_1+n_2} \right).$$

The sum of results in Propositions 6.3 and 6.4 gives us another expression for $Q_\gamma^0(a_1, a_2)$:

Corollary 6.5.

$$\begin{aligned} Q_\gamma^0(a_1, a_2) = & |\mathcal{X}_\gamma^{0,\xi}(\mathbf{F}_q)| + q^{2n_1+n_2} \left[\frac{1}{2}(3a_2 - 2 - 2n_1 - n_2)(3a_2 - 1 - 2n_1 - n_2) \right. \\ & - (2a_2 - a_1 - n_1 - 1)(2a_2 - a_1 - n_1) \\ & \left. - \frac{1}{2}(2a_2 - a_1 - n_2 - 1)(2a_2 - a_1 - n_2) \right] \\ & + q^{2n_1}(a_1 + a_2 - 2n_1 - 1) \left(n_2 q^{n_2} - \sum_{i=0}^{n_2-1} q^i \right) \\ & + 2q^{n_1+n_2}(a_1 + a_2 - n_1 - n_2 - 1) \left(n_1 q^{n_1} - \sum_{i=0}^{n_1-1} q^i \right) \\ & + (a_1 + a_2) \left(2q^{2n_1} \sum_{i=0}^{n_2-1} q^i + 4q^{n_1+n_2} \sum_{i=0}^{n_1-1} q^i + 3q^{2n_1+n_2} \right). \end{aligned}$$

In particular, this shows that $Q_\gamma^0(a_1, a_2)$ depends polynomially on $(a_1, a_2) \in \mathbf{N}^2$. As a corollary, the expression for $Q_\gamma^0(a_1, a_2)$ in Proposition 6.2 is also a polynomial in (a_1, a_2) , although it does not seem to be so.

6.1.3. Arthur's weighted orbital integral. Now we can compare the two expressions in Proposition 6.2 and Corollary 6.5 for $Q_\gamma^0(a_1, a_2)$. Look at their constant terms $Q_\gamma^0(0, 0)$. As $J_T^\xi(\gamma) = |\mathcal{X}_\gamma^{0,\xi}(\mathbf{F}_q)|$ in this case, we obtain the following:

Theorem 6.6. *Chaudouard and Laumon's weighted orbital integral for γ equals*

$$J_T^\xi(\gamma) = \sum_{i=1}^{n_1} i (q^{2i-1} + q^{2i-2}) + \sum_{i=n_1+n_2}^{2n_1+n_2-1} (4n_1 + 2n_2 - 4i - 3)q^i + (n_1^2 + 2n_1n_2)q^{2n_1+n_2}.$$

By Theorem 2.8 and Remark 2.3, we get Arthur's weighted orbital integral as well. For the orbital integral I_γ^G , as T is split, we can calculate it easily by equation (2.4):

$$I_\gamma^G = q^{2n_1+n_2}.$$

6.2. Calculation of $J_{M_1}(\gamma)$

We parametrize the Levi groups as before, with the further simplification $M_i := M_{\{i\}}$. Let $\gamma = \text{diag}(\gamma_1, \gamma_2, \gamma_3)$ and $\gamma' = \text{diag}(\gamma_1, \gamma_3, \gamma_2)$; notice that

$$J_{M_2}(\gamma) = J_{M_3}(\gamma').$$

Moreover, \mathcal{X}_γ and $\mathcal{X}_{\gamma'}$ have the same geometry, as they have the same root valuation (indeed, they have the same affine paving), which implies that

$$J_{M_2}(\gamma) = J_{M_3}(\gamma') = J_{M_3}(\gamma).$$

Hence it is enough to calculate $J_{M_1}(\gamma)$ and $J_{M_3}(\gamma)$. Notice that M_1 corresponds to the root α_2 and M_3 to the root α_1 .

As usual, we identify $X_*(T) \cong \mathbf{Z}^3$ and \mathfrak{a}_T^G with the hyperplane $x_1 + x_2 + x_3 = 0$ of $\mathfrak{a}_T = X_*(T) \otimes \mathbf{R} \cong \mathbf{R}^3$. The subspace $\mathfrak{a}_T^{M_1} \subset \mathfrak{a}_T^G$ becomes the line $\{(0, x, -x)\}$, and the subspace $\mathfrak{a}_{M_1}^G \subset \mathfrak{a}_T^G$ becomes $\{(-x, x/2, x/2)\}$. The lattice $\Lambda_{M_1}^0$ is identified with \mathbf{Z} by the mapping

$$\Lambda_{M_1}^0 \rightarrow \mathbf{Z} : (-(a+b), a, b) \mapsto a+b.$$

Its inclusion in $\mathfrak{a}_{M_1}^G$ is described by the mapping

$$\Lambda_{M_1}^0 \rightarrow \mathfrak{a}_{M_1}^G : (-(a+b), a, b) \mapsto (-(a+b), (a+b)/2, (a+b)/2).$$

We identify $\mathfrak{a}_{M_1}^G$ with \mathbf{R} by identifying $(-x, x/2, x/2)$ with x ; the inclusion $\Lambda_{M_1}^0 \subset \mathfrak{a}_{M_1}^G$ becomes the natural embedding $\mathbf{Z} \subset \mathbf{R}$. On the other hand, the discrete free abelian group $\Lambda \cong X_*(T)$ is naturally identified with \mathbf{Z}^3 , and the morphism $H_{M_1} : \Lambda \rightarrow \mathfrak{a}_{M_1}$ can be calculated to be

$$H_{M_1}(a_1, a_2, a_3) = \left(a_1, \frac{a_2 + a_3}{2}, \frac{a_2 + a_3}{2} \right).$$

Hence $\Lambda^{H_{M_1}}$ is freely generated by the element $\text{diag}(1, \epsilon, \epsilon^{-1})$.

According to Proposition 3.1, we can take Σ_γ^{G, M_1} to be the interval $[0, 2n_1]$ in $\mathfrak{a}_{M_1}^G \cong \mathbf{R}$. For $N \in \mathbf{N}$, $N \gg 0$, let Π_N be the interval $[-N, 2n_1 + N]$, regarded as a (G, M_1) -orthogonal family in $\mathfrak{a}_{M_1}^G$. We are going to calculate $(\Lambda^{H_{M_1}} \setminus \mathcal{X}_\gamma(\Pi_N))(\mathbf{F}_q)$ by the two approaches we have described.

In the Arthur–Kottwitz approach, we need to calculate

$$|(\Lambda^{H_{M_1}} \setminus F_\gamma^{G, M_1})(\mathbf{F}_q)| \quad \text{and} \quad |(\Lambda^{H_{M_1}} \setminus F_\gamma^{M_1, M_1})(\mathbf{F}_q)|.$$

Combining Proposition 3.8 and Corollary 3.10, we get

$$|(\Lambda^{H_{M_1}} \setminus F_\gamma^{G, M_1})(\mathbf{F}_q)| = |F_{\gamma, \mu}^{G, M_1}(\mathbf{F}_q)| = q^{2n_1 + n_2} + 2q^{n_1 + n_2} (1 + q + \cdots + q^{n_1 - 1}).$$

For the second calculation, since $F_\gamma^{M_1, M_1} = \mathcal{X}_\gamma^{M_1, (0)}$ and $\Lambda^{H_{M_1}} = \Lambda^{M_1}$, we have

$$|(\Lambda^{H_{M_1}} \setminus F_\gamma^{M_1, M_1})(\mathbf{F}_q)| = |(\Lambda^{M_1} \setminus \mathcal{X}_\gamma^{M_1, 0})(\mathbf{F}_q)| = q^{n_2}.$$

The reduction process is illustrated by a figure similar to [Figure 2](#). By [Corollary 3.7](#), we have the following:

Proposition 6.7.

$$\begin{aligned} |(\Lambda^{H_{M_1}} \setminus \mathcal{X}_\gamma^0(\Pi_N))(\mathbf{F}_q)| &= |(\Lambda^{H_{M_1}} \setminus F_\gamma^{G, M_1})(\mathbf{F}_q)| + 2Nq^{2n_1} |(\Lambda^{H_{M_1}} \setminus F_\gamma^{M_1, M_1})(\mathbf{F}_q)| \\ &= q^{2n_1+n_2} + 2q^{n_1+n_2} (1+q+\cdots+q^{n_1-1}) + 2Nq^{2n_1+n_2} \\ &= (2N+1)q^{2n_1+n_2} + 2q^{n_1+n_2} (1+q+\cdots+q^{n_1-1}). \end{aligned}$$

In the Harder–Narasimhan approach, we begin with counting points on the tail. By construction,

$$\begin{aligned} |(\Lambda^{H_{M_1}} \setminus {}^t\mathcal{X}_\gamma^0(\Pi_N))(\mathbf{F}_q)| &= 2 |(\Lambda^{H_{M_1}} \setminus F_\gamma^{G, M_1})(\mathbf{F}_q)| \\ &= 2 [q^{2n_1+n_2} + 2q^{n_1+n_2} (1+q+\cdots+q^{n_1-1})]. \end{aligned} \quad (6.16)$$

Then we calculate

$$|(\Lambda^{H_{M_1}} \setminus \mathcal{X}_\gamma^{M_1, 0, \xi^{M_1}})(\mathbf{F}_q)| = |(\Lambda^{H_{M_1}} \setminus \mathcal{X}_\gamma^{M_1, 0})(\mathbf{F}_q)| = q^{n_2}.$$

By [Theorem 4.8](#), this implies

$$\begin{aligned} |(\Lambda^{H_{M_1}} \setminus {}^m\mathcal{X}_\gamma^0(\Pi_N))(\mathbf{F}_q)| &= |(\Lambda^{H_{M_1}} \setminus \mathcal{X}_\gamma^{\xi, 0})(\mathbf{F}_q)| + [2N - (2n_1 + 1)] \cdot q^{2n_1} \\ &\quad \cdot |(\Lambda^{H_{M_1}} \setminus \mathcal{X}_\gamma^{M_1, 0, \xi^{M_1}})(\mathbf{F}_q)| \\ &= |(\Lambda^{H_{M_1}} \setminus \mathcal{X}_\gamma^{\xi, 0})(\mathbf{F}_q)| + [2N - (2n_1 + 1)] \cdot q^{2n_1+n_2}. \end{aligned} \quad (6.17)$$

Combining equations [\(6.16\)](#) and [\(6.17\)](#), we obtain the following:

Proposition 6.8.

$$\begin{aligned} |(\Lambda^{H_{M_1}} \setminus \mathcal{X}_\gamma^0(\Pi_N))(\mathbf{F}_q)| &= |(\Lambda^{H_{M_1}} \setminus \mathcal{X}_\gamma^{\xi, 0})(\mathbf{F}_q)| + [2N - (2n_1 + 1)] \cdot q^{2n_1+n_2} \\ &\quad + 2 [q^{2n_1+n_2} + 2q^{n_1+n_2} (1+q+\cdots+q^{n_1-1})]. \end{aligned}$$

Comparing [Propositions 6.7](#) and [6.8](#), we get the following:

Proposition 6.9.

$$|(\Lambda^{H_{M_1}} \setminus \mathcal{X}_\gamma^{\xi, 0})(\mathbf{F}_q)| = 2n_1q^{2n_1+n_2} - 2q^{n_1+n_2} (1+q+\cdots+q^{n_1-1}).$$

It remains to calculate the volume factor $\text{vol}_{dt}(\Lambda^{H_{M_1}} \setminus T(F)_{M_1}^1)$. By [equation \(2.5\)](#), it equals 1 because $S = T$ and the morphism $H_{M_1} : X_*(T) \rightarrow X_*(M)$ is surjective. The foregoing calculations can be summarized as follows:

Theorem 6.10. *We have*

$$J_{M_1}^\xi(\gamma) = |(\Lambda^{H_{M_1}} \setminus \mathcal{X}_\gamma^{\xi, 0})(\mathbf{F}_q)| = 2n_1q^{2n_1+n_2} - 2q^{n_1+n_2} (1+q+\cdots+q^{n_1-1}).$$

6.3. Calculation of $J_{M_3}(\gamma)$

We make identifications as before. The subspace $\mathfrak{a}_T^{M_3} \subset \mathfrak{a}_T^G$ becomes the line $\{(x, -x, 0)\}$, and the subspace $\mathfrak{a}_{M_3}^G \subset \mathfrak{a}_T^G$ becomes $\{(x/2, x/2, -x)\}$; they are identified with \mathbf{R} as before. The lattice $\Lambda_{M_3}^0$ is identified with \mathbf{Z} by the mapping

$$\Lambda_{M_3}^0 \rightarrow \mathbf{Z} : (a, b, -(a+b)) \mapsto a+b.$$

The inclusion $\Lambda_{M_3}^0 \subset \mathfrak{a}_{M_3}^G$ becomes again the natural embedding $\mathbf{Z} \subset \mathbf{R}$. Similarly, the group $\Lambda^{H_{M_3}}$ is freely generated by the element $\text{diag}(\epsilon, \epsilon^{-1}, 1)$.

By Proposition 3.1, we can take Σ_γ^{G, M_3} to be the interval $[0, n_1 + n_2]$ in $\mathfrak{a}_{M_3}^G \cong \mathbf{R}$. For $N \in \mathbf{N}$, $N \gg 0$, let Π_N be the interval $[-N, n_1 + n_2 + N]$, regarded as a (G, M_3) -orthogonal family in $\mathfrak{a}_{M_3}^G$. We calculate $(\Lambda^{H_{M_3}} \setminus \mathcal{X}_\gamma(\Pi_N))(\mathbf{F}_q)$ in two ways as before.

Using similar calculations as before, we get

$$|(\Lambda^{H_{M_3}} \setminus F_\gamma^{M_3, M_3})(\mathbf{F}_q)| = |(\Lambda^{M_3} \setminus \mathcal{X}_\gamma^{M_3, 0})(\mathbf{F}_q)| = q^{n_1}$$

and

$$\begin{aligned} |(\Lambda^{H_{M_3}} \setminus F_\gamma^{G, M_3})(\mathbf{F}_q)| &= |F_{\gamma, \mu}^{G, M_3}(\mathbf{F}_q)| = q^{2n_1+n_2} + q^{n_1+n_2} (1+q+\cdots+q^{n_1-1}) \\ &\quad + q^{2n_1} (1+q+\cdots+q^{n_2-1}). \end{aligned}$$

With Arthur–Kottwitz reduction, we obtain the following:

Proposition 6.11.

$$\begin{aligned} |(\Lambda^{H_{M_3}} \setminus \mathcal{X}_\gamma^0(\Pi_N))(\mathbf{F}_q)| &= |(\Lambda^{H_{M_3}} \setminus F_\gamma^{G, M_3})(\mathbf{F}_q)| + 2Nq^{n_1+n_2} |(\Lambda^{H_{M_3}} \setminus F_\gamma^{M_3, M_3})(\mathbf{F}_q)| \\ &= q^{2n_1+n_2} + q^{n_1+n_2} (1+q+\cdots+q^{n_1-1}) \\ &\quad + q^{2n_1} (1+q+\cdots+q^{n_2-1}) + 2Nq^{2n_1+n_2} \\ &= (2N+1)q^{2n_1+n_2} + q^{n_1+n_2} (1+q+\cdots+q^{n_1-1}) \\ &\quad + q^{2n_1} (1+q+\cdots+q^{n_2-1}). \end{aligned}$$

For the Harder–Narasimhan reduction, we count the points on the tail

$$\begin{aligned} |(\Lambda^{H_{M_3}} \setminus {}^t\mathcal{X}_\gamma^0(\Pi_N))(\mathbf{F}_q)| &= 2 |(\Lambda^{H_{M_3}} \setminus F_\gamma^{G, M_3})(\mathbf{F}_q)| \\ &= 2 [q^{2n_1+n_2} + q^{n_1+n_2} (1+q+\cdots+q^{n_1-1}) \\ &\quad + q^{2n_1} (1+q+\cdots+q^{n_2-1})] \end{aligned}$$

and the ξ -stable points

$$|(\Lambda^{H_{M_3}} \setminus \mathcal{X}_\gamma^{M_3, 0, \xi^{M_3}})(\mathbf{F}_q)| = |(\Lambda^{H_{M_3}} \setminus \mathcal{X}_\gamma^{M_3, 0})(\mathbf{F}_q)| = q^{n_1}.$$

Hence the points in the main body are

$$\begin{aligned} |(\Lambda^{H_{M_3}} \setminus {}^m\mathcal{X}_\gamma^0(\Pi_N))(\mathbf{F}_q)| &= |(\Lambda^{H_{M_3}} \setminus \mathcal{X}_\gamma^{\xi, 0})(\mathbf{F}_q)| + [2N - (n_1 + n_2 + 1)] \cdot q^{n_1+n_2} \\ &\quad \cdot |(\Lambda^{H_{M_3}} \setminus \mathcal{X}_\gamma^{M_3, 0, \xi^{M_3}})(\mathbf{F}_q)| \\ &= |(\Lambda^{H_{M_3}} \setminus \mathcal{X}_\gamma^{\xi, 0})(\mathbf{F}_q)| + [2N - (n_1 + n_2 + 1)] \cdot q^{2n_1+n_2}. \end{aligned}$$

Combining them gives us the following:

Proposition 6.12.

$$\begin{aligned} |(\Lambda^{H_{M_3}} \setminus \mathcal{X}_\gamma^0(\Pi_N))(\mathbf{F}_q)| &= |(\Lambda^{H_{M_3}} \setminus {}^m \mathcal{X}_\gamma^0(\Pi_N))(\mathbf{F}_q)| + |(\Lambda^{H_{M_3}} \setminus {}^t \mathcal{X}_\gamma^0(\Pi_N))(\mathbf{F}_q)| \\ &= |(\Lambda^{H_{M_3}} \setminus \mathcal{X}_\gamma^{\xi,0})(\mathbf{F}_q)| + [2N - (n_1 + n_2 + 1)] \cdot q^{2n_1+n_2} \\ &\quad + 2[q^{2n_1+n_2} + q^{n_1+n_2}(1+q+\cdots+q^{n_1-1}) \\ &\quad + q^{2n_1}(1+q+\cdots+q^{n_2-1})]. \end{aligned}$$

Comparing Propositions 6.11 and 6.12, we get the following:

Proposition 6.13.

$$\begin{aligned} |(\Lambda^{H_{M_3}} \setminus \mathcal{X}_\gamma^{\xi,0})(\mathbf{F}_q)| &= (n_1 + n_2)q^{2n_1+n_2} - q^{n_1+n_2}(1+q+\cdots+q^{n_1-1}) \\ &\quad - q^{2n_1}(1+q+\cdots+q^{n_2-1}). \end{aligned}$$

As before, the volume factor $\text{vol}_{dt}(\Lambda^{H_{M_3}} \setminus T(F)_{M_3}^1)$ equals 1, and so we have the following:

Theorem 6.14. *We have*

$$\begin{aligned} J_{M_3}^\xi(\gamma) &= |(\Lambda^{H_{M_3}} \setminus \mathcal{X}_\gamma^{\xi,0})(\mathbf{F}_q)| = (n_1 + n_2)q^{2n_1+n_2} - q^{n_1+n_2}(1+q+\cdots+q^{n_1-1}) \\ &\quad - q^{2n_1}(1+q+\cdots+q^{n_2-1}). \end{aligned}$$

7. Calculations for GL_3 -mixed case

Let $G = \text{GL}_3$ and let $\gamma \in \mathfrak{gl}_3(F)$ be a regular semisimple integral element. Assume that $T \cong F^\times \times \text{Res}_{E_2/F} E_2^\times$, with E_2 a separable totally ramified field extension over F of degree 2. As before, we can reduce to the case in which γ is a matrix of the form

$$\gamma = \begin{bmatrix} a & & \\ & b_0 \epsilon^n & \\ & b_0 \epsilon^{n+1} & \end{bmatrix}, \quad (7.1)$$

with $a \in \mathcal{O}, b_0 \in \mathcal{O}^\times$. Let $m = \text{val}(a)$.

Let P be the parabolic subgroup $P = B_0 \cup B_0 s_2 B_0$, and let $P = MN$ be the standard Levi decomposition. We identify $X_*(A) \cong \mathbf{Z}^3$ in the usual way. This gives us an identification $\Lambda_M \cong \mathbf{Z}^2$ and hence $\Lambda_M \otimes \mathbf{R} \cong \mathbf{R}^2$. We also identify \mathfrak{a}_M^G with the line $x+y=0$ in \mathbf{R}^2 , which can be further identified with \mathbf{R} by taking the coordinate x . Under these identifications, the moment polytope Σ_γ of the fundamental domain F_γ can be taken to be the closed interval

$$\Sigma_\gamma = [-n(\gamma, P, P^-), 0] \subset \mathbf{R} \cong \mathfrak{a}_M^G.$$

To simplify the notation, we abbreviate $n(\gamma, P, P^-)$ to n_γ . We have

$$n_\gamma = \min\{2m, 2n+1\}.$$




Figure 5. The (G, A) -orthogonal family Π_{n_1, n_2} and its extension to a triangle.

By definition, we can take

$$F_\gamma = \mathcal{X}_\gamma \cap \mathcal{X}^{n+1}(\Sigma_\gamma).$$

This can be refined a little bit. For $(n_1, n_2) \in \mathbf{N}^2$, let Π_{n_1, n_2} be the positive (G, A) -orthogonal family as indicated in Figure 5 (excluding the dashed part).

Consider the positive (G, A) -orthogonal families $\Pi_{n_\gamma, n+1}$. For $i \in \mathbf{Z}$, $-n_\gamma \leq i \leq 0$, let

$$\Pi_{n_\gamma, n+1}^i = \Pi_{n_\gamma, n+1} \cap \pi_M^{-1}(i),$$

where $i \in \mathbf{Z}$ is considered as an element of \mathfrak{a}_M^G by the identification $\mathbf{R} \cong \mathfrak{a}_M^G$. By Theorem 5.2, we have

$$\mathcal{X}_\gamma^{M, (i, n+1-i)} \subset \mathcal{X}^{M, (i, n+1-i)} \left(\Pi_{n_\gamma, n+1}^i \right), \quad \text{for } i = -n_\gamma, \dots, 0.$$

This implies that

$$F_\gamma = \mathcal{X}_\gamma \cap \mathcal{X}^{n+1}(\Sigma_\gamma) = \mathcal{X}_\gamma \cap \mathcal{X}^{n+1}(\Pi_{n_\gamma, n+1}).$$

It is possible, but quite hard, to construct an affine paving of F_γ and count the number of rational points with it. Instead, we take an indirect route. Let $\Delta_{n_\gamma, n+1}$ be the completion of $\Pi_{n_\gamma, n+1}$ into a triangle, as indicated in Figure 5. We can count the number of rational points on $\mathcal{X}_\gamma^{n+1}(\Delta_{n_\gamma, n+1})$ quite easily, using the affine pavings in [9, Proposition 3.6]. The complementary $\mathcal{X}_\gamma^{n+1}(\Delta_{n_\gamma, n+1}) \setminus F_\gamma$ can be treated by the Arthur–Kottwitz reduction. Taking their difference, we find $|F_\gamma(\mathbf{F}_q)|$.

We calculate the number of rational points on $\mathcal{X}_\gamma^{n+1}(\Delta_{n_\gamma, n+1})$. For $N \in \mathbf{N}$, let

$$I_N = \text{Ad}(\text{diag}(\epsilon^N, 1, 1)) I.$$

According to [9, Proposition 3.6], when $N \gg 0$, we have an affine paving

$$\mathcal{X}_\gamma^{n+1}(\Delta_{n_\gamma, n+1}) = \bigcup_{\epsilon^{\mathbf{a}} \in \mathcal{X}_\gamma^{n+1}(\Delta_{n_\gamma, n+1})^A} \mathcal{X}_\gamma^{n+1}(\Delta_{n_\gamma, n+1}) \cap I_N \epsilon^{\mathbf{a}} K / K.$$




Figure 6. Counting points in the nonequivalued case.




Figure 7. Counting points in the equivalued case.

The dimension of the affine paving can be calculated using [9, Lemma 3.1], together with Theorem 5.2. When $m \leq n$ – that is, γ is not equivalued – the dimension of the paving is

$$\min\{a_2, m\} + \min\{a_3, m\} + \begin{cases} a_2 - a_3 & \text{if } 0 \leq a_2 - a_3 \leq n, \\ a_3 - a_2 - 1 & \text{if } 1 \leq a_3 - a_2 \leq n + 1. \end{cases}$$

Otherwise, the intersection is empty. When $m \geq n + 1$ – that is, γ is equivalued – the dimension of the paving is

$$\min\{a_2, n\} + \min\{a_3, n + 1\} + \begin{cases} a_2 - a_3 & \text{if } 0 \leq a_2 - a_3 \leq n, \\ a_3 - a_2 - 1 & \text{if } 1 \leq a_3 - a_2 \leq n + 1. \end{cases}$$

Otherwise, the intersection is empty. We summarize the situation in Figures 6 and 7. The triangle is cut into four parts by the two long red lines, and the dimension of the fibration f_P restricted to the affine pavings in different parts are given by different formulas. The two dashed lines bound the region where $\mathcal{X}_\gamma^M \cap I_N \epsilon^a K/K$ is nonempty.

Proposition 7.1. Let γ be a matrix in the form of equation (7.1). When $\text{val}(a) = m \leq n$, we have

$$\begin{aligned} |\mathcal{X}_\gamma^{n+1}(\Delta_{2m,n+1})(\mathbf{F}_q)| &= \sum_{j=0}^{2m-1} \left(\left\lfloor \frac{j}{2} \right\rfloor + 1 \right) q^j + (2m+n+1)q^{2m} \\ &\quad + \sum_{j=2m+1}^{m+n} (4m+n+1-j)q^j + \sum_{j=m+n+1}^{2m+n} (3(2m+n-j)+1)q^j \\ &\quad + q^{2m} \sum_{j=0}^n q^j. \end{aligned}$$

In the summation, we use the convention that a summand is empty if its subscript is greater than its superscript.

Proof. Summing along the dotted blue lines in the four regions of Figure 6, we get

$$\begin{aligned} |\mathcal{X}_\gamma^{n+1}(\Delta_{2m,n+1})(\mathbf{F}_q)| &= \sum_{i=0}^m q^i (1+q+\cdots+q^i) + \sum_{i=m+1}^{2m} q^i (1+q+\cdots+q^{2m-i}) \\ &\quad + \sum_{i=0}^{m-1} q^{i+m} (q^{m+1-i} + \cdots + q^n) + \sum_{i=0}^{m-1} q^{i+m} (q^{m-i} + \cdots + q^n) \\ &\quad + q^{2m} \sum_{i=2m+1}^{2m+n} (1+q+\cdots+q^{i-2m}) + q^{2m} \sum_{i=0}^n q^i. \end{aligned}$$

After rearranging the summand, we get the proposition. \square

Proposition 7.2. Let γ be a matrix in the form of equation (7.1). When $\text{val}(a) = m > n$, we have

$$|\mathcal{X}_\gamma^{n+1}(\Delta_{2n+1,n+1})(\mathbf{F}_q)| = \sum_{j=0}^{2n} \left(\left\lfloor \frac{j}{2} \right\rfloor + 1 \right) q^j + \sum_{j=2n+1}^{3n+1} (3(3n-j+1)+1)q^j + q^{2n+1} \sum_{j=0}^n q^j.$$

Proof. Summing along the dotted blue lines in the four regions of Figure 7, we get

$$\begin{aligned} |\mathcal{X}_\gamma^{n+1}(\Delta_{2n+1,n+1})(\mathbf{F}_q)| &= \sum_{i=0}^n q^i (1+q+\cdots+q^i) + \sum_{i=n+1}^{2n+1} q^i (1+q+\cdots+q^{2n+1-i}) \\ &\quad + \sum_{i=1}^n q^{i+n} (q^{n+1-i} + \cdots + q^n) + \sum_{i=1}^{n-1} q^{i+n+1} (q^{n+1-i} + \cdots + q^n) \\ &\quad + q^{2n+1} \sum_{i=2n+2}^{3n+1} (1+q+\cdots+q^{i-(2n+1)}) + q^{2n+1} \sum_{i=0}^n q^i. \end{aligned}$$

After rearranging the summand, we get the proposition. \square

Now we calculate the number of rational points on the complement $\mathcal{X}_\gamma^{n+1}(\Delta_{n_\gamma, n+1}) \setminus F_\gamma$. For $i \in \mathbf{Z}$, $1 \leq i \leq n+1$, let

$$\Delta_{n_\gamma, n+1}^i = \Delta_{n_\gamma, n+1} \cap \pi_M^{-1}(i),$$

where $i \in \mathbf{Z}$ is considered as an element of \mathfrak{a}_M^G by the identification $\mathbf{R} \cong \mathfrak{a}_M^G$.

Proposition 7.3. *Let γ be a matrix in the form of equation (7.1). We have*

$$\mathcal{X}_\gamma^{n+1}(\Delta_{n_\gamma, n+1}) \setminus F_\gamma = \bigcup_{i=1}^{n+1} f_P^{-1} \left(\mathcal{X}_\gamma^{M, (i, n+1-i)} \left(\Delta_{n_\gamma, n+1}^i \right) \right) \cap \mathcal{X}_\gamma,$$

where $(i, n+1-i) \in \mathbf{Z}^2$ is regarded as an element in Λ_M by the identification $\mathbf{Z}^2 \cong \Lambda_M$. Its number of rational points over \mathbf{F}_q equals

$$q^{n_\gamma} \sum_{j=0}^n (1 + q + \cdots + q^j).$$

Proof. Observe that the second assertion is a direct consequence of the first one by Proposition 2.4. It is thus enough to show the first assertion.

Set $x \in \mathcal{X}_\gamma^{n+1}(\Delta_{n_\gamma, n+1})$, and notice that it does not belong to F_γ if and only if

$$H_P(x) \in [1, n+1] \subset \mathbf{R} \cong \mathfrak{a}_M^G, \quad (7.2)$$

because $H_{P^-}(x) \leq H_P(x)$. This implies that

$$\mathcal{X}_\gamma^{n+1}(\Delta_{n_\gamma, n+1}) \setminus F_\gamma = \bigcup_{i=1}^{n+1} f_P^{-1} \left(\mathcal{X}_\gamma^{M, (i, n+1-i)} \left(\Delta_{n_\gamma, n+1}^i \right) \right) \cap \mathcal{X}_\gamma(\Delta_{n_\gamma, n+1}).$$

To finish the proof, we only need to show that

$$f_P^{-1} \left(\mathcal{X}_\gamma^{M, (i, n+1-i)} \left(\Delta_{n_\gamma, n+1}^i \right) \right) \cap \mathcal{X}_\gamma(\Delta_{n_\gamma, n+1}) = f_P^{-1} \left(\mathcal{X}_\gamma^{M, (i, n+1-i)} \left(\Delta_{n_\gamma, n+1}^i \right) \right) \cap \mathcal{X}_\gamma,$$

for $i = 1, \dots, n+1$. The inclusion ' \subset ' is obvious; we only need to show its inverse. For any point $x \in f_P^{-1} \left(\mathcal{X}_\gamma^{M, (i, n+1-i)} \left(\Delta_{n_\gamma, n+1}^i \right) \right) \cap \mathcal{X}_\gamma$, the inclusion (7.2) holds. By Proposition 3.1, together with the fact that $\text{Ec}(x)$ is a positive (G, A) -orthogonal family, we have

$$\text{Ec}(x) \subset \Delta_{n_\gamma, n+1},$$

whence the equality we want. \square

Summarizing all the foregoing discussions, we get the following:

Theorem 7.4. *Let γ be a matrix in the form of equation (7.1). When $\text{val}(a) = m \leq n$, we have*

$$|F_\gamma(\mathbf{F}_q)| = \sum_{j=0}^{2m-1} \left(\left\lfloor \frac{j}{2} \right\rfloor + 1 \right) q^j + (2m+1) \sum_{j=2m}^{m+n} q^j \\ + \sum_{j=m+n+1}^{2m+n-1} (2(2m+n-j)+1) q^j + q^{2m+n}.$$

When $\text{val}(a) = m > n$, we have

$$|F_\gamma(\mathbf{F}_q)| = \sum_{j=0}^{2n} \left(\left\lfloor \frac{j}{2} \right\rfloor + 1 \right) q^j + \sum_{j=2n+1}^{3n} (2(3n+1-j)+1) q^j + q^{3n+1}.$$

Now it is easy to deduce the weighted orbital integral $J_M^\xi(\gamma)$. For $N \in \mathbf{N}$, $N \gg 0$, let

$$\Pi_N = [-n_\gamma - N, N] \subset \mathbf{R} \cong \mathfrak{a}_M^G.$$

We can count the number of rational points $|\mathcal{X}_\gamma^{n+1}(\Pi_N)(\mathbf{F}_q)|$ in two ways. By the Arthur–Kottwitz reduction, we have

$$|\mathcal{X}_\gamma^{n+1}(\Pi_N)(\mathbf{F}_q)| = |F_\gamma(\mathbf{F}_q)| + 2Nq^{n_\gamma} \cdot |F_\gamma^M(\mathbf{F}_q)|.$$

By the Harder–Narasimhan reduction, we have

$$|\mathcal{X}_\gamma^{n+1}(\Pi_N)(\mathbf{F}_q)| = 2|F_\gamma(\mathbf{F}_q)| + |\mathcal{X}_\gamma^{n+1, \xi}(\mathbf{F}_q)| \\ + (2N - n_\gamma - 1)q^{n_\gamma} \cdot |F_\gamma^M(\mathbf{F}_q)|,$$

where we use the fact that $\mathcal{X}_\gamma^{M, \nu, \xi} = F_\gamma^M$ for any $\nu \in \Lambda_M$ because γ is anisotropic in $\mathfrak{m}(F)$. The comparison of the two expressions implies

$$|\mathcal{X}_\gamma^{n+1, \xi}(\mathbf{F}_q)| = (n_\gamma + 1)q^{n_\gamma} \cdot |F_\gamma^M(\mathbf{F}_q)| - |F_\gamma(\mathbf{F}_q)|.$$

By Theorems 7.4 and 5.2, we have the following:

Theorem 7.5. *Let γ be a matrix in the form of equation (7.1). When $\text{val}(a) = m \leq n$, we have*

$$J_M^\xi(\gamma) = |\mathcal{X}_\gamma^{n+1, \xi}(\mathbf{F}_q)| = 2mq^{2m+n} + \sum_{j=m+n+1}^{2m+n-1} 2(j-m-n)q^j - \sum_{j=0}^{2m-1} \left(\left\lfloor \frac{j}{2} \right\rfloor + 1 \right) q^j.$$

When $\text{val}(a) = m > n$, we have

$$J_M^\xi(\gamma) = |\mathcal{X}_\gamma^{n+1, \xi}(\mathbf{F}_q)| = (2n+1)q^{3n+1} + \sum_{j=2n+1}^{3n} (2j-4n-1)q^j - \sum_{j=0}^{2n} \left(\left\lfloor \frac{j}{2} \right\rfloor + 1 \right) q^j.$$

By Theorem 2.8 and Remark 2.3, we get Arthur's weighted orbital integral. As before, the orbital integral I_γ^G can be calculated by equation (2.4):

$$I_\gamma^G = q^{n_\gamma} \sum_{i=0}^n q^i = \begin{cases} q^{2m} \sum_{i=0}^n q^i & \text{if } m \leq n, \\ q^{2n+1} \sum_{i=0}^n q^i & \text{if } m > n. \end{cases}$$

8. Calculations for GL_3 -anisotropic case

Let $G = \mathrm{GL}_3$ and $\gamma \in \mathfrak{gl}_3(F)$ be a regular semisimple integral element. Assume that $\mathrm{char}(k) > 3$ and $T \cong \mathrm{Res}_{E_3/F} E_3^\times$, with $E_3 = \mathbf{F}_q((\epsilon^{\frac{1}{3}}))$. As before, take the basis $\{\epsilon^{\frac{2}{3}}, \epsilon^{\frac{1}{3}}, 1\}$ of E_3 over F . We can assume that γ is of the form

$$\gamma = \begin{bmatrix} & b_0 \epsilon^{n_1} & c_0 \epsilon^{n_2} \\ c_0 \epsilon^{n_2+1} & & b_0 \epsilon^{n_1} \\ b_0 \epsilon^{n_1+1} & c_0 \epsilon^{n_2+1} & \end{bmatrix}, \quad (8.1)$$

with $b_0, c_0 \in \mathcal{O}^\times$ and $n_1, n_2 \in \mathbf{N}$. In this case, Arthur's weighted orbital integral is the same as the orbital integral, and both are equal to $|\mathcal{X}_\gamma^0(\mathbf{F}_q)|$. The matrix γ is equivalued of valuation $n_1 + \frac{1}{3}$ if $n_1 \leq n_2$, and equivalued of valuation $n_2 + \frac{2}{3}$ if $n_2 < n_1$. According to Goresky, Kottwitz, and MacPherson [16], the affine Springer fiber \mathcal{X}_γ admits affine paving

$$\mathcal{X}_\gamma = \bigcup_{\mathbf{a}=(a_1, a_2, a_3) \in \mathbf{Z}^3} \mathcal{X}_\gamma \cap I\epsilon^{\mathbf{a}}K/K.$$

Let $S_{\mathbf{a}}$ be the cell $\mathcal{X}_\gamma \cap I\epsilon^{\mathbf{a}}K/K$. Restricted to the connected component \mathcal{X}_γ^0 , we can calculate that $S_{\mathbf{a}}$ is nonempty if and only if

$$a_1 - a_2 \leq n_1, \quad a_2 - a_3 \leq n_1, \quad a_3 - a_1 \leq n_1 + 1, \quad (8.2)$$

and that it is of dimension

$$\left| \left\{ (m, \alpha) \in \mathbf{Z} \times \Phi(G, A) \mid 0 \leq m + \alpha(x) < n_1 + \frac{1}{3}, m + \alpha(y_{\mathbf{a}}) < 0 \right\} \right|,$$

with $x = (1, 2/3, 1/3)$, $y_{\mathbf{a}} = (-a_1, -a_2, -a_3) \in X_*(A) \otimes \mathbf{R}$. The results are summarized in Figure 8. Summing up, we get the following:

Theorem 8.1. *Let $\gamma \in \mathfrak{gl}_3(\mathcal{O})$ be the matrix in equation (8.1). Suppose that $n_1 \leq n_2$; it is then equivalued of valuation $n_1 + \frac{1}{3}$. The orbital integral associated to γ equals*

$$\begin{aligned} I_\gamma^G = |\mathcal{X}_\gamma^0(\mathbf{F}_q)| &= 1 + 2 \sum_{i=1}^{\lfloor \frac{n_1}{3} \rfloor} q^{2(3i-1)} (q^2 + q + 1) \\ &+ \sum_{i=1}^{n_1} \left(i - 2 \left\lfloor \frac{i}{3} \right\rfloor - 1 \right) q^{2i-3} (q^3 + 2q^2 + 2q + 1) \\ &+ \sum_{i=n_1+1}^{2n_1} \left(2n_1 - i - 2 \left\lceil \frac{2n_1-i}{3} \right\rceil + 1 \right) q^{i+n_1-1} (q+2) \\ &+ q^{2n_1-1} \left(\left\lceil \frac{2n_1-1}{3} \right\rceil - \left\lfloor \frac{n_1-2}{3} \right\rfloor - 1 \right) + 2 \sum_{i=\lceil \frac{2n_1-1}{3} \rceil}^{n_1-1} q^{3i+1}, \end{aligned}$$

where $\lfloor x \rfloor$ denotes the largest integer less than or equal to x , and $\lceil x \rceil$ denotes the smallest integer greater than or equal to x .




Figure 9. Counting points for ramified anisotropic $\gamma \in \mathfrak{gl}_3(\mathcal{O})$: Second case.

Theorem 8.2. Let $\gamma \in \mathfrak{gl}_3(\mathcal{O})$ be the matrix in equation (8.1). Suppose that $n_2 < n_1$; it is then equivalent of valuation $n_2 + \frac{2}{3}$. The orbital integral associated to γ equals

$$\begin{aligned}
 I_\gamma^G &= |\mathcal{X}_\gamma^0(\mathbf{F}_q)| = 1 + 2 \sum_{i=1}^{\lfloor \frac{n_2}{3} \rfloor} q^{2(3i-1)} (q^2 + q + 1) \\
 &\quad + \sum_{i=1}^{n_2} \left(i - 2 \left\lfloor \frac{i}{3} \right\rfloor - 1 \right) q^{2i-3} (q^3 + 2q^2 + 2q + 1) + \left(n_2 - 2 \left\lceil \frac{n_2 - 1}{3} \right\rceil \right) q^{2n_2-1} (1 + 2q^2) \\
 &\quad + \sum_{i=n_2+2}^{2n_2} \left(2n_2 - i - 2 \left\lceil \frac{2n_2 - i}{3} \right\rceil + 1 \right) q^{i+n_2-1} (1 + 2q) \\
 &\quad + 2 \sum_{i=0}^{\lfloor \frac{n_2-2}{3} \rfloor} q^{3(n_2-i)-2} (1 + q) + 2q^{2n_2} \left(\left\lceil \frac{2n_2 - 1}{3} \right\rceil - \left\lfloor \frac{n_2 - 2}{3} \right\rfloor - 1 \right) + q^{3n_2+1}.
 \end{aligned}$$

Acknowledgments. We want to thank Gérard Laumon for the discussions which led to this work, and we want to thank the anonymous referee for the careful reading and very helpful suggestions.

Competing Interest. The author(s) declare none.

References

- [1] J. ARTHUR, The characters of discrete series as orbital integrals, *Invent. Math.* **32** (1976), 205–261.
- [2] J. ARTHUR, The trace formula in invariant form, *Ann. of Math. (2)* **114** (1981), 1–74.
- [3] J. ARTHUR, A local trace formula, *Publ. Math. Inst. Hautes Études Sci.* **73** (1991), 5–96.
- [4] J. ARTHUR, *An introduction to the trace formula*, in *Harmonic Analysis, the Trace Formula, and Shimura Varieties*, Clay Mathematics Proceedings, **4**, pp. 1–263 (American Mathematical Society, Providence, RI, 2005).
- [5] R. BEZRUKAVNIKOV, The dimension of the fixed point set on affine flag manifolds, *Math. Res. Lett.* **3** (1996), 185–189.
- [6] M. BOROVoi, *Abelian Galois Cohomology of Reductive Groups*, Memoirs of the American Mathematical Society, **132**(626) (1998).
- [7] P.-H. CHAUDOUARD AND G. LAUMON, Sur l’homologie des fibres de Springer affines tronquées, *Duke Math. J.* **145**(3) (2008), 443–535.
- [8] P.-H. CHAUDOUARD AND G. LAUMON, Le lemme fondamental pondéré, I. Constructions géométriques, *Compos. Math.* **146**(6) (2010), 1416–1506.
- [9] Z. CHEN, Pureté des fibres de Springer affines pour GL_4 , *Bull. Soc. Math. France* **142**(2) (2014), 193–222.
- [10] Z. CHEN, The ξ -stability on the affine grassmannian, *Math. Z.* **280**(3-4) (2015), 1163–1184.
- [11] Z. CHEN, On the fundamental domain of affine Springer fibers, *Math. Z.* **286**(3-4) (2017), 1323–1356.
- [12] Z. CHEN, ‘Truncated affine grassmannians and truncated affine Springer fibers for GL_3 ’, Preprint, (2014), <http://arxiv.org/abs/1401.1930>.
- [13] M. GORESKEY, R. KOTTWITZ AND R. MACPHERSON, Equivariant cohomology, Koszul duality, and the localization theorem. *Invent. Math.* **131**(1) (1998), 25–83.
- [14] M. GORESKEY, R. KOTTWITZ AND R. MACPHERSON, Homology of affine Springer fibers in the unramified case, *Duke Math. J.* **121**(3) (2004), 509–561.
- [15] M. GORESKEY, R. KOTTWITZ AND R. MACPHERSON, Regular points in affine Springer fibers, *Michigan Math. J.* **53**(1) (2005), 97–107.
- [16] M. GORESKEY, R. KOTTWITZ AND R. MACPHERSON, Purity of equivalued affine Springer fibers, *Represent. Theory* **10** (2006), 130–146.
- [17] D. KAZHDAN AND G. LUSZTIG, Fixed point varieties on affine flag manifolds, *Israel. J. Math.* **62** (1988), 129–168.
- [18] R. E. KOTTWITZ, Isocrystals with additional structure. *Compos. Math.* **56**(2) (1985), 201–220.
- [19] R. E. KOTTWITZ, Isocrystals with additional structure, II, *Compos. Math.* **109**(3) (1997), 255–339.
- [20] B. C. NGÔ, Le lemme fondamental pour les algèbres de Lie, *Publ. Math. Inst. Hautes Études Sci.* **111** (2010), 1–169.