

INEQUALITY CONJECTURES ON DERIVATIONS OF LOCAL k -TH HESSAIN ALGEBRAS ASSOCIATED TO ISOLATED HYPERSURFACE SINGULARITIES

NAVEED HUSSAIN, STEPHEN S.-T. YAU, AND HUIAIQING ZUO

ABSTRACT. Let $(V, 0)$ be an isolated hypersurface singularity. We introduce a series of new derivation Lie algebras $L_k(V)$ associated to $(V, 0)$. Its dimension is denoted as $\lambda_k(V)$. The $L_k(V)$ is a generalization of the Yau algebra $L(V)$ and $L_0(V) = L(V)$. These numbers $\lambda_k(V)$ are new numerical analytic invariants of an isolated hypersurface singularity. In this article we compute $L_1(V)$ for fewnomial isolated singularities (Binomial, Trinomial) and obtain the formulas of $L_1(V)$. We also formulate a sharp upper estimate conjecture for the $L_k(V)$ of weighted homogeneous isolated hypersurface singularities and we prove this conjecture for large class of singularities. Furthermore, we formulate another conjecture: $\dots > \lambda^{(i+1)}(V) > \lambda^i(V) \dots > \lambda^0(V) \geq \lambda_l(V)$, $l = 1, 2, \dots, n$. We partially prove it for binomial and trinomial singularities.

Keywords. isolated hypersurface singularity, Lie algebra, moduli algebra.
MSC(2010). 14B05, 32S05.

1. INTRODUCTION

The algebra of germs of holomorphic functions at the origin of \mathbb{C}^n is denoted as \mathcal{O}_n . Clearly, \mathcal{O}_n can be naturally identified with the algebra of convergent power series in n indeterminates with complex coefficients. As a ring \mathcal{O}_n has a unique maximal ideal m , the set of germs of holomorphic functions which vanish at the origin. Let $\mathbb{C}[x_1, \dots, x_n]$ be the polynomial ring. For any $f \in \mathbb{C}[x_1, \dots, x_n]$, we denote by $V = V(f)$ the germ at the origin of \mathbb{C}^n of hypersurface $\{f = 0\} \subset \mathbb{C}^n$. In other words, if the origin is an isolated zero of the gradient of f , then V is a germ of isolated hypersurface singularity. According to Hilbert's Nullstellensatz for an isolated singularity $V = V(f) = \{f = 0\}$ the factor-algebra $A(V) = \mathcal{O}_n / (f, \frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n})$ is finite dimensional. This factor-algebra is called the moduli algebra of V and its dimension $\tau(V)$ is called Tyurina number. The Mather-Yau theorem stated that: Let V_1 and V_2 be two isolated hypersurface singularities and, $A(V_1)$ and $A(V_2)$ be the moduli algebra, then $(V_1, 0) \cong (V_2, 0) \iff A(V_1) \cong A(V_2)$.

Recall that the order of the lowest nonvanishing term in the power series expansion of f at 0 is called the multiplicity (denoted by $mult(f)$) of the singularity $(V, 0)$. A polynomial $f \in \mathbb{C}[x_1, \dots, x_n]$ is said to be weighted homogeneous if there exist positive rational numbers w_1, \dots, w_n (weights of x_1, \dots, x_n) and d such that, $\sum a_i w_i = d$ for each monomial $\prod x_i^{a_i}$ appearing in f with nonzero coefficient. The number d is called weighted homogeneous degree (w -degree) of f with respect to weights w_j . The weight type of f is denoted as $(w_1, \dots, w_n; d)$. Without loss of generality, we can assume that

Both Yau and Zuo were supported by NSFC Grants 11961141005 and 11531007. Zuo was supported by NSFC Grant 11771231. Yau was supported by Tsinghua University start-up fund and Tsinghua University Education Foundation fund (042202008). Naveed was supported by innovation team project of Humanities and Social Sciences in Colleges and universities of Guangdong Province (No.: 2020wcxtd008).

$w\text{-deg}f = 1$. According to ([YS], [Sae]) the weight types of 1 or 2-dimensional weighted homogeneous hypersurface singularities are topological invariants. The Milnor number of the isolated hypersurface singularity is defined by $\mu = \dim \mathcal{O}_n / (\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n})$. The Milnor number in case of weighted homogeneous hypersurface singularity is calculated by: $\mu = (\frac{1}{w_1} - 1)(\frac{1}{w_2} - 1) \cdots (\frac{1}{w_n} - 1)$ [MO]. In 1971, Saito was the first person who gave the necessary and sufficient numerical condition for V to be defined by a weighted homogeneous polynomial. His beautiful theorem says that f is a weighted homogeneous polynomial after a biholomorphic change of coordinates $\iff \mu = \tau$ [Sa].

An important class of weighted homogeneous isolated hypersurface singularity is fewnomial singularities which are defined by Elashvili and Khimshiashvili [EK]. A weighted homogeneous polynomial $f(x_1, \dots, x_n)$ is called fewnomial if number of variables coincides with number of monomials ([EK], [YZ2], [Kh], [Khi]). According to Ebeling and Takahashi [ET], the fewnomial singularity, which is defined by a fewnomial polynomial, is also called an invertible singularity.

It is well-known that for any isolated hypersurface singularity $(V, 0) \subset (\mathbb{C}^n, 0)$ where $V = V(f) = \{f = 0\}$, based on the Mather-Yau theorem [MY], one considers the Lie algebra of derivations of moduli algebra $A(V) := \mathcal{O}_n / (f, \frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n})$, i.e., $L(V) = \text{Der}(A(V), A(V))$. It is known that $L(V)$ is a finite dimensional solvable Lie algebra ([Ya2], [Ya3]). $L(V)$ is called the Yau algebra of V in [Yu] and [Khi] in order to distinguish from Lie algebras of other types appearing in singularity theory ([AVZ], [AM]). The Yau algebra plays an important role in singularities. Yau and his collaborators have been systematically studying the Lie algebras of isolated hypersurface singularities begin from eighties (see, e.g., [Ya1]-[Ya3], [BY], [XY], [SY], [YZ1, YZ2], [CYZ], [CCYZ], [HYZ1]-[HYZ7], [CHYZ], [Hu]).

In the theory of isolated singularities, one always wants to find invariants associated to the isolated singularities. Hopefully with enough invariants found, one can distinguish between isolated singularities. However, not many invariants are known. In this paper we introduce the new series new derivation Lie algebra associated to the isolated hypersurface singularity $(V, 0)$. For any isolated hypersurface singularity $(V, 0) \subset (\mathbb{C}^n, 0)$ defined by the holomorphic function $f(x_1, \dots, x_n)$, let $Hess(f)$ be the Hessian matrix (f_{ij}) of the second order partial derivatives of f and $h(f)$ be the Hessian of f , i.e. the determinant of this matrix $Hess(f)$. More generally, for each k satisfying $0 \leq k \leq n$ we denote by $h_k(f)$ the ideal in \mathcal{O}_n generated by all $k \times k$ -minors in the matrix $Hess(f)$. In particular, the ideal $h_n(f) = (h(f))$ is a principal ideal. For each k as above, consider the graded k -th Hessian algebra of the polynomial f defined by

$$H_k(f) = \mathcal{O}_n / (f + J(f) + h_k(f)).$$

In particular, $H_0(f)$ is exactly the well-known moduli algebra $A(V)$.

It is easy to check that the isomorphism class of the local k -th Hessian algebra $H_k(f)$ is contact invariant of f , i.e. depends only on the isomorphism class of the germ $(V, 0)$ [DS]. The dimension of $L_k(V)$ is denoted by $\lambda_k(V)$ which is new numerical analytic invariant of an isolated hypersurface singularity.

Recall that in [HYZ2], we also generalized the Yau algebra and introduced a new series of k -th Yau algebras $L^k(V)$ which are defined to be the Lie algebras of derivations of the moduli algebras $A^k(V) = \mathcal{O}_n / (f, m^k J(f))$, $k \geq 0$, i.e., $L^k(V) = \text{Der}(A^k(V), A^k(V))$ and

where m is the maximal ideal of \mathcal{O}_n . In particular, it is Yau algebra when $k = 0$. The dimension of $L^k(V)$ is denoted by $\lambda^k(V)$.

It is interesting to bound the Yau number with a number which depends on weight type. In [YZ2], Yau and Zuo firstly proposed the sharp upper estimate conjecture that bound the Yau number $\lambda(V)$. They also proved that this conjecture holds in case of binomial isolated hypersurface singularities. Furthermore, in [HYZ1], this conjecture was verified for trinomial singularities (the definitions of fewnomial, binomial, and trinomial singularities can be found in [YZ2]).

A natural interesting question is: whether one can give a sharp bound for the new introduced $\lambda_k(V)$ of an isolated hypersurface singularities $(V, 0)$. We proposed the following sharp upper estimate conjecture.

Conjecture 1.1. *For each $0 \leq k \leq n$, assume that $\lambda_k(\{x_1^{a_1} + \cdots + x_n^{a_n} = 0\}) = h_k(a_1, \cdots, a_n)$. Let $(V, 0) = \{(x_1, x_2, \cdots, x_n) \in \mathbb{C}^n : f(x_1, x_2, \cdots, x_n) = 0\}$, ($n \geq 2$) be an isolated singularity defined by the weighted homogeneous polynomial $f(x_1, x_2, \cdots, x_n)$ of weight type $(w_1, w_2, \cdots, w_n; 1)$ and $\text{mult}(f) \geq 4$. Then $\lambda_k(V) \leq h_k(1/w_1, \cdots, 1/w_n)$.*

Remark 1.1. *The inequality in Conjecture 1.1 holds true under the condition of $\text{mult}(f) = 3$ and $k = 0$ for binomial singularities [YZ2] and trinomial singularities [HYZ1]. However, in this paper, we will give some examples which show that the inequality does not hold for $k = 1$ and $\text{mult}(f) = 3$ for binomial and trinomial singularities (see remarks 3.2-3.8).*

In [HYZ2] we also formulate a similar inequality conjecture as above for the $\lambda^k(V)$ of weighted homogeneous isolated hypersurface singularities and we have also proven this conjecture for large class of singularities, i.e., for binomial and trinomial singularities. Another natural interesting question is: whether there is any relation between the numerical invariants $\lambda_k(V)$ and $\lambda^k(V)$ of isolated hypersurface singularities $(V, 0)$. We proposed the following an Inequality conjecture.

Conjecture 1.2. *With the above notations, let $(V, 0)$ be an isolated hypersurface singularity defined by $f \in \mathcal{O}_n$, $n \geq 2$, and $\text{mult}(f) \geq 3$. Then for each $i \geq 0$,*

$$\cdots > \lambda^{(i+1)}(V) > \lambda^i(V) \cdots > \lambda^0(V) \geq \lambda_l(V), \quad l = 1, 2, \cdots, n.$$

In [CHYZ], we studied $L_n(V)$ (note that we use a different notation $L^*(V)$ in stead of $L_n(V)$ there) and obtained the following result.

Theorem 1.1. ([CHYZ], Theorem D) *Let f be a weighted homogeneous polynomial in $\mathbb{C}[x_1, x_2, \cdots, x_n]$ ($2 \leq n \leq 4$) with respect to weight system $(w_1, w_2, \cdots, w_n; 1)$ and with $\text{mult}(f) \geq 3$. Suppose that f defines an isolated singularity $(V, 0)$, then*

$$\lambda_n(V) = \lambda(V).$$

Remark 1.2. *It follows from Theorem 1.1 that $\lambda^0(V) = \lambda_2(V)$ for binomial singularities and $\lambda^0(V) = \lambda_3(V)$ for trinomial singularities.*

The main purpose of this paper is to verify the Conjecture 1.1 and Conjecture 1.2 for binomial and trinomial singularities. We obtain the following main results.

Main Theorem A. Let $(V, 0) = \{(x_1, x_2, \dots, x_n) \in \mathbb{C}^n : x_1^{a_1} + \dots + x_n^{a_n} = 0\}$, ($n \geq 2$; $a_i \geq 3$, $1 \leq i \leq n$). Then

$$\lambda_1(V) = h_1(a_1, \dots, a_n) = \sum_{j=1}^n \frac{a_j - 3}{a_j - 2} \prod_{i=1}^n (a_i - 2).$$

Main Theorem B. Let $(V, 0)$ be a binomial singularity defined by the weighted homogeneous polynomial $f(x_1, x_2)$ (see corollary 2.1) with weight type $(w_1, w_2; 1)$ and $\text{mult}(f) \geq 4$. Then

$$\lambda_1(V) \leq h_1\left(\frac{1}{w_1}, \frac{1}{w_2}\right) = \sum_{j=1}^2 \frac{\frac{1}{w_j} - 3}{\frac{1}{w_j} - 2} \prod_{i=1}^2 \left(\frac{1}{w_i} - 2\right).$$

Main Theorem C. Let $(V, 0)$ be a fewnomial singularity defined by the weighted homogeneous polynomial $f(x_1, x_2, x_3)$ (see proposition 2.2) with weight type $(w_1, w_2, w_3; 1)$ and $\text{mult}(f) \geq 4$. Then

$$\lambda_1(V) \leq h_1\left(\frac{1}{w_1}, \frac{1}{w_2}, \frac{1}{w_3}\right) = \sum_{j=1}^3 \frac{\frac{1}{w_j} - 3}{\frac{1}{w_j} - 2} \prod_{i=1}^3 \left(\frac{1}{w_i} - 2\right).$$

Main Theorem D. Let $(V, 0)$ be a binomial singularity defined by the weighted homogeneous polynomial $f(x_1, x_2)$ (see corollary 2.1) with weight type $(w_1, w_2; 1)$ and $\text{mult}(f) \geq 3$. Then

$$\lambda^2 > \lambda^1 > \lambda^0 = \lambda_2 > \lambda_1.$$

Main Theorem E. Let $(V, 0)$ be a trinomial singularity defined by the weighted homogeneous polynomial $f(x_1, x_2, x_3)$ (see proposition 2.2) with weight type $(w_1, w_2, w_3; 1)$ and $\text{mult}(f) \geq 3$. Then

$$\lambda^2 > \lambda^1 > \lambda^0 = \lambda_3 > \lambda_1.$$

Remark 1.3. In main theorem E, the $\lambda_2(V)$ is extremely complicated to compute.

Main Theorem F. Let $(V_f, 0) \subset (\mathbb{C}^n, 0)$ and $(V_g, 0) \subset (\mathbb{C}^m, 0)$ be defined by weighted homogeneous polynomials $f(x_1, x_2, \dots, x_n) = 0$ of weight type $(w_1, w_2, \dots, w_n; 1)$ and $g(y_1, y_2, \dots, y_m) = 0$ of weight type $(w_{n+1}, w_{n+2}, \dots, w_{n+m}; 1)$ respectively. With the above notations. Then

$$L_1(V_{f+g}) = L_1(V_f) \otimes H_1(V_g) + H_1(V_f) \otimes L_1(V_g). \quad (1.1)$$

We obtain the following result as an immediately corollary of Theorem F.

Corollary 1.1. Let $(V_f, 0) \subset (\mathbb{C}^n, 0)$ and $(V_g, 0) \subset (\mathbb{C}^m, 0)$ be defined by weighted homogeneous polynomials $f(x_1, x_2, \dots, x_n) = 0$ of weight type $(w_1, w_2, \dots, w_n; 1)$ and $g(y_1, y_2, \dots, y_m) = 0$ of weight type $(w_{n+1}, w_{n+2}, \dots, w_{n+m}; 1)$ respectively. With the above notations, let $\mu_1(V_f)$, $\mu_1(V_g)$ be dimensions of the 1-st Hessian algebras $H_1(V_f)$ and $H_1(V_g)$ respectively. Then

$$\lambda_1(V_{f+g}) = \mu_1(V_f)\lambda_1(V_g) + \mu_1(V_g)\lambda_1(V_f). \quad (1.2)$$

2. GENERALITIES ON DERIVATION LIE ALGEBRAS OF ISOLATED SINGULARITIES

In this section we shall briefly defined the basic definitions and important results which are helpful to solve the problem. The following basic concepts and results will be used to compute the derivation Lie algebras of isolated hypersurface singularities.

Let A, B be associative algebras over \mathbb{C} . The subalgebra of endomorphisms of A generated by the identity element and left and right multiplications by elements of A is called multiplication algebra $M(A)$ of A . The centroid $C(A)$ is defined as the set of endomorphisms of A which commute with all elements of $M(A)$. Obviously, $C(A)$ is a unital subalgebra of $\text{End}(A)$. The following statement is a particular case of a general result from Proposition 1.2 of [Bl]. Let $S = A \otimes B$ be a tensor product of finite dimensional associative algebras with units. Then

$$\text{Der}S \cong (\text{Der}A) \otimes C(B) + C(A) \otimes (\text{Der}B).$$

We will only use this result for commutative associative algebras with unit, in which case the centroid coincides with the algebra itself and one has following result for commutative associative algebras A, B :

Theorem 2.1. ([Bl]) *For commutative associative algebras A, B ,*

$$\text{Der}S \cong (\text{Der}A) \otimes B + A \otimes (\text{Der}B). \quad (2.1)$$

We shall use this formula in the sequel.

Definition 2.1. *Let J be an ideal in an analytic algebra S . Then $\text{Der}_J S \subseteq \text{Der}_{\mathbb{C}} S$ is Lie subalgebra of all $\sigma \in \text{Der}_{\mathbb{C}} S$ for which $\sigma(J) \subset J$.*

We shall use the following well-known result to compute the derivations.

Theorem 2.2. ([YZ2]) *Let J be an ideal in $R = \mathbb{C}\{x_1, \dots, x_n\}$. Then there is a natural isomorphism of Lie algebras*

$$(\text{Der}_J R)/(J \cdot \text{Der}_{\mathbb{C}} R) \cong \text{Der}_{\mathbb{C}}(R/J).$$

Recall that a derivation of commutative associative algebra A is defined as a linear endomorphism D of A satisfying the Leibniz rule: $D(ab) = D(a)b + aD(b)$. Thus for such an algebra A one can consider the Lie algebra of its derivations $\text{Der}(A, A)$ with the bracket defined by the commutator of linear endomorphisms.

Definition 2.2. *Let $(V, 0)$ be an isolated hypersurface singularity. The series new derivation Lie algebra arising from the isolated hypersurface singularity $(V, 0)$ is defined as $L_k(V) := \text{Der}(H_k(f), H_k(f))$, $0 \leq k \leq n$ (where $H_k(f) = \mathcal{O}_{n+1}/(f + J(f) + h_k(f))$ and $h_k(f)$ is the ideal in \mathcal{O}_n generated by all $k \times k$ -minors in the matrix $\text{Hess}(f)$). Its dimension is denoted as $\lambda_k(V)$. In particular, $H_0(f)$ is exactly the well-known moduli algebra $A(V)$. Thus $L_k(V)$ is a generalization of Yau algebra $L(V)$ and $L_0(V) = L(V)$. These numbers $\lambda_k(V)$ are new numerical analytic invariants of an isolated hypersurface singularity.*

Definition 2.3. *A polynomial $f \in \mathbb{C}[x_1, x_2, \dots, x_n]$ is called quasi-homogeneous (or weighted homogeneous) if there exist positive rational numbers w_1, \dots, w_n (called weights of indeterminates x_j) and d such that, for each monomial $\prod x_j^{k_j}$ appearing in f with non-zero coefficient, one has $\sum w_j k_j = d$. The number d is called the quasi-homogeneous*

degree (w -degree) of f with respect to weights w_j and is denoted $\deg f$. The collection $(w; d) = (w_1, \dots, w_n; d)$ is called the quasi-homogeneity type (qh -type) of f .

Definition 2.4. An isolated hypersurface singularity in \mathbb{C}^n is *fewnomial* if it can be defined by a n -nomial in n variables and it is a *weighted homogeneous fewnomial isolated singularity* if it can be defined by a weighted homogeneous fewnomial. 3-nomial isolated hypersurface singularity is also called *trinomial singularity*.

Proposition 2.1. Let f be a weighted homogeneous fewnomial isolated singularity with $\text{mult}(f) \geq 3$. Then f is analytically equivalent to a linear combination of the following three series:

- Type A. $x_1^{a_1} + x_2^{a_2} + \dots + x_{n-1}^{a_{n-1}} + x_n^{a_n}$, $n \geq 1$,
- Type B. $x_1^{a_1}x_2 + x_2^{a_2}x_3 + \dots + x_{n-1}^{a_{n-1}}x_n + x_n^{a_n}$, $n \geq 2$,
- Type C. $x_1^{a_1}x_2 + x_2^{a_2}x_3 + \dots + x_{n-1}^{a_{n-1}}x_n + x_n^{a_n}x_1$, $n \geq 2$.

Proposition 2.1 has an immediate corollary.

Corollary 2.1. Each binomial isolated singularity is analytically equivalent to one from the three series: A) $x_1^{a_1} + x_2^{a_2}$, B) $x_1^{a_1}x_2 + x_2^{a_2}$, C) $x_1^{a_1}x_2 + x_2^{a_2}x_1$.

Wolfgang and Atsushi [ET] give the following classification of weighted homogeneous fewnomial singularities in case of three variables.

Proposition 2.2. ([ET]) Let $f(x_1, x_2, x_3)$ be a weighted homogeneous fewnomial isolated singularity with $\text{mult}(f) \geq 3$. Then f is analytically equivalent to following five types:

- Type 1. $x_1^{a_1} + x_2^{a_2} + x_3^{a_3}$,
- Type 2. $x_1^{a_1}x_2 + x_2^{a_2}x_3 + x_3^{a_3}$,
- Type 3. $x_1^{a_1}x_2 + x_2^{a_2}x_3 + x_3^{a_3}x_1$,
- Type 4. $x_1^{a_1} + x_2^{a_2} + x_3^{a_3}x_1$,
- Type 5. $x_1^{a_1}x_2 + x_2^{a_2}x_1 + x_3^{a_3}$.

3. PROOF OF MAIN THEOREMS

In order to prove the main theorems, we need to prove following propositions.

Proposition 3.1. Let $(V, 0)$ be a weighted homogeneous fewnomial isolated singularity which is defined by $f = x_1^{a_1} + x_2^{a_2} + \dots + x_n^{a_n}$ ($a_i \geq 3, i = 1, 2, \dots, n$) with weight type $(\frac{1}{a_1}, \frac{1}{a_2}, \dots, \frac{1}{a_n}; 1)$. Then

$$\lambda_1(V) = \sum_{j=1}^n \frac{a_j - 3}{a_j - 2} \prod_{i=1}^n (a_i - 2).$$

Proof. The generalized moduli algebra

$$H_1(V) = \mathbb{C}\{x_1, x_2, x_3, \dots, x_n\} / (\tilde{J}(f))$$

and where $\tilde{J}(f)$ is ideal of 2nd order partial derivative of f . It is note that $H_1(V)$ has dimension $\prod_{i=1}^n (a_i - 2)$ and has a monomial basis of the form

$$\{x_1^{i_1}x_2^{i_2} \dots x_n^{i_n}, 0 \leq i_1 \leq a_1 - 3, 0 \leq i_2 \leq a_2 - 3, \dots, 0 \leq i_n \leq a_n - 3\},$$

with following relations:

$$x_1^{a_1-2} = 0, x_2^{a_2-2} = 0, x_3^{a_3-2} = 0, \dots, x_n^{a_n-2} = 0. \quad (3.1)$$

In order to compute a derivation D of $H_1(V)$ it suffices to indicate its values on the generators x_1, x_2, \dots, x_n which can be written in terms of the monomial basis. Without loss of generality, we write

$$Dx_j = \sum_{i_1=0}^{a_1-3} \sum_{i_2=0}^{a_2-3} \cdots \sum_{i_n=0}^{a_n-3} c_{i_1, i_2, \dots, i_n}^j x_1^{i_1} x_2^{i_2} \cdots x_n^{i_n}, \quad j = 1, 2, \dots, n.$$

Using the above relations (3.1) one easily finds the necessary and sufficient conditions defining a derivation of $H_1(V)$ as follows:

$$\begin{aligned} c_{0, i_2, i_3, \dots, i_n}^1 &= 0; 0 \leq i_2 \leq a_2 - 3, 0 \leq i_3 \leq a_3 - 3, \dots, 0 \leq i_n \leq a_n - 3; \\ c_{i_1, 0, i_3, \dots, i_n}^2 &= 0; 0 \leq i_1 \leq a_1 - 3, 0 \leq i_3 \leq a_3 - 3, \dots, 0 \leq i_n \leq a_n - 3; \\ c_{i_1, i_2, 0, \dots, i_n}^3 &= 0; 0 \leq i_1 \leq a_1 - 3, 0 \leq i_2 \leq a_2 - 3, \dots, 0 \leq i_n \leq a_n - 3; \\ &\vdots \\ c_{i_1, i_2, i_3, \dots, i_{n-1}, 0}^n &= 0; 0 \leq i_1 \leq a_1 - 3, 0 \leq i_2 \leq a_2 - 3, \dots, 0 \leq i_{n-1} \leq a_{n-1} - 3. \end{aligned}$$

Therefore we obtain the following description of Lie algebras in question:

$$\begin{aligned} x_1^{i_1} x_2^{i_2} \cdots x_n^{i_n} \partial_1, & \quad 1 \leq i_1 \leq a_1 - 3, 0 \leq i_2 \leq a_2 - 3, 0 \leq i_3 \leq a_3 - 3, \dots, 0 \leq i_n \leq a_n - 3; \\ x_1^{i_1} x_2^{i_2} \cdots x_n^{i_n} \partial_2, & \quad 0 \leq i_1 \leq a_1 - 3, 1 \leq i_2 \leq a_2 - 3, 0 \leq i_3 \leq a_3 - 3, \dots, 0 \leq i_n \leq a_n - 3; \\ x_1^{i_1} x_2^{i_2} \cdots x_n^{i_n} \partial_3, & \quad 0 \leq i_1 \leq a_1 - 3, 0 \leq i_2 \leq a_2 - 3, 1 \leq i_3 \leq a_3 - 3, 0 \leq i_4 \leq a_4 - 3, \\ & \quad 0 \leq i_5 \leq a_5 - 3, 0 \leq i_6 \leq a_6 - 3, \dots, 0 \leq i_n \leq a_n - 3; \\ & \vdots \\ x_1^{i_1} x_2^{i_2} \cdots x_n^{i_n} \partial_n, & \quad 0 \leq i_1 \leq a_1 - 3, 0 \leq i_2 \leq a_2 - 3, 0 \leq i_3 \leq a_3 - 3, \dots, 1 \leq i_n \leq a_n - 3. \end{aligned}$$

Therefore we have the following formula

$$\lambda_1(V) = \sum_{j=1}^n \frac{a_j - 3}{a_j - 2} \prod_{i=1}^n (a_i - 2).$$

Q.E.D.

Remark 3.1. Let $(V, 0)$ be a weighted homogeneous fewnomial isolated singularity of type A which is defined by $f = x_1^{a_1} + x_2^{a_2}$ ($a_1 \geq 3, a_2 \geq 3$) with weight type $(\frac{1}{a_1}, \frac{1}{a_2}; 1)$. Then it follows from Proposition 3.1 that

$$\lambda_1(V) = 2a_1a_2 - 5(a_1 + a_2) + 12.$$

Proposition 3.2. Let $(V, 0)$ be a binomial isolated singularity of type B which is defined by $f = x_1^{a_1} + x_2^{a_2}x_1$ ($a_1 \geq 3, a_2 \geq 2$) with weight type $(\frac{1}{a_1}, \frac{a_1-1}{a_1a_2}; 1)$. Then

$$\lambda_1(V) = \begin{cases} 2a_1a_2 - 5(a_1 + a_2) + 15; & a_1 \geq 4, a_2 \geq 3 \\ a_2 - 2; & a_1 = 3, a_2 \geq 3 \\ 0; & a_1 \geq 3, a_2 = 2. \end{cases}$$

Furthermore, assuming that $\text{mult}(f) \geq 4$, we have

$$2a_1a_2 - 5(a_1 + a_2) + 15 \leq \frac{2a_1^2a_2}{a_1 - 1} - 5\left(\frac{a_1a_2}{a_1 - 1} + a_1\right) + 12.$$

Proof. It follows that the generalized moduli algebra

$$H_1(V) = \mathbb{C}\{x_1, x_2\} / \left(\frac{\partial^2 f}{\partial x_1^2}, \frac{\partial^2 f}{\partial x_1 \partial x_2}, \frac{\partial^2 f}{\partial x_2^2} \right)$$

has dimension $a_1a_2 - 2(a_1 + a_2) + 5$ and has a monomial basis of the form

$$\{x_1^{i_1}x_2^{i_2}, 0 \leq i_1 \leq a_1 - 3; 0 \leq i_2 \leq a_2 - 3; x_2^{a_2-2}\}, \quad (3.2)$$

with the following relations:

$$x_1^{a_1-2} = 0, \quad (3.3)$$

$$x_2^{a_2-1} = 0, \quad (3.4)$$

$$x_1x_2^{a_2-2} = 0. \quad (3.5)$$

In order to compute a derivation D of $H_1(V)$ it suffices to indicate its values on the generators x_1, x_2 which can be written in terms of the basis (3.2). Without loss of generality, we write

$$Dx_j = \sum_{i_1=0}^{a_1-3} \sum_{i_2=0}^{a_2-3} c_{i_1, i_2}^j x_1^{i_1} x_2^{i_2} + c_{0, a_2-2}^j x_2^{a_2-2}, \quad j = 1, 2.$$

Using the relations (3.3) - (3.5) one easily finds the necessary and sufficient conditions defining a derivation of $H_1(V)$ as follows:

$$c_{0,0}^1 = c_{0,1}^1 = \cdots = c_{0, a_2-3}^1 = 0; \quad (3.6)$$

$$c_{0,0}^2 = c_{1,0}^2 = \cdots = c_{a_1-4,0}^2 = 0. \quad (3.7)$$

Using (3.6) and (3.7) we obtain the following description of the Lie algebras in question. The following derivations form a basis of $\text{Der}H_1(V)$:

$$\begin{aligned} & x_1^{i_1}x_2^{i_2}\partial_1, 1 \leq i_1 \leq a_1 - 3, 0 \leq i_2 \leq a_2 - 3; x_1^{i_1}x_2^{i_2}\partial_2, 1 \leq i_1 \leq a_1 - 3, 1 \leq i_2 \leq a_2 - 3; \\ & x_2^{i_2}\partial_2; 1 \leq i_2 \leq a_2 - 2; x_1^{a_1-3}\partial_2. \end{aligned}$$

Therefore we have the following formula

$$\lambda_1(V) = 2a_1a_2 - 5(a_1 + a_2) + 15.$$

In case of $a_1 = 3, a_2 \geq 3$ we have following basis:

$$x_2^{i_2}\partial_2, 1 \leq i_2 \leq a_2 - 2.$$

Therefore we have

$$\lambda_1(V) = a_2 - 2.$$

It is follows from proposition 3.1 we have

$$h_1(a_1, a_2) = 2a_1a_2 - 5(a_1 + a_2) + 12.$$

After putting the weight type $(\frac{1}{a_1}, \frac{a_1-1}{a_1 a_2}; 1)$ of binomial isolated singularity of type B we have

$$h_1\left(\frac{1}{w_1}, \frac{1}{w_2}\right) = \frac{2a_1^2 a_2}{a_1 - 1} - 5\left(a_1 + \frac{a_1 a_2}{a_1 - 1}\right) + 12.$$

Finally we need to show that

$$2a_1 a_2 - 5(a_1 + a_2) + 15 \leq \frac{2a_1^2 a_2}{a_1 - 1} - 5\left(\frac{a_1 a_2}{a_1 - 1} + a_1\right) + 12. \quad (3.8)$$

After solving 3.8 we have $(a_1 - 5)(a_2 - 3) + a_1 a_2 - 12 \geq 0$.

Q.E.D.

Remark 3.2. Note that, for $\text{mult}(f) = 3$, the inequality in Proposition 3.2 does not hold true when $a_1 \geq 5$ and $a_2 = 2$.

Proposition 3.3. Let $(V, 0)$ be a binomial isolated singularity of type C which is defined by $f = x_1^{a_1} x_2 + x_2^{a_2} x_1$ ($a_1 \geq 2, a_2 \geq 2$) with weight type $(\frac{a_2-1}{a_1 a_2-1}, \frac{a_1-1}{a_1 a_2-1}; 1)$. Then

$$\lambda_1(V) = \begin{cases} 2a_1 a_2 - 5(a_1 + a_2) + 19; & a_1 \geq 5, a_2 \geq 5 \\ a_2 + 1; & a_1 = 3, a_2 \geq 3 \\ 3a_2 - 2; & a_1 = 4, a_2 \geq 5 \\ 9; & a_1 = 4, a_2 = 4 \\ 0; & a_1 = 2, a_2 \geq 2 \end{cases}$$

Furthermore, assuming that $\text{mult}(f) \geq 4$, we have

$$2a_1 a_2 - 5(a_1 + a_2) + 19 \leq \frac{2(a_1 a_2 - 1)^2}{(a_1 - 1)(a_2 - 1)} - 5(a_1 a_2 - 1) \left(\frac{a_1 + a_2 - 2}{(a_1 - 1)(a_2 - 1)}\right) + 12.$$

Proof. It follows that the generalized moduli algebra

$$H_1(V) = \mathbb{C}\{x_1, x_2\} / \left(\frac{\partial^2 f}{\partial x_1^2}, \frac{\partial^2 f}{\partial x_1 \partial x_2}, \frac{\partial^2 f}{\partial x_2^2}\right)$$

has dimension $a_1 a_2 - 2(a_1 + a_2) + 7$ and has a monomial basis of the form

$$\{x_1^{i_1} x_2^{i_2}, 0 \leq i_1 \leq a_1 - 3; 0 \leq i_2 \leq a_2 - 3; x_1^{a_1-2}; x_2^{i_2}, a_2 - 2 \leq i_2 \leq a_2 - 1\}, \quad (3.9)$$

with the following relations:

$$a_1 x_1^{a_1-1} + a_2 x_2^{a_2-1} = 0, \quad (3.10)$$

$$x_1^{a_1-2} x_2 = 0, \quad (3.11)$$

$$x_1 x_2^{a_2-2} = 0. \quad (3.12)$$

In order to compute a derivation D of $H_1(V)$ it suffices to indicate its values on the generators x_1, x_2 which can be written in terms of the basis (3.9). Without loss of generality, we write

$$Dx_j = \sum_{i_1=0}^{a_1-3} \sum_{i_2=0}^{a_2-3} c_{i_1, i_2}^j x_1^{i_1} x_2^{i_2} + c_{a_1-2, 0}^j x_1^{a_1-2} + \sum_{i_2=a_2-2}^{a_2-1} c_{0, i_2}^j x_2^{i_2}, \quad j = 1, 2.$$

Using the relations (3.10) - (3.12) one easily finds the necessary and sufficient conditions defining a derivation of $H_1(V)$ as follows:

$$c_{0,0}^1 = c_{0,1}^1 = \cdots = c_{0, a_2-4}^1 = 0; \quad (3.13)$$

$$c_{0,0}^2 = c_{1,0}^2 = \cdots = c_{a_1-4,0}^2 = 0; \quad (3.14)$$

$$-(a_1 - 1)a_2c_{1,0}^1 + a_1a_2c_{0,1}^2 = 0. \quad (3.15)$$

Using (3.13) - (3.15) we obtain the following description of the Lie algebras in question. The following derivations form a basis of $\text{Der}H_1(V)$:

$$\begin{aligned} & x_1^{i_1}x_2^{i_2}\partial_1, 1 \leq i_1 \leq a_1 - 3, 1 \leq i_2 \leq a_2 - 3; x_1^{i_1}x_2^{i_2}\partial_2, 1 \leq i_1 \leq a_1 - 3, 1 \leq i_2 \leq a_2 - 3; \\ & x_2^{i_2}\partial_1; a_2 - 3 \leq i_2 \leq a_2 - 1; x_1\partial_1 + \frac{a_1 - 1}{a_2 - 1}x_2\partial_2; x_1^{i_1}\partial_1, 2 \leq i_1 \leq a_1 - 2; x_2^{i_2}\partial_2, 2 \leq i_2 \leq a_2 - 1; \\ & x_1^{i_1}\partial_2, a_1 - 3 \leq i_1 \leq a_1 - 2. \end{aligned}$$

Therefore we have the following formula

$$\lambda_1(V) = 2a_1a_2 - 5(a_1 + a_2) + 19.$$

In case of $a_1 = 3, a_2 \geq 3$ we have following basis:

$$x_2^{a_2-2}\partial_1 + \frac{3x_1}{a_2}\partial_2; x_2^{a_2-1}\partial_1; x_1\partial_1 + \frac{2x_2}{a_2 - 1}\partial_2; x_2^{i_2}\partial_2, 2 \leq i_2 \leq a_2 - 1.$$

Therefore we have the following formula

$$\lambda_1(V) = a_2 + 1.$$

In case of $a_1 = 4, a_2 \geq 5$ we have following basis:

$$\begin{aligned} & x_2^{i_2}\partial_1; a_2 - 3 \leq i_2 \leq a_2 - 1; x_1\partial_1 + \frac{3x_2}{a_2 - 1}\partial_2; x_1x_2^{i_2}\partial_1, 1 \leq i_2 \leq a_2 - 3; \\ & x_1^2\partial_1; x_2^{i_2}\partial_2, 2 \leq i_2 \leq a_2 - 1; x_1^2\partial_2; x_1x_2^{i_2}\partial_2, 1 \leq i_2 \leq a_2 - 3. \end{aligned}$$

Therefore we have the following formula

$$\lambda_1(V) = 3a_2 - 2.$$

In case of $a_1 = 4, a_2 = 4$ we have following basis:

$$x_2^2\partial_1; x_2^3\partial_1; x_1x_2\partial_1; x_1^2\partial_1; x_2^2\partial_2; x_2^3\partial_2; x_1x_2\partial_2; x_1^2\partial_2.$$

Therefore we have the following formula

$$\lambda_1(V) = 9.$$

It is follows from proposition 3.1 and binomial isolated singularity of type C we have

$$h_1\left(\frac{1}{w_1}, \frac{1}{w_2}\right) = \frac{2(a_1a_2 - 1)^2}{(a_1 - 1)(a_2 - 1)} - 5\left(\frac{a_1a_2 - 1}{a_2 - 1} + \frac{a_1a_2 - 1}{a_1 - 1}\right) + 12.$$

Finally we need to show that

$$2a_1a_2 - 5(a_1 + a_2) + 19 \leq \frac{2(a_1a_2 - 1)^2}{(a_1 - 1)(a_2 - 1)} - 5\left(\frac{a_1a_2 - 1}{a_2 - 1} + \frac{a_1a_2 - 1}{a_1 - 1}\right) + 12. \quad (3.16)$$

After solving 3.16 we have

$$\begin{aligned} & a_1a_2^2[(a_2 - 2)(a_1 - 3) - a_1(a_2 - 5)] + a_2^3 + 3a_1^2a_2 + 11a_2^2(a_1 - 1) + 8a_1a_2(a_1 - 4) \\ & + 2a_1^2(a_2 - 4) + a_1a_2(a_1 - 4) + 17a_1 + 4(a_2 - 4) + (a_1 - 4) \geq 0. \end{aligned}$$

In case of $a_1 = 3, a_2 \geq 4$, we need to prove that:

$$a_2 + 1 \leq \frac{2(3a_2 - 1)^2}{2(a_2 - 1)} - 5\left(\frac{3a_2 - 1}{a_2 - 1} + \frac{3a_2 - 1}{2}\right) + 12. \quad (3.17)$$

After solving 3.17 we have

$$a_2^2 + 8a_2 - 15 \geq 0.$$

In case of $a_1 = 4, a_2 \geq 5$, we need to prove that:

$$3a_2 - 2 \leq \frac{2(4a_2 - 1)^2}{3(a_2 - 1)} - 5\left(\frac{4a_2 - 1}{a_2 - 1} + \frac{4a_2 - 1}{3}\right) + 12. \quad (3.18)$$

After solving 3.18 we have

$$a_2(a_2 - 2) - 10 \geq 0.$$

Q.E.D.

Remark 3.3. For $\text{mult}(f) = 3$, the inequality in Proposition 3.3 does not hold true when $a_1 = 2$ and $a_2 \geq 3$.

Remark 3.4. Let $(V, 0)$ be a fewnomial surface isolated singularity of type 1 (see Proposition 2.2) which is defined by $f = x_1^{a_1} + x_2^{a_2} + x_3^{a_3}$ ($a_1 \geq 3, a_2 \geq 3, a_3 \geq 3$) with weight type $(\frac{1}{a_1}, \frac{1}{a_2}, \frac{1}{a_3}; 1)$. Then it follows from Proposition 3.1 that

$$\lambda_1(V) = 3a_1a_2a_3 + 16(a_1 + a_2 + a_3) - 7(a_1a_2 + a_1a_3 + a_2a_3) - 36.$$

Proposition 3.4. Let $(V, 0)$ be a fewnomial surface isolated singularity of type 2 which is defined by $f = x_1^{a_1}x_2 + x_2^{a_2}x_3 + x_3^{a_3}$ ($a_1 \geq 2, a_2 \geq 2, a_3 \geq 3$) with weight type $(\frac{1-a_3+a_2a_3}{a_1a_2a_3}, \frac{a_3-1}{a_2a_3}, \frac{1}{a_3}; 1)$. Then

$$\lambda_1(V) = \begin{cases} 3a_1a_2a_3 - 7(a_1a_2 + a_1a_3 + a_2a_3) + 20(a_1 + a_3) + 16a_2 - 55; & a_1 \geq 4, a_2 \geq 4, a_3 \geq 4 \\ 2a_1a_3 - a_1 - 3a_3 - 1; & a_1 \geq 3, a_2 = 3, a_3 \geq 4 \\ 2a_2a_3 - 5a_2 - a_3 + 5; & a_1 = 3, a_2 \geq 4, a_3 \geq 4 \\ 2a_1a_2 - 3a_1 - 5a_2 + 10; & a_1 \geq 3, a_2 \geq 3, a_3 = 3 \\ a_3 - 3; & a_1 = 2, a_2 \geq 2, a_3 \geq 3 \\ a_1 - 3; & a_1 \geq 3, a_2 = 2, a_3 \geq 3. \end{cases}$$

Furthermore, assuming that $\text{mult}(f) \geq 4$, we have

$$\begin{aligned} 3a_1a_2a_3 - 7(a_1a_2 + a_1a_3 + a_2a_3) + 20(a_1 + a_3) + 16a_2 - 55 &\leq \frac{3a_1a_2^2a_3^3}{(1 - a_3 + a_2a_3)(a_3 - 1)} \\ &- 7\left(\frac{a_1a_2^2a_3^2}{(1 - a_3 + a_2a_3)(a_3 - 1)} + \frac{a_1a_2a_3^2}{1 - a_3 + a_2a_3} + \frac{a_2a_3^2}{a_3 - 1}\right) + 16\left(\frac{a_1a_2a_3}{1 - a_3 + a_2a_3} \right. \\ &\left. + \frac{a_2a_3}{a_3 - 1} + a_3\right) - 36. \end{aligned}$$

Proof. It is easy to see that the moduli algebra

$$H_1(V) = \mathbb{C}\{x_1, x_2, x_3\} / \left(\frac{\partial^2 f}{\partial x_1^2}, \frac{\partial^2 f}{\partial x_1 \partial x_2}, \frac{\partial^2 f}{\partial x_1 \partial x_3}, \frac{\partial^2 f}{\partial x_2^2}, \frac{\partial^2 f}{\partial x_3^2}, \frac{\partial^2 f}{\partial x_2 \partial x_3} \right)$$

has dimension $a_1 a_2 a_3 - 2(a_1 a_2 + a_1 a_3 + a_2 a_3) + 5(a_1 + a_3) + 4a_2 - 12$ and has a monomial basis of the form

$$\{x_1^{i_1} x_2^{i_2} x_3^{i_3}, 0 \leq i_1 \leq a_1 - 3; 0 \leq i_2 \leq a_2 - 3; 0 \leq i_3 \leq a_3 - 3; x_1^{i_1} x_3^{a_3 - 2}, 0 \leq i_1 \leq a_1 - 3; x_1^{a_1 - 2} x_3^{i_3}, 0 \leq i_3 \leq a_3 - 3\},$$

with the following relations:

$$x_1^{a_1 - 2} x_2 = 0, x_2^{a_2 - 2} x_3 = 0, x_3^{a_3 - 2} = 0, x_1^{a_1 - 1} = 0, x_2^{a_2 - 1} = 0. \quad (3.19)$$

In order to compute a derivation D of $H_1(V)$ it suffices to indicate its values on the generators x_1, x_2, x_3 which can be written in terms of the basis. Thus we can write

$$Dx_j = \sum_{i_1=0}^{a_1-3} \sum_{i_2=0}^{a_2-3} \sum_{i_3=0}^{a_3-3} c_{i_1, i_2, i_3}^j x_1^{i_1} x_2^{i_2} x_3^{i_3} + \sum_{i_1=0}^{a_1-3} c_{i_1, 0, a_3-2}^j x_1^{i_1} x_3^{a_3-2} + \sum_{i_3=0}^{a_3-3} c_{a_1-2, 0, i_3}^j x_1^{a_1-2} x_3^{i_3},$$

$j = 1, 2, 3.$

Using the relations (3.19) one easily finds the necessary and sufficient conditions defining a derivation of $H_1(V)$ as follows:

$$c_{0, i_2, i_3}^1 = 0; 0 \leq i_2 \leq a_2 - 4, 0 \leq i_3 \leq a_3 - 3; c_{0, a_2-3, 0}^1 = 0; \quad (3.20)$$

$$c_{i_1, 0, i_3}^2 = 0; 0 \leq i_1 \leq a_1 - 3, 0 \leq i_3 \leq a_3 - 4; c_{0, 0, a_3-2}^2 = 0; \quad (3.21)$$

$$c_{i_1, i_2, 0}^3 = 0; 0 \leq i_1 \leq a_1 - 3, 0 \leq i_2 \leq a_2 - 3; c_{a_1-2, 0, 0}^3 = 0; . \quad (3.22)$$

Using (3.20) - (3.22) we obtain the following description of Lie algebras in question:

$$\begin{aligned} & x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_1, 1 \leq i_1 \leq a_1 - 3, 0 \leq i_2 \leq a_2 - 3, 0 \leq i_3 \leq a_3 - 3; x_1^{a_1-2} x_3^{i_3} \partial_1, 0 \leq i_3 \leq a_3 - 3, \\ & x_2^{a_2-3} x_3^{i_3} \partial_1, 1 \leq i_3 \leq a_3 - 3; x_2^{a_2-2} \partial_1; x_1^{i_1} x_2^{a_2-2} \partial_1, 1 \leq i_1 \leq a_1 - 3, \\ & x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_2, 0 \leq i_1 \leq a_1 - 3, 1 \leq i_2 \leq a_2 - 3, 0 \leq i_3 \leq a_3 - 3; x_1^{a_1-2} x_3^{i_3} \partial_2, 0 \leq i_3 \leq a_3 - 3, \\ & x_1^{i_1} x_2^{a_2-2} \partial_2, 1 \leq i_1 \leq a_1 - 3; x_2^{a_2-2} \partial_2; x_1^{i_1} x_3^{a_3-3} \partial_2, 1 \leq i_1 \leq a_1 - 3, \\ & x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_3, 0 \leq i_1 \leq a_1 - 3, 0 \leq i_2 \leq a_2 - 3, 1 \leq i_3 \leq a_3 - 3, x_1^{i_1} x_2^{a_2-2} \partial_3, 1 \leq i_1 \leq a_1 - 3, \\ & x_2^{a_2-2} \partial_3; x_1^{a_1-2} x_3^{i_3} \partial_3, 1 \leq i_3 \leq a_3 - 3. \end{aligned}$$

Therefore we have

$$\lambda_1(V) = 3a_1 a_2 a_3 - 7(a_1 a_2 + a_1 a_3 + a_2 a_3) + 20(a_1 + a_3) + 16a_2 - 55.$$

In case of $a_1 \geq 3, a_2 = 3, a_3 \geq 4$, we obtain the following basis:

$$\begin{aligned} & x_1^{i_1} x_3^{i_3} \partial_1, 1 \leq i_1 \leq a_1 - 2, 0 \leq i_3 \leq a_3 - 3; x_1^{i_1} x_2 \partial_1, 1 \leq i_1 \leq a_1 - 3; x_2 \partial_1, \\ & x_1^{i_1} x_3^{a_3-3} \partial_2, 1 \leq i_1 \leq a_1 - 2; x_1^{i_1} x_2 \partial_2, 1 \leq i_1 \leq a_1 - 3; x_2 \partial_2, \\ & x_1^{i_1} x_3^{i_3} \partial_3, 0 \leq i_1 \leq a_1 - 2, 1 \leq i_3 \leq a_3 - 3; x_1^{i_1} x_2 \partial_3, 1 \leq i_1 \leq a_1 - 3. \end{aligned}$$

We have

$$\lambda_1(V) = 2a_1 a_3 - a_1 - 3a_3 - 1.$$

In case of $a_1 = 3, a_2 \geq 4, a_3 \geq 4$, we obtain the following basis:

$$\begin{aligned} & x_1 x_3^{i_3} \partial_1, 0 \leq i_3 \leq a_3 - 3; x_2^{a_2-3} x_3^{i_3} \partial_1, 1 \leq i_3 \leq a_3 - 3; x_2^{a_2-2} \partial_1, \\ & x_2^{i_2} x_3^{i_3} \partial_2, 1 \leq i_2 \leq a_2 - 3, 0 \leq i_3 \leq a_3 - 3; x_1 x_3^{i_3} \partial_2, 0 \leq i_3 \leq a_3 - 3; x_2^{a_2-2} \partial_2, \end{aligned}$$

$$x_2^{i_2} x_3^{i_3} \partial_3, 0 \leq i_2 \leq a_2 - 3, 1 \leq i_3 \leq a_3 - 3; x_1 x_3^{i_3} \partial_3, 1 \leq i_3 \leq a_3 - 3; x_2^{a_2-2} \partial_3.$$

We have

$$\lambda_1(V) = 2a_2 a_3 - 5a_2 - a_3 + 5.$$

In case of $a_1 \geq 3, a_2 \geq 3, a_3 = 3$, we obtain the following basis:

$$x_1^{i_1} x_2^{i_2} \partial_1, 1 \leq i_1 \leq a_1 - 3, 0 \leq i_2 \leq a_2 - 2; x_1^{i_1} x_2^{i_2} \partial_2, 0 \leq i_1 \leq a_1 - 3, 1 \leq i_2 \leq a_2 - 2; \\ x_2^{a_2-2} \partial_1; x_1^{a_1-2} \partial_1; x_1^{a_1-2} \partial_2.$$

We have

$$\lambda_1(V) = 2a_1 a_2 - 3a_1 - 5a_2 + 10.$$

In case of $a_1 = 2, a_2 \geq 2, a_3 \geq 3$, we obtain the following basis:

$$x_3^{i_3} \partial_3, 1 \leq i_3 \leq a_3 - 3.$$

In case of $a_1 \geq 3, a_2 = 2, a_3 \geq 3$, we obtain the following basis:

$$x_1^{i_1} \partial_1, 1 \leq i_1 \leq a_1 - 3.$$

Next, we need to show that when $a_1 \geq 4, a_2 \geq 4, a_3 \geq 4$, then

$$3a_1 a_2 a_3 - 7(a_1 a_2 + a_1 a_3 + a_2 a_3) + 20(a_1 + a_3) + 16a_2 - 55 \leq \frac{3a_1 a_2^2 a_3^3}{(1 - a_3 + a_2 a_3)(a_3 - 1)} \\ - 7\left(\frac{a_1 a_2^2 a_3^2}{(1 - a_3 + a_2 a_3)(a_3 - 1)} + \frac{a_1 a_2 a_3^2}{1 - a_3 + a_2 a_3} + \frac{a_2 a_3^2}{a_3 - 1}\right) + 16\left(\frac{a_1 a_2 a_3}{1 - a_3 + a_2 a_3} \right. \\ \left. + \frac{a_2 a_3}{a_3 - 1} + a_3\right) - 36.$$

After simplification we get

$$(a_3 - 1)^3 (a_2 - 3) a_1 + (a_1 - 1) a_2 a_3 ((a_3 - 1)(a_2 - 3) + (a_1 - 2)(a_3 - 2)) + a_2 (3a_3 - 2)(a_1 - 1) + a_2 (a_1 - 1) + 12 \geq 0.$$

We also need to show that when $a_1 \geq 3, a_3 \geq 4$, then

$$2a_1 a_3 - a_1 - 3a_3 - 1 \leq \frac{27a_1 a_3^3}{(1 + 2a_3)(a_3 - 1)} + 16\left(\frac{3a_1 a_3}{1 + 2a_3} + \frac{3a_3}{a_3 - 1} + a_3\right) \\ - 7\left(\frac{9a_1 a_3^2}{(1 + 2a_3)(a_3 - 1)} + \frac{3a_1 a_3^2}{1 + 2a_3} + \frac{3a_3^2}{a_3 - 1}\right) - 36.$$

After simplification we get

$$\frac{6a_1 a_3^3}{(1+2a_3)(a_3-1)} + \frac{a_3^2(a_1 a_3 - 3) + 3a_3^3(a_1 - 2)}{(1+2a_3)(a_3-1)} + \frac{17a_1 a_3^2(a_3 - 4)}{(1+2a_3)(a_3-1)} + \frac{5a_1 a_3^2}{(1+2a_3)(a_3-1)} + \frac{41a_1 a_3^2}{(1+2a_3)(a_3-1)} \\ + 16(a_3 - 3) + \frac{48a_3}{(a_3-1)} + 12 \geq 0.$$

Furthermore, we need to show that when $a_2 \geq 4, a_3 \geq 4$, then

$$2a_2 a_3 - 5a_2 - a_3 + 5 \leq \frac{9a_2^2 a_3^3}{(1 - a_3 + a_2 a_3)(a_3 - 1)} \\ - 7\left(\frac{3a_2^2 a_3^2}{(1 - a_3 + a_2 a_3)(a_3 - 1)} + \frac{3a_2 a_3^2}{1 - a_3 + a_2 a_3} + \frac{a_2 a_3^2}{a_3 - 1}\right) + 16\left(\frac{3a_2 a_3}{1 - a_3 + a_2 a_3}\right)$$

$$+ \frac{a_2 a_3}{a_3 - 1} + a_3) - 36.$$

It is easy to see that $a_2 \geq 4, a_3 \geq 4$ can be divided into three cases:

(1) : $a_2 \geq 4, a_3 \geq 5$, (2) : $a_2 \geq 5, a_3 \geq 4$, (3) : $a_2 = 4, a_3 = 4$.

We need to show that when $a_2 \geq 5, a_3 \geq 4$, then

$$\begin{aligned} 2a_2 a_3 - 5a_2 - a_3 + 5 &\leq \frac{9a_2^2 a_3^3}{(1 - a_3 + a_2 a_3)(a_3 - 1)} \\ &- 7\left(\frac{3a_2^2 a_3^2}{(1 - a_3 + a_2 a_3)(a_3 - 1)} + \frac{3a_2 a_3^2}{1 - a_3 + a_2 a_3} + \frac{a_2 a_3^2}{a_3 - 1}\right) + 16\left(\frac{3a_2 a_3}{1 - a_3 + a_2 a_3}\right. \\ &\left. + \frac{a_2 a_3}{a_3 - 1} + a_3\right) - 36. \end{aligned}$$

After simplification we get

$$\frac{a_2 a_3 [a_3 \{(a_2 - 5)a_3 + 5(a_3 - 3)\} + a_2(a_3 - 4)] + 35a_3(a_3 - 2) + 19a_3(a_3 - 1) + 16a_3 + 4a_2(a_3 - 1) + 10a_2 a_3 + 36}{(a_3 - 1)(1 - a_3 + a_2 a_3)} + (a_2 - 5) \geq 0.$$

Next we need to show that when $a_2 \geq 4, a_3 \geq 5$, then

$$\begin{aligned} 2a_2 a_3 - 5a_2 - a_3 + 5 &\leq \frac{9a_2^2 a_3^3}{(1 - a_3 + a_2 a_3)(a_3 - 1)} \\ &- 7\left(\frac{3a_2^2 a_3^2}{(1 - a_3 + a_2 a_3)(a_3 - 1)} + \frac{3a_2 a_3^2}{1 - a_3 + a_2 a_3} + \frac{a_2 a_3^2}{a_3 - 1}\right) + 16\left(\frac{3a_2 a_3}{1 - a_3 + a_2 a_3}\right. \\ &\left. + \frac{a_2 a_3}{a_3 - 1} + a_3\right) - 36. \end{aligned}$$

After simplification we get

$$\frac{(a_3 - 5)a_3^2 [5a_2 - 17] + a_2^2 a_3 [a_3 - 5] + (a_2 - 4)(21a_3 + 4a_2 a_3) + a_2(a_2 a_3^2 - 5) + a_3 [4a_2^2(a_3 - 5) + 2] + 41}{(a_3 - 1)(1 - a_3 + a_2 a_3)} \geq 0.$$

It is also noted that the following inequality

$$\begin{aligned} 2a_2 a_3 - 5a_2 - a_3 + 5 &\leq \frac{9a_2^2 a_3^3}{(1 - a_3 + a_2 a_3)(a_3 - 1)} \\ &- 7\left(\frac{3a_2^2 a_3^2}{(1 - a_3 + a_2 a_3)(a_3 - 1)} + \frac{3a_2 a_3^2}{1 - a_3 + a_2 a_3} + \frac{a_2 a_3^2}{a_3 - 1}\right) + 16\left(\frac{3a_2 a_3}{1 - a_3 + a_2 a_3}\right. \\ &\left. + \frac{a_2 a_3}{a_3 - 1} + a_3\right) - 36, \end{aligned}$$

satisfy for $a_2 = 4, a_3 = 4$.

Q.E.D.

Remark 3.5. For $\text{mult}(f) = 3$, the inequality in Proposition 3.4 is not true when $a_1 = 2, a_2 \geq 3$ and $a_3 \geq 3$.

Proposition 3.5. Let $(V, 0)$ be a fewnomial surface isolated singularity of type 3 which is defined by $f = x_1^{a_1} x_2 + x_2^{a_2} x_3 + x_3^{a_3} x_1$ ($a_1 \geq 2, a_2 \geq 2, a_3 \geq 2$) with weight type

$$\left(\frac{1 - a_3 + a_2 a_3}{1 + a_1 a_2 a_3}, \frac{1 - a_1 + a_1 a_3}{1 + a_1 a_2 a_3}, \frac{1 - a_2 + a_1 a_2}{1 + a_1 a_2 a_3}; 1\right).$$

Then

$$\lambda_1(V) = \begin{cases} 3a_1a_2a_3 + 20(a_1 + a_2 + a_3) - 7(a_1a_2 + a_1a_3 + a_2a_3) \\ -63; \\ a_3 - 2; \end{cases} \quad \begin{matrix} a_1 \geq 3, a_2 \geq 3, a_3 \geq 3 \\ a_1 = 2, a_2 \geq 2, a_3 \geq 2 \end{matrix}$$

Furthermore, assuming that $\text{mult}(f) \geq 4$, we have

$$\begin{aligned} & 3a_1a_2a_3 + 20(a_1 + a_2 + a_3) - 7(a_1a_2 + a_1a_3 + a_2a_3) - 63 \leq \frac{3(1+a_1a_2a_3)^3}{(1-a_3+a_2a_3)(1-a_1+a_1a_3)(1-a_2+a_1a_2)} \\ & + 16\left(\frac{1+a_1a_2a_3}{1-a_3+a_2a_3} + \frac{1+a_1a_2a_3}{1-a_1+a_1a_3} + \frac{1+a_1a_2a_3}{1-a_2+a_1a_2}\right) - 7\left(\frac{(1+a_1a_2a_3)^2}{(1-a_3+a_2a_3)(1-a_1+a_1a_3)} + \frac{(1+a_1a_2a_3)^2}{(1-a_1+a_1a_3)(1-a_2+a_1a_2)}\right. \\ & \left. + \frac{(1+a_1a_2a_3)^2}{(1-a_3+a_2a_3)(1-a_2+a_1a_2)}\right) - 36. \end{aligned}$$

Proof. It is easy to see that the moduli algebra

$$H_1(V) = \mathbb{C}\{x_1, x_2, x_3\} / \left(\frac{\partial^2 f}{\partial x_1^2}, \frac{\partial^2 f}{\partial x_1 \partial x_2}, \frac{\partial^2 f}{\partial x_1 \partial x_3}, \frac{\partial^2 f}{\partial x_2^2}, \frac{\partial^2 f}{\partial x_2^2}, \frac{\partial^2 f}{\partial x_2 \partial x_3} \right)$$

has dimension $a_1a_2a_3 - 2(a_1a_2 + a_1a_3 + a_2a_3) + 5(a_1 + a_2 + a_3) - 14$ and has a monomial basis of the form

$$\{x_1^{i_1} x_2^{i_2} x_3^{i_3}, 0 \leq i_1 \leq a_1 - 3; 0 \leq i_2 \leq a_2 - 3; 0 \leq i_3 \leq a_3 - 3; x_1^{a_1-2} x_3^{i_3}, 0 \leq i_3 \leq a_3 - 3; x_2^{i_2} x_3^{a_3-2}, 0 \leq i_2 \leq a_2 - 3; x_1^{i_1} x_2^{a_2-2}, 0 \leq i_1 \leq a_1 - 3\},$$

and in case of $a_1 = 2, a_2 \geq 2, a_3 \geq 2$ the monomial basis are $\{x_3^{i_3}, 0 \leq i_3 \leq a_3 - 2\}$ with the following relations:

$$x_1^{a_1-2} x_2 = 0, x_2^{a_2-2} x_3 = 0, x_3^{a_3-2} x_1 = 0, x_1^{a_1-1} = 0, x_2^{a_2-1} = 0, x_3^{a_3-1} = 0. \quad (3.23)$$

In order to compute a derivation D of $H_1(V)$ it suffices to indicate its values on the generators x_1, x_2, x_3 which can be written in terms of the basis. Thus we can write

$$\begin{aligned} Dx_j &= \sum_{i_1=0}^{a_1-3} \sum_{i_2=0}^{a_2-3} \sum_{i_3=0}^{a_3-3} c_{i_1, i_2, i_3}^j x_1^{i_1} x_2^{i_2} x_3^{i_3} + \sum_{i_1=0}^{a_1-3} c_{i_1, a_2-2, 0}^j x_1^{i_1} x_2^{a_2-2} + \sum_{i_3=0}^{a_3-3} c_{a_1-2, 0, i_3}^j x_1^{a_1-2} x_3^{i_3} \\ &+ \sum_{i_2=0}^{a_2-3} c_{0, i_2, a_3-2}^j x_2^{i_2} x_3^{a_3-2}, \quad j = 1, 2, 3. \end{aligned}$$

Using the relations (3.23) one easily finds the necessary and sufficient conditions defining a derivation of $H_1(V)$ as follows:

$$c_{0, i_2, i_3}^1 = 0; 0 \leq i_2 \leq a_2 - 4, 0 \leq i_3 \leq a_3 - 3, c_{0, a_2-3, 0}^1 = 0; \quad (3.24)$$

$$c_{i_1, 0, i_3}^2 = 0; 0 \leq i_1 \leq a_1 - 3, 0 \leq i_3 \leq a_3 - 4, c_{0, 0, a_3-3}^2 = 0; \quad (3.25)$$

$$c_{i_1, i_2, 0}^3 = 0; 0 \leq i_1 \leq a_1 - 4, 0 \leq i_2 \leq a_2 - 3, c_{a_1-3, 0, 0}^3 = 0. \quad (3.26)$$

Using (3.24) - (3.26) we obtain the following description of Lie algebras in question:

$$\begin{aligned} & x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_1, 1 \leq i_1 \leq a_1 - 3, 0 \leq i_2 \leq a_2 - 3, 0 \leq i_3 \leq a_3 - 3; x_2^{i_2} x_3^{a_3-2} \partial_1, 1 \leq i_2 \leq a_2 - 4, \\ & x_2^{a_2-3} x_3^{i_3} \partial_1, 1 \leq i_3 \leq a_3 - 2; x_1^{i_1} x_2^{a_2-2} \partial_1, 1 \leq i_1 \leq a_1 - 3; x_1^{a_1-2} x_3^{i_3} \partial_1, 0 \leq i_3 \leq a_3 - 3; \\ & x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_2, 0 \leq i_1 \leq a_1 - 3, 1 \leq i_2 \leq a_2 - 3, 0 \leq i_3 \leq a_3 - 3; x_1^{a_1-2} x_3^{i_3} \partial_2, 0 \leq i_3 \leq a_3 - 3; \end{aligned}$$

$$\begin{aligned}
& x_1^{i_1} x_2^{a_2-2} \partial_2, \quad 1 \leq i_1 \leq a_1 - 3; x_2^{a_2-2} \partial_2; x_1^{i_1} x_3^{a_3-3} \partial_2, \quad 1 \leq i_1 \leq a_1 - 3, \\
& x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_3, \quad 0 \leq i_1 \leq a_1 - 3, 0 \leq i_2 \leq a_2 - 3, 1 \leq i_3 \leq a_3 - 3; \\
& x_2^{a_2-2} \partial_3; x_1^{i_1} x_2^{a_2-2} \partial_3, \quad 1 \leq i_1 \leq a_1 - 3; x_3^{a_3-2} \partial_2; x_3^{a_3-2} \partial_3; x_2^{a_2-2} \partial_1; \\
& x_1^{a_1-3} x_2^{i_2} \partial_3, \quad 1 \leq i_2 \leq a_2 - 3; x_1^{a_1-2} x_3^{i_3} \partial_3, \quad 0 \leq i_3 \leq a_3 - 3; x_3^{a_3-2} \partial_1; \\
& x_2^{i_2} x_3^{a_3-2} \partial_3, \quad 1 \leq i_2 \leq a_2 - 3; x_2^{i_2} x_3^{a_3-2} \partial_2, \quad 1 \leq i_2 \leq a_2 - 3.
\end{aligned}$$

Therefore we have

$$\lambda_1(V) = 3a_1 a_2 a_3 + 20(a_1 + a_2 + a_3) - 7(a_1 a_2 + a_1 a_3 + a_2 a_3) - 63.$$

In case of $a_1 = 2, a_2 \geq 2, a_3 \geq 2$, we obtain the following basis:

$$x_3^{i_3} \partial_3, \quad 1 \leq i_3 \leq a_3 - 2.$$

Furthermore, we need to show that when $a_1 \geq 3, a_2 \geq 3, a_3 \geq 3$, then

$$\begin{aligned}
& 3a_1 a_2 a_3 + 20(a_1 + a_2 + a_3) - 7(a_1 a_2 + a_1 a_3 + a_2 a_3) - 63 \leq \frac{3(1+a_1 a_2 a_3)^3}{(1-a_3+a_2 a_3)(1-a_1+a_1 a_3)(1-a_2+a_1 a_2)} \\
& + 16\left(\frac{1+a_1 a_2 a_3}{1-a_3+a_2 a_3} + \frac{1+a_1 a_2 a_3}{1-a_1+a_1 a_3} + \frac{1+a_1 a_2 a_3}{1-a_2+a_1 a_2}\right) - 7\left(\frac{(1+a_1 a_2 a_3)^2}{(1-a_3+a_2 a_3)(1-a_1+a_1 a_3)} + \frac{(1+a_1 a_2 a_3)^2}{(1-a_1+a_1 a_3)(1-a_2+a_1 a_2)}\right. \\
& \left. + \frac{(1+a_1 a_2 a_3)^2}{(1-a_3+a_2 a_3)(1-a_2+a_1 a_2)}\right) - 36.
\end{aligned}$$

After simplification we get

$$\begin{aligned}
& 8(a_1 a_2 + a_2 a_3 + a_1 a_3) + a_1(a_2 - 3) + a_2(a_3 - 3) + a_3(a_1 - 3) + 6a_1^2[a_2(a_3 - 3) + a_3(a_2 - 3)] \\
& + 6a_2^2[a_1(a_3 - 3) + a_3(a_1 - 3)] + 6a_3^2[a_1(a_2 - 3) + a_2(a_1 - 3)] + 6(a_1^2 + a_2^2 + a_3^2) + 6(a_1^3 a_2 + a_2^3 a_3 \\
& + a_3^3 a_1) + 4a_1^2 a_2^2 a_3^2 + 8(a_1 a_2^2 a_3 + a_1 a_2 a_3^2) + 4a_1^2 a_2 a_3 + 2a_1 a_2 a_3 [3a_1 - 8] + a_1^3 a_2 a_3^2 (a_3 - 3)(a_2 - 3) \\
& + a_1^2 a_3^2 (a_3 - 3)(a_1 a_2 - 2) + 2a_1^2 a_2 a_3^2 (a_3 + a_2 - 6) + 6a_1 a_2 a_3^3 (a_1 - 3) + a_1^2 a_2^3 a_3 (a_3 - 3)(a_1 - 3) \\
& + a_1^2 a_2^2 (a_1 - 3)(a_2 a_3 - 2) + 2a_1^3 a_2 a_3 (a_2 - 3) + 2a_1^2 a_2^2 a_3 (a_1 - 3 + (a_3 - 3)) + a_1 a_2^2 a_3^3 (a_2 - 3)(a_1 \\
& - 3) + a_2^2 a_3^2 (a_2 - 3)(a_1 a_3 - 2) + 14 \geq 0. \quad \text{Q.E.D.}
\end{aligned}$$

Remark 3.6. For $\text{mult}(f) = 3$, the inequality in Proposition 3.5 is not true when $a_1 = 2, a_2 \geq 2$ and $a_3 \geq 3$.

Proposition 3.6. Let $(V, 0)$ be a fewnomial surface isolated singularity of type 4 which is defined by $f = x_1^{a_1} + x_2^{a_2} + x_3^{a_3} x_1$ ($a_1 \geq 3, a_2 \geq 3, a_3 \geq 2$) with weight type $(\frac{1}{a_1}, \frac{1}{a_2}, \frac{a_1-1}{a_1 a_3}; 1)$. Then

$$\lambda_1(V) = \begin{cases} 3a_1 a_2 a_3 + 16(a_1 + a_3) + 20a_2 - 7(a_1 a_2 + a_1 a_3 + a_2 a_3) & a_1 \geq 4, a_2 \geq 3, a_3 \geq 3 \\ -45; & a_1 = 3, a_2 \geq 3, a_3 \geq 2 \\ 2a_2 a_3 - 3a_2 - 5a_3 + 7; & a_1 \geq 4, a_2 \geq 3, a_3 = 2 \\ a_2 - 3; & \end{cases}$$

Furthermore, assuming that $\text{mult}(f) \geq 4$, we have

$$\begin{aligned}
& 3a_1 a_2 a_3 + 16(a_1 + a_3) + 20a_2 - 7(a_1 a_2 + a_1 a_3 + a_2 a_3) - 45 \leq \frac{3a_1^2 a_2 a_3}{a_1 - 1} + 16(a_1 + a_2 + \frac{a_1 a_3}{a_1 - 1}) \\
& - 7(a_1 a_2 + \frac{a_1 a_2 a_3}{a_1 - 1} + \frac{a_1^2 a_3}{a_1 - 1}) - 36.
\end{aligned}$$

Proof. It is easy to see that the moduli algebra

$$H_1(V) = \mathbb{C}\{x_1, x_2, x_3\} / \left(\frac{\partial^2 f}{\partial x_1^2}, \frac{\partial^2 f}{\partial x_1 \partial x_2}, \frac{\partial^2 f}{\partial x_1 \partial x_3}, \frac{\partial^2 f}{\partial x_2^2}, \frac{\partial^2 f}{\partial x_3^2}, \frac{\partial^2 f}{\partial x_2 \partial x_3} \right)$$

has dimension $a_1 a_2 a_3 - 2(a_1 a_2 + a_1 a_3 + a_2 a_3) + 4(a_1 + a_3) + 5a_2 - 10$ and has a monomial basis of the form

$$\{x_1^{i_1} x_2^{i_2} x_3^{i_3}, 0 \leq i_1 \leq a_1 - 3; 0 \leq i_2 \leq a_2 - 3; 0 \leq i_3 \leq a_3 - 3; x_2^{i_2} x_3^{a_3 - 2}, 0 \leq i_2 \leq a_2 - 3\},$$

with the following relations:

$$x_1^{a_1 - 2} = 0, x_2^{a_2 - 2} = 0, x_3^{a_3 - 2} = 0, x_3^{a_3 - 2} x_1 = 0. \quad (3.27)$$

In order to compute a derivation D of $H_1(V)$ it suffices to indicate its values on the generators x_1, x_2, x_3 which can be written in terms of the basis. Thus we can write

$$Dx_j = \sum_{i_1=0}^{a_1-3} \sum_{i_2=0}^{a_2-3} \sum_{i_3=0}^{a_3-3} c_{i_1, i_2, i_3}^j x_1^{i_1} x_2^{i_2} x_3^{i_3} + \sum_{i_2=0}^{a_2-3} c_{0, i_2, a_3-2}^j x_2^{i_2} x_3^{a_3-2}, \quad j = 1, 2, 3.$$

Using the relations (3.27) one easily finds the necessary and sufficient conditions defining a derivation of $H_1(V)$ as follows:

$$c_{0, i_2, i_3}^1 = 0; 0 \leq i_2 \leq a_2 - 3; 0 \leq i_3 \leq a_3 - 3, \quad (3.28)$$

$$c_{i_1, 0, i_3}^2 = 0; 0 \leq i_1 \leq a_1 - 3; 0 \leq i_3 \leq a_3 - 3, \quad (3.29)$$

$$c_{i_1, i_2, 0}^3 = 0; 0 \leq i_1 \leq a_1 - 4; 0 \leq i_2 \leq a_2 - 3. \quad (3.30)$$

Using (3.28) - (3.30) we obtain the following description of Lie algebras in question:

$$\begin{aligned} & x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_1, \quad 1 \leq i_1 \leq a_1 - 3, 0 \leq i_2 \leq a_2 - 3, 0 \leq i_3 \leq a_3 - 3; x_2^{i_2} x_3^{a_3 - 2} \partial_1, \quad 0 \leq i_2 \leq a_2 - 3, \\ & x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_2, \quad 0 \leq i_1 \leq a_1 - 3, 1 \leq i_2 \leq a_2 - 3, 0 \leq i_3 \leq a_3 - 3; x_2^{i_2} x_3^{a_3 - 2} \partial_2, \quad 1 \leq i_2 \leq a_2 - 3, \\ & x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_3, \quad 0 \leq i_1 \leq a_1 - 3, 0 \leq i_2 \leq a_2 - 3, 1 \leq i_3 \leq a_3 - 3, x_2^{i_2} x_3^{a_3 - 2} \partial_3, \quad 0 \leq i_2 \leq a_2 - 3, \\ & x_1^{a_1 - 3} x_2^{i_2} \partial_3, \quad 0 \leq i_2 \leq a_2 - 3. \end{aligned}$$

Therefore we have

$$\lambda_1(V) = 3a_1 a_2 a_3 + 16(a_1 + a_3) + 20a_2 - 7(a_1 a_2 + a_1 a_3 + a_2 a_3) - 45.$$

In case of $a_1 = 3, a_2 \geq 3, a_3 \geq 2$ we obtain we following basis:

$$x_2^{i_2} x_3^{i_3} \partial_2, \quad 1 \leq i_2 \leq a_2 - 3, 0 \leq i_3 \leq a_3 - 2; x_2^{i_2} x_3^{i_3} \partial_3, \quad 0 \leq i_2 \leq a_2 - 3, 1 \leq i_3 \leq a_3 - 2.$$

Therefore we have

$$\lambda_1(V) = 2a_2 a_3 - 3a_2 - 5a_3 + 7.$$

In case of $a_1 \geq 4, a_2 \geq 3, a_3 = 2$ we obtain we following basis:

$$x_2^{i_2} \partial_2, \quad 1 \leq i_2 \leq a_2 - 3.$$

Furthermore, we need to show that when $a_1 \geq 4, a_2 \geq 4, a_3 \geq 3$, then

$$\begin{aligned} & 3a_1 a_2 a_3 + 16(a_1 + a_3) + 20a_2 - 7(a_1 a_2 + a_1 a_3 + a_2 a_3) - 45 \leq \frac{3a_1^2 a_2 a_3}{a_1 - 1} + 16(a_1 + a_2 + \frac{a_1 a_3}{a_1 - 1}) \\ & - 7(a_1 a_2 + \frac{a_1 a_2 a_3}{a_1 - 1} + \frac{a_1^2 a_3}{a_1 - 1}) - 36. \end{aligned}$$

After solving above inequality, we get

$$\frac{a_2 a_3 (2a_1 - 7)}{a_1 - 1} + 5a_1 a_3 + a_3 (a_1 - 1) + \frac{15a_3}{a_1 - 1} + \frac{a_2 [a_1 (a_3 - 4) + 4]}{a_1 - 1} \geq 0.$$

Q.E.D.

Remark 3.7. For $\text{mult}(f) = 3$, the inequality in Proposition 3.6 is not true when $a_1 \geq 5$, $a_2 \geq 5$ and $a_3 = 2$.

Proposition 3.7. Let $(V, 0)$ be a fewnomial surface isolated singularity of type 5 which is defined by $f = x_1^{a_1}x_2 + x_2^{a_2}x_1 + x_3^{a_3}$ ($a_1 \geq 2, a_2 \geq 2, a_3 \geq 3$) with weight type $(\frac{a_2-1}{a_1a_2-1}, \frac{a_1-1}{a_1a_2-1}, \frac{1}{a_3}; 1)$. Then

$$\lambda_1(V) = \begin{cases} 3a_1a_2a_3 + 16(a_1 + a_2) + 26a_3 - 7(a_1a_2 + a_1a_3 + a_2a_3) - 59; & a_1 \geq 5, a_2 \geq 5, a_3 \geq 3 \\ a_3 - 3; & a_1 = 2, a_2 \geq 2, a_3 \geq 3 \\ 2a_2a_3 - 5a_2 + 2a_3 - 5; & a_1 = 3, a_2 \geq 3, a_3 \geq 3 \\ 6a_2a_3 - 14a_2 - 8a_3 + 17; & a_1 = 4, a_2 \geq 4, a_3 \geq 3 \end{cases}$$

Furthermore, assuming that $\text{mult}(f) \geq 4$, we have

$$3a_1a_2a_3 + 16(a_1 + a_2) + 26a_3 - 7(a_1a_2 + a_1a_3 + a_2a_3) - 59 \leq \frac{3a_3(a_1a_2-1)^2}{(a_2-1)(a_1-1)} + 16\left(\frac{a_1a_2-1}{a_2-1} + \frac{a_1a_2-1}{a_1-1} + a_3\right) - 7\left(\frac{(a_1a_2-1)^2}{(a_2-1)(a_1-1)} + \frac{a_3(a_1a_2-1)}{a_1-1} + \frac{a_3(a_1a_2-1)}{a_2-1}\right) - 36.$$

Proof. It is easy to see that the moduli algebra

$$H_1(V) = \mathbb{C}\{x_1, x_2, x_3\} / \left(\frac{\partial^2 f}{\partial x_1^2}, \frac{\partial^2 f}{\partial x_1 \partial x_2}, \frac{\partial^2 f}{\partial x_1 \partial x_3}, \frac{\partial^2 f}{\partial x_2^2}, \frac{\partial^2 f}{\partial x_3^2}, \frac{\partial^2 f}{\partial x_2 \partial x_3} \right)$$

has dimension $a_1a_2a_3 - 2(a_1a_2 + a_1a_3 + a_2a_3) + 4(a_1 + a_2) + 7a_3 - 14$ and has a monomial basis of the form

$$\{x_1^{i_1}x_2^{i_2}x_3^{i_3}, 0 \leq i_1 \leq a_1 - 3; 0 \leq i_2 \leq a_2 - 3; 0 \leq i_3 \leq a_3 - 3; x_1^{a_1-2}x_3^{i_3}, 0 \leq i_3 \leq a_3 - 3; x_2^{i_2}x_3^{i_3}, a_2 - 2 \leq i_2 \leq a_2 - 1; 0 \leq i_3 \leq a_3 - 3\},$$

with the following relations:

$$x_1^{a_1-2}x_2 = 0, x_2^{a_2-2}x_1 = 0, x_3^{a_3-2} = 0, a_1x_1^{a_1-1} + a_2x_2^{a_2-1} = 0. \quad (3.31)$$

In order to compute a derivation D of $H_1(V)$ it suffices to indicate its values on the generators x_1, x_2, x_3 which can be written in terms of the basis. Thus we can write

$$Dx_j = \sum_{i_1=0}^{a_1-3} \sum_{i_2=0}^{a_2-3} \sum_{i_3=0}^{a_3-3} c_{i_1, i_2, i_3}^j x_1^{i_1} x_2^{i_2} x_3^{i_3} + \sum_{i_3=0}^{a_3-3} c_{a_1-2, 0, i_3}^j x_1^{a_1-2} x_3^{i_3} + \sum_{i_2=a_2-2}^{a_2-1} \sum_{i_3=0}^{a_3-3} c_{0, i_2, i_3}^j x_2^{i_2} x_3^{i_3},$$

$$j = 1, 2, 3.$$

Using the relations (3.31) one easily finds the necessary and sufficient conditions defining a derivation of $H_1(V)$ as follows:

$$c_{0, i_2, i_3}^1 = 0; 0 \leq i_2 \leq a_2 - 4, 0 \leq i_3 \leq a_3 - 3; \quad (3.32)$$

$$c_{i_1, 0, i_3}^2 = 0; 0 \leq i_1 \leq a_1 - 4, 0 \leq i_3 \leq a_3 - 3; \quad (3.33)$$

$$c_{i_1, i_2, 0}^3 = 0; 0 \leq i_1 \leq a_1 - 3, 0 \leq i_2 \leq a_2 - 3; c_{a_1-2, 0, 0}^3 = 0; \quad (3.34)$$

$$c_{0, i_2, 0}^3 = 0; a_2 - 2 \leq i_2 \leq a_2 - 1; -a_2(a_1 - 1)c_{1, 0, i_3}^1 + a_2(a_2 - 1)c_{0, 1, i_3}^2 = 0; 0 \leq i_3 \leq a_3 - 3. \quad (3.35)$$

Using (3.32) - (3.34) we obtain the following description of Lie algebras in question:

$$x_1^{i_1}x_2^{i_2}x_3^{i_3}\partial_1, 2 \leq i_1 \leq a_1 - 3, 0 \leq i_2 \leq a_2 - 3, 0 \leq i_3 \leq a_3 - 3; x_1^{a_1-2}x_3^{i_3}\partial_1, 0 \leq i_3 \leq a_3 - 3,$$

$$\begin{aligned}
& x_1 x_3^{i_3} \partial_1 + \frac{a_1 - 1}{a_2 - 1} x_2 x_3^{i_3} \partial_2, \quad 0 \leq i_3 \leq a_3 - 3; x_2^2 x_3^{i_3} \partial_1, \quad a_2 - 3 \leq i_2 \leq a_2 - 1, 0 \leq i_3 \leq a_3 - 3, \\
& x_1 x_2^{i_2} x_3^{i_3} \partial_1, \quad 1 \leq i_2 \leq a_2 - 3, 0 \leq i_3 \leq a_3 - 3; x_2^2 x_3^{i_3} \partial_2, \quad 2 \leq i_2 \leq a_2 - 1, 0 \leq i_3 \leq a_3 - 3; \\
& x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_2, \quad 1 \leq i_1 \leq a_1 - 3, 1 \leq i_2 \leq a_2 - 3, 0 \leq i_3 \leq a_3 - 3; x_1^{a_1 - 2} x_3^{i_3} \partial_3, \quad 1 \leq i_3 \leq a_3 - 3, \\
& x_1^{i_1} x_3^{i_3} \partial_2, \quad a_1 - 3 \leq i_1 \leq a_1 - 4, 0 \leq i_3 \leq a_3 - 3; x_2^{i_2} x_3^{i_3} \partial_3, \quad 0 \leq i_2 \leq a_2 - 1, 1 \leq i_3 \leq a_3 - 3; \\
& x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_3, \quad 1 \leq i_1 \leq a_1 - 3, 0 \leq i_2 \leq a_2 - 3, 1 \leq i_3 \leq a_3 - 3.
\end{aligned}$$

Therefore we have

$$\lambda_1(V) = 3a_1 a_2 a_3 + 16(a_1 + a_2) + 26a_3 - 7(a_1 a_2 + a_1 a_3 + a_2 a_3) - 59.$$

In case of $a_1 = 4, a_2 \geq 4, a_3 \geq 3$, we obtain the following basis:

$$\begin{aligned}
& x_2^{i_2} x_3^{i_3} \partial_1, \quad 2 \leq i_2 \leq a_2 - 1, 0 \leq i_3 \leq a_3 - 3; x_1^2 x_3^{i_3} \partial_1, \quad 0 \leq i_3 \leq a_3 - 3, \\
& x_1^2 x_3^{i_3} \partial_2, \quad 0 \leq i_3 \leq a_3 - 3; x_1 x_2^{i_2} x_3^{i_3} \partial_1, \quad 0 \leq i_2 \leq a_2 - 3, 0 \leq i_3 \leq a_3 - 3, \\
& x_2^{i_2} x_3^{i_3} \partial_2, \quad 2 \leq i_2 \leq a_2 - 1, 0 \leq i_3 \leq a_3 - 3; x_1 x_2^{i_2} x_3^{i_3} \partial_2, \quad 1 \leq i_2 \leq a_2 - 3, 0 \leq i_3 \leq a_3 - 3, \\
& x_2^{i_2} x_3^{i_3} \partial_3, \quad 0 \leq i_2 \leq a_2 - 1, 1 \leq i_3 \leq a_3 - 3; x_1 x_2^{i_2} x_3^{i_3} \partial_3, \quad 0 \leq i_2 \leq a_2 - 3, 1 \leq i_3 \leq a_3 - 3, \\
& x_1^2 x_3^{i_3} \partial_3, \quad 1 \leq i_3 \leq a_3 - 3.
\end{aligned}$$

We have

$$\lambda_1(V) = 6a_2 a_3 - 14a_2 - 8a_3 + 17.$$

In case of $a_1 = 3, a_2 \geq 3, a_3 \geq 3$, we obtain the following basis:

$$\begin{aligned}
& x_2^{i_2} x_3^{i_3} \partial_1, \quad a_2 - 2 \leq i_2 \leq a_2 - 1, 0 \leq i_3 \leq a_3 - 3; x_1 x_3^{i_3} \partial_1, \quad 0 \leq i_3 \leq a_3 - 3, \\
& x_1 x_3^{i_3} \partial_3, \quad 1 \leq i_3 \leq a_3 - 3; x_2^{i_2} x_3^{i_3} \partial_2, \quad 2 \leq i_2 \leq a_2 - 1, 0 \leq i_3 \leq a_3 - 3, \\
& x_2^{i_2} x_3^{i_3} \partial_3, \quad 0 \leq i_2 \leq a_2 - 1, 1 \leq i_3 \leq a_3 - 3.
\end{aligned}$$

We have

$$\lambda_1(V) = 2a_2 a_3 - 5a_2 + 2a_3 - 5.$$

In case of $a_1 = 2, a_2 \geq 2, a_3 \geq 3$, we obtain the following basis:

$$x_3^{i_3} \partial_3, \quad 1 \leq i_3 \leq a_3 - 3.$$

We have

$$\lambda_1(V) = a_3 - 3.$$

Next, we need to show that when $a_1 \geq 5, a_2 \geq 5, a_3 \geq 4$, then

$$\begin{aligned}
& 3a_1 a_2 a_3 + 16(a_1 + a_2) + 26a_3 - 7(a_1 a_2 + a_1 a_3 + a_2 a_3) - 59 \leq \frac{3a_3(a_1 a_2 - 1)^2}{(a_2 - 1)(a_1 - 1)} + 16\left(\frac{a_1 a_2 - 1}{a_2 - 1} + \frac{a_1 a_2 - 1}{a_1 - 1}\right. \\
& \left. + a_3\right) - 7\left(\frac{(a_1 a_2 - 1)^2}{(a_2 - 1)(a_1 - 1)} + \frac{a_3(a_1 a_2 - 1)}{a_1 - 1} + \frac{a_3(a_1 a_2 - 1)}{a_2 - 1}\right) - 36.
\end{aligned}$$

After simplification, we get

$$\begin{aligned}
& a_1(a_1 - 5)(a_2 - 1)(a_3 + (a_1 - 1)a_2(a_2 - 5)a_3) + 2a_1^2(a_3 - 2)(a_2 - 3) + a_2^2 a_1 + 11a_1(a_2 - 5) \\
& + 11a_2(a_1 - 5) + 7a_3(a_1 - 3) + 22a_1 a_2 + 17a_1 a_3 + 6a_2 a_3 + 48 + 54a_2 + a_1 a_2(a_1 - 5) \\
& + 2(a_1 - 2)a_2(a_2 - 5)(a_3 - 3) + (a_1 - 2)(a_3 - 3) \geq 0.
\end{aligned}$$

Now we need to show that when $a_1 = 4, a_2 \geq 4, a_3 \geq 4$, then

$$6a_2a_3 - 14a_2 - 8a_3 + 17 \leq \frac{a_3(4a_2 - 1)^2}{(a_2 - 1)} + 16\left(\frac{4a_2 - 1}{a_2 - 1} + \frac{4a_2 - 1}{3} + a_3\right) - 7\left(\frac{(4a_2 - 1)^2}{3(a_2 - 1)} + \frac{a_3(4a_2 - 1)}{3} + \frac{a_3(4a_2 - 1)}{a_2 - 1}\right) - 36.$$

After simplification, we get

$$\frac{(4a_2-1)(3a_3-7)}{3(a_2-1)} + \frac{7a_2a_3+2a_2(a_3-5)+42(a_3-3)+14}{3} + \frac{(4a_2-1)(a_2(3a_3-14)+2a_2a_3+a_3)}{3(a_2-1)} \geq 0.$$

It is also easy to check that proposed conjecture also true for

$$a_1 = 3, a_2 \geq 3, a_3 \geq 4.$$

Q.E.D.

Remark 3.8. For $\text{mult}(f) = 3$, the inequality in Proposition 3.7 is not true when $a_1 = 2, a_2 \geq 3$ and $a_3 \geq 3$.

Proof of Main Theorem A.

Proof. It is an immediate corollary of Proposition 3.1.

Q.E.D.

Proof of Main Theorem B.

Proof. Let $f \in \mathbb{C}\{x_1, x_2\}$ be a weighted homogeneous fewnomial isolated singularity. Then f can be divided into the following three types:

Type A. $x_1^{a_1} + x_2^{a_2}$,

Type B. $x_1^{a_1}x_2 + x_2^{a_2}$,

Type C. $x_1^{a_1}x_2 + x_2^{a_2}x_1$.

The Main Theorem B is an immediate corollary of Remark 3.1, Proposition 3.2, and Proposition 3.3.

Q.E.D.

Proof of Main Theorem C.

Proof. Let $f \in \mathbb{C}\{x_1, x_2, x_3\}$ be a weighted homogeneous fewnomial isolated surface singularity. Then f can be divided into the following five types:

Type 1. $x_1^{a_1} + x_2^{a_2} + x_3^{a_3}$,

Type 2. $x_1^{a_1}x_2 + x_2^{a_2}x_3 + x_3^{a_3}$,

Type 3. $x_1^{a_1}x_2 + x_2^{a_2}x_3 + x_3^{a_3}x_1$,

Type 4. $x_1^{a_1} + x_2^{a_2} + x_3^{a_3}x_1$,

Type 5. $x_1^{a_1}x_2 + x_2^{a_2}x_1 + x_3^{a_3}$.

The Main Theorem C is an immediate corollary of Remark 3.4, Propositions 3.4, 3.5, 3.6, and 3.7.

Q.E.D.

Proof of Main Theorem D.

Proof. It is easy to see, from Remarks 1.2, 3.1, Propositions 3.2-3.3, Propositions 4.1-4.3 in [YZ2], Propositions 3.1-3.3 in [HYZ3], and Propositions 3.2-3.4 in [HYZ2], the inequality $\lambda^2 > \lambda^1 > \lambda^0 = \lambda_2 > \lambda_1$ holds true.

Q.E.D.

Proof of Main Theorem E.

Proof. It is easy to see, from Remarks 1.2, 3.4, Propositions 3.4-3.7, Proposition 4.1 in [YZ2], Propositions 3.1, 3.2 in [HYZ1], Propositions 3.4, 3.5 in [HYZ3], Propositions 3.5-3.9 in [HYZ2], and Propositions 3.4-3.8 in [HYZ5], the inequality $\lambda^2 > \lambda^1 > \lambda^0 = \lambda_3 > \lambda_1$ holds true.

Q.E.D.

Proof of Main Theorem F.

Proof.

$$\begin{aligned}
 L_1(V_{f+g}) &= \text{Der}\mathbb{C}\{x_1, \dots, x_n, y_1, \dots, y_m\} / \left(\frac{\partial^2(f+g)}{\partial x_1^2}, \dots, \frac{\partial^2(f+g)}{\partial y_m^2}, \right. \\
 &\quad \frac{\partial^2(f+g)}{\partial x_1 \partial x_2}, \dots, \frac{\partial^2(f+g)}{\partial x_1 \partial y_m}, \\
 &\quad \frac{\partial^2(f+g)}{\partial x_2 \partial x_3}, \dots, \frac{\partial^2(f+g)}{\partial x_2 \partial y_m}, \\
 &\quad \vdots \\
 &\quad \left. \frac{\partial^2(f+g)}{\partial y_{m-1} \partial y_m} \right) \\
 &= \text{Der}\mathbb{C}\{x_1, \dots, x_n, y_1, \dots, y_m\} / \left(\frac{\partial^2 f}{\partial x_1^2}, \dots, \frac{\partial^2 f}{\partial x_n^2}, \right. \\
 &\quad \frac{\partial^2 f}{\partial x_1 \partial x_2}, \dots, \frac{\partial^2 f}{\partial x_1 \partial x_n}, \\
 &\quad \frac{\partial^2 f}{\partial x_2 \partial x_3}, \dots, \frac{\partial^2 f}{\partial x_2 \partial x_n}, \\
 &\quad \vdots \\
 &\quad \frac{\partial^2 f}{\partial x_{n-1} \partial x_n}, \\
 &\quad \frac{\partial^2 g}{\partial x_1^2}, \dots, \frac{\partial^2 g}{\partial y_m^2}, \\
 &\quad \frac{\partial^2 g}{\partial x_1 \partial x_2}, \dots, \frac{\partial^2 g}{\partial x_1 \partial y_m}, \\
 &\quad \frac{\partial^2 g}{\partial x_2 \partial x_3}, \dots, \frac{\partial^2 g}{\partial x_2 \partial y_m}, \\
 &\quad \vdots \\
 &\quad \left. \frac{\partial^2 g}{\partial y_{m-1} \partial y_m} \right) \\
 &= \text{Der}[\mathbb{C}\{x_1, \dots, x_n\} / \left(\frac{\partial^2 f}{\partial x_1^2}, \dots, \frac{\partial^2 f}{\partial x_n^2}, \right. \\
 &\quad \left. \frac{\partial^2 f}{\partial x_1 \partial x_2}, \dots, \frac{\partial^2 f}{\partial x_1 \partial x_n}, \right)
 \end{aligned}$$

$$\begin{aligned}
& \left. \begin{aligned} & \frac{\partial^2 f}{\partial x_2 \partial x_3}, \dots, \frac{\partial^2 f}{\partial x_2 \partial x_n}, \\ & \vdots \\ & \frac{\partial^2 f}{\partial x_{n-1} \partial x_n} \end{aligned} \right) \\
& \otimes \mathbb{C}\{y_1, \dots, y_m\} \left/ \left(\begin{aligned} & \frac{\partial^2 g}{\partial y_1^2}, \dots, \frac{\partial^2 g}{\partial y_m^2}, \\ & \frac{\partial^2 g}{\partial y_1 \partial y_2}, \dots, \frac{\partial^2 g}{\partial y_1 \partial y_m}, \\ & \frac{\partial^2 g}{\partial y_2 \partial y_3}, \dots, \frac{\partial^2 g}{\partial y_2 \partial y_m}, \\ & \vdots \\ & \frac{\partial^2 g}{\partial y_{m-1} \partial y_m} \end{aligned} \right) \right] \\
= & \text{Der}[\mathbb{C}\{x_1, \dots, x_n\} \left/ \left(\begin{aligned} & \frac{\partial^2 f}{\partial x_1^2}, \dots, \frac{\partial^2 f}{\partial x_n^2}, \\ & \frac{\partial^2 f}{\partial x_1 \partial x_2}, \dots, \frac{\partial^2 f}{\partial x_1 \partial x_n}, \\ & \frac{\partial^2 f}{\partial x_2 \partial x_3}, \dots, \frac{\partial^2 f}{\partial x_2 \partial x_n}, \\ & \vdots \\ & \frac{\partial^2 f}{\partial x_{n-1} \partial x_n} \end{aligned} \right) \right] \\
& \otimes \mathbb{C}\{y_1, \dots, y_m\} \left/ \left(\begin{aligned} & \frac{\partial^2 g}{\partial y_1^2}, \dots, \frac{\partial^2 g}{\partial y_m^2}, \\ & \frac{\partial^2 g}{\partial y_1 \partial y_2}, \dots, \frac{\partial^2 g}{\partial y_1 \partial y_m}, \\ & \frac{\partial^2 g}{\partial y_2 \partial y_3}, \dots, \frac{\partial^2 g}{\partial y_2 \partial y_m}, \\ & \vdots \\ & \frac{\partial^2 g}{\partial y_{m-1} \partial y_m} \end{aligned} \right) \right) \\
& + \mathbb{C}\{x_1, \dots, x_n\} \left/ \left(\begin{aligned} & \frac{\partial^2 f}{\partial x_1^2}, \dots, \frac{\partial^2 f}{\partial x_n^2}, \\ & \frac{\partial^2 f}{\partial x_1 \partial x_2}, \dots, \frac{\partial^2 f}{\partial x_1 \partial x_n}, \end{aligned} \right) \right)
\end{aligned}$$

$$\begin{aligned} & \left(\frac{\partial^2 f}{\partial x_2 \partial x_3}, \dots, \frac{\partial^2 f}{\partial x_2 \partial x_n}, \right. \\ & \quad \vdots \\ & \quad \left. \frac{\partial^2 f}{\partial x_{n-1} \partial x_n} \right) \\ \otimes \operatorname{Der}[\mathbb{C}\{y_1, \dots, y_m\} / & \left(\frac{\partial^2 g}{\partial y_1^2}, \dots, \frac{\partial^2 g}{\partial y_m^2}, \right. \\ & \quad \frac{\partial^2 g}{\partial y_1 \partial y_2}, \dots, \frac{\partial^2 g}{\partial y_1 \partial y_m}, \\ & \quad \frac{\partial^2 g}{\partial y_2 \partial y_3}, \dots, \frac{\partial^2 g}{\partial y_2 \partial y_m}, \\ & \quad \vdots \\ & \quad \left. \frac{\partial^2 g}{\partial y_{m-1} \partial y_m} \right)] \\ = & L_1(V_f) \otimes H_1(V_g) + H_1(V_f) \otimes L_1(V_g). \end{aligned}$$

The first equality above comes from the fact that f, g are weighted homogeneous while the fourth equality follows from the Theorem 2.1. Q.E.D.

REFERENCES

[AM] A. G. Aleksandrov and B. Martin, *Derivations and deformations of Artin algebras*, Beitrage Zur Algebra und Geometrie 33 (1992), 115-130.

[AVZ] V. Arnold, A. Varchenko, and S. Gusein-Zade, *Singularities of differentiable mappings*, 2nd ed. (Russian) MCNMO, Moskva, 2004.

[BY] M. Benson and S. S.-T. Yau, *Lie algebra and their representations arising from isolated singularities: Computer method in calculating the Lie algebras and their cohomology*, Advance studies in Pure Mathematics 8, Complex Analytic Singularities, (1986), 3-58.

[Bl] R. Block, *Determination of the differentially simple rings with a minimal ideal*, Ann. Math. 90 (1969), 433-459.

[CCYZ] B. Chen, H. Chen, S. S.-T. Yau, and H. Zuo, *The non-existence of negative weight derivations on positive dimensional isolated singularities: generalized Wahl conjecture*, J. Differential Geom. 115(2) (2020), 195-224.

[CHYZ] B. Chen, N. Hussain, S. S.-T. Yau, and H. Zuo, *Variation of complex structures and variation of Lie algebras II: New Lie algebras arising from singularities*, J. Differential Geom. 115(3) (2020), 437-473.

[CYZ] H. Chen, S. S.-T. Yau, and H. Zuo, *Non-existence of negative weight derivations on positively graded Artinian algebras*, Trans. Amer. Math. Soc. 372(4) (2019), 2493-253.

[DS] A. Dimca and G. Sticlaru, *Hessian ideals of a homogeneous polynomial and generalized Tjurina algebras*, Documenta Math. 20 (2015), 689-705.

[EK] A. Elashvili and G. Khimshiashvili, *Lie algebras of simple hypersurface singularities*, J. Lie Theory 16(4) (2006), 621-649.

[ET] W. Ebeling and A. Takahashi, *Strange duality of weighted homogeneous polynomial*, J. Compositio Math. 147 (2011), 1413-1433.

[Hu] N. Hussain, *Survey on derivation Lie algebras of isolated singularities*, Methods Appl. Anal. 25(4) (2018), 307-322.

- [HYZ1] N. Hussain, S. S.-T. Yau, and H. Zuo, *On the derivation Lie algebras of fewnomial singularities*, Bull. Aust. Math. Soc. 98(1) (2018), 77-88.
- [HYZ2] N. Hussain, S. S.-T. Yau, and H. Zuo, *On the new k -th Yau algebras of isolated hypersurface singularities*, Math. Z. 294(1-2) (2020), 331-358.
- [HYZ3] N. Hussain, S. S.-T. Yau, and H. Zuo, *k -th Yau number of isolated hypersurface singularities and an inequality conjecture*, to appear, J. Aust. Math. Soc., published online: April 2019, doi:10.1017/S1446788719000132.
- [HYZ4] N. Hussain, S. S.-T. Yau, and H. Q. Zuo, *Generalized Cartan matrices arising from new derivation Lie algebras of isolated hypersurface singularities*, Pacific. J. Math. 305(1) (2020), 189-218.
- [HYZ5] N. Hussain, S. S.-T. Yau, and H. Zuo, *On two Inequality conjectures for the k -th Yau numbers of isolated hypersurface singularities*, to appear, Geometriae Dedicata, published online: July 2020, <https://doi.org/10.1007/s10711-020-00549-z>
- [HYZ6] N. Hussain, S. S.-T. Yau, and H. Zuo, *On The Generalized Cartan matrices arising from k -th Yau algebras of isolated hypersurface singularities*, 25pp. in ms, to appear, Algebras and Representation Theory.
- [HYZ7] N. Hussain, S. S.-T. Yau, and H. Zuo, *New k -th Yau algebras of isolated hypersurface singularities and weak Torelli-type theorem*, 26pp. in ms., to appear, Math. Research Lett.
- [Kh] A. Khovanski, *fewnomials*, American Mathematical Society, Providence, RI, 1991. Translated from the Russian by Smilka Zdravkovska.
- [Khi] G. Khimshiashvili, *yau algebras of fewnomial singularities*, preprint. <http://www.math.uu.nl/publications/preprints/1352.pdf>
- [MO] J. Milnor and P. Orlik, *Isolated singularities defined by weighted homogeneous polynomials*, Topology 9 (1970), 385-393.
- [MY] J. Mather and S. S.-T. Yau, *Classification of isolated hypersurface singularities by their moduli algebras*, Invent. Math. 69 (1982), 243-251.
- [Sa] K. Saito, *Quasihomogene isolierte Singularitäten von Hyperflächen*, Invent. Math. 14 (1971), 123-142.
- [Sae] O. Saeki, *Topological invariance of weights for weighted homogeneous isolated singularities in \mathbb{C}^3* , Proc. A.M.S. 103(3) (1988), 905-909.
- [SY] C. Seeley and S. S.-T. Yau, *Variation of complex structure and variation of Lie algebras*, Invent. Math. 99 (1990), 545-565.
- [XY] Y.-J. Xu and S. S.-T. Yau, *Micro-local characterization quasi-homogeneous singularities*, Amer. J. Math. 118(2) (1996), 389-399.
- [Ya1] S. S.-T. Yau, *Continuous family of finite-dimensional representations of a solvable Lie algebra arising from singularities*, Proc. Natl. Acad. Sci. USA 80 (1983), 7694-7696.
- [Ya2] S. S.-T. Yau, *Solvable Lie algebras and generalized Cartan matrices arising from isolated singularities*, Math. Z. 191 (1986), 489-506.
- [Ya3] S. S.-T. Yau, *Solvability of Lie algebras arising from isolated singularities and nonisolatedness of singularities defined by $sl(2, \mathbb{C})$ invariant polynomials*, Amer. J. Math. 113 (1991), 773-778.
- [YS] E. Yoshinaga and M. Suzuki, *Topological types of quasihomogeneous singularities in \mathbb{C}^2* , Topology 18(2) (1979), 113-116.
- [Yu] Y. Yu, *On Jacobian ideals invariant by reducible $sl(2; \mathbb{C})$ action*, Trans. Amer. Math. Soc. 348 (1996), 2759-2791.
- [YZ1] S. S.-T. Yau and H. Zuo, *Derivations of the moduli algebras of weighted homogeneous hypersurface singularities*, J. Algebra 457 (2016), 18-25.
- [YZ2] S. S.-T. Yau and H. Zuo, *A Sharp upper estimate conjecture for the Yau number of weighted homogeneous isolated hypersurface singularity*, Pure Appl. Math. Q. 12(1) (2016), 165-181.

SCHOOL OF DATA SCIENCES, HUASHANG COLLEGE GUANGDONG UNIVERSITY OF FINANCE AND ECONOMICS, GUANGZHOU GUANGDONG, 511300 P. R. CHINA

E-mail address: naveed1573@gmail.com

DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING, 100084, P. R. CHINA.

E-mail address: yau@uic.edu

DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING, 100084, P. R. CHINA

E-mail address: hqzuo@mail.tsinghua.edu.cn