

# QUASI-LOCAL ENERGY IN PRESENCE OF GRAVITATIONAL RADIATION

PO-NING CHEN, MU-TAO WANG, AND SHING-TUNG YAU

ABSTRACT. We discuss our recent work [4] in which gravitational radiation was studied by evaluating the Wang-Yau quasi-local mass of surfaces of fixed size at the infinity of both axial and polar perturbations of the Schwarzschild spacetime, à la Chandrasekhar [1].

We compute the Wang-Yau quasi-local mass [7, 8] of “spheres of unit size” at null infinity to capture the information of gravitational radiation. The set-up, following Chandrasekhar [1], is a gravitational perturbation of the Schwarzschild solution, which is governed by the Regge-Wheeler equation (see below). We take a sphere of a fixed areal radius and push it all the way to null infinity. The limit of the geometric data is that of a standard configuration and thus the optimal embedding equation [7, 8, 2] can be solved.

Let us first consider the axial perturbations. The metric perturbation is of the form:

$$-(1 - \frac{2m}{r})dt^2 + \frac{1}{1 - \frac{2m}{r}}dr^2 + r^2d\theta^2 + r^2\sin^2\theta(d\phi - q_2dr - q_3d\theta)^2.$$

The linearized vacuum Einstein equation is solved by a separation of variable Ansatz in which  $q_2$  and  $q_3$  are explicitly given by the Teukolsky function and the Legendre function.

In particular,

$$q_3 = \sin(\sigma t) \frac{C_\mu(\theta)}{\sin\theta} \frac{(r^2 - 2mr)}{\sigma^2 r^4} \frac{d}{dr}(rZ^{(-)})$$

for a solution of frequency  $\sigma$  and a separation of variable constant  $\mu$ . Here  $C_\mu(\theta)$  is related to the  $\mu$ -th Legendre function  $P_\mu$  by

$$C_\mu(\theta) = \sin\theta \frac{d}{d\theta} \left( \frac{1}{\sin\theta} \frac{dP_\mu(\cos\theta)}{d\theta} \right).$$

---

*Date:* March 30, 2016.

P.-N. Chen is supported by NSF grant DMS-1308164, M.-T. Wang is supported by NSF grant DMS-1405152, and S.-T. Yau is supported by NSF grants PHY-0714648 and DMS-1308244. This work was partially supported by a grant from the Simons Foundation (#305519 to Mu-Tao Wang). Part of this work was carried out when P.-N. Chen and M.-T. Wang were visiting the Department of Mathematics and the Center of Mathematical Sciences and Applications at Harvard University.

After the change of variable

$$r_* = r + 2m \ln\left(\frac{r}{2m} - 1\right),$$

$Z^{(-)}$  satisfies the Regge-Wheeler equation:

$$\left(\frac{d^2}{dr_*^2} + \sigma^2\right)Z^{(-)} = V^{(-)}Z^{(-)},$$

where

$$V^{(-)} = \frac{r^2 - 2mr}{r^5}[(\mu^2 + 2)r - 6m],$$

and  $\mu$  is a separation of variable constant.


On the Schwarzschild spacetime

$$-\left(1 - \frac{2m}{r}\right)dt^2 + \frac{1}{1 - \frac{2m}{r}}dr^2 + r^2d\theta^2 + r^2\sin^2\theta d\phi^2,$$

we consider an asymptotically flat Cartesian coordinate system  $(t, y_1, y_2, y_3)$  with  $y_1 = r \sin \theta \sin \phi$ ,  $y_2 = r \sin \theta \cos \phi$ ,  $y_3 = r \cos \theta$ . Given  $(d_1, d_2, d_3) \in \mathbb{R}^3$  with  $d^2 = \sum_{i=1}^3 d_i^2$ , consider the 2-surface

$$\Sigma_{t,d} = \left\{ (t, y_1, y_2, y_3) : \sum_{i=1}^3 (y_i - d_i)^2 = 1 \right\}.$$

We compute the quasi-local mass of  $\Sigma_{t,d}$  as  $d \rightarrow \infty$ .



Denote

$$A(r) = \frac{(r^2 - 2mr)}{\sigma^2 r^3} \frac{d}{dr} (rZ^{(-)}).$$

The linearized optimal embedding equation of  $\Sigma_{t,d}$  is reduced to two linear elliptic equations on the unit 2-sphere  $S^2$ :

$$\begin{aligned}\Delta(\Delta + 2)\tau &= [-A''(1 - Z_1^2) + 6A'Z_1 + 12A]Z_2Z_3 \\ (\Delta + 2)N &= (A'' - 2A'Z_1 + 4A)Z_2Z_3,\end{aligned}$$

where  $\tau$  and  $N$  are the respective time and radial components of the solution, and  $Z_1, Z_2, Z_3$  are the three standard first eigenfunctions of  $S^2$ .  $A'$  and  $A''$  are derivatives with respect to  $r$ , and  $r^2$  is substituted by  $r^2 = d^2 + 2Z_1 + 1$  in the above equations.

The quasi-local mass of  $\Sigma_{t,d}$  with respect to the optimal isometric embedding is then

$$E(\Sigma_{t,d}) = C^2\{\sin^2(\sigma t)E_1 + \sigma^2 \cos^2(\sigma t)E_2\} + O\left(\frac{1}{d^3}\right),$$

where  $E_1$  and  $E_2$  are two integrals on the standard unit 2-sphere, that depend on the solution  $\tau$  and  $N$  of the optimal isometric embedding equation. Explicitly,


$$E_1 = \int_{S^2} (1/2) [A^2Z_2^2(7Z_3^2 + 1) + 2AA'Z_1Z_3^2(3Z_2^2 - 1) - N(\Delta + 2)N]$$

$$E_2 = \int_{S^2} [A^2Z_2^2Z_3^2 - \tau\Delta(\Delta + 2)\tau].$$

In particular,

$$\partial_t E(\Sigma_{t,d}) = \frac{\sigma \sin(2\sigma t)C^2(\theta)}{d^2} \{E_1 - \sigma^2 E_2\} + O\left(\frac{1}{d^3}\right).$$

Let us compare the quasi-local mass on the small spheres  $\Sigma_{t,d}$  along a certain direction to the quasi-local mass of the large coordinate spheres  $S_{t,r}$ .



Naively, one may expect to recover  $\partial_t E(S_{t,r})$  by integrating the energy radiated away at all directions  $\partial_t E(\Sigma_{t,d})$ . However, our calculation indicates that there are nonlinear correction terms from the quasi-local energy that should be taken into account.

We can also consider the polar perturbation of the Schwarzschild space-time in which the metric coefficients  $g_{tt}$ ,  $g_{rr}$ ,  $g_{\theta\theta}$ , and  $g_{\phi\phi}$  are perturbed in

$$-(1 - \frac{2m}{r})dt^2 + \frac{1}{1 - \frac{2m}{r}}dr^2 + r^2d\theta^2 + r^2\sin^2\theta d\phi^2.$$

The gravitational perturbation is governed by the Zerilli equation

$$\left(\frac{d^2}{dr_*^2} + \sigma^2\right)Z^{(+)} = V^{(+)}Z^{(+)},$$

where

$$V^{(+)} = \frac{2(r^2 - 2mr)}{r^5(nr + 3m)^2} [n^2(n+1)r^3 + 3mn^2r^2 + 9m^2nr + 9m^3],$$

and  $n$  is the separation of variable constant. Again, we compute the quasi-local mass of spheres of unit-size at null infinity. The calculation is similar to the axial perturbation case but the result is different as the leading term is of the order  $\frac{1}{d}$  (as opposed to  $\frac{1}{d^2}$  for axial-perturbation) with nonzero coefficients. If such a linear perturbation can be realized as an actual perturbation of the Schwarzschild spacetime, the result would contradict the positivity of the quasi-local mass [6, 7, 8]. From this, we deduce the following conclusion: There does not exist any gravitational perturbation of the Schwarzschild spacetime that is of purely polar type in the sense of Chandrasekhar [1].

For an actual gravitational perturbation of the Schwarzschild solution, the vanishing of the  $\frac{1}{d}$  gives a limiting integrand that integrates to zero on the limiting 2-sphere at null infinity. In fact, the quasi-local mass density  $\rho$  (see [3, equation 2.2]) of  $\Sigma_{t,d}$  can be computed at the pointwise level. Up to an  $O(\frac{1}{d^3})$  term

$$\begin{aligned} \rho = & \left(K - \frac{1}{4}|H|^2\right) \\ & - \frac{(|H| - 2)^2}{4} + \frac{1}{d^2} \left\{ \frac{1}{2}|\nabla^2 N|^2 + ((\Delta + 2)N)^2 - \frac{1}{4}(\Delta N)^2 \right. \\ & \left. - \frac{1}{4}(\Delta\tau)^2 + \frac{1}{2}[\nabla^a \nabla^b (\tau_a \tau_b) - |\nabla\tau|^2 - \Delta|\nabla\tau|^2] \right\}, \end{aligned}$$

where  $K$  is the Gauss curvature of  $\Sigma_{t,d}$ . The first line, which integrates to zero, is of the order of  $\frac{1}{d}$  and is exactly the mass aspect function of the Hawking mass [5]. The  $\frac{1}{d^2}$  term of the quasi local mass  $\int_{\Sigma_d} \rho d\mu_{\Sigma_{t,d}}$  has contributions from the second and third lines (of the order of  $\frac{1}{d^2}$ ), the  $\frac{1}{d^2}$  term of the first line, and the  $\frac{1}{d}$  term of the area element  $d\mu_{\Sigma_{t,d}}$ . The

above integral formula is obtained after performing integrations by parts and applying the optimal embedding equation several times.

To each closed loop on the limiting 2-sphere at null infinity, we can thus associate a non-vanishing arc integral that is of the order of  $\frac{1}{d}$ , where  $d$  is the distance from the source. We expect the freedom in varying the shape of the loop can increase the detectability of gravitational waves.

## REFERENCES

- [1] S. Chandrasekhar, *The mathematical theory of black holes*, reprint of the 1992 edition, Oxford Classic Texts in the Physical Sciences, Oxford Univ. Press, New York.
- [2] P.-N. Chen, M.-T. Wang, and S.-T. Yau, *Evaluating quasi-local energy and solving optimal embedding equation at null infinity*, *Comm. Math. Phys.* **308** (2011), no.3, 845–863.
- [3] P.-N. Chen, M.-T. Wang, and S.-T. Yau, *Conserved quantities in general relativity: from the quasi-local level to spatial infinity*, *Comm. Math. Phys.* **338** (2015), no.1, 31–80.
- [4] P.-N. Chen, M.-T. Wang, and S.-T. Yau, *Quasi-local mass in the gravitational perturbations of black holes*, to appear.
- [5] S. W. Hawking, *Gravitational radiation in an expanding universe*, *J. Math. Phys.* **9**, 598 (1968).
- [6] C.-C. M. Liu and S.-T. Yau, *Positivity of quasi-local mass II*, *J. Amer. Math. Soc.* **19** (2006), no. 1, 181–204.
- [7] M.-T. Wang, and S.-T. Yau, *Quasi-local mass in general relativity*, *Phys. Rev. Lett.* **102** (2009), no. 2, no. 021101.
- [8] M.-T. Wang, and S.-T. Yau, *Isometric embeddings into the Minkowski space and new quasi-local mass*, *Comm. Math. Phys.* **288** (2009), no. 3, 919–942.

DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, NEW YORK, NY 10027,  
USA

*E-mail address:* mtwang@math.columbia.edu