

# Some extremal problems on graph theory

Chunhui Lai

Minnan Normal University

laich2011@msn.cn; laichunhui@mnnu.edu.cn

# Outline

- 1 Erdős Problem
- 2 Erdős conjecture
- 3 Hajós conjecture

# Outline

- 1 Erdős Problem
- 2 Erdős conjecture
- 3 Hajós conjecture

Let  $f(n)$  be the maximum number of edges in a graph on  $n$  vertices in which no two cycles have the same length. In 1975, Erdős raised the problem of determining  $f(n)$  (see J.A. Bondy and U.S.R. Murty [ Graph Theory with Applications (Macmillan, New York, 1976).], p.247, Problem 11). Y. Shi[ On maximum cycle-distributed graphs, Discrete Math. 71(1988) 57-71.] proved that

**Theorem 1.1 (Shi 1988)**

$$f(n) \geq n + [(\sqrt{8n - 23} + 1)/2]$$

for  $n \geq 3$ .

Y. Shi[The number of edges in a maximum cycle distributed graph, Discrete Mathematics, 104(1992), 205-209.] , Y. Shi [On simple MCD graphs containing a subgraph homomomorphic to  $K_4$ , Discrete Math , 126(1994), 325-338.], X. Jia[Some extremal problems on cycle distributed graphs, Congr. Number., 121(1996), 216-222.],

G. Chen, J. Lehel, M. S.Jacobson, and W. E.Shreve [ Note on graphs without repeated cycle lengths, J. Graph Theory, 29(1998),11-15.], C. Lai[On the maximum number of edges in a graph in which no two cycles have the same length. Combinatorics, graph theory, algorithms and applications (Beijing, 1993), 447–450, World Sci. Publ., River Edge, NJ, 1994.], C. Lai[A lower bound for the number of edges in a graph containing no two cycles of the same length, Electron. J. of Combin. 8(2001), #N9, 1 - 6.], S. Liu[Some extremal problems on the cycle length distribution of graphs. J. Combin. Math. Combin. Comput. 100 (2017), 155–171.] obtained some results.

E. Boros, Y. Caro, Z. Füredi and R. Yuster [Covering non-uniform hypergraphs, J. Combin. Theory Ser. B 82(2001), 270-284.] proved that

**Theorem 1.2 (Boros, Caro, Füredi and Yuster 2001)**

$$f(n) \leq n + 1.98\sqrt{n}(1 + o(1)).$$

C. Lai [On the size of graphs without repeated cycle lengths, Discrete Appl. Math. 232 (2017), 226-229.] improved the lower bound.

**Theorem 1.3 (Lai 2017)** Let  $t = 1260r + 169$  ( $r \geq 1$ ), then

$$f(n) \geq n + \frac{107}{3}t + \frac{7}{3}$$

for  $n \geq \frac{2119}{4}t^2 + 87978t + \frac{15957}{4}$ .



C. Lai [ On the size of graphs with all cycle having distinct length, Discrete Math. 122(1993) 363-364.] proposed the following conjecture:

**Conjecture 1.4 (Lai 1993)**

$$\liminf_{n \rightarrow \infty} \frac{f(n) - n}{\sqrt{n}} \leq \sqrt{3}.$$

It would be nice to prove that

$$\liminf_{n \rightarrow \infty} \frac{f(n) - n}{\sqrt{n}} \leq \sqrt{3 + \frac{3}{5}}.$$

$$\liminf_{n \rightarrow \infty} \frac{f(n) - n}{\sqrt{n}} \leq \sqrt{3 + \frac{1}{3}}.$$

C. Lai [On the number of edges in some graphs, Discrete Applied Mathematics 283 (2020), 751-755] construct a graph  $G$  having no two cycles with the same length which leads to the following result.

**Theorem 1.5 (Lai 2020)** Let  $t = 1260r + 169$  ( $r \geq 1$ ), then

$$f(n) \geq n + \frac{119}{3}t - \frac{26399}{3}$$

for  $n \geq \frac{1309}{2}t^2 - \frac{1349159}{6}t + \frac{6932215}{3}$ .

From Theorem 1.5, we have

$$\liminf_{n \rightarrow \infty} \frac{f(n) - n}{\sqrt{n}} \geq \sqrt{2 + \frac{40}{99}},$$

which is better than the previous bounds  $\sqrt{2}$  (see Shi 1988),  
 $\sqrt{2 + \frac{7654}{19071}}$  (see Lai 2017).

Combining this with Boros, Caro, Füredi and Yuster's upper bound, we get

$$1.98 \geq \limsup_{n \rightarrow \infty} \frac{f(n) - n}{\sqrt{n}} \geq \liminf_{n \rightarrow \infty} \frac{f(n) - n}{\sqrt{n}} \geq \sqrt{2 + \frac{40}{99}}.$$

The sequence  $(c_1, c_2, \dots, c_n)$  is the cycle length distribution of a graph  $G$  of order  $n$ , where  $c_i$  is the number of cycles of length  $i$  in  $G$ . Let  $f(a_1, a_2, \dots, a_n)$  denote the maximum possible number of edges which satisfies  $c_i \leq a_i$  where  $a_i$  is a nonnegative integer. Y. Shi posed the problem of determining  $f(a_1, a_2, \dots, a_n)$ , which extended the problem due to Erdős, it is clearly that  $f(n) = f(1, 1, \dots, 1)$  (see [Y. Shi, Some problems of cycle length distribution, J. Nanjing Univ. (Natural Sciences), Special Issue On Graph Theory, 27(1991), 233-234]).

Let  $g(n, m) = f(a_1, a_2, \dots, a_n)$ ,  $a_i = 1$  for all  $i/m$  be integer,  $a_i = 0$  for all  $i/m$  be not integer. It is clearly that  $f(n) = g(n, 1)$ .

From the Theorem 1.5, we have

$$\liminf_{n \rightarrow \infty} \frac{g(n, m) - n}{\sqrt{\frac{n}{m}}} \geq \sqrt{2 + \frac{40}{99}},$$

for all integer  $m$ .

We obtain the following result.



**Theorem 1.6 (Lai 2020)** Let  $m$  be even,  $s_1 > s_2$ ,  $s_1 + 3s_2 > k$ , then

$$g(n, m) \geq n + (k + s_1 + 2s_2 + 1)t - 1$$

for

$$n \geq \left(\frac{3}{4}mk^2 + \frac{1}{2}mks_1 + \frac{3}{2}mks_2 + \frac{1}{2}ms_1^2 + \frac{3}{2}ms_1s_2 + \frac{9}{4}ms_2^2 + mk + ms_1 + 3ms_2 + \frac{1}{2}m\right)t^2 + \left(\frac{1}{4}mk + \frac{1}{2}ms_1 + \frac{3}{4}ms_2 - k - s_1 - 2s_2 + \frac{1}{2}m - 1\right)t + 1.$$

From Theorem 1.6, we have

$$\liminf_{n \rightarrow \infty} \frac{g(n, m) - n}{\sqrt{\frac{n}{m}}} \geq$$

$$\sqrt{\frac{(k + s_1 + 2s_2 + 1)^2}{\left(\frac{3}{4}k^2 + \frac{1}{2}ks_1 + \frac{3}{2}ks_2 + \frac{1}{2}s_1^2 + \frac{3}{2}s_1s_2 + \frac{9}{4}s_2^2 + k + s_1 + 3s_2 + \frac{1}{2}\right)}},$$

for all even integer  $m$ .

Let  $s_1 = 28499066, s_2 = 4749839, k = 14249542$ , then

$$\liminf_{n \rightarrow \infty} \frac{g(n, m) - n}{\sqrt{\frac{n}{m}}} > \sqrt{2.444},$$

for all even integer  $m$ .

We make the following conjecture:

**Conjecture 1.7 (Lai 2020)**

$$\liminf_{n \rightarrow \infty} \frac{f(n) - n}{\sqrt{n}} > \sqrt{2.444}.$$

Let  $f_2(n)$  be the maximum number of edges in a 2-connected graph on  $n$  vertices in which no two cycles have the same length.

Y. Shi [On maximum cycle-distributed graphs, Discrete Math. 71(1988) 57-71.] proved that

**Theorem 1.8 (Shi 1988)** For every integer  $n \geq 3$ ,  
 $f_2(n) \leq n + \lfloor \frac{1}{2}(\sqrt{8n - 15} - 3) \rfloor$ .

G. Chen, J. Lehel, M. S.Jacobson, and W. E.Shreve [ Note on graphs without repeated cycle lengths, J. Graph Theory, 29(1998),11-15.] proved that

**Theorem 1.9 (Chen, Lehel, Jacobson, and Shreve 1998)**

$$f_2(n) \geq n + \sqrt{n/2} - o(\sqrt{n})$$

E. Boros, Y. Caro, Z. Füredi and R. Yuster [ Covering non-uniform hypergraphs, J. Combin. Theory Ser. B 82(2001), 270-284.] improved this lower bound significantly:

**Theorem 1.10 (Boros, Caro, Füredi and Yuster 2001)**

$$f_2(n) \geq n + \sqrt{n} - O(n^{\frac{9}{20}}).$$

Combining this with Shi's results we know:

**Corollary 1.11 (Boros, Caro, Füredi and Yuster 2011)**

$$\sqrt{2} \geq \limsup \frac{f_2(n) - n}{\sqrt{n}} \geq \liminf \frac{f_2(n) - n}{\sqrt{n}} \geq 1.$$

Boros, Caro, Füredi and Yuster [E. Boros, Y. Caro, Z. Füredi and R. Yuster, Covering non-uniform hypergraphs, J. Combin. Theory Ser. B 82(2001), 270-284.] made the following conjecture:

**Conjecture 1.12 (Boros, Caro, Füredi and Yuster 2001)**

$$\lim \frac{f_2(n) - n}{\sqrt{n}} = 1.$$



It is easy to see that Conjecture 1.12 implies the (difficult) upper bound in the Erdős Turan Theorem [P. Erdős , On a problem of Sidon in additive number theory and on some related problems. Addendum, J. Lond. Math. Soc. 19 (1944), 208.][P. Erdős and P. Turan, On a problem of Sidon in additive number theory, and on some related problems, J. Lond. Math. Soc. 16 (1941), 212–215.](see [E. Boros, Y. Caro, Z. Füredi and R. Yuster, Covering non-uniform hypergraphs, J. Combin. Theory Ser. B 82(2001), 270-284.]).

Jie Ma, Tianchi Yang [Non-repeated cycle lengths and Sidon sequences, arXiv:2007.12513 , 2020, Israel J. Math., to appear.] prove a conjecture of Boros, Caro, Füredi and Yuster on the maximum number of edges in a 2-connected graph without repeated cycle lengths, which is a restricted version of a longstanding problem of Erdős. Their proof together with the matched lower bound construction of Boros, Caro, Füredi and Yuster show that this problem can be conceptually reduced to the seminal problem of finding the maximum Sidon sequences in number theory.

**Theorem 1.13 (Ma, Yang 2020)**

Any  $n$ -vertex 2-connected graph with no two cycles of the same length contains at most  $n + \sqrt{n} + o(\sqrt{n})$  edges.

Let  $f_2(n, k)$  be the maximum number of edges in a graph  $G$  on  $n$  vertices in which no two cycles have the same length and  $G$  which consists of  $k$  blocks. A natural question is what is the maximum number of edges  $f_2(n, k)$ . It is clearly that  $f_2(n, 1) = f_2(n)$ .

K. Markström [ A note on uniquely pancyclic graphs, Australas. J. Combin. 44 (2009), 105-110.] raised the following problem:

**Problem 1.14 (Markström 2009)** Determine the maximum number of edges in a hamiltonian graph on  $n$  vertices with no repeated cycle lengths.

Let  $g(n)$  be the maximum number edges in an  $n$ -vertex, Hamiltonian graph with no repeated cycle length. J. Lee, C. Timmons [A note on the number of edges in a Hamiltonian graph with no repeated cycle length, Australas. J. Combin. 69(2)(2017), 286-291.] prove the following.

**Theorem 1.15 (Lee and Timmons 2017)** If  $q$  is a power of a prime and  $n = q^2 + q + 1$ , then

$$g(n) \geq n + \sqrt{n - 3/4} - 3/2$$

A simple counting argument shows that  $g(n) < n + \sqrt{2n} + 1$ .

It would be nice to determining  $g(n)$  for infinitely many  $n$ .

The lower bound  $f(0, 0, 2, \dots, 2)$  is given by J. Xu and Y. Shi [The maximum possible number of edges in a simple graph with at most two cycles having the same length, J. Shanghai Normal Univ. (Natural Sciences), 32(3)(2003), 26-32.].

**Theorem 1.16 (Xu and Shi 2003)**

For  $n \geq 3$ ,

$$f(0, 0, 2, \dots, 2) \geq n - 1 + [(\sqrt{11n - 20})/2],$$

and the equality holds when  $3 \leq n \leq 10$ .

Given a graph  $H$ , what is the maximum number of edges of a graph with  $n$  vertices not containing  $H$  as a subgraph? This number is denoted  $ex(n, H)$ , and is known as the Turan number.



We denote by  $m_i(n)$  the numbers of cycles of length  $i$  in the complete graph  $K_n$  on  $n$  vertices. Obviously,

$$\begin{aligned} & ex(n, C_k) \\ &= f(0, 0, m_3(n), \dots, \\ & m_{k-1}(n), 0, m_{k+1}(n), \dots, m_n(n)) \\ &= f(0, 0, 2^{\frac{n(n-1)}{2}}, \dots, \\ & 2^{\frac{n(n-1)}{2}}, 0, 2^{\frac{n(n-1)}{2}}, \dots, 2^{\frac{n(n-1)}{2}}). \end{aligned}$$

Therefore, finding  $ex(n, C_k)$  is a special case of determining  $f(a_1, a_2, \dots, a_n)$ .

There are not good sufficient and necessary condition of when a graph on  $n$  vertices in which no two cycles have the same length. There are also not good sufficient and necessary condition of when a 2-connected graph on  $n$  vertices in which no two cycles have the same length. It would be nice to give good sufficient and necessary condition of when a graph on  $n$  vertices in which no two cycles have the same length. It would be nice to give good sufficient and necessary condition of when a 2-connected graph on  $n$  vertices in which no two cycles have the same length.

The survey article on this problem can be found in

Tian, Feng[The progress on some problems in graph theory.  
(Chinese) Qufu Shifan Daxue Xuebao Ziran Kexue Ban 1986, no. 2,  
30–36. MR0865617 (87m:05160)],

Zhang, Ke Min[Progress of some problems in graph theory.  
(Chinese) J. Math. Res. Exposition 27 (2007), no. 3, 563–576.  
MR2349503] and

Chunhui Lai, Mingjing Liu[Some open problems on cycles,  
Journal of Combinatorial Mathematics and Combinatorial Computing  
91 (2014), 51-64. MR3287706]

Also see Douglas B. West [Introduction to Graph Theory (Second edition), Prentice Hall, 2001] ,p77, Exercises 2.1.41 , Douglas B. West [Combinatorial Mathematics, Cambridge University Press,Cambridge, 2020] ,p249, Exercise 5.4.27 , RMM 2019 - Romanian Master of Mathematics 2019 <http://rmms.lbi.ro/rmm2019/index.php?id=home>, The 11th Romanian Master of Mathematics Competition Problem 3.

The progress of all 50 problems in [J.A. Bondy and U.S.R. Murty, Graph Theory with Applications (Macmillan, New York, 1976)] can be find in Stephen C. Locke, Unsolved problems:  
<http://math.fau.edu/locke/Unsolved.htm>

# Outline

- 1 Erdős Problem
- 2 Erdős conjecture
- 3 Hajós conjecture

P. Erdős conjectured that there exists a positive constant  $c$  such that  $ex(n, C_{2k}) \geq cn^{1+1/k}$  (see P. Erdős, Some unsolved problems in graph theory and combinatorial analysis, Combinatorial Mathematics and its Applications (Proc. Conf., Oxford, 1969) , pp. 97–109, Academic Press, London, 1971).

P. Erdős [Extremal problems in graph theory, in Theory of graphs and its applications, Proc. Symp. ( Smolenice, 1963), (M. Fiedler, ed.), 29-36. New York: Academic Press, 1965.] and J.A Bondy, M. Simonovits [ Cycle of even length in graphs, J. Combin. Theory Ser. B,16(1974),97-105.] obtained that

**Theorem 2.1 (Erdős 1965, Bondy and Simonovits 1974)**

$$ex(n, C_{2k}) \leq ckn^{1+1/k}$$



R. Wenger [Extremal graphs with no  $C^4, s, C^6, s$ , or  $C^{10}, s$ ,, J. Combin. Theory Ser. B 52(1991), 113-116.] proved that

**Theorem 2.2 (Wenger 1991)**

$$ex(n, C_4) \geq \left(\frac{n}{2}\right)^{3/2},$$

$$ex(n, C_6) \geq \left(\frac{n}{2}\right)^{4/3},$$

$$ex(n, C_{10}) \geq \left(\frac{n}{2}\right)^{6/5}$$

Z. Füredi[Graph without quadrilaterals, J. Combin. Theory Ser. B 34(1983), 187-190.] proved that

**Theorem 2.3 (Füredi 1983)** If  $q$  is a power of 2, then

$$ex(q^2 + q + 1, C_4) = q(q + 1)^2/2$$

Z. Füredi [ On the number of edges of quadrilateral-free graphs, J. Combin. Theory Ser. B 68(1996), 1-6.] proved that

**Theorem 2.4 (Füredi 1996)** Let  $G$  be a quadrilateral-free graph with  $e$  edges on  $q^2 + q + 1$  vertices, and suppose that  $q \geq 15$ . Then  $e \leq q(q + 1)^2/2$ .

**Corollary 2.5 (Füredi 1996)** If  $q$  is a prime power greater than 13,  $n = q^2 + q + 1$ . Then

$$ex(n, C_4) = q(q + 1)^2/2.$$

Z. Füredi, A. Naor and J. Verstraete[ on the Turan number for the hexagon, Adv. Math. 203(2) (2006), 476-496.] proved that

**Theorem 2.6 (Füredi, Naor and Verstraete 2006)**

$$ex(n, C_6) > 0.5338n^{4/3}$$

for infinitely many  $n$  and

$$ex(n, C_6) < 0.6272n^{4/3}$$

if  $n$  is sufficiently large.

This refutes the Erdős-Simonovits conjecture in 1982 for hexagons(see[Füredi, Naor and Verstraete 2006]).

Firke, Frank A.; Kosek, Peter M.; Nash, Evan D.; Williford, Jason. [Extremal graphs without 4-cycles. J. Combin. Theory Ser. B 103 (2013), no. 3, 327–336. MR3048158] proved that

**Theorem 2.7 (Firke, Kosek, Nash, Williford 2013)**

For  $q$  even,

$$ex(q^2 + q, C_4) \leq q(q + 1)^2/2 - q.$$

M. Tait, C. Timmons [Sidon sets and graphs without 4-cycles, J. Comb. 5(2) (2014), 155 - 165.] proved that

**Theorem 2.8 (M. Tait, C. Timmons 2014)**

If  $q$  is an odd prime power, then

$$ex(q^2 - q - 2, C_4) \geq \frac{1}{2}q^3 - q^2 - O(q^{\frac{3}{4}}).$$

Jialin He, Jie Ma, Tianchi Yang [Stability and supersaturation of 4-cycles, arXiv:1912.00986v3 [math.CO]] proved that

**Theorem 2.9 (He, Ma, Yang 2019)**

Let  $q$  be even and  $G$  be a  $C_4$ -free graph on  $q^2 + q + 1$  vertices with at least  $\frac{1}{2}q(q+1)^2 - \frac{1}{2}q + o(q)$  edges. Then there exists a unique polarity graph of order  $q$ , which contains  $G$  as a subgraph.

The survey article on this Erdos conjecture can be found in  
F.R.K, Chung [Open problems of Paul Erdos in graph theory, J.  
Graph Theory 25 (1997), 3-36.]

Z.Furedi, M. Simonovits [The history of degenerate (bipartite)  
extremal graph problems. Erdos centennial, 169-264, Bolyai Soc.  
Math. Stud., 25, Janos Bolyai Math. Soc., Budapest, 2013.]

J. Verstraete [Extremal problems for cycles in graphs, Recent  
trends in combinatorics, 83-116, IMA Vol. Math. Appl., 159,  
Springer, Cham, 2016].

Chunhui Lai, Mingjing Liu [Some open problems on cycles,  
Journal of Combinatorial Mathematics and Combinatorial Computing  
91 (2014), 51-64. MR3287706]



Generalized Turán problems can be found in

N. Alon, R. Yuster [The Turán number of sparse spanning graphs. *J. Combin. Theory Ser. B* 103(3) (2013), 337 - 343.]

N. Alon, C. Shikhelman [Many  $T$  copies in  $H$ -free graphs. *J. Combin. Theory Ser. B* 121 (2016), 146 - 172.]

N. Alon, C. Shikhelman [H-free subgraphs of dense graphs maximizing the number of cliques and their blow-ups. *Discrete Math.* 342 (2019), 988 - 996.]

P. Loh, M. Tait, C. Timmons, R. M. Zhou [Induced Turán numbers. *Combin. Probab. Comput.* 27(2) (2018), 274 - 288.]

C. Palmer, M. Tait, C. Timmons, A. Z. Wagner [Turán numbers for Berge-hypergraphs and related extremal problems. *Discrete Math.* 342 (2019), no. 6, 1553-1563.]

Y. Caro, Z. Tuza [Regular Turán numbers, arXiv:1911.00109 [math.CO]]

Y. Lan, Y. Shi, Z. Song [Extremal theta-free planar graphs. *Discrete Math.* 342 (2019), no. 12, 111610, 8 pp.]

D. Gerbner, E. Gyria, A. Methuku, M. Vizer [Generalized Turán problems for even cycles. *J. Combin. Theory Ser. B* 145 (2020), 169 - 213.]

etc.

There are not good sufficient and necessary condition of when a graph on  $n$  vertices in which contains  $k$  cycle. For  $k = n$ , it is Hamiltonian problem, The survey article on Hamiltonian problem can be found in [Gould, Ronald J. Updating the Hamiltonian problem—a survey. *J. Graph Theory* 15 (1991), no. 2, 121–157. MR1106528 (92m:05128); Gould, Ronald J. Advances on the Hamiltonian problem—a survey. *Graphs Combin.* 19 (2003), no. 1, 7–52. MR1974368 (2004a:05092); Gould, Ronald J. Recent advances on the Hamiltonian problem: Survey III. *Graphs Combin.* 30 (2014), no. 1, 1–46. MR3143857 (Reviewed)]. It would be nice to give good sufficient and necessary condition of when a graph on  $n$  vertices in which contains  $k$  cycle for some  $k$ .

# Outline

- 1 Erdős Problem
- 2 Erdős conjecture
- 3 Hajós conjecture

# Hajós conjecture

An *eulerian graph* is a graph (not necessarily connected) in which each vertex has even degree. Let  $G$  be an eulerian graph. A *circuit decomposition* of  $G$  is a set of edge-disjoint circuits  $C_1, C_2, \dots, C_t$  such that  $E(G) = C_1 \cup C_2 \cup \dots \cup C_t$ . It is well known that every eulerian graph has a circuit decomposition. A natural question is to find the smallest number  $t$  such that  $G$  has a circuit decomposition of  $t$  circuits?

Such smallest number  $t$  is called the *circuit decomposition* number of  $G$ , denoted by  $cd(G)$ . For each edge  $xy \in E(G)$ , let  $m(xy)$  be the number of edges between  $x$  and  $y$ . The *multiple number* of  $G$  is defined by  $m(G) = \sum_{uv \in E(G)} (m(uv) - 1)$ . (See G. Fan and B. Xu [Hajós conjecture and projective graphs. Discrete Math. 252 (2002), no. 1-3, 91 - 101])

The following conjecture is due to Hajós(see L. Lovasz, On covering of graphs, in: P. Erdos, G.O.H. Katona (Eds.), Theory of Graphs, Academic Press, New York, 1968, pp. 231 - 236 ).

**Hajós conjecture:**  $cd(G) \leq \frac{|V(G)|}{2}$  for every simple eulerian graph  $G$ .

N. Dean [What is the smallest number of dicycles in a dicycle decomposition of an Eulerian digraph? *J. Graph Theory* 10 (1986), no. 3, 299–308.] proved that

**Theorem 3.1 (Dean 1986)** Hajós conjecture is equivalent to the following statement: If  $G$  is even, then  $cd(G) \leq \frac{|V(G)|-1}{2}$ .



L. Lovasz [On covering of graphs, in: P. Erdos, G.O.H. Katona (Eds.), Theory of Graphs, Academic Press, New York, 1968, pp. 231 - 236 ] proved that

**Theorem 3.2 (Lovasz 1968)** A graph of  $n$  vertices can be covered by  $\leq \lceil n/2 \rceil$  disjoint paths and circuits.

T. Jiang [On Hajós conjecture, J. China Univ. Sci. Tech. 14 (1984) 585 - 592 (in Chinese).] and K. Seyffarth [Hajós conjecture and small cycle double covers of planar graphs, Discrete Math. 101 (1992) 291 - 306.] proved that

**Theorem 3.3 (Jiang 1984; Seyffarth 1992)**  $cd(G) \leq \frac{|V(G)|-1}{2}$  for every simple planar eulerian graph  $G$ .

A. Granville, A. Moisiadis [On Hajós conjecture, in: Proceedings of the 16th Manitoba Conference on Numerical Mathematics and Computing, Congr. Numer. 56 (1987) 183 - 187.] and O. Favaron, M. Kouider [Path partitions and cycle partitions of eulerian graphs of maximum degree 4, Studia Sci. Math. Hungar. 23 (1988) 237 - 244.] proved that

**Theorem 3.4 (Granville and Moisiadis 1987; Favaron and Kouider 1988)** If  $G$  is an even multigraph of order  $n$ , of size  $m$ , with  $\Delta(G) \leq 4$ , then  $cd(G) \leq \frac{n+M-1}{2}$  where  $M = m - m^*$  and  $m^*$  is the size of the simple graph induced by  $G$ .

G. Fan and B. Xu [Hajós conjecture and projective graphs.  
Discrete Math. 252 (2002), no. 1-3, 91 - 101] proved that

**Theorem 3.5 (Fan and Xu 2002)** *If  $G$  is an eulerian graph with*

$$cd(G) > \frac{|V(G)| + m(G) - 1}{2},$$

*then  $G$  has a reduction  $H$  such that*

$$cd(H) > \frac{|V(H)| + m(H) - 1}{2}$$

*and the number of vertices of degree less than six in  $H$  plus  $m(H)$  is at most one.*

**Corollary 3.6 (Fan and Xu 2002)** *Hajós conjecture is valid for projective graphs.*

**Corollary 3.7 (Fan and Xu 2002)** *Hajós conjecture is valid for  $K_6^-$  minor free graphs.*

B. Xu [Hajós conjecture and connectivity of Eulerian graphs. *J. Syst. Sci. Complex.* 15 (2002), no. 3, 295 - 298. ] also proved the following two results:



**Theorem 3.8 (Xu 2002)** *If  $G$  is an eulerian graph with*

$$cd(G) > \frac{|V(G)| + m(G) - 1}{2}$$

*such that*

$$cd(H) \leq \frac{|V(H)| + m(H) - 1}{2}$$

*for each proper reduction of  $G$ , then  $G$  is 3-connected. Moreover, if  $S = \{x, y, z\}$  is a 3-cut of  $G$ , letting  $G_1$  and  $G_2$  be the two induced subgraph of  $G$  such that  $V(G_1) \cap V(G_2) = S$  and  $E(G_1) \cup E(G_2) = E(G)$ , then either  $S$  is not an independent set, or  $G_1$  and  $G_2$  are both eulerian graphs.*

**Corollary 3.9 (Xu 2002)** *To prove Hajós' conjecture, it suffices to prove*

$$cd(G) \leq \frac{|V(G)| + m(G) - 1}{2}$$

*for every 3-connected eulerian graph  $G$ .*

G. Fan [Covers of Eulerian graphs. J. Combin. Theory Ser. B 89 (2003), no. 2, 173 - 187.] proved that

**Theorem 3.10 (Fan 2003)** Every eulerian graph on  $n$  vertices can be covered by at most  $\lfloor \frac{n-1}{2} \rfloor$  circuits such that each edge is covered an odd number of times.

This settles a conjecture made by Chung in 1980(see[Fan 2003]).

B. Xu and L. Wang [Decomposing toroidal graphs into circuits and edges. Discrete Appl. Math. 148 (2005), no. 2, 147 - 159. ] give **Theorem 3.11 (Xu and Wang 2005)** *The edge set of each even toroidal graph can be decomposed into at most  $(n + 3)/2$  circuits in  $O(mn)$  time, where a toroidal graph is a graph embedable on the torus.*

**Theorem 3.12 (Xu and Wang 2005)** *The edge set of each toroidal graph can be decomposed into at most  $3(n-1)/2$  circuits and edges in  $O(mn)$  time.*

E. Fuchs, L. Gellert, I. Heinrich [Cycle decompositions of pathwidth-6 graphs, Journal of Graph Theory, 94(2)2020, 224-251] give

**Theorem 3.13 (Fuchs, Gellert, Heinrich 2020)** Every Eulerian graph  $G$  of pathwidth at most 6 satisfies Hajós conjecture.

## We do not think Hajós conjecture is true.

By the proof of Lemma 3.3 in N. Dean [What is the smallest number of dicycles in a dicycle decomposition of an Eulerian digraph? J. Graph Theory 10 (1986), no. 3, 299–308], if exists  $k$  vertices counterexample, then will exist  $ik - i + 1 (i = 2, 3, 4, \dots)$  vertices counterexamples.

The survey article on this problem can be found in

Pyber, L.[ Covering the edges of a graph by . . . . Sets, graphs and numbers (Budapest, 1991), 583 - 610, Colloq. Math. Soc. János Bolyai, 60, North-Holland, Amsterdam, 1992. MR1218220 ],

Fan, G. H.[ Integer flows and subgraph covers (in Chinese). Sci Sin Math, 2017, 47: 457 - 466, doi: 10.1360/N012016- 00177] and

Chunhui Lai, Mingjing Liu[Some open problems on cycles, Journal of Combinatorial Mathematics and Combinatorial Computing 91 (2014), 51-64. MR3287706]



# Acknowledgement:

Project supported by the National Science Foundation of China (No.61379021; No. 11401290), NSF of Fujian (2015J01018; 2018J01423; 2020J01795), Fujian Provincial Training Foundation for "Bai-Quan-Wan Talents Engineering", Project of Fujian Education Department (JZ160455), Fujian Key Laboratory of Granular Computing and Applications(Minnan Normal University), the Institute of Meteorological Big Data-Digital Fujian and Fujian Key Laboratory of Data Science and Statistics.

The authors would like to thank Professor Y. Caro, G. Fan, B. Xu, R. Yuster for their advice and sending some papers to us. The authors would like to thank Professor E. Boros, R. Gould, G.O.H. Katona, C. Zhao for their advice. The authors would like to thank Professor N. Alon, M. Tait, C. Timmons, T. Yang for their sending some papers to us.

The End

Thanks for your  
attention!