

# GRAPHS WITHOUT REPEATED CYCLE LENGTHS

Chunhui Lai \*

Department of Mathematics, Zhangzhou Teachers College, Zhangzhou,  
Fujian 363000, P. R. of CHINA; and Graph Theory and Combinatorics Laboratory,  
Institute of Systems Science, Academy of Mathematics and Systems Science,  
Chinese Academy of Sciences, Beijing 100080, P. R. of CHINA

`zjlaichu@public.zzptt.fj.cn`

MR Subject Classifications: 05C38, 05C35

Key words: graph, cycle, number of edges

Australasian Journal of Combinatorics 27(2003), 101-105.

## Abstract

In 1975, P. Erdős proposed the problem of determining the maximum number  $f(n)$  of edges in a graph of  $n$  vertices in which any two cycles are of different lengths. In this paper, it is proved that

$$f(n) \geq n + 36t$$

for  $t = 1260r + 169$  ( $r \geq 1$ ) and  $n \geq 540t^2 + \frac{175811}{2}t + \frac{7989}{2}$ . Consequently,  $\liminf_{n \rightarrow \infty} \frac{f(n)-n}{\sqrt{n}} \geq \sqrt{2 + \frac{2}{5}}$ , which is better than the previous bounds  $\sqrt{2}$  (see [2]),  $\sqrt{2 + \frac{2562}{6911}}$  (see [7]).

## 1 Introduction

Let  $f(n)$  be the maximum number of edges in a graph on  $n$  vertices in which no two cycles have the same length. In 1975, Erdős raised the problem of determining  $f(n)$  (see [1], p.247, Problem 11). Shi[2] proved that

$$f(n) \geq n + [(\sqrt{8n - 23} + 1)/2]$$

for  $n \geq 3$ . Lai[3,4,5,6,7] proved that for  $n \geq \frac{6911}{16}t^2 + \frac{514441}{8}t - \frac{3309665}{16}$ ,  $t = 27720r + 169$ ,

$$f(n) \geq n + 32t - 1,$$

---

\*Project Supported by NSF of Fujian(A96026), Science and Technology Project of Fujian(K20105) and Fujian Provincial Training Foundation for "Bai-Quan-Wan Talents Engineering".

and for  $n \geq e^{2m}(2m+3)/4$ ,

$$f(n) < n - 2 + \sqrt{n \ln(4n/(2m+3))} + 2n + \log_2(n+6).$$

Boros, Caro, Füredi and Yuster[8] proved that

$$f(n) \leq n + 1.98\sqrt{n}(1 + o(1)).$$

In this paper, we construct a graph  $G$  having no two cycles with the same length which leads the following result.

**Theorem.** Let  $t = 1260r + 169$  ( $r \geq 1$ ), then

$$f(n) \geq n + 36t$$

for  $n \geq 540t^2 + \frac{175811}{2}t + \frac{7989}{2}$ .

## 2 Proof of the theorem

**Proof.** Let  $t = 1260r + 169$ ,  $r \geq 1$ ,  $n_t = 540t^2 + \frac{175811}{2}t + \frac{7989}{2}$ ,  $n \geq n_t$ . We shall show that there exists a graph  $G$  on  $n$  vertices with  $n + 36t$  edges such that all cycles in  $G$  have distinct lengths.

Now we construct the graph  $G$  which consists of a number of subgraphs:  $B_i$ , ( $0 \leq i \leq 21t - 1$ ,  $27t \leq i \leq 28t + 64$ ,  $29t - 734 \leq i \leq 29t + 267$ ,  $30t - 531 \leq i \leq 30t + 57$ ,  $31t - 741 \leq i \leq 31t + 58$ ,  $32t - 740 \leq i \leq 32t + 57$ ,  $33t - 741 \leq i \leq 33t + 57$ ,  $34t - 741 \leq i \leq 34t + 52$ ,  $35t - 746 \leq i \leq 35t + 60$ ,  $36t - 738 \leq i \leq 36t + 60$ ,  $37t - 738 \leq i \leq 37t + 799$ ,  $i = 21t + 2j + 1$  ( $0 \leq j \leq t - 1$ ),  $i = 21t + 2j$  ( $0 \leq j \leq \frac{t-1}{2}$ ),  $i = 23t + 2j + 1$  ( $0 \leq j \leq \frac{t-1}{2}$ ), and  $i = 26t$ ).

Now we define these  $B_i$ 's. These subgraphs all have a common vertex  $x$ , otherwise their vertex sets are pairwise disjoint.

For  $0 \leq i \leq t - 1$ , let the subgraph  $B_{21t+2i+1}$  consist of a cycle

$$xu_i^1 u_i^2 \dots u_i^{25t+2i-1} x$$

and a path:

$$xu_{i,1}^1 u_{i,1}^2 \dots u_{i,1}^{(19t+2i-1)/2} u_{i,1}^{(23t+2i+1)/2}$$

Based the construction,  $B_{21t+2i+1}$  contains exactly three cycles of lengths:

$$21t + 2i + 1, 23t + 2i, 25t + 2i.$$

For  $0 \leq i \leq \frac{t-3}{2}$ , let the subgraph  $B_{21t+2i}$  consist of a cycle

$$xv_i^1 v_i^2 \dots v_i^{25t+2i} x$$

and a path:

$$xv_{i,1}^1v_{i,1}^2\dots v_{i,1}^{9t+i-1}v_i^{12t+i}$$

Based the construction,  $B_{21t+2i}$  contains exactly three cycles of lengths:

$$21t + 2i, 22t + 2i + 1, 25t + 2i + 1.$$

For  $0 \leq i \leq \frac{t-3}{2}$ , let the subgraph  $B_{23t+2i+1}$  consist of a cycle

$$xw_i^1w_i^2\dots w_i^{26t+2i+1}x$$

and a path:

$$xw_{i,1}^1w_{i,1}^2\dots w_{i,1}^{(21t+2i-1)/2}w_i^{(25t+2i+1)/2}$$

Based the construction,  $B_{23t+2i+1}$  contains exactly three cycles of lengths:

$$23t + 2i + 1, 24t + 2i + 2, 26t + 2i + 2.$$

For  $58 \leq i \leq t - 742$ , let the subgraph  $B_{27t+i-57}$  consist of a cycle

$$C_{27t+i-57} = xy_i^1y_i^2\dots y_i^{132t+11i+893}x$$

and ten paths sharing a common vertex  $x$ , the other end vertices are on the cycle  $C_{27t+i-57}$ :

$$\begin{aligned} & xy_{i,1}^1y_{i,1}^2\dots y_{i,1}^{(17t-1)/2}y_i^{(37t-115)/2+i} \\ & xy_{i,2}^1y_{i,2}^2\dots y_{i,2}^{(19t-1)/2}y_i^{(57t-103)/2+2i} \\ & xy_{i,3}^1y_{i,3}^2\dots y_{i,3}^{(19t-1)/2}y_i^{(77t+315)/2+3i} \\ & xy_{i,4}^1y_{i,4}^2\dots y_{i,4}^{(21t-1)/2}y_i^{(97t+313)/2+4i} \\ & xy_{i,5}^1y_{i,5}^2\dots y_{i,5}^{(21t-1)/2}y_i^{(117t+313)/2+5i} \\ & xy_{i,6}^1y_{i,6}^2\dots y_{i,6}^{(23t-1)/2}y_i^{(137t+311)/2+6i} \\ & xy_{i,7}^1y_{i,7}^2\dots y_{i,7}^{(23t-1)/2}y_i^{(157t+309)/2+7i} \\ & xy_{i,8}^1y_{i,8}^2\dots y_{i,8}^{(25t-1)/2}y_i^{(177t+297)/2+8i} \\ & xy_{i,9}^1y_{i,9}^2\dots y_{i,9}^{(25t-1)/2}y_i^{(197t+301)/2+9i} \\ & xy_{i,10}^1y_{i,10}^2\dots y_{i,10}^{(27t-1)/2}y_i^{(217t+305)/2+10i}. \end{aligned}$$

As a cycle with  $d$  chords contains  $\binom{d+2}{2}$  distinct cycles,  $B_{27t+i-57}$  contains exactly 66 cycles of lengths:

$27t + i - 57,$	$28t + i + 7,$	$29t + i + 210,$	$30t + i,$
$31t + i + 1,$	$32t + i,$	$33t + i,$	$34t + i - 5,$
$35t + i + 3,$	$36t + i + 3,$	$37t + i + 742,$	$38t + 2i - 51,$
$38t + 2i + 216,$	$40t + 2i + 209,$	$40t + 2i,$	$42t + 2i,$
$42t + 2i - 1,$	$44t + 2i - 6,$	$44t + 2i - 3,$	$46t + 2i + 5,$
$46t + 2i + 744,$	$48t + 3i + 158,$	$49t + 3i + 215,$	$50t + 3i + 209,$
$51t + 3i - 1,$	$52t + 3i - 1,$	$53t + 3i - 7,$	$54t + 3i - 4,$
$55t + 3i - 1,$	$56t + 3i + 746,$	$59t + 4i + 157,$	$59t + 4i + 215,$
$61t + 4i + 208,$	$61t + 4i - 2,$	$63t + 4i - 7,$	$63t + 4i - 5,$
$65t + 4i - 2,$	$65t + 4i + 740,$	$69t + 5i + 157,$	$70t + 5i + 214,$
$71t + 5i + 207,$	$72t + 5i - 8,$	$73t + 5i - 5,$	$74t + 5i - 3,$
$75t + 5i + 739,$	$80t + 6i + 156,$	$80t + 6i + 213,$	$82t + 6i + 201,$
$82t + 6i - 6,$	$84t + 6i - 3,$	$84t + 6i + 738,$	$90t + 7i + 155,$
$91t + 7i + 207,$	$92t + 7i + 203,$	$93t + 7i - 4,$	$94t + 7i + 738,$
$101t + 8i + 149,$	$101t + 8i + 209,$	$103t + 8i + 205,$	$103t + 8i + 737,$
$111t + 9i + 151,$	$112t + 9i + 211,$	$113t + 9i + 946,$	$122t + 10i + 153,$
$122t + 10i + 952,$	$132t + 11i + 894.$		

$B_0$  is a path with an end vertex  $x$  and length  $n - n_t$ . Other  $B_i$  is simply a cycle of length  $i$ .

Then  $f(n) \geq n + 36t$ , for  $n \geq n_t$ . This completes the proof.

From the above theorem, we have

$$\liminf_{n \rightarrow \infty} \frac{f(n) - n}{\sqrt{n}} \geq \sqrt{2 + \frac{2}{5}},$$

which is better than the previous bounds  $\sqrt{2}$  (see [2]),  $\sqrt{2 + \frac{2562}{6911}}$  (see [7]).

Combining this with Boros, Caro, Füredi and Yuster's upper bound, we get

$$1.98 \geq \limsup_{n \rightarrow \infty} \frac{f(n) - n}{\sqrt{n}} \geq \liminf_{n \rightarrow \infty} \frac{f(n) - n}{\sqrt{n}} \geq \sqrt{2.4}.$$

We make the following conjecture:

**Conjecture.**

$$\lim_{n \rightarrow \infty} \frac{f(n) - n}{\sqrt{n}} = \sqrt{2.4}.$$

## Acknowledgment

The author thanks Prof. Yair Caro and Raphael Yuster for sending me references[8]. The author thanks Prof. Genghua Fan and Cheng Zhao for their valuable suggestions.

## References

- [1] J.A. Bondy and U.S.R. Murty, Graph Theory with Applications (Macmillan, New York, 1976).
- [2] Y. Shi, On maximum cycle-distributed graphs, Discrete Math. 71(1988) 57-71.
- [3] Chunhui Lai, On the Erdős problem, J. Zhangzhou Teachers College(Natural Science Edition) 3(1)(1989) 55-59.
- [4] Chunhui Lai, Upper bound and lower bound of  $f(n)$ , J. Zhangzhou Teachers College(Natural Science Edition) 4(1)(1990) 29,30-34.
- [5] Chunhui Lai, On the size of graphs with all cycle having distinct length, Discrete Math. 122(1993) 363-364.
- [6] Chunhui Lai, The number of edges in a graph in which no two cycles have the same length, J. Zhangzhou Teachers College (Natural Science Edition) 8(4)(1994), 30-34.
- [7] Chunhui Lai, A lower bound for the number of edges in a graph containing no two cycles of the same length, The Electronic J. of Combinatorics 8(2001), #N9.
- [8] E. Boros, Y. Caro, Z. Füredi and R. Yuster, Covering non-uniform hypergraphs (submitted, 2000).