

A GEOMETRIC CONSTRUCTION OF REPRESENTATIONS OF THE BEREZIN-TOEPLITZ QUANTIZATION

KWOKWAI CHAN, NAICHUNG CONAN LEUNG, AND QIN LI

ABSTRACT. For a Kähler manifold X equipped with a prequantum line bundle L , we give a geometric construction of a family of representations of the Berezin-Toeplitz deformation quantization algebra $(C^\infty(X)[[\hbar]], \star_{BT})$ parametrized by points $z_0 \in X$. The key idea is to use peak sections to suitably localize the Hilbert spaces $H^0(X, L^{\otimes m})$ around z_0 .

1. INTRODUCTION

Quantization plays important roles in both physics and in mathematics. Two outstanding approaches are the deformation quantization ([1–3, 11, 20, 24]) and geometric quantization [18, 21, 29, 30, 32]. This paper is an attempt to understand the intriguing relationship between these two quantization schemes.

To begin with, let us consider a symplectic vector space $X = \mathbb{R}^{2n}$ equipped with the standard symplectic form $\omega = \sum_{j=1}^n dx_j \wedge dy_j$. A complex polarization (i.e., complex structure) identifies X with \mathbb{C}^n with coordinates $z_j = x_j + iy_j$'s. Then geometric quantization gives the *Bargmann-Fock space* $\mathcal{H}L^2(\mathbb{C}^n, \mu_{\hbar})$, elements of which are L^2 integrable entire holomorphic functions with respect to the density $\mu_{\hbar}(z) := (\pi\hbar)^{-n} e^{-|z|^2/\hbar}$. A smooth function $f = f(z, \bar{z}) \in C^\infty(X)$ acts on $\mathcal{H}L^2(\mathbb{C}^n, \mu_{\hbar})$ as a Toeplitz operator T_f defined by setting $T_{z_j} = m_{z_j}$ (i.e., multiplication by z_j) and $T_{\bar{z}_j} = \hbar \frac{\partial}{\partial z_j}$. The composition of these operators defines a *star product* via the formula $T_f \circ T_g = T_{f \star g}$, which is given explicitly by

$$f \star g := \exp \left(-\hbar \sum_{i=1}^n \frac{\partial}{\partial z_i} \frac{\partial}{\partial \bar{w}_i} \right) (f(z, \bar{z})g(w, \bar{w}))|_{z=w}.$$

This endows $C^\infty(X)[[\hbar]]$ with a noncommutative algebra structure, or a deformation quantization of (X, ω) , and $\mathcal{H}L^2(\mathbb{C}^n, \mu_{\hbar})$ is naturally its representation.

Note that the space of polynomials $\mathbb{C}[z_1, \dots, z_n]$ is preserved under this action, and we have the decomposition $\mathbb{C}[z_1, \dots, z_n] = \sum_m \mathbb{C}[z_1, \dots, z_n]_m$, where $\mathbb{C}[z_1, \dots, z_n]_m$ is the space of degree m homogeneous polynomials on \mathbb{C}^n which can be regarded as the space of holomorphic sections of $L^{\otimes m}$ with L being the trivial line bundle.

2010 *Mathematics Subject Classification.* 53D55 (58J20, 81T15, 81Q30).

Key words and phrases. Deformation quantization, geometric quantization, Berezin-Toeplitz star product, Toeplitz operator, peak section.

In general, we are interested in a compact Kähler manifold (X, ω, J) with integral $[\omega]$, so that there exists a prequantum line bundle L whose curvature F_L satisfies $\frac{\sqrt{-1}}{2\pi}F_L = \omega$. Geometric quantization of $(X, m\omega)$ gives the Hilbert space $H^0(X, L^{\otimes m})$, the space of holomorphic sections of $L^{\otimes m}$. To a smooth function $f \in C^\infty(X)$, we can similarly associate the Toeplitz operator

$$T_{f,m} := \Pi_m \circ m_f : H^0(X, L^{\otimes m}) \rightarrow H^0(X, L^{\otimes m}),$$

where m_f is multiplication by f and Π_m denotes the orthogonal projection from the space of L^2 sections $L^2(X, L^{\otimes m})$ to $H^0(X, L^{\otimes m})$.

An important result in the Berezin-Toeplitz quantization is that this gives rise to a star product \star_{BT} , and hence the *Berezin-Toeplitz deformation quantization algebra* $(C^\infty(X)[[\hbar]], \star_{BT})$ [3, 17, 28]:

$$f \star_{BT} g := \sum_{i \geq 0} \hbar^i C_i(f, g),$$

where $C_i(-, -)$ are bi-differential operators, $\|\cdot\|$ is the operator norm, and $K_N(f, g)$ is independent of m such that the following estimates hold:

$$(1.1) \quad \|T_{f,m} \circ T_{g,m} - \sum_{i=0}^{N-1} \left(\frac{1}{m}\right)^i T_{C_i(f,g),m}\| \leq K_N(f, g) \left(\frac{1}{m}\right)^N.$$

Unlike the flat case, however, the estimate (1.1) says that the difference $T_{f,m} \circ T_{g,m} - T_{f \star_{BT} g, m}$ is only *asymptotically* zero when m tends to infinity. So $(C^\infty(X)[[\hbar]], \star_{BT})$ does not quite act on $H^0(X, L^{\otimes m})$; we do not even expect a representation of $(C^\infty(X)[[\hbar]], \star_{BT})$ on the product $\prod_m H^0(X, L^{\otimes m})$.

On the other hand, as $m \rightarrow \infty$, physically speaking we are scaling X to the *large volume limit*. We would expect the physical system to behave like that on a flat space around any given point $z_0 \in X$. We are going to show that this is indeed the case. To be more precise, we will use *peak sections* $S_{m,p,r}$ of $H^0(X, L^m)$ to appropriately localize the Hilbert spaces around z_0 and produce a representation H_{z_0} of the Berezin-Toeplitz deformation quantization algebra $(C^\infty(X)[[\hbar]], \star_{BT})$.

In a suitably chosen coordinates (and frame of L) around z_0 , $S_{m,p,r}$ is equal to $z_1^{p_1} \cdots z_n^{p_n}$ up to order $2r - 1$. Because of the error terms, the peak sections in $\prod_m H^0(X, L^m)$ with a fixed r behave in a compatible way with the actions of $T_{z_j} = m_{z_j}$ and $T_{\bar{z}_j} = \hbar \frac{d}{dz_j}$ around z_0 only up to order $2r - 1$, which is *not* enough to produce a representation of $(C^\infty(X)[[\hbar]], \star_{BT})$. To construct our representation H_{z_0} , we need to find a clever way to increase the order r of Taylor expansions at z_0 to infinity when the choices of peak sections $S_{m,p,r}$'s are changing in r .

To achieve this, we consider $\sum_m \alpha_{m,r} \in \prod_m H^0(X, L^{\otimes m})$, which represents a sum of peak sections of various tensor powers $L^{m'}$'s and where we are actually only keeping track of its Taylor expansion around z_0 up to order $2r - 1$. A key observation is that $\{\sum_m \alpha_{m,r}\}_{r=1}^\infty$, or simply a double sequence $\{\alpha_{m,r}\}$, will define elements in $\prod_m H^0(X, L^{\otimes m})$ with more and

more terms of their Taylor expansions around z_0 being identical if the following condition holds: there exists a sequence of complex numbers $\{a_{p,k}\}_{p,k \geq 0}$ such that, for each fixed $r > 0$, we have the following estimates:

$$(1.2) \quad \langle \alpha_{m,r} - \sum_{2k+|p| \leq r} a_{p,k} \cdot \frac{1}{m^k} \cdot S_{m,p,r+1}, S_{m,q,r+1} \rangle_m = O\left(\frac{1}{m^{r+1}}\right),$$

for any multi-index q with $|q| \leq r$.

We call such $\{\alpha_{m,r}\}$ an *admissible sequence at z_0* , and they span a linear subspace $V_{z_0} \subset \prod_r (\prod_m H^0(X, L^{\otimes m}))$. In fact, only the coefficients $\{a_{p,k}\}$ of $\{\alpha_{m,r}\}$ would record the whole Taylor expansions at z_0 . Thus this defines an equivalence relation \sim on V_{z_0} and the desired vector space can be constructed as

$$H_{z_0} := V_{z_0} / \sim.$$

Theorem 1.1 (=Theorem 3.16). *The vector space H_{z_0} is a representation of the Berezin-Toeplitz deformation quantization algebra $(C^\infty(X)[[\hbar]], \star_{BT})$.*

We will prove that this representation possesses various nice properties, as expected from the physical point of view. First of all, it is *local*, namely, for any smooth function $f \in C^\infty(X)$, the action of the Toeplitz operator T_f on H_{z_0} depends only on the infinite jets of f at z_0 ; this will be proved in Theorem 3.18. Also, for every real-valued function f , the operator T_f on H_{z_0} is self-adjoint; see Proposition 3.14. Last but not the least, it is irreducible in a suitable sense, as we will see in Theorem 3.21.

Remark 1.2. The space H_{z_0} has the structure of a formal Hilbert space with inner products taking values in $\mathbb{C}[[\hbar]]$, as in the work [5] of Bordemann-Waldmann, where they gave some algebraic constructions of representations of deformation quantization algebras.

On the other hand, our results are closely related to the work [27] of Reshetikhin-Takhtajan, where they related the star product to formal Feynman-Laplace expansions of formal integrals.

Acknowledgement.

We thank Si Li and Siye Wu for useful discussions. The first named author thanks Martin Schlichenmaier and Siye Wu for inviting him to attend the conference GEOQUANT 2019 held in September 2019 in Taiwan, in which he had stimulating and very helpful discussions with both of them as well as Jørgen Ellegaard Andersen, Motohico Mulase, Georgiy Sharygin and Steve Zelditch.

K. Chan was supported by grants of the Hong Kong Research Grants Council (Project No. CUHK14302617 & CUHK14303019) and direct grants from CUHK. N. C. Leung was supported by grants of the Hong Kong Research Grants Council (Project No. CUHK14301117 & CUHK14303518) and direct grants from CUHK. Q. Li supported by a grant from National Natural Science Foundation of China for young scholars (Project No. 11501537).

2. THE FEYNMAN-LAPLACE THEOREM AND PERTURBATIONS OF THE BARGMANN-FOCK SPACE

In this section, we perform the local computations needed for proving Theorem 1.1.

2.1. The Bargmann-Fock space and Wick algebra.

Recall that in the flat case when $X = \mathbb{C}^n$ (and prequantum line bundle L is trivial), the Hilbert space on which the Toeplitz operators act is the well-known *Bargmann-Fock space* $\mathcal{HL}^2(\mathbb{C}^n, \mu_{\hbar})$ consisting of L^2 integrable entire holomorphic functions with respect to the density $\mu_{\hbar}(z) = (\pi\hbar)^{-n}e^{-|z|^2/\hbar}$; here \hbar is regarded as a positive real number. It is easy to see, by direct computations, that the holomorphic polynomials

$$\frac{z^I}{\sqrt{I!\hbar^{|I|}}},$$

where I runs over all multi-indices, form an orthonormal basis of $\mathcal{HL}^2(\mathbb{C}^n, \mu_{\hbar})$.

We will only need the Toeplitz operators associated to polynomials, i.e., multiplying by a polynomial $f \in \mathbb{C}[z, \bar{z}]$ which is in general non-holomorphic and then projecting back to the holomorphic subspace. For example, when $n = 1$, we have

$$T_z = m_z, \quad T_{\bar{z}} = \hbar \frac{d}{dz}, \quad T_{f_1(z)f_2(\bar{z})} = f_2 \left(\hbar \frac{d}{dz} \right) \circ m_{f_1(z)},$$

Let $f, g \in \mathbb{C}[z, \bar{z}]$. Then there is a formula for the composition of Toeplitz operators: $T_f \circ T_g = T_{f \star g}$, where

$$f \star g := \exp \left(-\hbar \sum_{i=1}^n \frac{\partial}{\partial z_i} \frac{\partial}{\partial \bar{w}_i} \right) (f(z, \bar{z})g(w, \bar{w}))|_{z=w}.$$

This product can be extended to formal power series, giving the definition of the Wick algebra:

Definition 2.1. The *Wick algebra* is $\mathcal{W}_{\mathbb{C}^n} := \mathbb{C}[[y, \bar{y}]][[\hbar]]$ equipped with the multiplication:

$$(2.1) \quad f \star g := \exp \left(-\hbar \sum_{i=1}^n \frac{\partial}{\partial y_i} \frac{\partial}{\partial \bar{y}'_i} \right) (f(y, \bar{y})g(y', \bar{y}'))|_{y=y'}.$$

Remark 2.2. Here we use y, \bar{y} instead of z, \bar{z} in order to distinguish functions on \mathbb{C}^n and Wick algebra in later sections.

There is an action of the Wick algebra on the Bargmann-Fock space

$$\mathcal{F}_{\mathbb{C}^n} := \mathbb{C}[[y_1, \dots, y_n]][[\hbar]],$$

which we call the *Bargmann-Fock action*, defined as follows: let $y_{i_1} \cdots y_{i_k} \bar{y}_{j_1} \cdots \bar{y}_{j_l}$ be any monomial in the Wick algebra, then we turn it to a differential operator on $\mathcal{F}_{\mathbb{C}^n}$ by

$$(2.2) \quad y_{i_1} \cdots y_{i_k} \bar{y}_{j_1} \cdots \bar{y}_{j_l} \mapsto \left(\hbar \cdot \frac{\partial}{\partial y_{j_1}} \right) \circ \cdots \circ \left(\hbar \cdot \frac{\partial}{\partial y_{j_l}} \right) \circ m_{y_{i_1} \cdots y_{i_k}};$$

in other words, holomorphic polynomials are mapped to creation operators and anti-holomorphic ones are mapped to annihilation operators. This assignment is also known as the *Wick normal ordering*.

Remark 2.3. There is another Fock representation on $\mathcal{W}_{\mathbb{C}^n}$ where $y^{i'}$'s act as annihilators and $\bar{y}^{j'}$'s act as creators. Explicitly, $\mathcal{F}'_{\mathbb{C}^n} := \mathbb{C}[[\bar{y}_1, \cdots, \bar{y}_n]][[\hbar]]$, and the operators associated to a monomial is given by

$$y_{i_1} \cdots y_{i_k} \bar{y}_{j_1} \cdots \bar{y}_{j_l} \mapsto m_{\bar{y}_{j_1} \cdots \bar{y}_{j_l}} \circ \left(-\hbar \cdot \frac{\partial}{\partial \bar{y}_{i_1}} \right) \circ \cdots \circ \left(-\hbar \cdot \frac{\partial}{\partial \bar{y}_{i_k}} \right)$$

This Fock representation is equivalent to the GNS representation on \mathbb{C}^n in [5].

Notation 2.4. Throughout this paper, we will use the following notation for multi-indices: let $I = (i_1, \cdots, i_n)$ and $J = (j_1, \cdots, j_m)$ then we set

$$y^I := y_1^{i_1} \cdots y_n^{i_n}, \quad \bar{y}^J := \bar{y}_1^{j_1} \cdots \bar{y}_n^{j_m}.$$

We also use the notations: $|I| := i_1 + \cdots + i_n$ and $I! := i_1! \cdots i_n!$.

Let I, J be as above. We assign a \mathbb{Z} -grading on $\mathcal{W}_{\mathbb{C}^n}$ by letting the monomial $\hbar^k y^I \bar{y}^J$ to have degree $2k + |I| + |J|$. There is an associated decreasing filtration on $\mathcal{W}_{\mathbb{C}^n}$ given by the set $(\mathcal{W}_{\mathbb{C}^n})_k$ of power series in $\mathcal{W}_{\mathbb{C}}$ whose terms are all of degree $\geq k$. In a similar way, we can define a grading and filtration on both $\mathcal{F}_{\mathbb{C}^n}$ and $\mathbb{C}[[\hbar]]$. Note that this grading is preserved by both the Wick product and the Bargmann-Fock action.

We will also need the following extension of the Wick algebra:

Definition 2.5. The extension $\mathcal{W}_{\mathbb{C}^n}^+$ of the Wick algebra $\mathcal{W}_{\mathbb{C}^n}$ is defined as follows:

- Elements of $\mathcal{W}_{\mathbb{C}^n}^+$ are given by power series, possibly with negative powers of \hbar ,
- The degrees of terms of an element $U \in \mathcal{W}_{\mathbb{C}^n}^+$ have a uniform lower bound which could be negative; equivalently, there exists $k \geq 0$ so that every term in $\hbar^k \cdot U$ has non-negative degree.
- For an element $U \in \mathcal{W}_{\mathbb{C}^n}^+$, there exists a finite number of terms for any given non-negative total degree.

Remark 2.6. The definition of $\mathcal{W}_{\mathbb{C}^n}^+$ here is *different* from the one in [11, p. 224]; in that definition, monomials in $\mathcal{W}_{\mathbb{C}^n}^+$ must have non-negative total degrees. Our extension will be important later for proving the irreducibility of our representation.

Note that $\mathcal{W}_{\mathbb{C}^n}^+$ is closed under the Wick product. One of the motivations of this extension is to allow the following exponentials:

Example 2.7. Let $H \in (\mathcal{W}_{\mathbb{C}^n})_3$, i.e., every term in H is of degree at least 3, then the following classical and quantum exponentials both live in $\mathcal{W}_{\mathbb{C}^n}^+$:

$$\begin{aligned}\exp(H/\hbar) &= 1 + \frac{H}{\hbar} + \frac{1}{2!} \frac{H \cdot H}{\hbar^2} + \cdots, \\ \exp^*(H/\hbar) &= 1 + \frac{H}{\hbar} + \frac{1}{2!} \frac{H \star H}{\hbar^2} + \cdots\end{aligned}$$

Notation 2.8. In this paper, we will use the notation $e^{H/\hbar}$ to denote the classical exponential of H/\hbar .

We can define $\mathcal{F}_{\mathbb{C}^n}^+$ in a similar way. It is clear that there is a naturally extended Bargmann-Fock action of $\mathcal{W}_{\mathbb{C}^n}^+$ on $\mathcal{F}_{\mathbb{C}^n}^+$. The following lemma shows that the subspace $\mathcal{F}_{\mathbb{C}^n} \subset \mathcal{F}_{\mathbb{C}^n}^+$ is closed under the action of elements in $\mathcal{W}_{\mathbb{C}^n}^+$ of a special form.

Lemma 2.9. Suppose $H = \sum_{k, |I| \geq 0, |J| \geq 0} \hbar^k a_{k, I, J} \cdot y^I \bar{y}^J \in (\mathcal{W}_{\mathbb{C}^n})_3$ has no purely holomorphic terms, i.e., $a_{k, I, 0} = 0$. Then $\mathcal{F}_{\mathbb{C}^n}$ is closed under the action of $\exp(H/\hbar)$ and $\exp^*(H/\hbar)$.

Proof. Each \bar{y}^j in every monomial of H/\hbar acts as $\hbar \frac{\partial}{\partial y^j}$, and the \hbar in this differential operator will cancel with the \hbar in the denominator. So the output can only have nonnegative powers of \hbar . \square

Lemma 2.10. The classical exponential $\exp(H/\hbar)$, where $H \in (\mathcal{W}_{\mathbb{C}^n})_3$, can also be written as a quantum exponential:

$$\exp(H/\hbar) = \exp^*(H'/\hbar),$$

with $H' \in \mathcal{W}_{\mathbb{C}^n}^+$. In particular, $\exp(H/\hbar)$ is invertible in $\mathcal{W}_{\mathbb{C}^n}^+$ and

$$(\exp(H/\hbar))^{-1} = \exp^*(-H'/\hbar).$$

Proof. Let $A = \exp(H/\hbar) - 1 \in (\mathcal{W}_{\mathbb{C}^n})_1$. Then H' is defined via the following formal logarithm with respect to the quantum product \star :

$$H' = \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k} A^k,$$

where $A^k = A \star A \star \cdots \star A$ denotes the k -th power with respect to the quantum product. The fact that $A \in (\mathcal{W}_{\mathbb{C}^n})_1$ implies that each term of H' is of positive degree, and in each degree there are only finitely many terms in H' , i.e., $H' \in \mathcal{W}_{\mathbb{C}^n}^+$. \square

2.2. Formal Hilbert spaces.

In the Kähler geometry setting, we cannot in general reduce to the local model of the Bargmann-Fock space, and will need to consider a more general situation. For this purpose, we need the following theorem:

Theorem 2.11 (Feynman-Laplace). *Let X be a compact n -dimensional manifold (possibly with boundary), and let f be a smooth function attaining a unique minimum on X at an interior point $x_0 \in X$, and assume that the Hessian of f is non-degenerate at x_0 ; also, let $\mu = \alpha(x) \cdot e^{g(x)} d^n x$ be a top-degree form. Then the integral*


$$I(\hbar) := \int_X \mu e^{-\frac{1}{\hbar}f(x)} = \int_X \alpha(x) \cdot e^{-\frac{f(x)+\hbar g(x)}{\hbar}} dx_1 \cdots dx_n,$$

has the following asymptotic expansion as $\hbar \rightarrow 0^+$:

$$I(\hbar) \sim \sum_{k \geq 0} a_k \cdot \hbar^k,$$

where each coefficient a_k is a sum of Feynman weights which depends only on the infinite jets of the functions f, g at the point x_0 .

More explicitly, each a_k is a sum over connected graphs of genus k . Recall that the genus of a graph γ is the sum of the genera of the vertices in γ (in our situation, each vertex has genus either 0 or 1, labeled by $f(x)$ and $g(x)$ respectively, since the integrand is $e^{-\frac{f(x)+\hbar g(x)}{\hbar}}$), and $k = 1 - \chi(\gamma)$ where $\chi(\gamma)$ denotes the Euler characteristic of γ . The propagator in the Feynman weights is given by the inverse of the Hessian of f at x_0 . The following picture shows a Feynman graph:



Here every vertex labeled by f must be at least trivalent, and every vertex labeled by $\hbar \cdot g$ must be univalent. For more details on the Feynman-Laplace Theorem, we refer the readers to Pavel Mnev's excellent exposition in [25], and for a detailed exposition of Feynman graph computations, we refer the readers to [7].

We will mainly apply the Feynman-Laplace Theorem to a function $f(z, \bar{z})$ on a closed disk $\mathbb{D}^{2n} \subset \mathbb{C}^n$ such that the origin $0 \in \mathbb{D}^{2n}$ is the unique minimum of f and $f(0) = 0$. So, by taking an appropriate complex coordinate system $z = (z_1, \dots, z_n)$ centered at 0, the Taylor expansion of f at the origin is given by

$$f(z, \bar{z}) = |z|^2 + O(|z|^3).$$

Theorem 2.11 gives an asymptotic expansion of the following integral:

$$\frac{(\sqrt{-1})^n}{\hbar^n} \int_{\mathbb{D}^{2n}} h(z, \bar{z}) e^{-\frac{f(z, \bar{z}) + \hbar g(z, \bar{z})}{\hbar}} dz_1 d\bar{z}_1 \cdots dz_n d\bar{z}_n.$$

Remark 2.12. The above integral clearly depends on the radius of \mathbb{D}^{2n} , but its asymptotic expansion is actually independent of the radius.

Theorem 2.11 implies that the asymptotic expansion of the above integral depends only on the Taylor expansions of the functions f, g and h at the origin. We can thus replace these functions by formal power series in $\mathcal{W}_{\mathbb{C}^n}$, and define a formal integral:

Definition 2.13. For $\phi(y, \bar{y}) \in (\mathcal{W}_{\mathbb{C}^n})_3$ and $h(y, \bar{y}) \in \mathcal{W}_{\mathbb{C}^n}$, we define the following formal integral:

$$\frac{1}{\hbar^n} \int h(y, \bar{y}) \cdot e^{\frac{-|y|^2 + \phi(y, \bar{y})}{\hbar}} \in \mathbb{C}[[\hbar]]$$

via the Feynman rule in Theorem 2.11.

Remark 2.14. We omit the standard differential form $(\sqrt{-1})^n dz_1 d\bar{z}_1 \cdots dz_n d\bar{z}_n$ in the notation of formal integral.

Lemma 2.15. *The formal integral preserves the decreasing filtration on $\mathcal{W}_{\mathbb{C}^n}$ and $\mathbb{C}[[\hbar]]$; more precisely, if $h(y, \bar{y}) \in (\mathcal{W}_{\mathbb{C}^n})_k$, then the formal integral lies in $(\mathbb{C}[[\hbar]])_k$.*

Proof. The leading term of the formal integral

$$\frac{1}{\hbar^n} \int h(y, \bar{y}) \cdot e^{\frac{-|y|^2 + \phi(y, \bar{y})}{\hbar}} \in \mathbb{C}[[\hbar]]$$

is the same as that in the free case, i.e., when $\phi = 0$. Thus the leading term of the integral have the same degree as the leading degree of $h(y, \bar{y})$. \square

Using this formal integral, we can define a *Hilbert space in the formal sense*, namely, its inner product takes values in the formal Laurent series $\mathbb{C}((\sqrt{\hbar}))$:

Definition 2.16. On the $\mathbb{C}((\sqrt{\hbar}))$ -vector space $\mathcal{W}_{\mathbb{C}^n} \otimes_{\mathbb{C}[[\hbar]]} \mathbb{C}((\sqrt{\hbar}))$, we define a complex conjugation by extending the complex conjugation on polynomials in \mathbb{C}^n :

$$(\sqrt{\hbar})^k a_{I,J} y^I \bar{y}^J \mapsto (\sqrt{\hbar})^k \bar{a}_{I,J} \bar{y}^I y^J.$$

Fix $\phi(y, \bar{y}) \in (\mathcal{W}_{\mathbb{C}^n})_3$. Then for $f, g \in \mathcal{W}_{\mathbb{C}^n}((\sqrt{\hbar}))$, we define their *formal inner product* as the following formal integral:

$$(2.3) \quad \langle f, g \rangle := \frac{1}{\hbar^n} \cdot \int f \bar{g} \cdot e^{\frac{-|y|^2 + \phi(y, \bar{y})}{\hbar}},$$

which is in turn defined using Feynman graph expansions as in Definition 2.13 and takes value in $\mathbb{C}((\sqrt{\hbar}))$.

The following are some simple properties of this formal inner product.

Lemma 2.17. *Suppose that ϕ is real, i.e., $\phi = \bar{\phi}$. Then the formal inner product (2.3) is Hermitian, namely, $\langle f, g \rangle = \overline{\langle g, f \rangle}$.*

Lemma 2.15 implies the following

Corollary 2.18. *The formal inner product of $f, g \in \mathcal{W}_{\mathbb{C}^n}^+$ is a formal power series in \hbar , i.e., $\langle f, g \rangle \in \mathbb{C}[[\hbar]]$.*

Remark 2.19. We allow ϕ to have \hbar -dependence. In particular, the fact that ϕ is of at least degree 3 guarantees that the graph expansion of (2.3) is valid. In the Kähler geometry setting, f will be given by the logarithm of the norm of a local holomorphic frame of the prequantum line bundle, and g will be the logarithm of the volume form.

The following lemma explains the reason for considering an extension of $\mathcal{W}_{\mathbb{C}^n}$ by $\mathbb{C}((\hbar))$:

Lemma 2.20. *The holomorphic polynomials*

$$\frac{y^I}{\sqrt{I!\hbar^{|I|}}}$$

form a basis of the formal Hilbert space, which is orthonormal modulo \hbar , i.e.,

$$\left\langle \frac{y^I}{\sqrt{I!\hbar^{|I|}}}, \frac{y^J}{\sqrt{J!\hbar^{|J|}}} \right\rangle = \delta_{I,J} + O(\hbar).$$

Proof. The proof for the cases where $I = J$ is obvious since the computation of the leading term is the same as that in the Bargmann-Fock space. For the cases where $I \neq J$, the terms $y^I \bar{y}^J$ cannot be fully contracted using the quadratic part $-|y|^2/\hbar$. The “interaction” part $e^{\phi/\hbar}$ needs to come in so that we can get a full contraction which takes value in $\mathbb{C}[[\hbar]]$. Notice that this contraction preserves the filtration induced by the grading on $\mathcal{W}_{\mathbb{C}^n}$ and $\mathbb{C}[[\hbar]]$. Now $\frac{y^I}{\sqrt{I!\hbar^{|I|}}}, \frac{\bar{y}^J}{\sqrt{J!\hbar^{|J|}}}$ have degree 0, and all terms in $e^{\phi/\hbar}$ have degrees strictly greater than 0. The result follows. \square

Corollary 2.21. *Given two different multi-indices $I \neq J$, we have the following asymptotics:*

$$\hbar^{-n} \int y^I \bar{y}^J e^{\frac{-|y|^2 + \phi(y, \bar{y})}{\hbar}} = O(\hbar^{\max\{|I|, |J|\}}).$$

Let $\phi = \sum_{k,I,J} \hbar^k \phi_{k,I,J} y^I \bar{y}^J$, and suppose ϕ satisfies the property that $\phi_{0,I,J} = 0$ if either $|I| = 1$ or $|J| = 1$. Then we further have the refinement:

$$\hbar^{-n} \int y^I \bar{y}^J e^{\frac{-|y|^2 + \phi(y, \bar{y})}{\hbar}} = o(\hbar^{\max\{|I|, |J|\}}).$$

Proof. Let $K = (k_1, \dots, k_n)$ be the multi-index given by $k_l = \max\{i_l, j_l\}, 1 \leq l \leq n$. The worst scenario is when $y^I \bar{y}^J$ together with terms in $e^{\phi/\hbar}$ form a multiple of $y^K \bar{y}^K$ so that we get full contraction. These terms coming from $e^{\phi/\hbar}$ must be a multiple of

$$\hbar^{-l} \cdot y^{K-I} \bar{y}^{K-J}$$

for some l . Thus the leading term of the integral in the statement is $O(\hbar^{|K|-l})$. If $l \leq 0$, then there is nothing to show since $|K| - l \geq |K| \geq \max\{|I|, |J|\}$. Thus we assume that $l > 0$. Since every monomial in ϕ/\hbar contains at least one $y^{i'}$ s, it follows that

$$(2.4) \quad l \leq |K - I|.$$

From Lemma 2.20, we know that

$$\hbar^{-n} \int y^I \bar{y}^J \cdot \frac{1}{\hbar^l} y^{K-I} \bar{y}^{K-J} = \hbar^{-n} \int \frac{y^K \bar{y}^K}{\hbar^l} = O\left(\hbar^{|K|-l}\right).$$

The statement follows since $|K| - l \geq |K| - |K - I| = |I|$, and also $|K| - l \geq |J|$ by a similar argument. For the refinement under the additional condition on ϕ , we only need to notice that the inequality (2.4) can be refined to

$$l \leq \frac{|K - I|}{2} < |K - I|.$$

□

We want to define the notion of orthogonal projection and formal Toeplitz operators using this formal inner product. To do so, we need the following technical theorem (which is also important in the sequel [6] to this paper):

Theorem 2.22. *Suppose $\phi \in (\mathcal{W}_{\mathbb{C}^n})_3$ contains no purely holomorphic monomials. For any $f \in \mathcal{W}_{\mathbb{C}^n}$, there exists a unique $O_f \in \mathcal{W}_{\mathbb{C}^n}$ such that*

- (1) *For any $s \in \mathcal{F}_{\mathbb{C}^n}$, the element $T_{O_f}(s) \in \mathcal{F}_{\mathbb{C}^n}$ satisfies the following equalities:*

$$\langle T_{O_f}(s), y^I \rangle = \langle f \cdot s, y^I \rangle,$$

for every multi-index I ; here T_{O_f} denotes the Bargmann-Fock action by O_f .

- (2) *IF f is a monomial, then the leading term of O_f is exactly f , i.e.,*

$$O_f = f + \text{higher order terms.}$$

Proof. Given $s \in \mathcal{F}_{\mathbb{C}^n}$, suppose there exists $s' \in \mathcal{F}_{\mathbb{C}^n}$ such that

$$(2.5) \quad T_{f \cdot e^{\phi/\hbar}}(s) = T_{e^{\phi/\hbar}}(s').$$

There is the following straightforward computation:

$$\begin{aligned} \int \left(e^{\phi/\hbar} \cdot s' \right) \cdot \bar{y}^I \cdot e^{-\frac{|y|^2}{\hbar}} &= \int T_{e^{\phi/\hbar}}(s') \cdot \bar{y}^I \cdot e^{-\frac{|y|^2}{\hbar}} \\ &= \int T_{f \cdot e^{\phi/\hbar}}(s) \cdot \bar{y}^I \cdot e^{-\frac{|y|^2}{\hbar}} \\ &= \int \left(f \cdot e^{\phi/\hbar} \cdot s \right) \cdot \bar{y}^I \cdot e^{-\frac{|y|^2}{\hbar}} \end{aligned}$$

for any purely antiholomorphic monomial \bar{y}^I ; here the first equality follows from the fact that $T_{e^{\phi/\hbar}}(s')$ is the orthogonal projection of $e^{\phi/\hbar} \cdot s'$ with respect to the standard Gaussian measure. So for every multi-index I , we have

$$\int s' \cdot \bar{y}^I \cdot e^{-\frac{|y|^2 + \phi(y, \bar{y})}{\hbar}} = \int f s \cdot \bar{y}^I \cdot e^{-\frac{|y|^2 + \phi(y, \bar{y})}{\hbar}}.$$

Hence we only need to solve the equation (2.5) for s' .

By Lemma 2.10, $e^{\phi/\hbar}$ is invertible under the Wick product and its inverse is:

$$\left(e^{\phi/\hbar}\right)^{-1} = \exp^* \left(- \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k} (e^{\phi/\hbar} - 1)^k \right).$$

From the following expansion

$$- \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k} (e^{\phi/\hbar} - 1)^k = - \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k} \left(\frac{\phi}{\hbar} + \frac{1}{2!} \frac{\phi^2}{\hbar^2} + \dots \right)^k,$$

it is easy to see that each monomial in the expansion of $(e^{\phi/\hbar})^{-1}$ satisfies the following property: in every \hbar^{-k} term, the antiholomorphic components must have degrees at least k . By a similar argument as in the proof of Lemma 2.9, we see that there is a well-defined action of $(e^{\phi/\hbar})^{-1}$ on $\mathcal{F}_{\mathbb{C}^n}$.

Therefore we get the following explicit description of s' :

$$(2.6) \quad s' = T_{(e^{\phi/\hbar})^{-1}} \circ T_{f \cdot e^{\phi/\hbar}}(s).$$

The next step is to look at the term $T_{f \cdot e^{\phi/\hbar}}(s)$ more closely. According to (2.1), we have

$$f \cdot e^{\phi/\hbar} = e^{\phi/\hbar} \star f - \sum_{k \geq 1} \hbar^k C_k \left(e^{\phi/\hbar}, f \right).$$

Since $C_k(-, -)$ is a bi-differential operator, the term $C_k(e^{\phi/\hbar}, f)$ is still of the form $e^{\phi/\hbar} \cdot g_k(y, \bar{y})$ for some $g_k \in \mathcal{W}_{\mathbb{C}^n}$ and satisfies the condition that

$$\text{leading degree of } g_k - \text{leading degree of } f \geq k.$$

Thus by an induction on the degree, this procedure can be iterated, and we can find $O_f \in \mathcal{W}_{\mathbb{C}^n}$ whose first terms are exactly f such that $f \cdot e^{\phi/\hbar} = e^{\phi/\hbar} \star O_f$. This implies that

$$(2.7) \quad T_{f \cdot e^{\phi/\hbar}}(s) = (T_{e^{\phi/\hbar}} \circ T_{O_f})(s).$$

In particular, we see that $s' = T_{O_f}(s)$. □

By the first statement of this theorem, we have the following:

Definition 2.23. The *orthogonal projection operator*

$$(2.8) \quad \pi_\phi : \mathcal{W}_{\mathbb{C}^n} \rightarrow \mathcal{F}_{\mathbb{C}^n} = \mathbb{C}[[y_1, \dots, y_n]][[\hbar]]$$

is defined by requiring that

$$\langle f, y^I \rangle = \langle \pi_\phi(f), y^I \rangle$$

for all multi-indices I ; here $\langle -, - \rangle$ is the inner product defined by equation (2.3).

We can also define the formal Toeplitz operators:

Definition 2.24. The *formal Toeplitz operator* $T_{\phi, f}$ associated to $f \in \mathcal{W}_{\mathbb{C}^n}$ is defined as the composition of multiplication by f and the projection π_ϕ :

$$T_{\phi, f} := \pi_\phi \circ m_f.$$

Theorem 2.22 gives an explicit algorithm to compute $T_{\phi,f}$: we only need to find $O_f \in \mathcal{W}_{\mathbb{C}^n}$ associated to f , and then $T_{\phi,f} = T_{O_f}$. A simple observation is that if $f = f(y)$ is a holomorphic power series, then T_f is simply the multiplication m_f since then $O_f = f$.

Here we give a description of the adjoint operator of a formal Toeplitz operator:

Lemma 2.25. *Suppose $\phi \in \mathcal{W}_{\mathbb{C}^n}$ is real, i.e., $\phi = \bar{\phi}$. Then for any $f \in \mathcal{W}_{\mathbb{C}^n}$, the adjoint of the formal Toeplitz operator $T_{\phi,f}$ is given by $T_{\phi,\bar{f}}$. In particular, $T_{\phi,f}$ is self-adjoint if and only if $f \in \mathcal{W}_{\mathbb{C}^n}$ is real.*

Proof. According to the definition of formal Toeplitz operators, for any elements $s_1, s_2 \in \mathcal{F}_{\mathbb{C}^n}$, we have

$$\begin{aligned} \langle T_{\phi,f}(s_1), s_2 \rangle &= \int T_{\phi,f}(s_1) \cdot \bar{s}_2 \cdot e^{\frac{-|y|^2 + \phi(y,\bar{y})}{\hbar}} \\ &= \int f \cdot s_1 \cdot \bar{s}_2 \cdot e^{\frac{-|y|^2 + \phi(y,\bar{y})}{\hbar}} \\ &= \int s_1 \cdot \overline{\bar{f}} \cdot s_2 \cdot e^{\frac{-|y|^2 + \phi(y,\bar{y})}{\hbar}} \\ &= \langle s_1, T_{\phi,\bar{f}}(s_2) \rangle. \end{aligned}$$

□

2.3. Local asymptotics via formal Hilbert space.

Let $\mathbb{D}^{2n} \subset \mathbb{C}^n$ be a ball centered at 0, with $dvol_{\mathbb{D}^{2n}} := (\sqrt{-1})^n e^{\psi(z,\bar{z})} dz_1 d\bar{z}_1 \cdots dz_n d\bar{z}_n$ the volume form. For every smooth function f on \mathbb{D}^{2n} , we will let $J_f \in \mathcal{W}_{\mathbb{C}^n}$ denote the Taylor expansion of f at the origin:

$$J_f := \sum_{I,J \geq 0} \frac{1}{I!J!} \frac{\partial^{|I|+|J|} f}{\partial z^I \partial \bar{z}^J}(0) y^I \bar{y}^J.$$

The previous algebraic computations together with the Feynman-Laplace Theorem 2.11 give the following asymptotics as $\hbar \rightarrow 0^+$:

Theorem 2.26. *Suppose $\varphi(z,\bar{z})$ is a smooth function on \mathbb{D}^{2n} which attains its unique minimum at the origin. Let f, φ, s be functions on \mathbb{D}^{2n} such that $\bar{\partial}s = 0$, φ has a unique minimum at the origin and satisfies*

$$(2.9) \quad J_\varphi = |y|^2 + \sum_{I,J \geq 2} \frac{1}{I!J!} \frac{\partial^{|I|+|J|} \varphi}{\partial z^I \partial \bar{z}^J}(0) y^I \bar{y}^J.$$

There exist complex numbers $a_{k,I}$ so that for every fixed multi-index J , we have the following asymptotics as $\hbar \rightarrow 0$:

$$(2.10) \quad \frac{1}{\hbar^n} \int_{\mathbb{D}^{2n}} \left(f \cdot s - \sum_{2k+|I| \leq r} \frac{1}{\hbar^k} a_{k,I} z^I \right) \cdot \bar{z}^J e^{-\frac{\varphi(z,\bar{z})}{\hbar}} dvol_{\mathbb{D}^{2n}} = O(\hbar^{r+1}).$$

In particular, these $a_{k,I}$'s only depend on the Taylor expansions of f, s, φ and ψ at the origin.

Proof. We define a function $\phi = |z|^2 - \varphi(z, \bar{z}) + \hbar\psi(z, \bar{z})$. Then we define $a_{k,I}$'s via the following equation:

$$\sum_{k, |I| \geq 0} a_{k,I} \hbar^k \cdot z^I = T_{(e^{J\phi/\hbar})^{-1}} \circ T_{J_f \cdot e^{J\phi/\hbar}}(J_s).$$

From Theorem 2.11, we have, for any \bar{z}^J , the following equality of asymptotic \hbar -expansions

$$\frac{1}{\hbar^n} \cdot \int_{\mathbb{D}^{2n}} f \cdot s \cdot \bar{z}^J e^{-\frac{\varphi(z, \bar{z})}{\hbar}} d\text{vol}_{\mathbb{D}^{2n}} = \frac{1}{\hbar^n} \int J_f \cdot J_s \cdot \bar{y}^J e^{-\frac{|y|^2 + J\phi}{\hbar}}.$$

On the other hand, there is the following identity by Theorem 2.22:

$$\frac{1}{\hbar^n} \int J_f \cdot J_s \cdot \bar{y}^J e^{-\frac{|y|^2 + J\phi}{\hbar}} = \frac{1}{\hbar^n} \int \left(\sum_{k, |I| \geq 0} a_{k,I} \hbar^k \cdot y^I \right) \cdot \bar{y}^J e^{-\frac{|y|^2 + J\phi}{\hbar}}.$$

Now equation (2.10) follows from Corollary 2.21 since the truncated higher order terms will only contribute to integrals of type $o(\hbar^{r+1})$. \square

3. GEOMETRIC REPRESENTATIONS OF THE BEREZIN-TOEPLITZ QUANTIZATION

In this section, we construct a family of representations of the Berezin-Toeplitz deformation quantization parametrized by the points in the Kähler manifold X , and describe its basic properties such as locality and irreducibility.

The organization of this section is as follows: In Section 3.1, we study \mathbb{C}^n as the motivating example to illustrate the idea behind the general definition of admissible sequences. In Section 3.2, we construct representations of the Berezin-Toeplitz deformation quantization using peak sections (whose properties are reviewed in Section A), which reduce the proof of Theorem 1.1 to the local computations done in Section 2. In Section 3.3, we prove locality and (modified) irreducibility of our representations.

3.1. Admissible sequences on \mathbb{C}^n .

Recall that the prequantum line bundle L on \mathbb{C}^n is trivialized by a global holomorphic frame $\mathbb{1}$. The Hermitian inner product of $L^{\otimes m}$ under this trivialization is given by

$$h^m(\mathbb{1}^{\otimes m}, \mathbb{1}^{\otimes m}) = e^{-m \cdot |z|^2}.$$

We would like to define an action of $C^\infty(\mathbb{C}^n)[[\hbar]]$ on $\mathcal{F}_{\mathbb{C}^n}$ such that its restriction to polynomials is exactly the Bargmann-Fock action. To do this, we first apply asymptotic analysis to give an equivalent description of $\mathcal{F}_{\mathbb{C}^n}$.

First of all, we consider the vector space

$$V := \prod_{r \geq 0} \left(\prod_{m \geq 0} H^0(\mathbb{C}^n, L^{\otimes m}) \right),$$

an element of which is a double sequence $\alpha = (\alpha_{m,r})$ with $\alpha_{m,r} \in H^0(\mathbb{C}^n, L^{\otimes m})$.

We also consider the map

$$(3.1) \quad F : \mathcal{F}_{\mathbb{C}^n} \rightarrow V, \quad a = \sum_{k,I} a_{k,I} \hbar^k z^I \mapsto \alpha = \{\alpha_{m,r}\},$$

defined by setting

$$\alpha_{m,r} := \left(\sum_{2k+|I| \leq r} m^{-k} \cdot a_{k,I} z^I \right) \otimes \mathbb{1}^m \in H^0(\mathbb{C}^n, L^{\otimes m}).$$

If the element in $\mathcal{F}_{\mathbb{C}^n}$ is of the form $\sum_I a_I z^I$, i.e., it does not include \hbar , then each $\alpha_{m,r}$ is a holomorphic section of $L^{\otimes m}$ which is a polynomial of degree $\leq r$ truncated from α under the trivialization $\mathbb{1}^{\otimes m}$. For general elements, \hbar^k is mapped to $1/m^k$ in the corresponding components of the double sequence.

We now consider an action of smooth functions on the image of the map (3.1). Let f be any smooth function on \mathbb{C}^n . It is clear that the Toeplitz operator $T_{f,m}$ is in general not well-defined since \mathbb{C}^n is noncompact. We apply the asymptotic analysis in Section 2.3 to obtain the following

Proposition 3.1. *Let f be any smooth function on \mathbb{C}^n , and let $F(a) = \{\alpha_{m,r}\}$ be defined as above. Then there exists $b = \sum_{k,I} \hbar^k b_{k,I} z^I \in \mathcal{F}_{\mathbb{C}^n}$ with $F(b) = \{\beta_{r,m}\}$, satisfying the following asymptotics as $m \rightarrow \infty$:*

$$(3.2) \quad m^n \cdot \int_{\mathbb{C}^n} (f \cdot \alpha_{m,r} - \beta_{m,r}) \cdot \bar{z}^J \cdot e^{-m \cdot |z|^2} = O\left(\frac{1}{m^{r+1}}\right)$$

for every fixed $r \geq 0$ and multi-index J . In particular, the formal power series $b \in \mathcal{F}_{\mathbb{C}^n}$ is uniquely determined by α and the Taylor expansion of f at the origin $0 \in \mathbb{C}^n$.

Proof. Explicitly, we need to prove the following:

$$m^n \cdot \int_{\mathbb{C}^n} (f \cdot \alpha_{m,r} - \sum_{2k+|I| \leq r} \frac{1}{m^k} \cdot b_{k,I} z^I) \cdot \bar{z}^J \cdot e^{-m \cdot |z|^2} = O\left(\frac{1}{m^{r+1}}\right).$$

Since $\alpha_{m,r}$'s are truncated from the same formal power series, we see that $\{f \cdot \alpha_{m,r}\}$ has the same property. Thus the result follows from Theorem 2.26. \square

Example 3.2. Let us consider the simplest example where $n = 1$, and let $a = z$. Then the double sequence $\{\alpha_{m,r}\}$ is explicitly defined by $\alpha_{m,r} := z \otimes \mathbb{1}^m$. Let $f = \bar{z}$, then a simple computation of the Wick ordering gives $\hbar \frac{\partial}{\partial \bar{z}}(z) = \hbar$. It follows that

$$b_{k,I} = \begin{cases} 1, & (k, I) = (1, 0), \\ 0, & \text{otherwise.} \end{cases}$$

We can interpret Proposition 3.1 as follows. Let $T_{f,m}$ denote the Toeplitz operators on $H^0(\mathbb{C}^n, L^{\otimes m})$ associated to the function f , then the double sequence $\{T_{f,m}(\alpha_{m,r})\} \in V$ can be ‘‘approximated’’ by a vector in the image of F . To make this precise, we define the subspace of *admissible sequences* $V_0 \subset V$:

Definition 3.3. We call $\alpha = \{\alpha_{m,r}\} \in V$ an *admissible sequence* if there exists $b = \sum_{k,I} b_{k,I} \hbar^k z^I \in \mathcal{F}_{\mathbb{C}^n}$ with $F(b) = \{\beta_{r,m}\}$ such that, for every fixed $r > 0$, we have

$$m^n \cdot \int_{\mathbb{C}^n} (\alpha_{m,r} - \beta_{m,r}) \cdot \bar{z}^J \cdot e^{-m \cdot |z|^2} = O\left(\frac{1}{m^{r+1}}\right)$$

for any multi-index J .

It is clear that there are inclusions $F(V) \subset V_0 \subset V$. Furthermore, there is a natural equivalence relation \sim on V : If $\alpha_i \in V_0, i = 1, 2$ are two admissible sequences with b_i their corresponding elements in $\mathcal{F}_{\mathbb{C}^n}$ respectively, then we say α_1 is equivalent to α_2 if $b_1 = b_2 \in \mathcal{F}_{\mathbb{C}^n}$.

3.2. Construction of the representation.

For a general compact Kähler manifold X equipped with a prequantum line bundle L , we define

$$V := \prod_{r \geq 0} \left(\prod_{m \geq 0} H^0(X, L^{\otimes m}) \right)$$

as before. The situation is more complicated than \mathbb{C}^n because L is non-trivial and there are no obvious global holomorphic sections analogous to $z^I \otimes \mathbb{1}^m$ on \mathbb{C}^n . The best replacement (or approximation) for the polynomial sections on \mathbb{C}^n are given by so-called *peak sections*, with which we can define double sequences of holomorphic sections with asymptotic properties similar to equation (3.2):

Definition 3.4. For every point $z_0 \in X$, we fix a set $\{S_{m,p,r}\}$ of normalized peak sections centered at z_0 , as introduced in Section A. A sequence of holomorphic sections $\alpha = \{\alpha_{m,r} \in H^0(X, L^{\otimes m})\}$, regarded as an element in V , is called an *admissible sequence at z_0* if it satisfies the following two conditions:

- (1) For every fixed r , the norm of the sequence $\{\alpha_{m,r}\}_{m>0}$ has a uniform bound:

$$\|\alpha_{m,r}\|_m \leq C_r.$$

- (2) There is a sequence of complex numbers $\{a_{p,k}\}_{p,k \geq 0}$ such that, for each fixed $r > 0$, we have

$$(3.3) \quad \langle \alpha_{m,r} - \sum_{2k+|p| \leq r} a_{p,k} \cdot \frac{1}{m^k} \cdot S_{m,p,r+1}, S_{m,q,r+1} \rangle_m = O\left(\frac{1}{m^{r+1}}\right),$$

for any multi-index q with $|q| \leq r$.

We define the subspace $V_{z_0} \subset V$ as the \mathbb{C} -linear span of admissible sequences at z_0 .

Equation (3.3) is the analogue of equation (3.2) in the flat case. According to Lemma A.9, the coefficients $a_{p,k}$ are uniquely determined.

Remark 3.5. The index r in admissible sequences corresponds to the degree in the Wick algebra and Bargmann-Fock space.

The complex numbers $\{a_{p,k}\}$ are called the *coefficients* of the admissible sequence α . Note that they are independent of either the tensor power m and the *weight index* r . The coefficients define a natural equivalence relation \sim on V_{z_0} , namely, α is equivalent to β (denoted as $\alpha \sim \beta$) if the coefficients of $\alpha - \beta$ are all 0.

Remark 3.6. It follows from this definition that, for each fixed r , even if we change finitely many terms of the double sequence $\{\alpha_{m,r}\}$, its equivalence class remains the same (cf. direct limits).

The vector space we would like to construct is then simply the quotient by this equivalence relation:

$$H_{z_0} := V_{z_0} / \sim .$$

It follows from asymptotics of inner products of peak sections $S_{m,p,r}$'s as $m \rightarrow \infty$ and equation (3.3) that H_{z_0} is a formal Hilbert space.

Remark 3.7. The vector space H_{z_0} is defined as a sub-quotient, instead of just as a linear span of peak sections. This is because in general the Toeplitz operators do not preserve the space of peak sections.

We now give some examples of admissible sequences:

Example 3.8. Suppose we fix any multi-index q . Then we define an admissible sequence α as follows: we let $\alpha_{m,r} := 0$ if m is too small and there is no (normalized) peak section of $L^{\otimes m}$ corresponding to the index r , and let $\alpha_{m,r} = S_{m,q,r+1}$ be simply the normalized peak section. It is then easy to see that this is indeed an admissible sequence with coefficients $a_{p,r} = 0$ if $(p,r) \neq (q,0)$ and $a_{q,0} = 1$.

Example 3.9. We give an example of equivalent admissible sequences. Let α be the admissible sequence as in the previous example. We now construct a sequence also consisting of normalized peak sections similar to α but with a higher order error term. Namely, we let $\beta = \{\beta_{m,r}\}$ be the admissible sequence given by $\beta_{m,r} = S_{m,q,r+2}$. It is easy to show that $\alpha \sim \beta$.

It is not difficult to see that the admissible sequences in Example 3.8 form a basis of V_{z_0} as a \mathbb{C} -vector space, and thus every vector can be written as $\{a_{p,k}\}$. More precisely, we have the following lemma:

Lemma 3.10. *We have the following isomorphism of \mathbb{C} -vector spaces:*

$$(3.4) \quad H_{z_0} \cong \mathbb{C}[[y_1, \dots, y_n]][[\hbar]].$$

Proof. The above isomorphism is given by $\{a_{p,k}\} \mapsto \sum_{p,k} a_{p,k} \cdot \hbar^k y^p$. □

Remark 3.11. The isomorphism in the above lemma depends on a choice of K -coordinates centered at z_0 , and thus is unique only up to a $U(n)$ -transformation.

Now for every sequence of operators $\{A_m\}_{m \geq 0}$, where $A_m \in \text{End}(H^0(X, L^{\otimes m}))$, we have an obvious action on V :

$$\{\alpha_{m,r}\} \mapsto \{A_m(\alpha_{m,r})\}.$$

We apply this to the sequence of Toeplitz operators $\{T_{f,m}\}_{m \geq 0}$ associated to any given smooth function $f \in C^\infty(X)$.

Lemma 3.12. *Suppose that $\alpha = \{\alpha_{m,r}\}$ is an admissible sequence. Then $\{T_{f,m}(\alpha_{m,r})\}$ is also an admissible sequence for any smooth function f .*

Proof. We consider the sequence $T_f(\alpha) := \{T_{f,m}(\alpha_{m,r})\}_{m,r \geq 0}$. First of all, since the operator norm of $T_{f,m}$ is bounded by $\|f\|_\infty$ for any fixed $r > 0$, the sequence $\{T_{f,m}(\alpha_{m,r})\}_{m > 0}$ has bounded norm.

To show that $T_f(\alpha)$ satisfies the second asymptotic property, we split the integral which defines that property into two parts: one inside the disk $\{\rho(z) < 1\}$ and the other outside:

$$\begin{aligned} & m^n \int_X h^m(T_{f,m}(\alpha_{m,r}), S_{m,p,r+1}) \cdot dV_g \\ &= m^n \int_X h^m(f \cdot \alpha_{m,r}, S_{m,p,r+1}) \cdot dV_g \\ &= m^n \int_{X \setminus \{\rho(z) < 1\}} h^m(f \cdot \alpha_{m,r}, S_{m,p,r+1}) \cdot dV_g + m^n \int_{\{\rho(z) < 1\}} h^m(f \cdot \alpha_{m,r}, S_{m,p,r+1}) \cdot dV_g, \end{aligned}$$

where the first equality follows from the fact that $T_{f,m}(\alpha_{m,r})$ is the orthogonal projection of $f \cdot \alpha_{m,r}$ to the space of holomorphic sections, and that $S_{m,p,r+1}$ are all holomorphic sections. For the integral outside the disk, we have the following estimate:

$$\begin{aligned} & \left| m^n \int_{X \setminus \{\rho(z) < 1\}} h^m(f \cdot \alpha_{m,r}, S_{m,p,r+1}) \cdot dV_g \right| \\ & \leq m^n \left(\int_{X \setminus \{\rho(z) < 1\}} \|f \cdot \alpha_{m,r}\|_{h^m}^2 \cdot dV_g \right)^{1/2} \cdot \left(\int_{X \setminus \{\rho(z) < 1\}} \|S_{m,p,r+1}\|_{h^m}^2 \cdot dV_g \right)^{1/2} \\ & \leq \|f\|_\infty \cdot C_r \cdot O\left(\frac{1}{m^{(2r+2+|p|)/2}}\right) = O\left(\frac{1}{m^{r+1}}\right). \end{aligned}$$

Here the multi-index p satisfies $|p| \leq r$, and the constant C_r is given by the upper bound of the sequence $\{\alpha_{m,r}\}_{m > 0}$. For the second inequality, we have used boundedness of $\{\alpha_{m,r}\}_{m > 0}$, and equation (A.9).

Hence it remains to consider the integral inside the disk $\{\rho(z) < 1\}$. In this local setting, the computation is the same as the asymptotics of the Gaussian integral on \mathbb{D}^{2n} , and also that in the formal Hilbert space. So the statement follows from Theorem 2.26. \square

This lemma shows that for any smooth function f , the sequence of Toeplitz operators $\{T_{f,m}\}_{m > 0}$ gives a well-defined linear operator $T_f : V_{z_0} \rightarrow V_{z_0}$, $\alpha \mapsto T_f(\alpha)$.

Lemma 3.13. *Suppose that two admissible sequences are equivalent, i.e., $\alpha \sim \beta$. Then for any smooth function f , we have $T_f(\alpha) \sim T_f(\beta)$.*

Proof. We only need to show that if the coefficients $a_{k,I}$ of $\alpha = \{\alpha_{m,r}\}$ vanish, then $T_f(\alpha)$ has the same property. Notice that the condition $a_{k,I} = 0$ is equivalent to the following equalities for all indices r, q , as $m \rightarrow \infty$:

$$m^n \int_X h^m(\alpha_{m,r}, S_{m,q,r+1}) \cdot dV_g = O\left(\frac{1}{m^{r+1}}\right).$$

Similar to the argument of Lemma 3.12, there is

$$\begin{aligned} \left| m^n \int_X h^m(T_{f,m}(\alpha_{m,r}), S_{m,q,r+1}) \cdot dV_g \right| &= \left| m^n \int_X h^m(f \cdot \alpha_{m,r}, S_{m,q,r+1}) \cdot dV_g \right| \\ &\leq m^n \cdot \|f\|_\infty \left| \int_X h^m(\alpha_{m,r}, S_{m,q,r+1}) \cdot dV_g \right| \\ &= O\left(\frac{1}{m^{r+1}}\right). \end{aligned}$$

□

Hence, for every smooth function f on X , the sequence of Toeplitz operators $\{T_{f,m}\}_{m>0}$ gives a well-defined linear operator T_f on the vector space H_{z_0} . We can further extend it to an action of $C^\infty(X)[[\hbar]]$ on H_{z_0} by letting $\hbar^k \cdot f$ act as

$$T_{\hbar^k \cdot f} : \{\alpha_{m,r}\} \mapsto \left\{ \frac{1}{m^k} \cdot T_{f,m}(\alpha_{m,r}) \right\}.$$

Lemma 2.25 implies the following:

Proposition 3.14. *Suppose $f \in C^\infty(X)$ is a real function. Then for every $z_0 \in X$, the operator T_f on H_{z_0} is self-adjoint.*

Lemma 3.15. *Let $A = \{A_m\}_{m \geq 0}$ and $B = \{B_m\}_{m \geq 0}$ be two sequences of bounded operators preserving asymptotic sequences and satisfying the condition that*

$$(3.5) \quad \|A_m - B_m\| = O\left(\frac{1}{m^{k+1}}\right).$$

Then for any admissible sequence α , the two admissible sequences $A(\alpha)$ and $B(\alpha)$ have the same coefficients up to weight k .

Proof. Equation (3.5) implies that the operators $\{A_m - B_m\}$ will increase the weight of α by $k + 1$. The lemma follows. □

Here is our main theorem:

Theorem 3.16. *Let $z_0 \in X$ be any point. The action of $C^\infty(X)[[\hbar]]$ on the vector space H_{z_0} satisfies the following relation:*

$$(3.6) \quad T_f \circ T_g = T_{fg} + \sum_{k \geq 1} \hbar^k \cdot T_{C_k(f,g)}, \quad f, g \in C^\infty(X),$$

where $C^k(-, -)$ are the bi-differential operators which appear in the Berezin-Toeplitz quantization. Therefore, H_{z_0} is a representation of the Berezin-Toeplitz deformation quantization algebra $(C^\infty(X)[[\hbar]], \star_{BT})$.

Proof. We first recall the property of Toeplitz operators:

$$\|T_{f,m} \circ T_{g,m} - (T_{fg,m} + \sum_{k=1}^n \frac{1}{m^k} \cdot T_{C_k(f,g),m})\| = O\left(\frac{1}{m^{n+1}}\right).$$

We apply Lemma 3.15 by putting $A_m := T_{f,m} \circ T_{g,m}$ and $B_m := T_{fg} + \sum_{k=1}^n \frac{1}{m^k} \cdot T_{C_k(f,g),m}$. Then $A(\alpha)$ and $B(\alpha)$ have the same coefficients up to order n . The theorem follows by letting $n \rightarrow \infty$. \square

Remark 3.17. The representation and also the isomorphism (3.4) are independent of the choice of the set of peak sections because for every multi-index p and $p' > |p|$, different choices of peak sections only differ by higher order terms.

3.3. Locality and irreducibility of the representation.

3.3.1. *Locality.* We will give an explicit formula of our representation under the isomorphism (3.4). Given any K -coordinates (z_1, \dots, z_n) centered at z_0 , we define $J_{f,z_0} \in \mathcal{W}_{\mathbb{C}^n}$ by

$$(3.7) \quad J_{f,z_0} := \sum_{|I|, |J| \geq 0} \frac{1}{I!J!} \frac{\partial^{|I|+|J|} f}{\partial z^I \partial \bar{z}^J}(z_0) y^I \bar{y}^J,$$

where the sum is over all multi-indices.

Theorem 3.18. *Let f be any smooth function on X , and J_{f,z_0} be defined as above. We define $O_{f,z_0} \in \mathcal{W}_{\mathbb{C}^n}$ as the unique solution of the following equation:*

$$J_{f,z_0} \cdot e^{\Phi/\hbar} = e^{\Phi/\hbar} \star O_{f,z_0}.$$

Then the action of T_f on $\alpha \in H_{z_0}$ is given by

$$T_f(\alpha) = O_{f,z_0} \star \alpha.$$

In particular, this implies that the representation H_{z_0} is local in $f \in C^\infty(X)$, i.e., it only depends on the infinite jets of f at z_0 .

Proof. The representation H_{z_0} depends on an arbitrarily small neighborhood of z_0 . So the result follows from the computation of the formal Toeplitz operators on \mathbb{C}^n in Theorem 2.22. \square

As a straightforward corollary, we have:

Corollary 3.19. *Let $f, g \in C^\infty(X)$ be smooth functions on X . Then*

$$O_{f \star_{BT} g, z_0} = O_{f, z_0} \star O_{g, z_0}.$$

This gives an algorithm for computing $f \star_{BT} g$: for every $z_0 \in X$, in order to find $(f \star_{BT} g)(z_0)$, we only need to compute the Wick product $O_{f, z_0} \star O_{g, z_0}$ and then collect all the constant terms in the Wick algebra.

3.3.2. Irreducibility. We now consider irreducibility of our representation. The first observation is that the Bargmann-Fock space is *not* an irreducible representation of $\mathcal{W}_{\mathbb{C}^n}$: for every $f \in \mathcal{W}_{\mathbb{C}^n}$, we have the invariant subspaces

$$T_f((\mathcal{F}_{\mathbb{C}^n})_k) \subset (\mathcal{F}_{\mathbb{C}^n})_k.$$

But this is the *only* reason why the representation fails to be irreducible. So to have a suitable notion of irreducibility, we make use of the extended algebra $\mathcal{W}_{\mathbb{C}^n}^+$, which allows terms with negative degrees, and the corresponding extension $\mathcal{F}_{\mathbb{C}^n}^+$. It is then quite easy to check that we indeed obtain an irreducible representation.

We now define an extension of $C^\infty(X)[[\hbar]]$, which is the geometric analogue of $\mathcal{W}_{\mathbb{C}^n}^+$.

Definition 3.20. For every smooth function $f \in C^\infty(X)$, let $\deg_{z_0}(f)$ be the vanishing order of f at z_0 . Then let $(C^\infty(X)[[\hbar]])_{z_0}^+$ be the extension of $C^\infty(X)[[\hbar]]$ which consists of formal functions:

$$\sum_{i \in \mathbb{Z}} \hbar^i \cdot f_i,$$

where $f_i \in C^\infty(X)$ are smooth functions on X satisfying the conditions that

- the sum $\deg_{z_0}(f_i) + 2i$ has a uniform lower bound for all i , and
- for every degree k , the following expression is a finite sum:

$$\sum_{2i + \deg_{z_0}(f_i) = k} \hbar^i \cdot f_i.$$

In the same way we can define the extension $H_{z_0} \subset H_{z_0}^+$. It is easy to check that the extension $(C^\infty(X)[[\hbar]])_{z_0}^+$ is closed under the star product \star_{BT} , and acts on $H_{z_0}^+$. Furthermore, the map $f \mapsto O_{f, z_0}$ can be extended to

$$(3.8) \quad (C^\infty(X)[[\hbar]])_{z_0}^+ \rightarrow \mathcal{W}_{\mathbb{C}^n}^+.$$

Theorem 3.21. *For every $z_0 \in X$, the representation $H_{z_0}^+$ of $(C^\infty(X)[[\hbar]])_{z_0}^+$ is irreducible.*

Proof. Let W be a sub-representation of $H_{z_0}^+$. We choose any non-zero $a \in W$, which can be written as:

$$a = \sum_{2i + |I| \geq k} a_{i, I} \hbar^i \cdot y^I.$$

Since for a local holomorphic function f , we have

$$O_{f,z_0} = J_{f,z_0},$$

which consists of only creators in $(H_{z_0})^+$, we only need to find $f \in (C^\infty(X)[[\hbar]])_{z_0}^+$ such that $T_f(a) = \hbar^l$ for some l , and the result will follow.

We choose a non-zero term in a of leading order $a_{i_0, I_0} \hbar^{i_0} \cdot z^{I_0}$, $2i_0 + |I_0| = k$, such that i_0 is the least possible. Let $f_0 \in (C^\infty(X)[[\hbar]])_{z_0}^+$ be a formal function which is $a_{i_0, I_0}^{-1} \bar{z}^{I_0}$ near z_0 . So the leading term of the image of f_0 under the map (3.8) is $a_{i_0, I_0}^{-1} \bar{y}^{I_0}$, and the degree of the function f_0 is exactly $|I_0|$. We have

$$T_{f_0} \left(\sum_{2i+|I| \geq k} a_{i,I} \hbar^i \cdot y^I \right) = \hbar^{i_0+|I_0|} + \sum_{2i+|I|=2(i_0+|I_0|)+1} b_{i,I} \hbar^i y^I + \text{higher degree terms.}$$

The next step is to find a formal function $f_1 \in (C^\infty(X)[[\hbar]])_{z_0}^+$, so that

$$(T_{f_0} + T_{f_1}) \left(\sum_{2i+|I| \geq k} a_{i,I} \hbar^i \cdot y^I \right) = \hbar^{i_0+|I_0|} + \text{higher degree terms.}$$

Let g be a formal function which equals $-\frac{1}{\hbar^{i_0+|I_0|}} \sum_{2j+|J|=2(i_0+|I_0|)+1} b_{j,J} \hbar^j z^J$ near z_0 . It is easy to see that the total degree of g is 1, and we have

$$(T_{f_0} + T_g \circ T_{f_0}) \left(\sum_{2i+|I| \geq k} a_{i,I} \hbar^i \cdot y^I \right) = \hbar^{i_0+|I_0|} + (\text{terms of degree } \geq 2(i_0 + |I_0|) + 2).$$

Although $T_g \circ T_{f_0} = \sum_{i \geq 0} \hbar^i T_{C_i(g, f_0)}$ is an infinite sum, but those high enough \hbar will map terms in a to terms of high degree. More precisely, we can simply let $f_1 = \sum_{i=0}^N \hbar^i C_i(g, f_0)$, where $N = i_0 + |I_0| - k/2 + 2$. Then we have

$$(T_{f_0} + T_{f_1}) \left(\sum_{2i+|I| \geq k} a_{i,I} \hbar^i \cdot y^I \right) = \hbar^{i_0+|I_0|} + (\text{terms of degree } \geq 2(i_0 + |I_0|) + 2).$$

From the formula of \star_{BT} , it is easy to see that the total degree of $\hbar^i C_i(g, f_0)$ is no less than the sum of the degree of g_0 and f which equals $1 + |I_0|$. This implies that the degree of terms in f_1 is strictly greater than the degree of f_0 . This procedure can be repeated and we obtain the desired formal function $f = \sum_{i \geq 0} f_i \in (C^\infty(X)[[\hbar]])_{z_0}^+$. \square

APPENDIX A. PEAK SECTIONS IN KÄHLER GEOMETRY

In this appendix, we briefly review of the notion of peak sections, which was introduced in [31] and plays an important role in the study of asymptotic expansions of the Bergmann kernel [33] with applications to balanced embeddings and constant scalar curvature metrics as well as in the theory of the geometric quantization. For our purpose, we

need to introduce a normalized version of peak sections and describe their basic properties. We first recall the notion of K -coordinates and K -frame of the prequantum line bundle L .

Definition A.1. Let e_{L,z_0} be a holomorphic frame of the prequantum line bundle L in a neighborhood of a point $z_0 \in X$, and let (z_1, \dots, z_n) be a holomorphic coordinate system centered at z_0 . Let $\varphi(z) := -\log \|e_{L,z_0}\|$. We say that (z_1, \dots, z_n) are K -coordinates with K -frame e_{L,z_0} if the Taylor expansion of $\varphi(z)$ at z_0 is of the following form:

$$(A.1) \quad \varphi(z) \sim |z|^2 + \sum a_{JK} z^J \bar{z}^K, \quad |J| \geq 2, |K| \geq 2.$$

For Kähler manifolds with real analytic Kähler form, the existence of K -coordinates and K -frames was shown by Bochner. For Kähler manifolds with only smooth Kähler form, such a coordinate system and frame do not exist in general; then we may consider a weaker K -coordinates and K -frames of finite order. But to avoid further technical complications, let us assume that the Kähler manifolds in this paper always admit K -coordinates and K -frames.

It is obvious that this local holomorphic frame e_{L,z_0} is unique up to a multiplication by a complex number of modulus 1. In particular, the leading term of the Taylor expansion of φ with degree at least 3 is given by the curvature:

$$(A.2) \quad \varphi(z, \bar{z}) = |z|^2 + \sum_{i,j,k,l} R_{i\bar{j}k\bar{l}} z_i z_k \bar{z}_j \bar{z}_l + O(|z|^5).$$

Also note that equation (2.9) is satisfied.

Lemma A.2. Suppose the volume form is $(\sqrt{-1})^n \cdot e^{\psi(z, \bar{z})} \cdot dz^1 \dots dz^n d\bar{z}^1 \dots d\bar{z}^n = \omega^n$. Then the purely (anti-)holomorphic derivatives of $\psi(z, \bar{z})$ vanish at z_0 under the K -coordinates, i.e.,

$$\frac{\partial^{|I|} \psi}{\partial z^I}(z_0) = \frac{\partial^{|J|} \psi}{\partial \bar{z}^J}(z_0) = 0$$

for all multi-indices with $|I|, |J| > 0$.

Proof. We will only prove the vanishing of purely holomorphic derivatives at z_0 ; the proof for antiholomorphic ones is the same. It suffices to show that the statement is valid for functions $\omega_{i_1 \bar{j}_1} \dots \omega_{i_n \bar{j}_n}$, where

$$\omega_{i\bar{j}} = \frac{\partial^2 \varphi}{\partial z^i \partial \bar{z}^j},$$

and φ is Kähler potential. But then equation (A.1) implies that

$$\frac{\partial^{|I|+1} \varphi}{\partial z^I \partial \bar{z}^j}(z_0) = 0, \quad |I| \geq 2,$$

from which the statement follows easily. □

Equation (A.1) together with Lemma A.2 tell us that the Taylor expansions of $\varphi(z, \bar{z})$ and $\psi(z, \bar{z})$ satisfy the technical conditions in Theorem 2.26. This allows us to apply the algebraic computations in the formal setting in Section 2.2.

We now recall the following proposition in Tian's paper [31] and, in particular, the definition of peak sections. Let (z_1, \dots, z_n) be a K -coordinate with K -frame e_{L, z_0} at $z_0 \in X$, and consider the function $\rho(z) := \sqrt{|z_1|^2 + \dots + |z_n|^2}$.

Proposition A.3 (Lemma 1.2 in [31]). *For an n -tuple of integers $p = (p_1, \dots, p_n) \in \mathbb{Z}_+^n$ and an integer $r > |p| = p_1 + \dots + p_n$, there exists an $m_0 > 0$ such that, for $m > m_0$, there is a holomorphic global section S , called a peak section, of the line bundle $L^{\otimes m}$, satisfying*

$$(A.3) \quad \int_X \|S\|_{h^m}^2 dV_g = 1, \quad \int_{X \setminus \{\rho(z) \leq \frac{\log m}{\sqrt{m}}\}} \|S\|_{h^m}^2 dV_g = O\left(\frac{1}{m^{2r}}\right),$$

and locally at z_0 ,

$$(A.4) \quad S(z) = \lambda_{m,p} \cdot \left(z_1^{p_1} \cdots z_n^{p_n} + O(|z|^{2r}) \right) e_{L, z_0}^m \left(1 + O\left(\frac{1}{m^{2r}}\right) \right),$$

where $\|\cdot\|_{h^m}$ is the norm on $L^{\otimes m}$ given by h^m , and $O\left(\frac{1}{m^{2r}}\right)$ denotes a quantity dominated by C/m^{2r} with the constant C depending only on r and the geometry of X , moreover

$$(A.5) \quad \lambda_{m,p}^{-2} = \int_{\rho(z) \leq \log m / \sqrt{m}} |z_1^{p_1} \cdots z_n^{p_n}|^2 \cdot e^{-m \cdot \varphi(z)} dV_g,$$

where $dV_g = \det(g_{i\bar{j}}) (\sqrt{-1} / (2\pi))^n dz_1 \wedge d\bar{z}_1 \wedge \dots \wedge dz_n \wedge d\bar{z}_n$ is the volume form.

Here we use the same notation as in the introduction of K -frame: $h^m(e_{L, z_0}, e_{L, z_0}) = e^{-m \cdot \varphi(z)}$. Geometrically, a peak section is, roughly speaking, a global holomorphic section of a high enough tensor power of L whose norm is 1 and is concentrated around a given point z_0 on the Kähler manifold.

We want to define a section $S_{m,p,r}$ of the line bundle $L^{\otimes m}$ by normalizing the peak section $S(z)$ in Proposition A.3 so that its Taylor expansion at z_0 under the K -frame $e_{L, z_0}^{\otimes m}$ is exactly equal to $z_1^{p_1} \cdots z_n^{p_n}$ up to order $2r - 1$. This forces $S_{m,p,r}$ to be of the form:

$$(A.6) \quad S_{m,p,r} := \lambda_{m,p}^{-1} \cdot \left(1 + O\left(\frac{1}{m^{2r}}\right) \right) \cdot S(z).$$

We now give an estimate of $\lambda_{m,p}^{-1}$. We have, for $m \gg 0$,

$$\begin{aligned} \lambda_{m,p}^{-2} &= \int_{\rho(z) \leq \log m / \sqrt{m}} |z_1^{p_1} \cdots z_n^{p_n}|^2 \cdot e^{-m \cdot \rho(z)} dV_g \\ &\leq \int_{\rho(z) \leq 1} |z_1^{p_1} \cdots z_n^{p_n}|^2 \cdot e^{-m \cdot \rho(z)} dV_g = O\left(\frac{1}{m^{|p|+n}}\right), \end{aligned}$$

where the estimate follows from Theorem 2.11. In particular, there is a constant C_p , depending only on the point z_0 and the multi-index p , such that

$$\lambda_{m,p}^{-1} \cdot \left(1 + O\left(\frac{1}{m^{2r}}\right)\right) \leq C_p \cdot \left(\frac{1}{m^{\frac{|p|+n}{2}}}\right).$$

We define a normalized version of the inner product of sections of $L^{\otimes m}$:

Definition A.4. Let s_1, s_2 be (smooth) sections of $L^{\otimes m}$. Their (normalized) inner product is defined as

$$(A.7) \quad \langle s_1, s_2 \rangle_m := m^n \cdot \int_X h^m(s_1, s_2) dV_g,$$

where $n = \dim_{\mathbb{C}} X$, and we let $\|s\|_m$ be the norm of a section s under this inner product.

Remark A.5. An explanation of the normalization factor m^n is the following: consider \mathbb{C}^n with the volume form

$$\left(\frac{\sqrt{-1}}{2\pi}\right)^n e^{-m \cdot |z|^2} dz_1 \wedge d\bar{z}_1 \wedge \cdots \wedge dz_n \wedge d\bar{z}_n,$$

then the factor m^n normalizes the volume to 1 under this volume form.

We summarize the properties of $S_{m,p,r}$ as follows:

$$(A.8) \quad \|S_{m,p,r}\|_m^2 \leq m^n \cdot C \cdot \lambda_{m,p}^{-2} = O\left(\frac{1}{m^{|p|}}\right);$$

$$(A.9) \quad m^n \cdot \int_{M \setminus \{\rho(z) \leq 1\}} \|S_{m,p,r}\|_{h^m}^2 dV_g \leq m^n \cdot \int_{M \setminus \{\rho(z) \leq \log m / \sqrt{m}\}} \|S_{m,p,r}\|_{h^m}^2 dV_g \\ \leq m^n \cdot C \cdot \lambda_{m,p}^{-2} \cdot O\left(\frac{1}{m^{2r}}\right) = O\left(\frac{1}{m^{2r+|p|}}\right).$$

Remark A.6. The constant C in the above estimates comes from the number $1 + O\left(\frac{1}{m^{2r}}\right)$ in equation (A.6).

Locally around z_0 , we have

$$(A.10) \quad S_{m,p,r}(z) = \left(z_1^{p_1} \cdots z_n^{p_n} + O(|z|^{2r})\right) \cdot e_L^m.$$

The first property (A.8) implies that for fixed p, r , the sequence $\{S_{m,p,r}\}$ is bounded for all m . The second property (A.9) is saying that the sections $S_{m,p,r}$ are asymptotically ‘‘concentrated’’ around the point z_0 . The third property (A.10) is saying that asymptotically, $S_{m,p,r}$ has an assigned leading term of the Taylor expansion at the point z_0 .

Remark A.7. The third property of $S_{m,p,r}$ is the reason for calling it a *normalized peak section*: its Taylor expansion at z_0 has leading term exactly equal to the monomial $z^p \cdot e_L^m$ corresponding to the multi-index p .

Remark A.8. According to [31], for every fixed p, r , peak sections exist only when m is big enough. We will adopt the follows two conventions

- $S_{m,p,r} := 0$ for small m ,
- $S_{m,p,r} := 0$ if $r \leq |p|$.

There is the following estimate of the inner product between peak sections:

Lemma A.9. *Given two normalized peak sections $S_{m,p^1,r}, S_{m,p^2,r}$, we have the following asymptotic expansion of their inner product up to order r :*

$$(A.11) \quad \langle S_{m,p^1,r}, S_{m,p^2,r} \rangle_m - \sum_{k=1}^{p'-1} a_k \cdot \frac{1}{m^k} = O\left(\frac{1}{m^{p'}}\right),$$

where the coefficients a_k 's are the same as those in the expansion of the following formal integral:

$$\int z^{p^1} \bar{z}^{p^2} e^{-\frac{|z|^2 + \phi(z, \bar{z})}{\hbar}} = \sum_{k \geq 1} a_k \cdot \hbar^k.$$

Thus, for fixed multi-indices p^1, p^2 , the a_i 's are independent of $r \gg 0$. In particular, the leading term of the asymptotic expansion of $\|S_{m,p,r}\|_m^2$ is given by

$$\left(\frac{1}{m}\right)^{|p|} p!,$$

which is non-zero.

Proof. We split the integral defining the inner product to two parts:

$$\frac{1}{m^n} \int_{\{\rho(z) < 1\}} h^m(S_{m,p^1,r}, S_{m,p^2,r}) \cdot dV_g + \frac{1}{m^n} \int_{X \setminus \{\rho(z) < 1\}} h^m(S_{m,p^1,r}, S_{m,p^2,r}) \cdot dV_g,$$

where the second part is $O\left(\frac{1}{m^{2r + \frac{|p^1| + |p^2|}{2}}}\right)$ by using Cauchy-Schwarz inequality and equation (A.9). Thus to show equation (A.11), the integral outside the disk $\{\rho(z) < 1\}$ can be ignored. For the integral over the disk, we can apply Theorem 2.11 to obtain the desired asymptotic expansion. In particular, the coefficients a_k 's are the same as those coming from the formal integral. \square

As an immediate corollary, we have the following:

Corollary A.10. *Let p^1, p^2 be multi-indices, and let $r > \max\{|p^1|, |p^2|\}$, then we have the following estimate of the inner product between $S_{m,p^1,r}$ and $S_{m,p^2,r}$:*

$$\langle S_{m,p^1,r}, S_{m,p^2,r} \rangle_m = \begin{cases} O\left(\frac{1}{m^{|p^1|}}\right), & p^1 = p^2 \\ o\left(\frac{1}{m^{\max\{|p^1|, |p^2|\}}}\right), & p^1 \neq p^2. \end{cases}$$

Proof. The case where $p_1 = p_2$ is given by equation (A.8). For $p_1 \neq p_2$, we need to estimate an integral. For the integral inside the disk $\{\rho(z) < 1\}$, this estimate is given by Corollary 2.21 where the technical condition on ϕ is implied by the existence of K -coordinates and K -frame. For the estimate of the integral outside the disk, we use Cauchy-Schwarz inequality:

(A.12)

$$m^n \cdot \int_{X \setminus \{\rho(z) < 1\}} h^m(S_{m,p_1,r}, S_{m,p_2,r}) \cdot dV_g = O\left(\frac{1}{m^{\frac{4r+|p_1|+|p_2|}{2}}}\right) = o\left(\frac{1}{m^{\max\{|p_1|, |p_2|\}}}\right).$$

□

REFERENCES

- [1] F. Bayen, M. Flato, C. Fronsdal, A. Lichnerowicz, and D. Sternheimer, *Deformation theory and quantization. I. Deformations of symplectic structures*, Ann. Physics **111** (1978), no. 1, 61–110.
- [2] ———, *Deformation theory and quantization. II. Physical applications*, Ann. Physics **111** (1978), no. 1, 111–151.
- [3] M. Bordemann, E. Meinrenken, and M. Schlichenmaier, *Toeplitz quantization of Kähler manifolds and $\mathfrak{gl}(N)$, $N \rightarrow \infty$ limits*, Comm. Math. Phys. **165** (1994), no. 2, 281–296.
- [4] M. Bordemann and S. Waldmann, *A Fedosov star product of the Wick type for Kähler manifolds*, Lett. Math. Phys. **41** (1997), no. 3, 243–253.
- [5] ———, *Formal GNS construction and states in deformation quantization*, Comm. Math. Phys. **195** (1998), no. 3, 549–583.
- [6] K. Chan, N. C. Leung, and Q. Li. in preparation.
- [7] K. Costello, *Renormalization and effective field theory*, Mathematical Surveys and Monographs, vol. 170, American Mathematical Society, Providence, RI, 2011.
- [8] ———, *A geometric construction of the Witten genus, II*, available at [arXiv:1111.4234](https://arxiv.org/abs/1111.4234) [math.QA].
- [9] ———, *Renormalization and the Batalin-Vilkovisky formalism*, available at [arXiv:0706.1533](https://arxiv.org/abs/0706.1533) [math.QA].
- [10] S. K. Donaldson, *Planck’s constant in complex and almost-complex geometry*, XIIIth International Congress on Mathematical Physics (London, 2000), 2001, pp. 63–72.
- [11] B. V. Fedosov, *A simple geometrical construction of deformation quantization*, J. Differential Geom. **40** (1994), no. 2, 213–238.
- [12] ———, *Deformation quantization and index theory*, Mathematical Topics, vol. 9, Akademie Verlag, Berlin, 1996.
- [13] M. Kapranov, *Rozansky-Witten invariants via Atiyah classes*, Compositio Math. **115** (1999), no. 1, 71–113.
- [14] A.V. Karabegov, *Deformation quantizations with separation of variables on a Kähler manifold*, Comm. Math. Phys. **180** (1996), no. 3, 745–755.
- [15] ———, *On Fedosov’s approach to deformation quantization with separation of variables*, Conférence Moshé Flato 1999, Vol. II (Dijon), 2000, pp. 167–176.
- [16] ———, *A formal model of Berezin-Toeplitz quantization*, Comm. Math. Phys. **274** (2007), no. 3, 659–689.
- [17] A.V. Karabegov and M. Schlichenmaier, *Identification of Berezin-Toeplitz deformation quantization*, J. Reine Angew. Math. **540** (2001), 49–76.
- [18] A. A. Kirillov, *Geometric quantization*, Current problems in mathematics. Fundamental directions, Vol. 4, 1985, pp. 141–178, 291.
- [19] M. Kontsevich, *Feynman diagrams and low-dimensional topology*, First European Congress of Mathematics, Vol. II (Paris, 1992), 1994, pp. 97–121.
- [20] ———, *Deformation quantization of Poisson manifolds*, Lett. Math. Phys. **66** (2003), no. 3, 157–216.

- [21] B. Kostant, *Quantization and unitary representations. I. Prequantization*, Lectures in modern analysis and applications, III, 1970, pp. 87–208. Lecture Notes in Math., Vol. 170.
- [22] Z. Lu and B. Shiffman, *Asymptotic expansion of the off-diagonal Bergman kernel on compact Kähler manifolds*, J. Geom. Anal. **25** (2015), no. 2, 761–782.
- [23] X. Ma and G. Marinescu, *Toeplitz operators on symplectic manifolds*, J. Geom. Anal. **18** (2008), no. 2, 565–611.
- [24] ———, *Berezin-Toeplitz quantization on Kähler manifolds*, J. Reine Angew. Math. **662** (2012), 1–56.
- [25] P. Mnev, *Lectures on Batalin-Vilkovisky formalism and its applications in topological quantum field theory*, available at [arXiv:1707.08096](https://arxiv.org/abs/1707.08096) [math-ph].
- [26] N. Neumaier, *Universality of Fedosov’s construction for star products of Wick type on pseudo-Kähler manifolds*, Rep. Math. Phys. **52** (2003), no. 1, 43–80.
- [27] N. Reshetikhin and L. A. Takhtajan, *Deformation quantization of Kähler manifolds*, L. D. Faddeev’s Seminar on Mathematical Physics, 2000, pp. 257–276.
- [28] M. Schlichenmaier, *Deformation quantization of compact Kähler manifolds by Berezin-Toeplitz quantization*, Conférence Moshé Flato 1999, Vol. II (Dijon), 2000, pp. 289–306.
- [29] J. Śniatycki, *Geometric quantization and quantum mechanics*, Applied Mathematical Sciences, vol. 30, Springer-Verlag, New York-Berlin, 1980.
- [30] J.-M. Souriau, *Structure des systèmes dynamiques*, Maîtrises de mathématiques, Dunod, Paris, 1970.
- [31] G. Tian, *On a set of polarized Kähler metrics on algebraic manifolds*, J. Differential Geom. **32** (1990), no. 1, 99–130.
- [32] N. M. J. Woodhouse, *Geometric quantization*, Second, Oxford Mathematical Monographs, The Clarendon Press, Oxford University Press, New York, 1992. Oxford Science Publications.
- [33] S. Zelditch, *Szego kernels and a theorem of Tian*, Internat. Math. Res. Notices **6** (1998), 317–331.

DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, HONG KONG
Email address: kwchan@math.cuhk.edu.hk

THE INSTITUTE OF MATHEMATICAL SCIENCES AND DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, HONG KONG
Email address: leung@math.cuhk.edu.hk

DEPARTMENT OF MATHEMATICS, SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY, SHENZHEN, CHINA
Email address: liqin@sustech.edu.cn