A Review of Mathematical Modeling, Simulation and Analysis of Membrane Channel Charge Transport

Duan Chen¹*and Guo-Wei Wei^{2,3}[†], ¹Department of Mathematics and Statistics University of North Carolina at Charlotte, NC 28223, USA ²Department of Mathematics Michigan State University, MI 48824, USA ³Department of Biochemistry and Molecular Biology Michigan State University, MI 48824, USA

January 31, 2017

Abstract

The molecular mechanism of ion channel gating and substrate modulation is elusive for many voltage gated ion channels, such as eukaryotic sodium ones. The understanding of channel functions is a pressing issue in molecular biophysics and biology. Mathematical modeling, computation and analysis of membrane channel charge transport have become an emergent field and give rise to significant contributions to our understanding of ion channel gating and function. This review summarizes recent progresses and outlines remaining challenges in mathematical modeling, simulation and analysis of ion channel charge transport. One of our focuses is the Poisson-Nernst-Planck (PNP) model and its generalizations. Specifically, the basic framework of the PNP system and some of its extensions, including size effects, ion-water interactions, coupling with density functional theory and relation to fluid flow models. A reduced theory, the Poisson-Boltzmann-Nernst-Planck (PBNP) model, and a differential geometry based ion transport model are also discussed. For proton channel, a multiscale and multiphysics Poisson-Boltzmann-Kohn-Sham (PBKS) model is presented. We show that all of these ion channel models can be cast into a unified variational multiscale framework with a macroscopic continuum domain of the solvent and a microscopic discrete domain of the solute. The main strategy is to construct a total energy functional of a charge transport system to encompass the polar and nonpolar free energies of solvation and chemical potential related energies. Using the Euler-Lagrange variation, the coupled PNP equations and other transport equations are derived, whose solutions lead to the minimization of the total free energy and explicit profiles of electrostatic potential and densities of charge species. Current computational algorithms and tools for numerical simulations and results from mathematical analysis of ion channel systems are also surveyed. As this review will be frequently updated, to help improve it, please do not hesitate to send corrections and suggestions to the authors.

Key words: Variational multiscale models, Charge transport, Ion channels, Laplace-Beltrami equation, Poisson-Boltzmann equation, Nernst-Planck equation,

^{*}E-mail: Duan.Chen@uncc.edu

[†]E-mail: Wei@math.msu.edu

Contents

I	Introduction	2
11	The Poisson-Nernst-Planck model II.A Governing equations II.B Gramicidin channel: A showcase for the PNP model	5 6 7
111	Generalized PNP models III.A lon-size effects III.B Classical density functional theory (cDFT) based PNP (cDFT-PNP) III.C Fluid flow and chemical reactions III.D lon-water interactions III.E Poisson-Boltzmann-Nernst-Planck model	9 10 11 11 12 14
IV	Computational algorithms and implementation IV.A Finite difference based methods IV.B Finite element based methods IV.C Other computational methods available for ion channels IV.D Iterative schemes for coupled systems	16 16 17 17 18
v	Mathematical analysis	19
VI	Other mathematical models for ion channel transport VI.A Poisson-Boltzmann-Kohn-Sham model for proton transport VI.B Differential geometry (DG) based charge transport models Variational solute-solvent interface Generalized Laplace-Beltrami (LB) equation Generalized Poisson-Nernst-Planck equation Generalized correlation	20 23 24 25 26 26
vı	I Concluding remarks	28

I Introduction

Membrane charge transport is one of the most important biological processes in lives, as it facilitates signal transduction, action potential, cardiac rhythms, muscle contraction, T-cell activation, etc. Transporting charges could be small proteins, mobile ions, and dipoles in solvent environment, and they are conducted by membrane channels such as ion channel, proton pumps, or general transporter proteins. In a specific but critical manner, ion channels are a type of transporter proteins embedded in cell membranes. They have tube-like water pores in the middle that facilitate selected ion permeation and maintain proper cellular ion compositions. Phospholipid bilayer of cells provides a hydrophobic barrier to the passage of charged ions in extra- and intracellular environment, but strongly polar or even charged amino acids of channel proteins provide a conducting pathway across the hydrophobic interior of the membrane bilayer (Ikezu & Gendelman (2008), Karniadakis et al. (2005)). Resulting ionic flux plays a key role in almost many physiological phenomena from nerve and muscle excitation, human sensory transduction, to cell volume and blood pressure regulation, etc. More critically, ion channels are prominent factors to human health. One example is in cancer research: glioblastoma multiforme exhibits abnormal upregulation of gBK potassium (K^+) and CIC-3 chloride (CI⁻) channels, which aid glioblastoma cells changing cellular volume very rapidly, thus help extremely aggressive invasive behavior of the tumor cells. Another example is the M2 proton (H^+) channel in influenza A virus. The M2 proton channel conducts protons into the virion core, acidifies the virus interior, and leads viral ribo nucleo protein (RNP) complexes release and start viral replication. It has been intensively studied that ion channels are frequent targets in research of new drugs for human diseases (Cross et al. (2012), Dong et al. (2013), Fermini & Priest (2008), Miao et al. (2015), Zhou (2011a)). Advanced giga-seal patch-clamp technique has made the measurement of ionic flowing through a single channel possible (Chung & Kuyucak (2002)), and experimental



Figure 1: Homology structure of a sodium channel constructed from a mosquito genetic data. (a) Top view of a mosquito voltage gated sodium channel. Four homologous domains (I through IV) are illustrated by using four different colors. In each domain, there are six transmembrane segments (S1 through S6). Segments S1-S4 of the channel constitute the voltage-sensing domain (VSD), which is away from the pore, while segments S5 and S6, together with the membrane-reentrant pore loop, form the pore. Image credit: Zixuan Cang. (b) The sodium channel viewed in a cell membrane. Image credit: Christopher Opron.

discoveries are available for structures and functions of potassium channels (Brohawn et al. (2014), Labro et al. (2015)), sodium channels (Catterall (2012), Payandeh et al. (2012)), calcium channels (Fu et al. (2014)), and proton channels (Hondares et al. (2014)). These developments set the stage for theoretical/mathematical modeling to simulate charge transport, to reproduce experimental data, to predict new phenomena, and to offer direction in ion channel-targeted drug design (Boiteuxa et al. (2014), Lacroix et al. (2013)).

There is quite good understanding of the molecular basis of voltage gated potassium channels across various species, such as bacteria, insects, and mammals, due to the availability of many high quality X-ray crystallographic structures (Gutman et al. (2005), Long et al. (2005)), and detailed theoretical analysis (Jensen et al. (2012b)). Voltage gated potassium channels typically are simple, single-domain proteins that assemble to form functional homotetramers. However, eukaryotic voltage gated sodium channel (Nav) channels, such as those of mosquitoes and humen, are complex and four homologous domain proteins assembled into pseudotetrameric structures, with no X-ray crystallographic structure, see Fig. 1. The only existing high resolution X-ray crystallographic structures for Nav channels are from bacteria (Bagneris et al. (2013), McCusker et al. (2012), Payandeh et al. (2012), Shaya et al. (2014)). Similarly to voltage gated potassium channels, bacterial Nav channels have four identical subunits arranged to form a functional channel. The molecular mechanism of eukaryotic Nav channel gating transitions, between voltage-sensing, activation, and deactivation as well as their coupling and interaction with drugs and/or insecticides, is largely unknown or unclear, partially due to limited experimental means, theoretical models and computational power (Dong et al. (2014), Du et al. (2013, in press 2015)). Such a gap in our understanding severely hinders our ability to design effective mosquito insecticides as well as qualified drugs for epilepsy, irregular cardiac arrhythmias, hyperalgesia, myotonia, and anesthesia.

Modern ion channel models are based on the biological understanding of the function and gating mechanism of ion channels, which heavily depend on their molecular structures. A channel protein usually consists of several hundreds to thousands amino-acid residues. Some of them have simpler structures, for example, the Gramicidin A (GA) channel obtained from the soil bacterial species Bacillus brevis is just a dimer that consists of two head-to-head β -helical parts in a bilayer membrane. All of the residues form a narrow pore of about 4Å in diameter and 25Å in length that simply conducts monovalent cations, binds bivalent cations, while rejects anions. On the contrast, the KcsA (potassium crystallographically-sited activation) channel has a relatively complicated structure and hence complete functions. It is comprised of around 560 residues that form four

identical subunits, each containing two alpha-helices connected by a loop of approximately 30 amino acids. There are three primary functioning sections of the KcsA channel: the opening pore on the cytoplasmic side of the cell interior, a small cavity filled with water and a mix of sodium (Na^+) and potassium (K^+) ions, and the selectivity filter (Egwolf, B. & Roux, B. (2010)). The KcsA channel is specialized to facilitate and regulate the conduction of K⁺ ions in particular (Egwolf, B. & Roux, B. (2010)). Further, the technology of homology provides ability to discover more ion channel three dimensional (3D) structures from their genetic data. Figure 1 shows the homology structure of a sodium channel constructed from a mosquito genetic data. In fact, voltage gated potassium and sodium channels are homologous to each other. Therefore, a full description of of ion channels at *atomic level* is required in order to investigate their function and gating comprehensively. On the other hand, a major characteristics of an ion channel system is its heterogeneity. It is inhomogeneous in terms of materials: as shown in Fig. 2(a), an ion channel exists in an extremely complicated environment including cell membrane, water molecules, mobile ions and other molecular components. All of these components are subject to intensive mutual long-range (e.g., electrostatics) and short-range (e.g. Lennard-Jones) interactions. It is also inhomogeneous in terms of functions: an ion channel may have both water-rich, or bulk solvent regions where ions are fully hydrated, and a region where water molecules can only form a single file and ionwater clusters need to be rearranged. Additionally, an ion channel may have different function domains that are responsible for voltage sensing, gating, conducting efficiency and selectivity. In fact, some of these domains may extend away from the channel pore, as in the case of voltage gated sodium channels (Catterall (2012), Payandeh et al. (2012)). Furthermore, ion channel gating can be modulated by ligand, toxicity, substrate, et cetera, which is often the basis for anesthesia and insect control. Therefore, theoretical modeling and simulations of the molecular mechanism for charge transport in ion channels often involve an excessively large number of degrees of freedom and encounter enormous challenges (Wei (2010, 2013), Wei et al. (2012)).

One of the pioneering works about the functions of ion channels is the Hodgkin-Huxley model derived by Hodgkin & Huxley (1952), which is used to study action potentials initiated and propagated in neurons by modeling the ensemble of voltage dependent channels in nerve fibers using nonlinear ordinary differential equations. This model is still a basis for many present interesting ion channel studies. In this review, we focus on models for the functions of a single ion channel. Among mathematical/theoretical models of ion channels developed in past decades, molecular dynamics (MD) stands among top of the hierarchy in terms of accuracy. As a type of all-atom channel model, it treats channel protein, cell membrane, solvent and ions explicitly with positions and velocities propagated by the Newtonian dynamics. MD simulation provides a way to investigate the ultimate details of how structures move and which motions may be linked to biological functions. From an MD simulation trajectory, a variety of thermodynamic (e.g., the free energy changes associated with solvation or ion binding inside a channel pore) or kinetic (e.g., the rate of ion passing through a channel) quantities can be calculated. Computer tools based on MD have been extensively developed by Engels et al. (1995), Klapper et al. (1986), MacKerell et al. (1998), Madura et al. (1995), Perlman et al. (1995), Prabhu et al. (2008) and simulations have been widely used to compute energetics (potential of mean forces PMFs) of ion or water transport through biological channels. Further, MD is a suitable model to study some specific function unit of ion channels such as selectivity mechanism in the local selectivity filters. However, the major drawback of the explicit method is the extremely large number of degree of freedom for the system, so the computation remains very expensive even with contemporary computer powers due to the necessarily small time step (10^{-15}) seconds) versus the ion permeation time scale (10^{-6} seconds). Some important ion channel models in MD are referred to Baker et al. (2016), Dai & Zhou (2014), Heymann et al. (2013), Im & Roux (2002), Lin et al. (2013), Marx & Hutter (2000), Roux (2002), Roux et al. (2004), Schumaker et al. (2000). Specifically, gating motion of ion channels, and their interactions between transmembrane were studied in Dai et al. (2015), Dai & Zhou (2014), Heymann et al. (2013), Jensen et al. (2012a). Meanwhile, some fast algorithms were developed for all-atoms models to study protein folding and binding Qin & Zhou (2013, 2014).

Brownian dynamics (BD) fades out the molecular details of membrane bilayer and water molecules, while only treats mobile ions and channel protein explicitly. The motion of target ions is governed by the Langevin equation. In this model, the forces acting on the ions include frictional/random forces from the surrounding solvent, and the total electrostatic forces due to other mobile ions, fixed charges in the channel protein, solvent polarized electrostatic field and/or applied transmembrane potential that are determined by solving the Poisson equation. Many BD based algorithms are developed and because of the significant reduction in the number of degrees of freedom. The BD simulations are highly efficient. The trajectories of mobile ions can be simulated at usually microsecond scale, from which the single channel current can be derived by counting the number of ions that move across the channel. A series of work has been established to study functions of channel proteins via BD by Cheng & Coalson (2005), Cheng et al. (2010), Coalson & Kurnikova (2005), Gordon et al. (2009).

One of simplest ion channel models is the *Poisson-Nernst-Planck* (PNP) theory, which is a mean-field approach with a low resolution of ion channels, but offers high efficiency. Using a continuum approximation, the PNP model treats the ion flow as the averaged ion concentration driven by the electrostatic potential force and ion concentration gradient. Meanwhile, it incorporates the static atomistic charge description of channel proteins. Thus, it hybrids the macroscopic/continuum description of ionic channel flows with the microscopic/discrete representation of protein electrostatic charge sources, see Fig 2(b). For a 1:1 electrolyte, the PNP system (without physical parameters) is essentially a system of coupled partial differential equations (PDEs):

$$\begin{aligned} & -\epsilon \Delta \phi = \rho_f + p - n, \\ & \frac{\partial p}{\partial t} = \nabla \cdot (\nabla p + p \nabla \phi) \\ & \frac{\partial n}{\partial t} = \nabla \cdot (\nabla n - n \nabla \phi), \end{aligned}$$
(1)

where $\phi(\mathbf{r})$, $p(\mathbf{r}, t)$ and $n(\mathbf{r}, t)$ are electrostatics, concentrations of positive mobile ions (cations) and negative mobile ions (anions), receptively. In this approach, both water and cell membrane are approximated by dielectric function ϵ , and the structure of channel protein is modeled by static point charges or atomic charge density ρ_f . Under this framework, concentrations of the ions through the channel follow the Ohm's and Fick's law, and form two drift-diffusion equations in the same structure. It is worthwhile to point out that under non-flux boundary conditions, the PNP model at stationary state reduces to the Poisson-Boltzmann (PB) model Wei et al. (2012). The PNP model was introduced to the field of molecular biophysics in early 1990s by Barcilon et al. (1992), Chen et al. (1995), Eisenberg (1996), Eisenberg & Chen (1993), Eisenberg et al. (2010), Singer et al. (2008) from a similar approach, called drift-diffusion equations in electronic devices community and widely used in ion channel simulations afterwards, as one of the current major workhorses (Allen et al. (2001), Cardenas et al. (2000), Chen et al. (1997), Choudhary et al. (2010), Constantin & Siwy (2007), Corry et al. (2003), Dyrka et al. (2008), Gillespie et al. (2002), Graf et al. (2004), Hwang et al. (2006), Kurnikova et al. (1999), Levitt (1999), Mamonov et al. (2006), Simakov & Kurnikova (2010), Singer et al. (2008), Zhou, Lu, Huber, Holst & McCammon (2008)).

But the reality of ion channels is far more complex than three PDEs. The starting point of the fundamental PNP system in Eq. (1) is the assumption of volume-less point charge approximation of mobile ions embedded in a structureless and homogeneous continuum model of water molecules. Thus, the original model neglects the steric effects of ions in significant geometric confinement in channel pore, ion-water interactions, polarization of water molecules, channel motion, and many other interactions that are not directly and exclusively relate to electrostatics, but critically contribute to complicated functions of ion channels such as selectivity and activation. Some theoretical approaches have been proposed towards the direction of improving the traditional Poisson-Boltzmann or PNP theory, such as in Antypov et al. (2005), Bazant et al. (2011, 2004), Ben-Yaakov et al. (2011), Schmuck & Bazant (2015), Vlachy (1999), Wei (2010), Wei et al. (2012). Development, analysis, and computation of PNP-based models for ion channel transport have also attracted much attention in the community of applied mathematics.

In this review, we first review the fundamental PNP model and its application on a simple ion channel in Section II. The recent progresses on new models development, computational methods, and mathematical analysis are surveyed in Sections III, IV, and V, respectively. Finally, we discuss some other advanced mathematical models in Section VI, including the Poisson-Boltzmann-Kohn-Sham model for proton channels and a differential geometry based multiscale model for a comprehensive understanding of solvation and charge transport.

II The Poisson-Nernst-Planck model

The variational derivation of Poisson-Boltzmann (PB) equation was given in early 1990s by Sharp & Honig (1990). Similarly, the system of PNP equations can be derived by the variation of a total energy functional



Figure 2: Illustration of an ion channel system and the multiscale approach. (a) Atomic view of the Gramicidin A channel in the membrane and aqueous environment; (b) A cross section of the multiscale representation of the system.

(Fogolari & Briggs (1997)). However, unlike the PB equation or the Poisson equation, which can derived entirely from total variation, the derivation of the Nernst-Planck equation follows two steps, namely, using the energy variation to obtain the chemical potential and then using Fick's laws of diffusion to attain the Nernst-Planck equation. A somewhat more rigorous derivation from the conservation laws has been given recently by Wei (2010). This approach allows the coupling to flow velocity and potential chemical reactions, such as protonation and deprotonation of amino acids, which occur very often during ion channel permeation.

In this review, we follow a simple derivation as in Fogolari & Briggs (1997), Zheng & Wei (2011). Assuming there are multiple ionic species in an ion channel system and ρ_{α} is the concentration of the α -th ion species, the total free energy for the system can be described in terms of the electrostatic potential Φ and the concentration ρ_{α} as the following:

$$G_{\text{total}}^{\text{PNP}}[\Phi, \{\rho_{\alpha}\}] = \int \left\{ -\frac{\epsilon_m}{2} |\nabla \Phi|^2 + \Phi \ \rho_m - \frac{\epsilon_s}{2} |\nabla \Phi|^2 + \Phi \sum_{\alpha} \rho_{\alpha} q_{\alpha} + \sum_{\alpha} \left[(\mu_{\alpha}^0 - \mu_{\alpha 0})\rho_{\alpha} + k_B T \rho_{\alpha} \ln \frac{\rho_{\alpha}}{\rho_{\alpha 0}} - k_B T (\rho_{\alpha} - \rho_{\alpha 0}) + \lambda_{\alpha} \rho_{\alpha} \right] \right\} d\mathbf{r}.$$
(2)

The first row of Eq. (2) is the electrostatic free energies of the system. Here, the ion channel protein is modeled as the fixed charge density ρ_m in atomic details with dielectric constant ϵ_m . In contrast, the solvent is modeled by the ionic density ρ_{α} and water molecules are treated as a dielectric continuum with dielectric constant ϵ_s . Here q_{α} is the charge of α th ion species. The second row includes chemical potential related energy and entropy of mobile ions, where $\mu_{\alpha 0}$ is the chemical potential and μ_{α}^0 is the reference chemical potential of the α th species at which the associated reference concentration is $\rho_{\alpha 0}$. k_BT is the thermal energy with k_B being the Boltzmann constant and T being the temperature. At last, a Lagrange multiplier λ_{α} is used to ensure appropriate physical properties at equilibrium (Fogolari & Briggs (1997)).

By applying the variational principle, governing equations for the variables Φ and ρ_{α} of the system can be obtained.

II.A Governing equations

First, the Poisson equation can be derived by taking the variation with respect to the electrostatic potential Φ , i.e,

$$\frac{\delta G_{\text{total}}^{\text{PNP}}}{\delta \Phi} \Rightarrow \nabla \cdot (\epsilon \nabla \Phi) + \rho_m + \sum_{\alpha} \rho_{\alpha} q_{\alpha} = 0.$$
(3)

Then it yields

$$-\nabla \cdot (\epsilon \nabla \Phi) = \rho_m + \sum_{\alpha} \rho_{\alpha} q_{\alpha}.$$
(4)

In many multiscale models such as in Chen (2016), Zheng et al. (2011), the Poisson equation is defined in the whole computational domain Ω , which consists of the solute domain Ω^+ and the solvent domain Ω^- , on which the dielectric function ϵ is defined as a piecewise constant function

$$\epsilon = \begin{cases} \epsilon_m, & \mathbf{r} \in \Omega^+, \\ \epsilon_s, & \mathbf{r} \in \Omega^- \end{cases}$$
(5)

At this moment domains Ω^+ and Ω^- are assumed to be divided by a given molecular surface Γ , i.e., $\Omega = \Omega^+ \cup \Omega^-$ and $\Gamma = \Omega^+ \cap \Omega^-$. Naturally, the Nernst-Planck equation (and ρ_α) is confined in the domain Ω^- . Note that the dividing surfaces for dielectric constant ϵ and ρ_α are not necessary to be the same, as suggested in Kirkwood (1934). However, this review, we assume the same surface.

The derivation of the Nernst-Planck equation follows two steps. First, the variation of the total free energy functional with respect to ion concentration ρ_{α} is the relative generalized potential $\mu_{\alpha}^{\text{gen}}$ of species α , i.e.:

$$\frac{\delta G_{\text{total}}^{\text{PNP}}}{\delta \rho_{\alpha}} \Rightarrow \mu_{\alpha}^{\text{gen}} = \mu_{\alpha}^{0} - \mu_{\alpha 0} + k_{B}T \ln \frac{\rho_{\alpha}}{\rho_{\alpha 0}} + q_{\alpha}\Phi + \lambda_{\alpha}, \tag{6}$$

where

$$\lambda_{\alpha} = -\mu_{\alpha}^{0} \quad , \quad \rho_{\alpha} = \rho_{\alpha 0} e^{-\frac{q_{\alpha} \Phi - \mu_{\alpha 0}}{k_{B} T}}, \tag{7}$$

and therefore

$$\mu_{\alpha}^{\text{gen}} = k_B T \ln \frac{\rho_{\alpha}}{\rho_{\alpha 0}} + q_{\alpha} \Phi - \mu_{\alpha 0}.$$
(8)

This quantity vanishes at the system equilibrium, while in a non-equilibrium situation, the ion flux density \mathbf{J}_{α} is given through the gradient of the relative generalized potential, i.e., $\mathbf{J}_{\alpha} = -D_{\alpha}\rho_{\alpha}\nabla \frac{\mu_{\alpha}^{\text{gen}}}{k_{B}T}$, where D_{α} is the diffusion coefficient of species α . If stream velocity and chemical reaction are neglected (Wei (2010)), the the mass conservation of species α gives $\frac{\partial \rho_{\alpha}}{\partial t} = -\nabla \cdot \mathbf{J}_{\alpha}$, i.e.,

$$\frac{\partial \rho_{\alpha}}{\partial t} = \nabla \cdot \left[D_{\alpha} \left(\nabla \rho_{\alpha} + \frac{\rho_{\alpha} q_{\alpha}}{k_B T} \nabla \Phi \right) \right],\tag{9}$$

Equation (9) is only defined in the solvent domain Ω^- but forms a coupled system with Eq. (4) for describing the charge concentrations ρ_{α} and the electrostatic potential Φ . The solutions of these equations need to be pursued self-consistently. Based on the ionic flux density, the general formulation to calculate an important physical observable-ionic current, as the total flux through a cross section at the steady state, is

$$I = \sum_{\alpha=1}^{N_c} q_\alpha \int_{L_x, L_y} D_\alpha \left(\frac{\partial \rho_\alpha}{\partial z} + \frac{\rho_\alpha q_\alpha}{k_B T} \frac{\partial \Phi}{\partial z} \right) dx dy.$$
(10)

II.B Gramicidin channel: A showcase for the PNP model

As a showcase, electrostatic profiles of the GA channel, calculated from the PNP system, is mapped on the protein surface and shown in Fig. 3. Overall the GA is neutral in terms of charges, but its surface electrostatic potential is mostly negative near the channel mouth as indicated by the red color in the graph. Also, the inner wall of the channel pore is also intensively negatively charged as shown in Fig. 3(b). This fact indicates the obvious selectivity of the GA channel — it selects cations and suppresses anions. The electrostatics of the channel system greatly depends on the dielectric constants used in Eq. (5). However, the choice of this key model parameter is very subtle and nontrivial because rotation and polarization of water molecules in the narrow channel pore are significantly different from those in bulk solvent. The true dielectric constants have been



Figure 3: 3D illustration of the electrostatic profile of the Gramicidin A (GA) channel. The red and blue colors represent negative and positive electrostatics, respectively. (a) Top view of the GA channel; (b) Side view of the GA channel.



Figure 4: Concentration profiles in the GA channel. (a) with reference concentration $\rho_0 = n_0 = 0.1$ M and variable transmembrane voltage Φ_0 (b) with $\Phi_0 = 50$ mV and variable ρ_0 . Two dashed vertical lines represent the entrance and exist of the channel.

explored in order to obtain a reasonable prediction in Chen & Wei (2013). There is a general agreement that ϵ_m could be taken as a constant that slightly greater than 2, which is the value used in the solvation study. While the dielectric constant ϵ_s for the solvent should be position dependent. The dielectric constant $\epsilon_{\text{bath}} = 80$ is the value widely accepted in the literature for the bath water region.

Concentration profiles of mobile ions in a 1:1 electrolyte, such as KCl or NaCl, in the GA channel are illustrated in Fig 4, where the ion concentrations of K^+ and Cl^- in the GA channel are displayed, against transmembrane voltage Φ_0 and reference concentration $\rho_0 = n_0$, respectively. In these figures, the concentration of K^+ is dominant over the concentration of Cl^- in the channel region, so these simulations agree with the fact that GA channel only conducts cations but reject anions. When the reference $\rho_0 = n_0$ is fixed and the transmembrane potential difference Φ_0 is zero, see Fig. 4(a), distribution of K^+ tends to the right end of the channel and then ion current is generated. Fig. 4 shows concentration of K^+ in channel increases as ρ_0

increases, while it is a constant in the bulk solvent due to the neutral electrostatics.



Figure 5: A comparison of simulated I-V curves and experimental data from Cole et al. (2002) for Gramicidin A channel.

One way to validate the proposed model is to compare the simulated current-voltage (IV) curves with experimental results. In electrophysiology, the voltage refers to the voltage across a membrane, and the current is the flow of charged ions across protein pore. Some experimental results of I-V curves of the GA channel for KCl were reported by Cole et al. (2002). For the bulk diffusion coefficients of K⁺ and Cl⁻, the experimental data are used, i.e., $D_{\rm K} = 1.96 \times 10^{-5} {\rm cm}^2 {\rm /s}$ and $D_{\rm Cl} = 2.03 \times 10^{-5} {\rm cm}^2 {\rm /s}$ for K⁺ and Cl⁻, respectively (Zheng et al. (2011)). However, the diffusion coefficients in the channel pore are not known in general. In order to match experimental results, smaller diffusion coefficients are usually used in the channel region due to the restricted diffusion in most ion channels. Figure 5 presents the reasonable match between the IV curve simulated by the PNP model and experimental data from Cole et al. (2002), with a diffusion coefficient in the channel 25 times smaller than the bulk coefficient.

III Generalized PNP models

In the charge dynamics modeled by the traditional NP equation (9), mobile ions are treated as volume-less point charges. This is a reasonable assumption for bulk or diluted solvents, but it could be problematic for crowded ionic population in a narrow channel pore. As shown in Fig. 6, ionic sizes of mobile ions are comparable to geometry configuration in the narrow channel pore, and they play significant roles in interactions with water molecules and selectivity of ion channels. Many generalizations of the original PNP framework have been proposed. Mathematical models for the finite size effects in ionic solutions were proposed by introducing an energy term that represents the hard sphere repulsion of ions under the PNP framework, as in Hyon et al. (2010, 2011), Lin & Eisenberg (2014). The total energy then is made of the entropic energy, electrostatic potential energy, and the repulsive potential energy. As a result, modifications of the Poisson-Nernst-Planck (PNP) equations were derived, including the effects of the finite size of ions that are so important in the concentrated solutions near electrodes, active sites of enzymes, and selectivity filters of proteins. More recently, a nonlinear



Figure 6: (a) A schematic drawing of steric effects of ions and ion-water interactions in channel pore. Molecular surface Γ partitions the whole region into the solute part (Ω^+) and the solvent part (Ω^-). In the extremely narrow channel pore, ion-water clusters need to re-arrange their original configurations in bulk solvent in order to go through the channel region. In this situation, ion size effects and polarization of water dipoles affected in the vicinity of an ion play significant roles in ion transport dynamics.

Poisson model, a Poisson-Nernst-Planck-Fermi model, and an ionic concentration and size dependent dielectric permittivity Poisson-Boltzmann model were proposed to study the water molecules in solvent as heterogeneous media in the mean-field theory (Hu & Wei (2012), Li & Lu (2014), Liu & Eisenberg (2014)). Other work of modification of PNP theory can be found in Adalsteinsson et al. (2008), Burger et al. (2012), Hyon et al. (2011), Kilic et al. (2007), Li et al. (2010), Qiao et al. (2016, 2014), Xu, Ma & Liu (2014).

III.A Ion-size effects

A specific mathematical model for the finite size, specially, the repulsive effects of mobile ions has been introduced in Hyon et al. (2010) for ion channels. An appropriate energy term that represents the hard sphere repulsion of ions was built into the total energy consisting of the entropic and electrostatic potential energies, then variational approach leads to a modified NP equation for ionic species ρ_{α} as follows:

$$\frac{\partial \rho_{\alpha}}{\partial t} = \nabla \cdot \left\{ D_{\alpha} \left[\nabla \rho_{\alpha} + \frac{\rho_{\alpha}}{k_{B}T} \left(q_{\alpha} \nabla \Phi - \int \frac{12\varepsilon_{\alpha,\alpha}(a_{\alpha} + a_{\alpha})^{12}(\mathbf{r} - \mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|^{14}} \rho_{\alpha}(\mathbf{r}') d\mathbf{r}' \right. \right. \tag{11}$$

$$- \sum_{\beta,\beta \neq \alpha} \int \frac{6\varepsilon_{\alpha,\beta}(a_{\alpha} + a_{\beta})^{12}(\mathbf{r} - \mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|^{14}} \rho_{\beta}(\mathbf{r}') d\mathbf{r}' \right] \right\}.$$

Comparing to Eq. (9), this model involves two integral terms with the Lennard-Jones hard-core repulsion kernels to model the repulsing energy between ion ρ_{α} themselves and with all other ion ρ_{β} , where $\varepsilon_{\alpha,\beta}$ are empirically chosen energy constants for the repulsive interactions between ionic concentration ρ_{α} and ρ_{β} with the radii of the two species are a_{α} and a_{β} , respectively.

In another work by Lu & Zhou (2011), the finite size effects of ions were modeled based on the Borukhov

model, in which an ideal-gas-like solvent entropy term is included in the total functional, to represent the unfavorable energy of over-packing or crowding of ions in narrow channels. After variation, the new NP equation reads:

$$\frac{\partial \rho_{\alpha}}{\partial t} = \nabla \cdot \left\{ D_{\alpha} \left[\nabla \rho_{\alpha} + \frac{\rho_{\alpha}}{k_B T} \left(q_{\alpha} \nabla \Phi + \frac{k_{\alpha} \rho_{\alpha} \sum_{\beta} a_{\beta}^3 \nabla \rho_{\beta}}{1 - \sum_{\beta} a_{\beta}^3 \rho_{\beta}} \right) \right] \right\},\tag{12}$$

where a_{β} is the radius of the β th ionic species and k_{α} is the ratio between a_{α} and the radius of water molecules.

In the work of Burger et al. (2012), a modified PNP system was established as for size effects in confined geometry of channels

$$\frac{\partial \rho_{\alpha}}{\partial t} = \nabla \cdot \left\{ D_{\alpha} \left[(1-c)\nabla \rho_{\alpha} + \rho_{\alpha}\nabla c + \frac{\rho_{\alpha}q_{\alpha}}{k_{B}T} (1-c)\nabla \Phi + \mu_{\alpha}\rho_{\alpha}(1-c)\nabla W_{\alpha}^{0} \right] \right\},\tag{13}$$

where μ_{α} is the entropy variable, $c = \sum_{\alpha} \rho_{\alpha}$, and W_{α}^0 is an external potential. The nonlinear mobilities described by the model was derived based on a discrete lattice-based hopping model with volume exclusion, with the investigation of the system entropy in time.

In these models, the ion-ion (repulsive) interaction terms are all in terms of either concentration or gradient of concentration of mobile ions. This is the so-called density functional approach, which could be unified in a framework of density functional theory (DFT). In the differential geometry based model discussed in Section VI.B, the size effects of ions can be included in a unified term called generalized correlations, which include not only ion-ion interactions, but also short-range interactions among ions, ion-water, and ion-proteins.

III.B Classical density functional theory (cDFT) based PNP (cDFT-PNP)

Another strategy to include the above discussed size effects or other interactions beyond electrostatics is to include them uniformly by the classical density functional theory (cDFT). In this approach, the cDFT-PNP can be written as (Meng et al. (2014)):

$$\frac{\partial \rho_{\alpha}}{\partial t} = \nabla \cdot \left\{ D_{\alpha} \left[\nabla \rho_{\alpha} + \frac{\rho_{\alpha}}{k_B T} \left(q_{\alpha} \nabla \Phi + \nabla \mu_{\alpha}^{\rm id}(\mathbf{r}) + \nabla \mu_{\alpha}^{\rm ex}(\mathbf{r}) \right) \right] \right\}.$$
(14)

Besides the electrostatics energy, the ideal chemical potential energy $\mu_{\alpha}^{id}(\mathbf{r})$ and excess chemical potential energy $\mu_{\alpha}^{ex}(\mathbf{r})$ are included in the ion dynamics. Both of the two terms can be expressed as functionals of ionic densities, i.e.,

$$\mu_{\alpha}^{\rm id}(\mathbf{r}) = -\ln\left[\gamma_{\alpha}\rho_{\alpha}(\mathbf{r})/\rho_{\alpha}^{0}\right],\tag{15}$$

where γ_{α} is the activity coefficient described by the extended Debye-Huckel theory. Meanwhile,

$$\mu_{\alpha}^{\text{ex}}(\mathbf{r}) = \frac{\delta F^{\text{ex}}(\{\rho_{\alpha}(\mathbf{r})\})}{\delta \rho_{\alpha}(\mathbf{r})}$$
(16)

and the excess free energy functional $F^{\text{ex}}(\{\rho_{\alpha}(\mathbf{r})\})$ includes hard-sphere components, short-range interactions, Coulomb interactions and electrostatic correlations, i.e.,

$$F^{\text{ex}}(\{\rho_{\alpha}(\mathbf{r})\}) = F^{\text{ex}}_{\text{hs}} + F^{\text{ex}}_{\text{sh}} + F^{\text{ex}}_{\text{C}} + F^{\text{ex}}_{\text{el}}, \qquad (17)$$

where the expression of each term can be found in Ref. Meng et al. (2014).

III.C Fluid flow and chemical reactions

When fluid flows play a crucial role in the density distribution of charge spices and electrostatic properties, the PNP equations can be coupled with the Navier-Stokes equations.

In Eisenberg et al. (2010), Wang, Liu & Tan (2016), Xu, Sheng & Liu (2014), the Poisson-Nernst-Planck-Navier-Stokes (PNPNS) equations were derived from the energetic variational approach (EnVarA) for a 1:1 electrolyte

$$E^{\text{total}} = \int_{\Omega} \left\{ \frac{c}{2} |\mathbf{v}|^2 + k_B T \left(n \ln \frac{n}{n_0} + p \ln \frac{p}{p_0} \right) + \frac{ze}{2\epsilon} (p-n) \int_{\Omega} G(\mathbf{r}, \mathbf{r}') (n-p) (\mathbf{r}') d(\mathbf{r}') \right\} d\mathbf{r},$$
(18)

where \mathbf{v} is the flow stream velocity and $c = \sum_{\alpha} \rho_{\alpha}$ is the total solvent mass density . Note that the energy components for electrostatics of the system is expressed by the Green's function $G(\mathbf{r}, \mathbf{r}')$. As results, the derived PNPNS equations are

$$-\epsilon\Delta\phi = ze(p-n)$$

$$\frac{\partial p}{\partial t} + \nabla \cdot (p\mathbf{r}) = \nabla \cdot \left(D_p \nabla p + \frac{ze}{k_B T} D_p p \nabla \phi\right)$$

$$\frac{\partial n}{\partial t} + \nabla \cdot (n\mathbf{r}) = \nabla \cdot \left(D_n \nabla n + \frac{ze}{k_B T} D_n n \nabla \phi\right)$$

$$\rho \left(\frac{\partial \mathbf{v}}{\partial t} + (\mathbf{v} \cdot \nabla) \mathbf{v}\right) = \eta \Delta \mathbf{v} - \nabla \Pi + (n-p) ze \nabla \phi$$

$$\nabla \cdot \mathbf{v} = 0$$
(19)

with the detailed formulation of Π in Ref. Xu, Sheng & Liu (2014).

In another work by Wei (2010), a total action functional was proposed as

$$G_{\text{total}}^{\text{NS-PNP}}[\Phi, \{\rho_{\alpha}\}] = \int \int \left\{ -\frac{\epsilon_{m}}{2} |\nabla \Phi|^{2} + \Phi \ \rho_{m} - \frac{\epsilon_{s}}{2} |\nabla \Phi|^{2} + \Phi \sum_{\alpha} \rho_{\alpha} q_{\alpha} + \sum_{\alpha} \left[\left(\mu_{\alpha}^{0} - \mu_{\alpha 0} \right) \rho_{\alpha} + k_{B} T \rho_{\alpha} \ln \frac{\rho_{\alpha}}{\rho_{\alpha 0}} - k_{B} T \left(\rho_{\alpha} - \rho_{\alpha 0} \right) + \lambda_{\alpha} \rho_{\alpha} \right] - \left[c \frac{\mathbf{v}^{2}}{2} - p + \frac{\mu_{f}}{8} \int^{t} \left(\frac{\partial \mathbf{v}_{i}}{\partial \mathbf{r}_{j}} + \frac{\partial \mathbf{v}_{j}}{\partial \mathbf{r}_{i}} \right)^{2} dt' \right] \right\} d\mathbf{r} dt,$$
(20)

where η is the viscosity of the fluid. The Einstein summation convention is used in the viscosity term. The first two rows in Eq. (20) have been discussed in the earlier sections. The last row in Eq. (20) describes the Lagrangian of an incompressible viscous flow with the kinetic energy, potential energy and viscous energy lost due to friction. Then the new NP equation derived from Eq. (20) is

$$\frac{\partial \rho_{\alpha}}{\partial t} + \mathbf{v} \cdot \nabla \rho_{\alpha} = \nabla \cdot D_{\alpha} \left[\nabla \rho_{\alpha} + \frac{\rho_{\alpha}}{k_B T} \nabla \left(q_{\alpha} \Phi - \frac{\mathbf{v}^2}{2} \right) \right] + \sum_{j} \bar{\nu}_{\alpha j} J^j, \tag{21}$$

where $\bar{\nu}_{\alpha j} J^j$ is the density production of α species per unit volume in the *j*th chemical reaction. Consequently, the Navier-Stokes equation results as

$$c\left(\frac{\partial \mathbf{v}}{\partial t} + \mathbf{v} \cdot \nabla \mathbf{v}\right) = -\nabla p + \nabla \cdot \mathbb{T} + \mathbf{F}_{\mathrm{E}},\tag{22}$$

where $\mathbf{F}_{\rm E} = \sum_{\alpha} \rho_{\alpha} q_{\alpha} \nabla \Phi$ and \mathbb{T} is the flow stress tensor Wei (2010)

$$\mathbb{T} = \frac{\eta}{2} \left(\frac{\partial \mathbf{v}_i}{\partial \mathbf{r}_j} + \frac{\partial \mathbf{v}_j}{\partial \mathbf{r}_i} \right) = \frac{\eta}{2} \left[\nabla \mathbf{v} + (\nabla \mathbf{v})^T \right].$$
(23)

Here the electrostatic potential is governed by the Poisson equation

$$-\nabla \cdot (\epsilon \nabla \Phi) = \rho_m + \sum_{\alpha} \rho_{\alpha} q_{\alpha}.$$
(24)

Note that chemical reactions do not contribute to the total mass and velocity transport due to the conservation. **III.D Ion-water interactions**

Another view of generalize PNP theory is to model the heterogeneous property of water molecules as a continuum. Experimental observations concluded that that dielectric response of water decreases as ionic concentration increases. A possible explanation is that water molecules form a hydration shell around a

solvated ion, and when away from the ion-water cluster, the orientation of water molecules generally follow the external electrostatic field. In contrast, the motions of those dipoles in the hydration shell are greatly restricted in the vicinity of an ion. They are oriented immediately along the field line generated by the cation or anion, as shown in Fig. 6, leading to an overall decrease in the dielectric response to the electrostatic field.

Based on this observation, a PNP model involving ion-water interactions was proposed by Chen (2016), by modeling water molecules in the solvent as a medium with dielectric function depending on the concentration of mobile ions, i.e., $\epsilon(\{\rho_{\alpha}\})$. In this work, the following total energy is considered:

$$G[\Phi, \{\rho_{\alpha}\}] = \int_{\Omega} \left[k_{B}T \sum_{\alpha} \rho_{\alpha} \ln \frac{\rho_{\alpha}}{\rho_{\alpha 0}} - \frac{\epsilon(\mathbf{r})}{2} |\nabla \Phi|^{2} + \Phi \rho_{m} + \Phi \sum_{\alpha} \rho_{\alpha} q_{\alpha} \right] d\mathbf{r}$$

+
$$\int_{\Omega} \sum_{\alpha} \left[k_{B}T(\rho_{\alpha 0} - \rho_{\alpha}) + (\mu_{\alpha}^{0} - \mu_{\alpha 0})\rho_{\alpha} + \lambda_{\alpha}\rho_{\alpha} \right] d\mathbf{r},$$
(25)

in which the whole domain Ω has been divided into the solute domain Ω^+ and the solvent domain Ω^- and

$$\epsilon(\mathbf{r}) = \begin{cases} \epsilon_m, & \mathbf{r} \in \Omega^+, \\ \epsilon(\{\rho_\alpha\}), & \mathbf{r} \in \Omega^-. \end{cases}$$
(26)

By a similar variation process, a new PNP system with ion-water interaction is obtained.

$$\begin{cases} -\nabla \cdot (\epsilon(\{\rho_{\alpha}\})\nabla\Phi) = \rho_{m} + \sum_{\alpha} \rho_{\alpha} q_{\alpha} \\ \frac{\partial \rho_{\alpha}}{\partial t} = \nabla \cdot \left\{ D_{\alpha} \left[\nabla \rho_{\alpha} + \rho_{\alpha} \nabla \left(\frac{q_{\alpha}}{k_{B}T} \Phi + U_{\alpha} \right) \right] \right\}, \end{cases}$$
(27)

The model of Chen (2016) inherits the structure of the PNP equation and introduces an extra potential energy term:

$$U_{\alpha} = -\frac{\delta\epsilon}{\delta\rho_{\alpha}} \frac{|\nabla\Phi|^2}{2k_B T}.$$
(28)

This energy depends on ionic species, ionic concentration and the electrostatics, and thus it is called the ion-water interaction (IWI) energy. Since $\frac{\delta\epsilon}{\delta\rho_{\alpha}} < 0$, the IWI energy is always positive regardless of the charge of ions, so it is an energy barrier for all ionic species. It offers the ability of the model to *distinguish different ions of the same charges*. In the conventional PNP, the overall potential energy for the dynamics of the α th ion is $\Phi(\mathbf{r})q_{\alpha}/k_{B}T$, i.e., as long as two ionic species have the same valence (e.g., Na⁺ and K⁺), they will have the identical transport dynamics, which is not realistic for some ion channels. In contrast, since $\delta\epsilon/\delta n_{\alpha}$ depends on specific ion types, the IWI energy $-\frac{\delta\epsilon}{\delta\rho_{\alpha}}\frac{|\nabla\Phi|^{2}}{2k_{B}T}$ will be obviously different for ions even with the same valence. This property can be used to study the selectivity of ion channels. Experimental results from Chandra (2000) suggest that the dependence of local dielectric response ϵ of water molecules to the K⁺ is different from that of Na⁺. In other words, the value $\frac{\delta\epsilon}{\delta\rho_{K^{+}}}$ in Eq. (28) is different from $\frac{\delta\epsilon}{\delta\rho_{Na^{+}}}$, where $\rho_{K^{+}}$ and $\rho_{Na^{+}}$ are local concentrations of K⁺ and Na⁺, respectively. Consequently, the overall transport potential energy, given by Eq. (27), takes different values for K⁺ and Na⁺.

Figures 7 (a) and (b) show the energy components in the KcsA channel, for K^+ and Na^+ , respectively. The blue curves in both figures indicate that the electrostatic potential energies are identical for Na^+ and K^+ (although they look differently due to the scaling). As shown by the green curve in Fig. 7 (a), the IWI energy of K^+ is up to 2 k_BT and the overall, effective transport energy (red curve) is alleviated through the channel, but is still negative. On the other hand, Na^+ ions experience the IWI energy as high as 8 k_BT , so the overall potential energy in the KcsA channel becomes an energy barrier. It is worth to pointing out that the dielectric functions and values of the parameter β for K^+ and Na^+ are taken from experiment results. Therefore, the proposed formulation is able to model the selectivity of the KcsA channel, although the reality is much more complicated. Noskov et al. (2004) investigated the selectivity of KcsA channel by comparing the free energy of K^+ and Na^+ in the pore and in the bulk solution through molecular dynamics. It was claimed that the carbonyl groups in the selectivity filter can compensate for the desolvation of K^+ with larger radius while not for the Na^+ of smaller radius. Therefore, both the all-atom explicit and the implicit continuum models demonstrate that ion-water interaction could be a critical factor for ion selectivity.



Figure 7: Profiles of potential energies of the KcsA channel simulated by the PNP-IWI model. Two dashed vertical lines separate the whole channel into three domains: the pore, cavity and filter regions. "Overall" means the sum of the electrostatics and ion-water interaction energies.

In Fig. 8 the current-voltage relations of KcsA channel are compared for the original PNP model and the PNP model with IWI modification. As shown by blue curves, the PNP model predicts similar magnitudes of ionic currents for K^+ and Na^+ ; the only difference is from the diffusion coefficients of two ion species $(K^+: 1.96 \times 10^{-9}m^2/s \text{ and } Na^+: 1.33 \times 10^{-9}m^2/s \text{ Kuyucak et al. (2001)})$. This is against the experimental observation that the conductance of K^+ in the KcsA channel is dominant over Na^+ . Secondly, it is widely believed that PNP model generally overestimates ionic current and a phenomenologically tuned diffusion coefficient is used to match experimental results. The green and red curves are the current-voltage generated by the PNP-IWI model for K^+ and Na^+ , respectively. From these curves, one can see that the simulated ionic current of K^+ is significantly higher than that of Na^+ . Additionally, diffusion coefficients do not need to be reduced in the PNP-IWI model.

III.E Poisson-Boltzmann-Nernst-Planck model

The PNP model provides good descriptions of each ionic species in a non-equilibrium system for charge transport phenomena. However, the computational cost will be extremely high when there is a large number of ionic types in the system because each charge species is governed by one 3D Nernst-Planck equation that needs to be numerically solved. To address this issue, a Poisson-Boltzmann Nernst-Planck (PBNP) model was introduced by Zheng & Wei (2011), in which only concentrations of the target ions (ions of interests) are modeled by the Nernst-Planck equation while those of other ions are described by the Boltzmann distribution. This approach is especially reasonable for charge transport modeling: an ion channel usually has selectivity and it only conducts some specific ion species. For the rest of ions, they can be considered in a quasi-equilibrium state in the bulk solvent. The validity and usefulness of the PBNP formulation were confirmed by independent researchers such as Kiselev et al. (2011).

Assume the total number of ion species in the system is N_c , among which ρ_{α} ($\alpha = 1, \dots, N_{\rm NP}$) are denoted as the densities of the target charge species, or ions of interests, thus $N_{\rm NP}$ is the total number of charge species treated by using the non-equilibrium Nernst-Planck (NP) equation. On the other hand, the densities of the remaining charge species in the system are ρ_{β} ($\beta = N_{\rm NP} + 1, \dots, N_c$) and $N_{\rm BD} = N_c - N_{\rm NP}$ is the total number of the remaining charge species which are represented by the equilibrium Boltzmann distribution.



Figure 8: Simulated current-voltage relations of $\rm K^+$ and $\rm Na^+$ in the KcsA channel.

Based on this consideration, the total free energy functional can be expressed by

$$G_{\text{total}}^{\text{PBNP}}[\Phi, \{\rho_{\alpha}\}] = \int \left\{ -\frac{\epsilon_{m}}{2} |\nabla \Phi|^{2} + \Phi \ \rho_{m} - \frac{\epsilon_{s}}{2} |\nabla \Phi|^{2} + \Phi \sum_{\alpha=1}^{N_{\text{NP}}} \rho_{\alpha} q_{\alpha} - \sum_{\beta=N_{\text{NP}}+1}^{N_{c}} k_{B} T \rho_{\beta 0} \left(e^{-\frac{q_{\beta} \Phi - \mu_{\beta 0}}{k_{B} T}} - 1 \right) + \sum_{\alpha=1}^{N_{\text{NP}}} \left[\left(\mu_{\alpha}^{0} - \mu_{\alpha 0} \right) \rho_{\alpha} + k_{B} T \rho_{\alpha} \ln \frac{\rho_{\alpha}}{\rho_{\alpha 0}} - k_{B} T \left(\rho_{\alpha} - \rho_{\alpha 0} \right) + \lambda_{\alpha} \rho_{\alpha} \right] \right\} d\mathbf{r}.$$

$$(29)$$

Equation (29) includes the same energy components as Eq. (2) does, but note two different treatments for charged ion species are taken for the charge source terms in the polar solvation free energy functional.

It is then a standard procedure to derive the PBNP equations from the total energy functional (29).

$$\begin{cases} -\nabla \cdot (\epsilon \nabla \Phi) = \rho_m + \left(\sum_{\alpha=1}^{N_{\rm NP}} q_\alpha \rho_\alpha + \sum_{\beta=N_{\rm NP}+1}^{N_c} q_\beta \rho_{\beta 0} e^{-\frac{q_\beta \Phi - \mu_{\beta 0}}{k_B T}}, \right). \\ \frac{\partial \rho_\alpha}{\partial t} = \nabla \cdot \left[D_\alpha \left(\nabla \rho_\alpha + \frac{\rho_\alpha q_\alpha}{k_B T} \nabla \Phi \right) \right], \end{cases}$$
(30)

In practical applications, the Nernst-Planck equation is only needed for the ions of interests ρ_{α} and usually $N_{\rm NP} << N_c$. This treatment can significantly reduce computational costs in simulations for a given level of modeling accuracy. For an electrolyte that contains several ionic species, it is strategically useful to focus on the ion of major interests and then use the PBNP equations, instead of the full PNP equations, in order to reduce model complexity and computational costs. The ability of quasi-equilibrium PBNP model to recover the full predictions of the non-equilibrium PNP model was tested in Wei et al. (2012).

Figure 9 provides the comparison of the cross sections of electrostatic potential and concentration profiles obtained from PNP and PBNP models for the GA channel (Zheng & Wei (2011)). The external voltage is set to

 $\Phi_0 = 100 \text{mV}$ and the salt (KCl) concentration is $n_0 = 0.5 \text{M}$. Concentration of Cl⁻, $n_{\text{Cl}^-}(\mathbf{r})$ is represented by using the Boltzmann distribution, while solve the Nernst-Planck equation for K⁺ density $n_{\text{K}^+}(\mathbf{r})$. In figures, solid curves are simulation results from the PBNP equation, while the dots are from the PNP equations. Electrostatic potential computed by the reduced LB-PBNP model agree quite well with the full PNP model. For the density profile, reduced PBNP model also does an excellent job in the channel region, which is the region of main interest.



Figure 9: Comparison of cross sections of electrostatic potential and concentration profiles from the PNP and PBNP models. Transmembrane voltage $\Phi_0 = 100 \text{mV}$, reference concentration $n_0 = 0.5 \text{ M}$. (a) Electrostatic potential profiles; (b) Concentration profiles.

IV Computational algorithms and implementation

In order to understand the realistic chemical, physical, and biological properties in ion transport processes and to predict reliable results, many algorithms and computational tools have been developed to obtain highly accurate and efficient numerical solutions of various proposed models.

IV.A Finite difference based methods

The PNP and related system can be discretized by finite difference method (FDM). In Chen, Chen, Chen, Geng & Wei (2011), Zheng et al. (2011), a set of FDM based algorithms was developed for the second-order convergence solutions of the PNP equations with 3D *realistic and complicated* solvent excluded surfaces, in which the matched interface and boundary method (MIB) developed in Xia & Wei (2014), Xia et al. (2012, 2011, 2014), Yu & Wei (2007), Yu et al. (2007), Zhao & Wei (2004), Zhou, Feig & Wei (2008), Zhou & Wei (2006), Zhou et al. (2006) was used to handle the discontinuous property of dielectric constants on solvent and solute domain, and Dirichlet to Neumann mapping technique (Geng et al. (2007)) was applied to rigorously treat Dirac delta singularities of fixed charges the protein channels. This PNP algorithm was used to construct a molecular level prototype for mechanoelectrical transducers in mammalian hair cells (Park & Wei (2014)). Finally, many PB solvers, as those in Baker et al. (2001), Bertonati et al. (2007), Im et al. (1998), Rocchia et al. (2001), Wang et al. (2012), can be incorporated to PNP systems.

Finite-difference methods for solving 1D and 2D time-dependent PNP equations was developed by Flavell et al. (2014), Liu & Wang (2014), with second-order accurate solutions in both space and time. These works focus on conservation of total ions, correct rates of energy dissipation, and positivity of the ion concentrations. A set of sufficient conditions on the step sizes of the numerical method were discussed to assure positivity of the ion concentrations solutions over long time scales (Flavell et al. (2014)). Relatively simple and easy-to-implement conservative schemes were established to preserve equilibrium solutions, and they were proved to satisfy the total exact concentrations, preserving positivity of the chemical concentrations under a mild Courant-Friedrichs-Lewy (CFL) condition, and the free energy dissipation law at the semi-discrete level (Liu & Wang (2014)). An energy preserving finite difference scheme was developed in He & Pan (2016).

Efficient numerical algorithms for solving 3D steady-state PNP equations with excess chemical potentials described by the classical density functional theory (cDFT) (Meng et al. (2014)). In these algorithms, the NP equations were transformed into Laplace equations through the Slotboom transformation. The algebraic multigrid method was applied to efficiently solve the system and excess chemical potentials was calculated through fast Fourier transforms with computational complexity of $O(N \log N)$, where N is the number of grid points.

Pseudo-time-coupled nonlinear models was proposed for biomolecular surface representation and solvation analysis by Zhao (2012). Recently, this type of approaches have been further extended to operator splitting alternating direction implicit (ADI) schemes for pseudo-time coupled nonlinear solvation simulations (Zhao (2014)). More recently, a fast ADI algorithm has been developed for geometric flow equations in biomolecular surface generations in Tian & Zhao (2014).

IV.B Finite element based methods

Finite element method (FEM) is also developed to solve the PNP equation (Chaudhry et al. (2014), Lu et al. (2010), Lu & Zhou (2011), Metti et al. (2016), Sun et al. (2016*a*), Tu et al. (2013)) and error analysis is available (Sun et al. (2016*b*), Yang & Lu (2013), Zinsl (2015)). An FEM based solver was developed for a modified form of the PNP equations that includes steric effects of mobile ions by Chaudhry et al. (2014). The algorithm in this work combines Newton's method to the nonlinear Galerkin form of the equations, which are augmented with stabilization terms to appropriately handle the drift-diffusion processes. Periodic boundary conditions of the PNP equations were used to conserve the number of ions in the solution domain and to make comparison with particle-based simulations possible.

A stable regularization scheme was applied to remove the singular component of the electrostatic potential induced by the permanent charges inside biomolecules, and then regular, well-posed weak form of PNP equations were formulated by Lu et al. (2010), Lu & Zhou (2011). For the steady-state problems, an inexact-Newton method was used to solve the coupled nonlinear elliptic equations, and for time integration for the non-steady-state electrodiffusion, the Adams-Bashforth-Crank-Nicolson method was devised. These computational algorithms were also generalized to a size-modified Poisson-Nernst-Planck (SMPNP) model that is able to treat nonuniform particle sizes by the Borukhov model.

A method of lines approach was proposed for the FEM discretization to approximately solve the PNP equations in Metti et al. (2016). This discretization scheme assures positivity of the numerical solutions for particle density. A discrete energy estimate was also established and extended to the FE solutions of an electrokinetic model, which couples the PNP system with the incompressible Navier-Stokes equations. A parallel FE simulator for ion transport was developed in Tu et al. (2013) and error analysis of the FEM for the PNP equations is available in Jerome & Kerkhoven (1990), Sun et al. (2016*a*). A free energy satisfying discontinuous Galerkin method was developed in Liu & Wang (2017).

IV.C Other computational methods available for ion channels

There are many other intelligent computational algorithms, some of which are hybrid methods, and some are currently developed for the equilibrium PB equation but could be extended to study non-equilibrium PNP-like equations in the future.

Hybrid models: Baker et al. (2016), Lin et al. (2013) developed an image-charge solvation method (ICSM) combined with molecular dynamics simulations to investigated the selectivity of the KcsA channel. In this hybrid model, all particles including the channel protein, water molecules and mobile ions were described at the atomic level with molecular dynamics in a small neighborhood around the channel, while the reaction field effect of the continuum approximation of the background was computed by the image method. The ICSM is able to demonstrate the function of the selectivity filter of the KcsA channel when potassium and sodium ions are considered. In Jung et al. (2009), a PNP with explicit resident ion, or ERIPNP model, was developed to study biting sites of K^+ ions in the KcsA channel. In this algorithms, the continuum PNP equations are accompanied by explicitly described individual ions with finite size in the selectivity filter of the channel. The ERINP model reproduced the experimental results with a realistic set of parameters and also reduced CPU costs.

Boundary integral method: A boundary integral equation program was provided for calculation of electrostatics in the Poisson Poisson-Boltzmann modeling of an ion channel in layered dielectric/electrolyte media in Lin et al. (2013), Ziner & Cai (2016). A layered media Green's function was used in order to accurately model the inhomogeneous background, including different physical properties and extra/intra cellular environments, cell membrane, and the cylindrical shape of ion channels. A series of parallel, treecode, or GPU-accelerated boundary integral equation methods were developed to calculate electrostatics of solvated biomolecules by Geng (2013), Geng & Jacob (2013), Geng & Krasny (2013).

Nonlocal algorithms: A nonlocal dielectric model and the associated computational package was developed for protein in ionic solvent, taking into account the polarization correlation among water molecules in Xie et al. (2013), Xie & Zhou (2007). Using solution splitting and reformulation techniques, the solution of the nonlocal dielectric model was shown to be uniquely obtained from solving two well-defined PDE systems and one Poisson-like boundary value problem. Additionally, a nonlocal linearized PB equation with uniform ionic size effect was also proposed and numerically tested on three protein molecules.

Stochastic methods: Stochastic walk-on-spheres (WOS) algorithms for solving the linearized Poisson-Boltzmann equation (LPBE) provide several attractive features not available in traditional deterministic solvers (Simonov et al. (2007)): Gaussian error bars can be computed easily, the algorithm is readily parallelized and requires minimal memory and multiple solvent environments can be accounted for by reweighting trajectories. Numerical optimizations that can make the computational time of Monte Carlo LPBE solvers competitive with deterministic methods was introduced by Mackoy et al. (2013). In the optimization techniques, each atom's contribution to the variance of the electrostatic solvation free energy was assured to be the same, and the bias-generating parameters in the algorithm were optimized, and an epsilon-approximate rather than exact nearest-neighbor search was utilized when determining the size of the next step in the Brownian motion.

Levitt approach, rate-equation approach and their equivalence: It was investigated by Levitt (1991a,b) that ion permeation can be modeled as continuous diffusion with the rate of ion transport being obtained from solving the steady-state diffusion equation. The calculation involves determining the potential of mean force and ion diffusion constant, which can be obtained from MD simulations. While in the rate-equation approach Kolomeisky (2007), Läuger (1985), the translocation of the permeant ion through a channel pore is modeled by a hopping mechanism along a discrete set of internal binding sites, and the rate of ion transport is obtained from solving a set of steady-state rate equations. These two approaches are related and the equivalence of them is proved by Zhou et al. (2011).

IV.D Iterative schemes for coupled systems

Regardless computational methods, governing equations in all models are coupled and need to be solved iteratively. To illustrate the iterative process, consider a coupled generalized PNP model as

$$\frac{\partial S^{k+1}}{\partial t} = |\nabla S^{k+1}| \left[\nabla \cdot \left(\gamma \frac{\nabla S^{k+1}}{|\nabla S^{k+1}|} \right) + V_{\text{LB}}(\Phi^k, \{\rho_\alpha^k\}) \right],\tag{31}$$

$$-\nabla \cdot \left(\epsilon(S^k)\nabla\Phi^k\right) = S^k \rho_m + (1 - S^k) \sum_{\alpha} \rho_{\alpha}^k q_{\alpha},\tag{32}$$

$$\nabla \cdot \left[D_{\alpha} \left(\nabla \rho_{\alpha}^{k} + \frac{\rho_{\alpha}^{k}}{k_{B}T} \nabla (q_{\alpha} \Phi^{k} + U_{\alpha}) \right) \right] = 0,$$
(33)

where the function $S(\mathbf{r})$ is a characteristic function identifying the solvent or solute domain. Derivation of the governing equation of $S(\mathbf{r})$, or Eq. (31) is discussed in detail in Section VI.B. In this case, solving Eq. (31) is considered as the outer loop, in which the function S^{k+1} in the (k+1)th step is solved from the electrostatics Φ^k and ρ^k_{α} in the kth step. On the other hand, solving for Φ^k and ρ^k_{α} with an available S^k is also an iterative process and it is called the inner loop. Namely, start with a certain definition of molecular surface S^0 , Φ^0 and ρ^0_{α} are solved self-consistently with this chosen S^0 till convergent. In this case, $\Phi^{0,0}$ is taken as the solution of the corresponding Poisson-Boltzmann equation, and it is used in Eq. (33) to solve for $\rho_{\alpha}^{0,1}$. Then $\rho^{0,1}$ is substituted in Eq. (32) to solve for $\Phi^{0,1}$. After *m*th step $\Phi^{0,m}$ and $\rho_{\alpha}^{0,m}$ are convergent, they are renamed as Φ^0 and ρ^0_{α} and used in the outer loop Eq.(31) to solve S^1 . In the *m*th inner loop for computing $\Phi^{k,m}$ and $\rho^{k,m}_{\alpha}$, the successive over relaxation scheme is utilized (

Chen et al. (2010))

$$\Phi^{k,m} = \zeta_1 \Phi^{k,m} + (1-\zeta_1) \Phi^{k,m-1} \rho^{k,m}_{\alpha} = \zeta_2 \rho^{k,m}_{\alpha} + (1-\zeta_2) \rho^{k,m-1}_{\alpha},$$
(34)

where $0 \le \zeta_1 \le 1$ and $0 \le \zeta_2 \le 1$ are relaxation factors. Larger values of ζ_1 and ζ_2 will lead to slower

convergence, while smaller values may cause instability. Alternatively, the Gummel iteration proposed by de Falco et al. (1964) can also be used to handle this type of problems.

The overall self-consistent process, including inner and outer iterations are summarized as follows.

Step 1: Initial atomic position and partial charge generation. The initial atomic positions of a protein are taken from the Protein Data Bank (PDB) (www.pdb.org), and the partial charge prescription is obtained by the software PDB2PQR Dolinsky et al. (2007, 2004), which provides \mathbf{r}_j and Q_j values in the formulation.

Step 2: Given initial guesses of Φ and ρ_{α} , the surface function S is obtained by the initial value problem Eq. (47). After the surface function S is determined, an isosurface is extracted for the interface Γ .

Step 3: Based on the interface Γ , normal direction **n** is computed by $\frac{\nabla S}{|\nabla S|}$ on the isosurface; the coupled Eqs. (32) and (33) are solved iteratively by above mentioned schemes.

Step 4: Go to Steps 2 and 3 for updating S, Φ and ρ_{α} until a convergence is reached based on a given tolerance. Noticed that in the *l*th outer loop for updating S, we use $S^{l+1} = \lambda_3 S^l + (1 - \lambda_3) S^{l+1}$. In each outer loop, the total free energy functional is evaluated for checking the convergence criteria.

V Mathematical analysis

There has been much effort in the mathematical analysis of the PNP equation and related systems in the past few decades. The nontrivial mathematics and important applications have made this analysis an attractive research topic. Qualitative properties of the steady state of PNP systems in ion channels, including existence and uniqueness of solutions upon boundary conditions, were studied in Park & Jerome (1997). Afterwards, mathematical analysis established for the PNP and related models in past decades are majorly in two aspects: one is the asymptotic behavior of the PNP model and the other is the multiple solutions of modified PNP equations Other analyses are referred to Cartailler et al. (2017), Deng & Li (2013), Hadjadj et al. (2010), Hineman & Ryham (2015).

The PNP system was studied as a singularly perturbed system, with the assumption that the Debye length is small relative to the diameter of the narrow ion channel, by Abaid et al. (2008), Eisenberg & Liu (2006), Liu (2005), Singer et al. (2008), Singer & Norbury (2009), Wang et al. (2014). Explicit derivation of higher order terms in the asymptotic expansions was obtained from special structures of the zeroth order inner and outer systems. Various current-voltage relations of ion channels were described in the case of zero permanent charges in a channel protein with electro-neutrality condition enforced at the ends of the channel (Abaid et al. (2008), Zhang (2015)). Effects of small fixed protein charges and channel geometry on ionic flow was studied in Ji et al. (2015). Ion channels involving two types of ions with three regions of piecewise constant permanent charge were studied by geometric singular perturbation theory, which gives rise to the existence and (local) uniqueness of the solution of the singular boundary value problem near each singular orbit (Eisenberg & Liu (2006), Liu (2009)). By this technique, multiple solutions of the system were discovered and they might explain a variety of multiple valued phenomena in biological channels, such as gating or some kinds of active transport. Liu (2005) studied the global behavior in terms of limiting fast and slow systems, in which three different time scales were indicated in the singularly perturbed PNP system.

Solutions of modified PNP system may explain more physical observation. The ability of PNP system is analyzed to study gating mechanism of ion channels by Lin & Eisenberg (2015). Discovered by experimental measurements, single protein channels produces unstable currents: nearly zero or a definite level of currents. One reason may be the spontaneous stochastic gating process. Existence of multiple solutions of steady state PNP-steric equations were studied to check whether it can describe this two levels of current. Indeed, two steady state solutions of PNP-steric equations were proved for three types of ion species (two types of cations and one type of anion) and four types of ion species (two types of cations and their counter-ions) with specific assumptions on permanent charges in channel proteins. In Jia et al. (2016), a quasi-one-dimensional steady-state PNP model modified with size effect was studied as a singularly perturbed differential system, with fixed boundary ion concentrations and electric potentials. The existence of solutions to the boundary value problem for small ion sizes was investigated with the ion sizes as small parameters. This analysis provided dependencies of current-voltage relations on boundary concentrations, diffusion coefficients and ion sizes.

An open problem in mathematical analysis is how to deal with the free boundary associated with ionic flow. A differential geometry based PNP system has been proposed where the interface between the ion channel and the solvent is governed by the Laplace-Beltrami equation as described in the next section. There is a pressing to analyze this more challenging system.

VI Other mathematical models for ion channel transport

VI.A Poisson-Boltzmann-Kohn-Sham model for proton transport

Proton transport is one special type of charge transport through membranes and plays a critical role in many biochemical processes (Chen, Chen & Wei (2012), Chen & Wei (2012)). For example, generation and conduction of large proton concentration gradients are required in energy transduction in a bioenergetic system: chemical energy is stored as proton gradient that drives the ATP generation in mitochondria of animal cells. While for plants, light energy is transducted into a proton gradient to create ATP in chloroplasts (Decoursey (2003)). Studies of proton transport are also important for public health: for example, the M2 proton (H⁺) channel in influenza A virus. Conducting protons into the virion core and thus acidifying the virus interior, the M2 proton channel leads viral ribo nucleo protein (RNP) complexes release and start viral replication (Schnell & Chou (2008), Sharma et al. (2010)). Also, it has been intensively studied that proton channels are frequent targets in the research of new drugs for human diseases such as cancers (Harguindey et al. (2009)).

However, the mechanism of proton transport is significantly different from that of regular ions such as sodium or potassium. The reason is that proton has the lightest mass among all ions and its effective radius at least 10^5 smaller because the H⁺ has no electron (Decoursey (2003)). The light mass and tiny size greatly facilitate proton transfer reaction and electrostatic interactions with surrounding molecules (Mitchell (1976)). Due to these unique physical properties, the mobility of protons in bulk solution is about fivefold higher than that of other cations (Bernal & Fowler (1933)). Major studies indicated that transport of protons is characterized as a succession of hops in the hydrogen-bond network and is described by the Grotthuss theory in Nagle & Morowitz (1978). Specifically, mechanism of proton transport for the influenza virus M2 channel was studied by Zhou (2011*b*,*c*), in which quantitative calculations have been implemented on the effort of pH, transmembrane voltage, and H₂O to D₂O exchange on the transport rate, and the rate-limiting was concluded to be binding to and unbinding from the His37 tetrad of protons.

Proton transport needs to be treated by quantum mechanical formulations, according to Nagle & Morowitz (1978), Pomes & Roux (2002*b*). Many algorithms, such as multistate empirical valence bond (MS-EVB) approach in Schmitt & Voth (1999), were developed by Schmitt and Voth, to compute dynamics of protons in bulk phase water with an emphasis on a quantum dynamical treatment. Additionally, using Feynman path integral dynamical simulations in Pomes & Roux (1996, 2002*a*,*b*), Roux and his coworkers investigated single file of water molecules in the Gramicidin A channel, which functions as a proton wire. Although not a governing factor, the nuclear quantum effect has a significant impact to proton transfer in equilibrium conditions (Pomes & Roux (1996, 2002*a*,*b*)). Especially under nonequilibrium conditions, such as the presence of an external electrostatic field (Drukker et al. (1998)), or hydrogen-bonding partners are greatly restricted for the displacements of water molecules in narrow channel pores (Decornez et al. (1999)), nuclear tunneling and nonadiabatic transitions need to be accounted in the proton translocation (Bothma et al. (2010), Pomes & Roux (2002*a*)). Other important works are included in Alexov & Gunner (1999), Cukier (2004*a*,*a*,*b*,*b*), Shepherd & Morrison (2010), Wang et al. (2006), Yan et al. (2007) and all these studies are based on a full-atom fashion.

In recent work by Chen & Wei (2013), a multiscale/multiphysics model was developed for the understanding of the molecular mechanism of proton transport in transmembrane proteins via continuum, atomic and quantum descriptions. As shown in Fig. 10, quantum dynamics of proton is expressed in terms of proton concentrations, combined with classical implicit solvent modeling. Further, in order to reduce the number of degrees of freedom, a new quantum density functional theory was constructed, based on the Boltzmann statistics to describe proton dynamics quantum mechanically, while numerous solvent molecules are implicitly treated as a dielectric continuum. A new density functional like formalism is introduced to represent protein density according to the Boltzmann statistics. Additionally, generalized correlations that model interactions among all the ions, and between ions and proteins are explored in detail in Chen & Wei (2012). In this model, proton kinetic and potential energies, the free energy of all other ions, and the polar and nonpolar energies of the whole system are integrated in a multiscale framework on an equal footing. In these models, we consider the



$\Omega_s \bigcup \Omega_m$: Generalized Laplace-Beltrami equation $\Omega_s \bigcup \Omega_m$: Generalized Poisson-Boltzmann equation Ω_s : Generalized Kohn-Sham equation

Figure 10: Illustration of the quantum dynamics in continuum model for proton channels. Dynamics of other ions are modeled by classical Boltzmann approximation in bulk solvent Ω_s , while dynamics of protons is modeled in quantum mechanics but in a mean-field approach.

energy functional:

$$G_{\text{Total}}[\Phi, n, \{\rho_{\beta}\}] = \int \left\{ -\frac{\epsilon_m}{2} |\nabla \Phi|^2 + \Phi \rho_m - \frac{\epsilon_s(\mathbf{r})}{2} |\nabla \Phi|^2 + \Phi n(\mathbf{r})q - k_B T \sum_{\beta} \rho_{\beta}^0 \left(e^{-\frac{q_{\beta} \Phi + U_{\beta} - \mu_{\beta0}}{k_B T}} - 1 \right) \right. \\ \left. + \int \frac{\hbar^2 e^{-(E-\mu_p)/k_B T}}{2m(\mathbf{r})} |\nabla \Psi_E(\mathbf{r})|^2 dE + U_{\text{GC}}[n] + U_{\text{Ext}}[n] \right. \\ \left. + \lambda \left[\frac{N_p}{V_{\Omega}} - \int e^{-\frac{E-\mu_p}{k_B T}} |\Psi_E(\mathbf{r})|^2 dE \right] \right\} d\mathbf{r}.$$
(35)

This energy functional follows the same pattern of Eq. (2) or (29), but in this case proton is the major ion of interests and its concentration is denoted as $n(\mathbf{r})$. Other ions, with concentration ρ_{β} , are modeled by the equilibrium Boltzmann distribution as in Eq. (29).

Proton is described in a quantum mechanical formulation in terms of kinetic and potential energies in the third line of Eq. (35), where \hbar is the reduced Planck constant, and $m(\mathbf{r})$ is the effective mass of the proton. Under this framework, the kinetic energy is represented by the gradient of the proton wavefunction $\Psi_E(\mathbf{r})$. The potential energy of protons includes electrostatic energy $\Phi n(\mathbf{r})q$, generalized correlation $U_{\text{GC}}[n]$, and external $U_{\text{Ext}}[n]$, which are approximated as functionals of the proton concentration. The wavefunction and number density of protons have the following relation:

$$n(\mathbf{r}) = \int |\Psi_E(\mathbf{r})|^2 e^{-\frac{E-\mu_P}{k_B T}} dE,$$
(36)

where μ_p is the general electrochemical potential of protons.

The total energy functional (35) represents a multiphysical and multiscale framework that contains the continuum approximation for the solvent, while explicitly takes into account the channel protein in discrete atomic details. More importantly, it puts the classical theory of electrostatics and the quantum mechanical description of protons on an equal footing. Similar energy frameworks have been developed for nano-electronic devices in Chen & Wei (2010).

The governing equation for the electrostatic potential can be derived by the variation of energy functional (35) with respect to electrostatic potential Φ

$$\frac{\delta G_{\text{Total}}[\Phi, n]}{\delta \Phi} = 0 \Longrightarrow -\nabla \cdot (\epsilon \nabla \Phi(\mathbf{r})) - \sum_{\beta}^{N'_c} \rho_{\beta}^0 e^{-\frac{q_{\beta} \Phi(\mathbf{r}) - \mu_{\beta}}{k_B T}} = n(\mathbf{r})q + \rho_f(\mathbf{r}).$$
(37)

In the present multiphysics model, the proton number density $n(\mathbf{r})$ in Eq. (37) is related to the wavefunction $\Psi_E(\mathbf{r})$, which is governed by the generalized Kohn-Sham equation. This equation is obtained by the variation of the total free energy functional (35) with respect to wavefunction Ψ_E^{\dagger}

$$\frac{\delta G_{\text{Total}}[\Phi, n]}{\delta \Psi_E^*} = 0 \Longrightarrow -\nabla \cdot \frac{\hbar^2}{2m(\mathbf{r})} \nabla \Psi_E(\mathbf{r}) + V(\mathbf{r}) \Psi_E(\mathbf{r}) = E \Psi_E(\mathbf{r}), \tag{38}$$

where we set the Lagrange multiplier $\lambda = E$. The total Hamiltonian of the proton is given by

$$H = -\nabla \cdot \frac{\hbar^2}{2m(\mathbf{r})} \nabla + V(\mathbf{r}), \tag{39}$$

in which the total effective potential energy

$$V(\mathbf{r}) = q\Phi(\mathbf{r}) + V_{\rm GC}(\mathbf{r}) + V_{\rm Ext}(\mathbf{r})$$
(40)

consists of electrostatic, generalized correlation and external contributions. The external potential can be omitted for a closed system without external fields.

It is important to note that generalized Kohn-Sham equation (38) is fundamentally different from the normal Kohn-Sham equation for electronic structures. The Kohn-Sham operator in Eq. (38) has an absolutely continuous spectrum and invokes the Boltzmann statistics for proton scattering. Whereas the normal Kohn-Sham operator has a discrete spectrum and assumes the Fermi Dirac statistics for electron occupations (bound states). Eqs. (37) and (38) form the Poisson-Boltzmann-Kohn-Sham (PBKS) equations.

From the above quantities, the proton current can be defined by standard probability flux, whose practical expression is the following

$$I = \frac{q}{h} \operatorname{Tr} \int G(E) V_{\text{intra}}^{ah} G^{\dagger}(E) V_{\text{extra}}^{ah} \left[e^{-\frac{E-\mu_{\text{extra}}}{k_B T}} - e^{-\frac{E-\mu_{\text{intra}}}{k_B T}} \right] dE.$$
(41)

where Tr is the trace operation, G is the Green's operator

$$G(E) = (E - H)^{-1}, (42)$$

and μ_{extra} and μ_{intra} are the external electrical field energies at extracellular and intracellular electrodes, respectively. Here V_{extra}^{ah} and V_{intra}^{ah} are the anti-Hermitian components of the external potentials (Chen & Wei (2013)).

Figures 11 (a)-(b) display the electrostatic potential energy and the generalized correlations of protons in the GA channel, respectively. The electrostatic potential energy in this model is very similar to the profiles in the PNP or PBNP model, because the electrostatics is determined mainly by the molecular structure of the GA channel. The dielectric constant of water molecules in the channel region is taken as a model parameter and three values, $\epsilon_{ch} = 20$, $\epsilon_{ch} = 40$, and $\epsilon_{ch} = 80$ are used in Fig. 11 (a). Two electrostatic potential wells present near the entrance and exist of the channel, corresponding to the two binding sites. Fig. 11(b) shows the generalized correlation of protons, including protein-proton, proton-water, and proton-ion interactions,



Figure 11: Potential energy components of proton transport through the GA channel along the z-axis. (a) Electrostatic potential energy. (b) Generalized correlations

modeled by density functionals. In this case, generalized correlations are all energy barriers that protons need to overcome during the transport, and the sum of electrostatics and generalized correlation is the overall potential energy of protons.

Figure 12 displays simulated results of proton conductance through Eq. (41), compared with experimental data from Schumaker et al. (2000) of the GA channel. The blue dots in each figure represent available experimental observations for certain voltage biases, while the red curves are from the PBKS model predictions calculated with sufficiently many voltage samples. The model parameters are chosen to match the experimental data but all of the choices are taken within the range of physical measurements.



Figure 12: Voltage-current relation of proton translocation of GA at different concentrations. Blue dots: experimental data of Eisenman et al. (1980); Red solid curves: QDC model prediction.

VI.B Differential geometry (DG) based charge transport models

Like other biological processes, charge transport takes place in an aqueous environment because 65%-90% of cell mass is water. Thus, understanding the solvation process of channel proteins and mobile ions in solvent has equal importance as ionic dynamics. In terms of modeling, a solvation process may include the creation of a solute cavity in the solvent, the hydrogen bond breaking and formation at the solvent-solute interface, the surface reconstruction of the solute molecule, and/or the entropy effect due to solvent-solute mixing

(Mukhopadhyay et al. (2012), Onufriev et al. (2002)). Physically, this process involves a variety of solventsolute interactions, such as the electrostatic, dipolar, induced dipolar, and van der Waals interactions between the solvent and the solute (Baker et al. (2006), Baker & McCammon (2003), Ren et al. (2013), Rocchia et al. (2001, 2002)). In the past several years, intensive investigations for modeling ion channel systems based on the PNP theory but with solvation processes through multiscale, multiphysics variational approach were carried out, which were based on a differential geometry based multiscale paradigm for large chemical and biological systems, such as fuel cells, nanofluidics (Park et al. (2015)), ion channels, molecular motors, and viruses (Chen, Chen & Wei (2012), Chen & Wei (2012, 2013), Chen et al. (2010), Chen, Baker & Wei (2011), Chen & Wei (2011), Wei (2010, 2013), Wei et al. (2012)). With abundant water molecules and atomic details in cell membrane, it is critical to perform dimensionality reduction and manifold contraction by multiscale approaches. The essential ingredient is to use the differential geometry theory and analysis for surfaces and geometric measure as a natural technique to distinguish macroscopic domain for solvent and membrane, from the microscopic domain for channel proteins. At the same time, the differential geometry based model couples the continuum mechanical description of the aquatic environment with the discrete atomistic description of ion channels.

These differential geometry based multiscale models are intensively investigated and practically implemented for various types of ion channel problems (Chen, Chen & Wei (2012), Chen & Wei (2012), Chen et al. (2010), Chen, Baker & Wei (2011), Chen & Wei (2011)). There are two representations, the Eulerian formulation (Chen et al. (2010), Chen & Wei (2011), Chen, Zhao, Chun, Thomas, Baker, Bates & Wei (2012), Wang & Wei (2015)) and the Lagrangian formulation (Chen, Baker & Wei (2011)) for the key element in the models, the solvent-solute interface (Bates et al. (2009, 2008)). For the former, the interface is described as a hypersurface function which is evolved according to the derived governing equations (Chen et al. (2010)), while for the latter, interface elements are directly evolved according to governing equations which prescribe a set of rules. The Lagrangian representation of a molecular surface can be obtained from the projection of the hypersurface function by using an isosurface extraction procedure. The Eulerian formulation is mathematically simple and computationally robust, while the Lagrangian formalism is straightforward for force prescription (Bates et al. (2009)) and is computationally efficient, but usually encounters difficulties in handling the geometric breakup and/or surface merging. Validation and equivalence of these two formulations are tested by the solvation analysis for biological and chemical compounds (Chen et al. (2010), Chen, Baker & Wei (2011)). These models were shown to deliver excellent solvation predictions of experimental data (Chen, Zhao, Chun, Thomas, Baker, Bates & Wei (2012), Wang & Wei (2015)).

Variational solute-solvent interface The first ingredient of the DG model is the definition of molecular surface. Implicit solvent models require a given solvent-solute interface, or molecular surface, to distinguish different domains with the corresponding physical features, e.g., dielectric functions and diffusion constants, and to separate appropriate computational domains. In many models, simple *ad hoc* molecular surfaces, such as the van der Waals surface, the solvent excluded surface (Richards (1977)), or the solvent accessible surface are often utilized in applications of protein-protein interactions (Crowley & Golovin (2005)), protein folding (Spolar & Record Jr. (1994)), DNA binding and bending (Dragan et al. (2004)). In Wei et al. (2005), the first PDE-based approach to construct biomolecular surfaces via curvature driven geometric flows was introduced. Later on, the first variational formulation of molecular surfaces is developed, and the resulting molecular surface, called the minimal molecular surface (MMS), was constructed by the mean curvature flow (Bates et al. (2006, 2008)). Physically, the new definition of molecular surface satisfies the physical requirement of free energy minimization of the ion channel system. Computationally, it avoids artificially geometric defects such as cusps or self-intersecting surfaces, which could lead to computational instabilities.

To develop the variational solute-solvent interface, a solute characteristic function $S(\mathbf{r})$ is introduced in the total free energy. Fig. 13 offers a 1D representation, in which $S(\mathbf{r})$ takes values one and zero in the solute and solvent domains, respectively but with a transient region. Correspondingly, the solvent characteristic function, $1 - S(\mathbf{r})$, represents the solvent domain. With this setup, the energy in Eq. (2) can be rewritten as



Figure 13: Illustration of surface characteristic function S and solvent characteristic function 1-S in a 1D setting.

$$G_{\text{total}}^{\text{PNP}}[S, \Phi, \{\rho_{\alpha}\}] = \int \{\gamma |\nabla S| + pS + (1 - S)U + S\left[-\frac{\epsilon_{m}}{2}|\nabla \Phi|^{2} + \Phi \rho_{m}\right] + (1 - S)\left[-\frac{\epsilon_{s}}{2}|\nabla \Phi|^{2} + \Phi \sum_{\alpha}\rho_{\alpha}q_{\alpha}\right] + (1 - S)\sum_{\alpha}\left[\left(\mu_{\alpha}^{0} - \mu_{\alpha0}\right)\rho_{\alpha} + k_{B}T\rho_{\alpha}\ln\frac{\rho_{\alpha}}{\rho_{\alpha0}} - k_{B}T\left(\rho_{\alpha} - \rho_{\alpha0}\right) + \lambda_{\alpha}\rho_{\alpha}\right]\right\}d\mathbf{r}.$$
(43)

The first row of Eq. (43) lists the nonpolar solvation free energy of the system, with γ being the surface tension and p being the hydrodynamic pressure, respectively. The solvent-solute interaction U is originally approximated by Chen et al. (2010), Chen, Baker & Wei (2011), the Lennard-Jones potential, i.e.,

$$U = \sum_{j}^{Na} U_j^{LJ}(\mathbf{r}) \tag{44}$$

where Na atoms are assumed for the channel protein. The Weeks-Chandler-Andersen (WCA) decomposition based on the original WCA theory from Weeks et al. (1971) is utilized to split the Lennard-Jones potential into attractive and repulsive parts. The WCA potential was used to account for the attractive dispersion interaction Chen et al. (2010), Chen, Baker & Wei (2011), but can be extended as the generalized correlation in terms of ionic concentrations, i.e.,

$$U = \sum_{\alpha} \rho_{\alpha} U_{\alpha}.$$
 (45)

Details of the formation and explanation of the generalized correlation U_{α} is given in the later Section.

The second and third rows of Eq. (43) follows the same concepts in Eq. (2), except that the solvent and solute characteristic functions are associated correspondingly.

Generalized Laplace-Beltrami (LB) equation By applying the variational principle, the governing equation for the function $S(\mathbf{r})$ is

$$\frac{\delta G_{\text{total}}^{\text{PNP}}}{\delta S} \Rightarrow -\nabla \cdot \left(\gamma \frac{\nabla S}{|\nabla S|}\right) + p - U - \frac{\epsilon_m}{2} |\nabla \Phi|^2 + \Phi \rho_m + \frac{\epsilon_s}{2} |\nabla \Phi|^2 - \Phi \sum_{\alpha} \rho_{\alpha} q_{\alpha} - \sum_{\alpha} \left[-\mu_{\alpha 0} \rho_{\alpha} + k_B T \rho_{\alpha} \ln \frac{\rho_{\alpha}}{\rho_{\alpha 0}} - k_B T \left(\rho_{\alpha} - \rho_{\alpha 0}\right)\right] = 0,$$
(46)

where Eq. (7) is used in the derivation. It is easier to pursue the solution of Eq. (46) by the following parabolic equation with an artificial time (Bates et al. (2009), Wei (2010), Zhao (2011)):

$$\frac{\partial S}{\partial t} = |\nabla S| \left[\nabla \cdot \left(\gamma \frac{\nabla S}{|\nabla S|} \right) + V_1 \right], \tag{47}$$

where the LB potential V_1 is

$$V_{1} = -p + U + \frac{\epsilon_{m}}{2} |\nabla \Phi|^{2} - \Phi \rho_{m} - \frac{\epsilon_{s}}{2} |\nabla \Phi|^{2} + \Phi \sum_{\alpha} \rho_{\alpha} q_{\alpha}$$

$$+ \sum_{\alpha} \left[k_{B}T \left(\rho_{\alpha} \ln \frac{\rho_{\alpha}}{\rho_{\alpha 0}} - \rho_{\alpha} + \rho_{\alpha 0} \right) - \mu_{\alpha 0} \rho_{\alpha} \right].$$

$$(48)$$

Figures 14 (a)-(b) show the molecular surface generated from the generalized LB equation (47). For comparison, the solvent exclusive surface (SES) of the GA generated via the MSMS package developed by Sanner et al. (1996) is also displayed in Figs. 14 (c)-(d). The SES only depends on the parameter used in the generating software package (water probe radius 1.4 Å and density 10). Once generated, it keeps fixed and is independent of the physiological conditions such as ion concentration or transmembrane voltage differences in simulations. On the other hand, generation of the surface from the generalized LB equation is an iteration process as stated in the earlier sections. As included in the LB potential $V_1(\Phi, \{\rho_\alpha\})$, evaluation of the characteristic functions $S(\mathbf{r})$ depends on the electrostatics and ionic concentration. The results showed in Figs. 14(a) and (b) are calculated under the transmembrane potential of 0.2 mV and with the ionic concentration of 0.1M.

Comparing Figs. 14(a) and (b) and Figs. 14(c) and (d), the SES commits geometric singularities, such as cusps and self-intersecting surfaces Zheng et al. (2011). These singularities may bring computational difficulties to the designed interface schemes and are unphysical in the solvent-solute interface. In contrast, the surface from the LB equation has a smoother appearance because of the diffusion mechanism, which gives less-intensive sharp changes near the solvent-solute boundary. More importantly, generations of the new molecular surface incorporate the interactions with the external ionic condition as well as transmembrane voltages. All of these characteristics make more physical sense, and many good results in applications are obtained in Chen et al. (2010), Chen, Baker & Wei (2011).

Generalized Poisson-Nernst-Planck equation Under this differential geometry based model, the generalized Poisson-Nernst-Planck equation can be derived by taking the variation with respect to the electrostatic potential Φ , i.e,

$$-\nabla \cdot (\epsilon(S)\nabla\Phi) = S\rho_m + (1-S)\sum_{\alpha} \rho_{\alpha}q_{\alpha},$$
(49)

where $\epsilon(S) = (1 - S)\epsilon_s + S\epsilon_m$ is an interface-dependent dielectric function. Eq. (49) depends on ion concentration ρ_{α} and the solute characteristic function S.

$$\frac{\partial \rho_{\alpha}}{\partial t} = \nabla \cdot \left[D_{\alpha} \left(\nabla \rho_{\alpha} + \frac{\rho_{\alpha}}{k_B T} \nabla (q_{\alpha} \Phi + U_{\alpha}) \right) \right], \tag{50}$$

This is the generalized Nernst-Planck equation where $q_{\alpha}\Phi + U_{\alpha}$ is a mean-field approach of all potentials. Equation (50) reduces to the standard Nernst-Planck equation in Eq. (9) when the solvent-solute interactions are taken as the original definition in Eq. (44). The differential geometry (DG) based model can also be applied to the PBNP and PBKS systems similarly.

Generalized correlation Many improvements for the PNP system, such as the size effects or ion-water interactions discussed earlier, can be adopted in the differential geometry based model, termed as generalized correlation, which are modeled in a mean-field approach.

In this approach, the generalized correlation is modeled by extending the term

$$\int (1-S)Ud\mathbf{r} \tag{51}$$



Figure 14: Surface representations of the GA channel. (a)-(b) Surface extracted from the generalized LB equation with S = 0.5. (c)-(d) MSMS surface with probe radius 1.4 and density 10.

in Eq. (43) as the free energy functional of the local ionic concentration ρ_{α} and its gradient $\nabla \rho_{\alpha}$ for all ions, i.e.,

$$G_{\rm GC}[S, \{\rho_{\alpha}\}] = \int (1 - S(\mathbf{r})) \sum_{\alpha} U_{\rm GC}[\{\rho_{\alpha}\}, \{\nabla\rho_{\alpha}\}] d\mathbf{r}.$$
(52)

With the assumption that the $\nabla \rho_{\alpha}$ dependence is omitted as a first order approximation, it takes

$$G_{\rm GC}[S, \{\rho_{\alpha}\}] = \sum_{\beta} \left(\frac{1}{2}\right)^{\delta_{\alpha\beta}} \int \int (1 - S(\mathbf{r}))\rho_{\alpha}(\mathbf{r})\rho_{\beta}(\mathbf{r}')K_{\beta}(|\mathbf{r} - \mathbf{r}'|)d\mathbf{r}' d\mathbf{r}$$
(53)

+
$$\int \int (1 - S(\mathbf{r}))\rho_{\alpha}(\mathbf{r})n_{w}(\mathbf{r}')K_{w}(|\mathbf{r} - \mathbf{r}'|)d\mathbf{r}'d\mathbf{r}$$
(54)

+
$$\sum_{j=1}^{N_a} \int (1 - S(\mathbf{r})) \rho_\alpha(\mathbf{r}) n_j(\mathbf{r}') K_j(|\mathbf{r} - \mathbf{r}'|) d\mathbf{r},$$
 (55)

where $\delta_{\alpha\beta}$ is the Kronecker delta function with $\delta_{\alpha\beta} = 1$ if $\alpha = \beta$ and $\delta_{\alpha\beta} = 0$ when $\alpha \neq \beta$. The kernels $K_{\beta}(|\mathbf{r} - \mathbf{r}'|)$, $K_w(|\mathbf{r} - \mathbf{r}'|)$, and $K_i(|\mathbf{r} - \mathbf{r}'|)$ model interactions among the α -th and β -th ions, water molecules (in sense of density $n_w(\mathbf{r})$), and protein atom distribution $n_i(\mathbf{r}')$, respectively.

Finally, by taking variation with respect to the concentration ρ_{α} , one obtains the generalize correlation potential

$$U_{\alpha}(\mathbf{r}) = \sum_{\beta} \rho_{\beta}(\mathbf{r}) * K_{\beta}(\mathbf{r}) + n_{w}(\mathbf{r}) * K_{w}(\mathbf{r}) + \sum_{j=1}^{N_{a}} n_{j}(\mathbf{r}) * K_{j}(\mathbf{r}),$$
(56)

where * represents the convolution operation.

VII Concluding remarks

Phenomena of charge transport present not only in naturally-designed devices such as ion channels, deoxyribonucleic acid (DNA) nanowires, ATPases, or neuron synapses, but also widely exist in human-made devices including solar cells, fuel cells, battery cells, molecular switches, nanotubes, field effect transistors, nanofibers, thin films, etc (Wei et al. (2012)). Thus, understanding of mechanism and dynamics of charge transport in these nano-scale biological or industrial devices is prerequisite to study their functions for human health or developing modern technologies. Quantitative modeling and simulation of charge transport have emerged as a new field in applied mathematics, which offers mathematical models, algorithms and analysis to reproduce experimental data, to predict new phenomena and hence to shed light on new directions of research for these nano-bio systems. Due to complexity of charge transport increases dramatically at nanoscale, where a large amount of components, both macroscopic and microscopic, interact with each other in multiphysical principles in a heterogeneous environment, Mathematical modeling and computation of these complex systems encounter formidable challenges. Fully atomic description for the whole system will offer the highest accuracy, but it is prohibited by the intractable large number of degree of freedom. Multiscale and multiphysics modeling that retains the atomistic description of the channel protein while treats the membrane and solvent as dielectric continuum gives rise to efficient approaches to ion channel charge transport.

In this paper we review recent progresses in mathematical modeling, algorithm and analysis of charge transport in ion channels. A major emphasis is the development in Poisson-Nernst-Planck (PNP) based models. The PNP equations can be derived from the total energy functional of an ion channel system and include a Poisson equation for electrostatic environment for the whole system and a series of Nernst-Planck equations for dynamics of mobile ions. Modified PNP models were developed under this framework in order to include more detailed physical properties such as ion size effects or ion-water interactions, or to reduce model complexity in multi-ion species systems. All the modification and improvement of PNP equations can be achieved by adding additional energy components in the total energy functional. On the other side, proton transport is a special type of charge transport due to the properties of hydrogen ions, and thus quantum dynamics is involved in modeling proton channels. There are quite a few literature about quantum models for proton transport in the literature, but it is the first time that a quantum dynamics in continuum model was established as a mean-field method. Finally, a differential geometry (DG) based multiscale model was developed, which includes the solvation process of ion channels in a solvent. In the multiscale treatment of the ion channel system, the DGPNP model generates a more reliable and robust definition of molecular surface, which takes into account the mutual interactions with electrostatics and ionic dynamics. Various computational algorithms and mathematical analysis are also reviewed in this work.

Although numerous efforts and work have been devoted in this area each year worldwide, including the availability of molecular structures of new ion channels, resolving a complete picture of molecular mechanism

of channel gating and charge transport remains a challenging task. On the one hand, structure determination of ion channel proteins and in general, G-protein-coupled receptors (GPCRs), is typically more difficult than that of global proteins. On the other hand, even with available structures, biophysical understanding of functioning principles for many existing ion channels are not completely clear. In particular, the gating mechanism of voltage gated eukaryotic sodium channels is still elusive.

Future mathematical modeling and simulation of ion channels will address the pressing needs in the understanding of their molecular mechanism. Much attention will be paid to the homology modeling of ion channel structures, such as the structure presented in Fig. 1 and their improvement. Additionally, the interactions of ion channels and drugs, which hold the key for the drug discovery for epilepsy, irregular cardiac arrhythmias, hyperalgesia, myotonia, and anesthesia, will be studied by molecular docking. An important component in docking analysis is the prediction of protein-ligand/drug binding affinities, which can be achieved by machine learning algorithms Wang, Zhao & Wei (submitted 2016). Moreover, the blind prediction of mutation impacts to ion channel current-voltage (I-V) curves (Cang & Wei (2017b)) and drug resistance will be a new topic in mathematical molecular bioscience and biophysics (Wei (2016)). In such predictions, machine learning methods, including deep learning, can be empowered by using mathematical features from differential geometry, algebraic topology, graph theory and partial differential equations (Cang & Wei (2017a), Wang, Zhao & Wei (submitted 2016), Xia & Wei (submitted 2016)). Finally, the PNP type of models will be coupled with molecular mechanism and chemical kinetics to address the conformational changes and protonation (or deprotonation) during the ion permeation. Given the importance of the physical and chemical phenomena of charge transport to both biological systems and nano-device engineering, theoretical modeling, numerical algorithms, mathematical analysis, and realistic applications of ion channel charge transport will be a focus of mathematical research in the future.

Acknowledgments

The work of GWW was supported in part by NSF Grant IIS- 1302285 and MSU Center for Mathematical Molecular Biosciences Initiative. The work of the author DC is funded by the Faculty Research Grant 2015-2017 provided by the University of North Carolina at Charlotte.

Change History

This article is initially submitted by Duan Chen and Guowei Wei. No change has been made so far.

References

- Abaid, N., Eisenberg, B. S. & Liu, W. S. (2008), 'Asymptotic expansions of I-V relations via a Poisson-Nernst-Planck system', *SIAM Journal on Applied Dynamical Systems* **7**(4), 1507–1526.
- Adalsteinsson, H., Debusschere, B. J., Long, K. R. & Najm, H. N. (2008), 'Components for atomistic-tocontinuum multiscale modeling of flow in micro- and nanofluidic systems', *Scient. Program.* **16**, 297–313.
- Alexov, E. & Gunner, M. R. (1999), 'Calculated Protein and Proton Motions Coupled to Electron Transfer: Electron Transfer from Q_A^- to Q_B in Bacterial Photosynthetic Reaction Centers', *Biochemistry* **38**, 8253–8270.
- Allen, R., Hansen, J.-P. & Melchionna, S. (2001), 'Electrostatic potential inside ionic solutions confined by dielectrics: a variational approach', *Phys Chem Chem Physics* **3**, 4177–4186.
- Antypov, D., Barbosa, M. C. & Holm, C. (2005), 'Incorporation of excluded-volume correlations into poissonboltzmann theory', *Physical Review E* 71, 061106.
- Bagneris, C., DeCaen, P. G., Hall, B. A., Naylor, C. E., Clapham, D. E., Kay, C. W. M. & Wallace, B. A. (2013), 'Role of the C-terminal domain in the structure and function of tetrameric sodium channels', *Nature Communications* 4, 2465.
- Baker, K., Chen, D. & Cai, W. (2016), 'Investigating the Selectivity of KcsA Channel by an Image Charge Solvation Method (ICSM) in Molecular Dynamics Simulations.', *Communications in Computational Physics* 19, 927–943.
- Baker, N. A., Bashford, D. & Case, D. A. (2006), Implicit solvent electrostatics in biomolecular simulation, in B. Leimkuhler, C. Chipot, R. Elber, A. Laaksonen, A. Mark, T. Schlick, C. Schutte & R. Skeel, eds, 'New Algorithms for Macromolecular Simulation', Springer.
- Baker, N. A. & McCammon, J. A. (2003), Electrostatic interactions, in P. Bourne & H. Weissig, eds, 'Structural Bioinformatics', John Wiley & Sons, Inc., New York, pp. 427–440.
- Baker, N. A., Sept, D., Holst, M. J. & Mccammon, J. A. (2001), 'The adaptive multilevel finite element solution of the Poisson-Boltzmann equation on massively parallel computers', *IBM Journal of Research and Development* 45(3-4), 427–438. URL: http://www.research.ibm.com/journal/rd/453/baker.html
- Barcilon, V., Chen, D. & Eisenberg, B. S. (1992), 'Ion flow through narrow membrane channels: Part ii', SIAM J. Appl. Math 52, 1405–1425.
- Bates, P. W., Chen, Z., Sun, Y. H., Wei, G. W. & Zhao, S. (2009), 'Geometric and potential driving formation and evolution of biomolecular surfaces', *J. Math. Biol.* **59**, 193–231.
- Bates, P. W., Wei, G. W. & Zhao, S. (2006), 'The minimal molecular surface', arXiv:q-bio/0610038v1 [q-bio.BM].
- Bates, P. W., Wei, G. W. & Zhao, S. (2008), 'Minimal molecular surfaces and their applications', Journal of Computational Chemistry 29(3), 380–91.
- Bazant, M. Z., Storey, B. D. & Kornyshev, A. A. (2011), 'Double layer in ionic liquids: Overscreening versus crowding', *Physical Review Letters* 106, 046102.
- Bazant, M. Z., Thornton, K. & Ajdari, A. (2004), 'Diffuse-charge dynamics in electrochemical systems', *Physical Review E* 70, 021506.
- Ben-Yaakov, D., Andelman, D., Podgornik, R. & Podgornik, R. (2011), 'Ion-specific hydration effects: Extending the Poisson-Boltzmann theory', Current Opinion in Colloid and Interface Science 16, 542–550.

- Bernal, J. & Fowler, R. (1933), 'A theory of water and ionic solution, with particular reference to hydrogen and hydroxyl ions', *J Chem Phys* 1, 515–548.
- Bertonati, C., Honig, B. & Alexov, E. (2007), 'Poisson-Boltzmann calculations of nonspecific salt effects on protein-protein binding free energy', *Biophysical Journal* 92, 1891–1899.
- Boiteuxa, C., Vorobyovb, I., Frenchc, R. J., Frenchd, C., Yarov-Yarovoye, V. & Allena, T. W. (2014), 'Local anesthetic and antiepileptic drug access and binding to a bacterial voltage-gated sodium channel', *Proc. Natl. Acad. Sci. USA* 36, 13057–13062.
- Bothma, J., Gilmore, J. & McKenzie, R. (2010), 'The role of quantum effects in proton transfer reactions in enzymes: quantum tunneling in a noisy environment?', New Journal of Physics 12(055002).
- Brohawn, S., Campbell, E. & MacKinnon, R. (2014), 'Physical mechanism for gating and mechanosensitivity of the human TRAAK K⁺ channel.', *Nature* **516**, 126–30.
- Burger, M., Schlake, B. & Wolfram, M. T. (2012), 'Nonlinear poisson-nernst-planck equations for ion flux through confined geometries', *Nonlinearity* 25, 961–990.
- Cang, Z. X. & Wei, G. W. (2017*a*), 'Feature functional theory-mutation predictor (FFT-MP) for the blind prediction of protein mutation energy changes', *Submitted*.
- Cang, Z. X. & Wei, G. W. (2017b), 'Topological fingerprints reveal protein-ligand binding mechanism ', Submitted .
- Cardenas, A. E., Coalson, R. D. & Kurnikova, M. G. (2000), 'Three-dimensional Poisson-Nernst-Planck theory studies: Influence of membrane electrostatics on Gramicidin A channel conductance', *Biophysical Journal* **79**, 80–93.
- Cartailler, J., Schuss, Z. & Holcman, D. (2017), 'Analysis of the Poisson–Nernst–Planck equation in a ball for modeling the Voltage–Current relation in neurobiological microdomains', *Physica D: Nonlinear Phenomena* 339, 39–48.
- Catterall, W. A. (2012), 'Voltage-gated sodium channels at 60: structure, function and pathophysiology', J. *Physiol.* **590**, 2577–2589.
- Chandra, A. (2000), 'Static dielectric constant of aqueous electrolyte solutions: Is there any dynamic contribution?', *Journal of Chemical Physics* **113**, 903–905.
- Chaudhry, J. H., Comer, J., Aksimentiev, A. & Olson, L. N. (2014), 'A Stabilized Finite Element Method for Modified Poisson-Nernst-Planck Equations to Determine Ion Flow Through a Nanopore', *Communications* in Computational Physics 15, 93–125.
- Chen, D. (2016), 'A New Poisson-Nernst-Planck Model with Ion-Water Interactions for Charge Transport in Ion Channels', *Bulletin of Mathematical Biology* p. accepted.
- Chen, D., Chen, Z., Chen, C., Geng, W. H. & Wei, G. W. (2011), 'MIBPB: A software package for electrostatic analysis', J. Comput. Chem. 32, 657 670.
- Chen, D., Chen, Z. & Wei, G. W. (2012), 'Quantum dynamics in continuum for proton transport II: Variational solvent-solute intersurface', *International Journal for Numerical Methods in Biomedical Engineering* 28, 25– 51.
- Chen, D., Lear, J. & Eisenberg, B. S. (1997), 'Permeation through an open channel: Poisson-Nernst-Planck theory of a synthetic ionic channel.', *Biophys J.* **72**(1), 97–116.
- Chen, D. P., Eisenberg, R. S., Jerome, J. W. & Shu, C. W. (1995), 'Hydrodynamic model of temperature change in open ionic channels', *Biophys. J.* 69, 2304 2322.

- Chen, D. & Wei, G. W. (2010), 'Modeling and simulation of electronic structure, material interface and random doping in nano-electronic devices', J. Comput. Phys. 229, 4431–4460.
- Chen, D. & Wei, G. W. (2012), 'Quantum dynamics in continuum for proton transport III: Generalized correlation', *J Chem. Phys.* **136**, 134109.
- Chen, D. & Wei, G. W. (2013), 'Quantum dynamics in continuum for proton transport I: Basic formulation', *Commun. Comput. Phys.* **13**, 285–324.
- Chen, Z., Baker, N. A. & Wei, G. W. (2010), 'Differential geometry based solvation models I: Eulerian formulation', J. Comput. Phys. 229, 8231-8258.
- Chen, Z., Baker, N. A. & Wei, G. W. (2011), 'Differential geometry based solvation models II: Lagrangian formulation', *J. Math. Biol.* **63**, 1139–1200.
- Chen, Z. & Wei, G. W. (2011), 'Differential geometry based solvation models III: Quantum formulation', J. Chem. Phys. 135, 194108.
- Chen, Z., Zhao, S., Chun, J., Thomas, D. G., Baker, N. A., Bates, P. B. & Wei, G. W. (2012), 'Variational approach for nonpolar solvation analysis', *Journal of Chemical Physics* **137**(084101).
- Cheng, M. H. & Coalson, R. D. (2005), 'An accurate and efficient empirical approach for calculating the dielectric self-energy and ion-ion pair potential in continuum models of biological ion channels', J Phys Chem B 109(1), 488–98.
- Cheng, M. H., Coalson, R. D. & Tang, P. (2010), 'Molecular dynamics and brownian dynamics investigation of ion permeation and anesthetic halothane effects on a proton-gated ion channel', J Am Chem Soc 132(46), 16442–9.
- Choudhary, O. P., Ujwal, R., Kowallis, W., Coalson, R., Abramson, J. & Grabe, M. (2010), 'The electrostatics of VDAC: implications for selectivity and gating', *J Mol Biol* **396(3)**, 580–92.
- Chung, S.-H. & Kuyucak, S. (2002), 'Recent advances in ion channel research', *Biochimica et Biophysica Acta* **1565**, 267–286.
- Coalson, R. D. & Kurnikova, M. G. (2005), 'Poisson-Nernst-Planck theory approach to the calculation of current through biological ion channels', *IEEE Trans Nanobioscience* **4(1)**, 81–93.
- Cole, C. D., Frost, A. S., Thompson, N., Cotten, M., Cross, T. A. & Busath, D. D. (2002), 'Noncontact dipole effects on channel permeation. vi. 5f- and 6F-Trp Gramicidin channel currents', *Biophysical Journal* 83, 1974–1986.
- Constantin, D. & Siwy, Z. S. (2007), 'Poisson-Nernst -Planck model of ion current rectification through a nanofluidic diode', *Phys. Rev. E* 76, 041202.
- Corry, B., Kuyucak, S. & Chung, S.-H. (2003), 'Dielectric self-energy in Poisson-Boltzmann and Poisson-Nernst-Planck models of ion channels', *Biophysical Journal* 84(6), 3594–3606.
- Cross, T. A., Dong, H., Sharma, M., Busath, D. D. & Zhou, H.-X. (2012), 'M2 protein from influenza A: from multiple structures to biophysical and functional insights', *Curr. Opin. Virol.* **2**, 128–133.
- Crowley, P. & Golovin, A. (2005), 'Cation-pi interactions in protein-protein interfaces', *Proteins Struct. Func. Bioinf.* **59**, 231–239.
- Cukier, R. I. (2004*a*), 'Quantum molecular dynamics simulation of proton transfer in cytochrome c oxidase', *Biochimica Et Biophysica Acta-Bioenergetics* **1656**, 189 – 202.
- Cukier, R. I. (2004*b*), 'Theory and simulation of proton-coupled electron transfer, hydrogen-atom transfer, and proton translocation in proteins', *Biochimica Et Biophysica Acta-Bioenergetics* **1655**, 37–44.

- Dai, J., Wollmuth, Lonnie, P. & Zhou, H.-X. (2015), 'Mechanism-Based Mathematical Model for Gating of Ionotropic Glutamate Receptors.', J. Phys. Chem. B 119, 10934–10940.
- Dai, J. & Zhou, H.-X. (2014), 'General rules for the arrangements and gating motions of pore-lining helices in homomeric ion channels.', *Nat. Commun.* **5**, 4641.
- de Falco, C., Jerome, J. W. & Sacco, R. (1964), 'A self-consitent iterative scheme for the one-dimensional steady-state transistor calculations', *IEEE Trans. Ele. Dev.* **11**, 455–465.
- Decornez, H., Drukker, K. & Hammes-Schiffer, H. (1999), 'Solvation and hydrogen-bonding effects on proton wires', J. Phys. Chem. A 103, 2891–2898.
- Decoursey, T. (2003), 'Voltage-gated proton channels and other proton transfer pathways', *Physiol Rev* **83**, 475–579.
- Deng, C. & Li, C. (2013), 'Endpoint bilinear estimates and applications to the two-dimensional Poisson– Nernst–Planck system', Nonlinearity 26(11), 2993.
- Dolinsky, T. J., Czodrowski, P., Li, H., Nielsen, J. E., Jensen, J. H., Klebe, G. & Baker, N. A. (2007), 'Pdb2pqr: Expanding and upgrading automated preparation of biomolecular structures for molecular simulations', *Nucleic Acids Res* 35, W522–525.
- Dolinsky, T. J., Nielsen, J. E., McCammon, J. A. & Baker, N. A. (2004), 'Pdb2pqr: an automated pipeline for the setup, execution, and analysis of Poisson-Boltzmann electrostatics calculations', *Nucleic Acids Res* 32, W665–W667.
- Dong, H., Yi, M., Cross, T. & Zhou, H.-X. (2013), 'Ab initio calculations and validation of the pH-dependent structures of the His37-Trp41 quartet, the heart of acid activation and proton conductance in the M2 protein of Influenza A virus', *Chem. Sci.* **4**, 2776–2787.
- Dong, K., Du, Y., Rinkevich, F. D., Nomura, Y., Xu, P., Wang, L., Silver, K., & Zhorov, B. S. (2014), 'Molecular biology of insect sodium channels and pyrethroid resistance', *Insect Biochem.* **50**, 1–17.
- Dragan, A., Read, C., Makeyeva, E., Milgotina, E., Churchill, M., Crane-Robinson, C. & Privalov, P. (2004), 'Dna binding and bending by hmg boxes: Energetic determinants of specificity', *J. Mol. Biol.* **343**, 371–393.
- Drukker, K., de Leeuw, S. H. & Hammes-Schiffer, H. (1998), 'Proton transport along water chains in an electric field', J. Chem. Phys. 108, 6799 –6808.
- Du, Y., Nomura, Y., Satar, G., Hu, Z., Nauen, R., He, S. Y., Zhorov, B. S. & Dong, K. (2013), 'Molecular evidence for dual pyrethroid-receptor sites on a mosquito sodium channel', *Proc. Nat. Acad. Sci.* 110, 11785– 11790.
- Du, Y., Nomura, Y., Zhorov, B. S. & Dong, K. (in press 2015), 'Rotational Symmetry of two pyrethroid receptor sites in the mosquito sodium channel', *Mol. Pharmacol.* .
- Dyrka, W., Augousti, A. T. & Kotulska, M. (2008), 'Ion flux through membrane channels: An enhanced algorithm for the Poisson-Nernst-Planck model', *J. Comput Chem* **29**, 1876–1888.
- Egwolf, B. & Roux, B. (2010), 'Ion selectivity of the kcsa channel: A perspective from multi-ion free energy landscapes', J. Chem. Phys. 401, 831–842.
- Eisenberg, B. S. (1996), 'Computing the field in proteins and channels', J. Membrane Biol. 150, 1-25.
- Eisenberg, B. S. & Chen, D. (1993), 'Poisson-Nernst-Planck (PNP) theory of an open ionic channel', *Bio-physical Journal* **64**, A22.
- Eisenberg, B. S., Hyon, Y. K. & Liu, C. (2010), 'Energy variational analysis of ions in water and channels: Field theory for primitive models of complex ionic fluids', *Journal of Chemical Physics* **133**, 104104.

- Eisenberg, B. S. & Liu, W. S. (2006), 'Poisson-Nernst-Planck systems for ion channels with permanent charges', SIAM Journal on Applied Dynamical Systems **38**(6), 1932–1966.
- Eisenman, G., Enos, B., Hagglund, J. & Sandbloom, J. (1980), 'Gramicidin as an example of a single-filing ion channel', Ann. N.Y. Acad. Sci. 339, 8–20.
- Engels, M., Gerwert, K. & Bashford, D. (1995), 'Computational studies on bacteriorhodopsin: Conformation and proton transfer energetics', *Biophys. Chem.* 56, 95.
- Fermini, B. & Priest, B. T. (2008), Ion Channels, Springer-Verlag.
- Flavell, A., Machen, M., Eisenberg, B., Kabre, J., Liu, C. & Li, X. (2014), 'A Conservative Finite Difference Scheme for Poisson-Nernst-Planck Equations', *Journal of Computational Electronics* 13, 235–249.
- Fogolari, F. & Briggs, J. M. (1997), 'On the variational approach to Poisson-Boltzmann free energies', Chemical Physics Letters 281, 135–139.
- Fu, Y., Westenbroek, R., Scheuer, T. & Catterall, W. (2014), 'Basal and β-adrenergic regulation of the cardiac calcium channel Ca_V1.2 requires phosphorylation of serine 1700.', *Proc Natl Acad Sci USA* 111, 16598– 16603.
- Geng, W. H. (2013), 'Parallel higher-order boundary integral electrostatics computation on molecular surfaces with curved triangulation', *J. Comput. Phys.* **241**, 253 –265.
- Geng, W. H. & Jacob, F. (2013), 'A gpu-accelerated direct-sum boundary integral poisson-boltzmann solver', Comput. Phys. Commun. 184, 1490 –1496.
- Geng, W. H. & Krasny, R. (2013), 'A treecode-accelerated boundary integral Poisson-Boltzmann solver for continuum electrostatics of solvated biomolecules', J. Comput. Phys. 247, 62–87.
- Geng, W., Yu, S. & Wei, G. W. (2007), 'Treatment of charge singularities in implicit solvent models', *Journal* of Chemical Physics **127**, 114106.
- Gillespie, D., Nonner, W. & Eisenberg, B. S. (2002), 'Coupling poisson-nernst-planck and density functional theory to calculate ion flux', *Journal of Physics: Condensed Matter* **14**(46), 12129–12145.
- Gordon, D., Krishnamurthy, V. & Chung, S. (2009), 'Generalized langevin models of molecular dynamics simulations with applications to ion channels', *J. Chem. Phys.* **131**(134102).
- Graf, P., Kurnikova, M. G., Coalson, R. D. & Nitzan, A. (2004), 'Comparison of dynamic lattice Monte Carlo simulations and the dielectric self-energy Poisson-Nernst-Planck continuum theory for model ion channels', J. Phys. Chem. B 108, 2006–2015.
- Gutman, G. A., Chandy, K. G., Grissmer, S., Lazdunski, M., McKinnon, D., Pardo, L. A., Robertson, G. A., Rudy, B., Sanguinetti, M. C., Stuhmer, W. & Wang, X. (2005), 'International Union of Pharmacology. LIII. Nomenclature and molecular relationships of voltage-gated potassium channels', *Pharmacol Rev* 57, 473– 508.
- Hadjadj, L., Hamdache, K. & Hamroun, D. (2010), 'Solutions to a nonlinear Poisson-Nernst-Planck system in an ionic channel', *Mathematical Methods in the Applied Sciences* **33**(15), 1794–1806.
- Harguindey, S., Arranz, J., Wahl, M., Orives, G. & Reshkin, S. (2009), 'Proton Transport Inhibitors as Potentially Selective Anticancer Drugs', *Anticancer Research* 29, 2127–2136.
- He, D. & Pan, K. (2016), 'An energy preserving finite difference scheme for the Poisson-Nernst-Planck system', *Applied Mathematics and Computation* **287**, 214–223.
- Heymann, G., Dai, J., Li, M., Silberberg, S. D., Zhou, H.-X. & Swartz, K. J. (2013), 'Inter- and intrasubunit interactions between transmembrane helices in the open state of P2X receptor channels.', *Proc. Natl. Acad. Sci. USA* **110**, E4045–E4054.

- Hineman, J. L. & Ryham, R. J. (2015), 'Very weak solutions for Poisson-Nernst-Planck system', *Nonlinear Analysis: Theory, Methods & Applications* **115**, 12–24.
- Hodgkin, A. L. & Huxley, A. F. (1952), 'A quantitative description of membrane current and its application to conduction and excitation in nerve.', *The Journal of Physiology* **117**, 500–544.
- Hondares, E., Brown, M. A., Musset, B., Morgan, D., Cherny, V. V., Taubert, C., Bhamrah, M. K., Coe, D., Marelli-Berg, F., Gribben, J., Dyer, M. J., DeCoursey, T. E. & Capasso, M. (2014), 'Enhanced activation of an amino-terminally truncated isoform of the voltage-gated proton channel HVCN1 enriched in malignant B cells.', Proc Natl Acad Sci USA. 111, 18078–83.
- Hu, L. & Wei, G. W. (2012), 'Nonlinear poisson equation for heterogeneous media', Biophys. J. .
- Hwang, H., Schatz, G. C. & Ratner, M. A. (2006), 'Ion current calculations based on three dimensional Poisson-Nernst-Planck theory for a cyclic peptide nanotube', *Journal of Physical Chemistry B* **110**, 6999– 7008.
- Hyon, Y., Eisenberg, B. & Liu, C. (2010), 'A mathematical model of the hard sphere repulsion in ionic solutions', *Communications in Mathematical Sciences* **9**, 459–475.
- Hyon, Y., Eisenberg, B. S. & Liu, C. (2011), 'A mathematical model for the hard sphere repulsion in ionic solution', *Commun. Math. Sci.* 9, 459 475.
- Ikezu, T. & Gendelman, H. E. (2008), Neuroimmune Pharmacology, Springer.
- Im, W., Beglov, D. & Roux, B. (1998), 'Continuum solvation model: electrostatic forces from numerical solutions to the Poisson-Boltzmann equation', *Computer Physics Communications* 111(1-3), 59–75. URL: http://dx.doi.org/10.1016/S0010-4655(98)00016-2
- Im, W. & Roux, B. (2002), 'Ion permeation and selectivity of ompf porin:a theoretical study based on molecular dynamics, Brownian dynamics, and continuum electrodiffusion theory', J. Mol. Biol. 322, 851–869.
- Jensen, M., Jogini, V., Borhani, D., Leffler, A., Dror, R. & Shaw, D. (2012a), 'Mechanism of Voltage Gating in Potassium Channels.', Science 336, 229–233.
- Jensen, M. O., Jogini, V., Borhani, D. W., Leffler, A. E., Dror, R. O. & Shaw, D. E. (2012b), 'Mechanism of Voltage Gating in Potassium Channels', *Science* **336**, 229–232.
- Jerome, J. & Kerkhoven, T. (1990), 'A Finite Element Approximation Theory for the Drift Diffusion Semiconductor Model', SIAM J. Numer. Anal 28, 403–422.
- Ji, S., Liu, W. & Zhang, M. (2015), 'Effects of (Small) Permanent Charge and Channel Geometry on Ionic Flows via Classical Poisson–Nernst–Planck Models', SIAM Journal on Applied Mathematics 75(1), 114–135.
- Jia, Y., Liu, W. & Zhang, M. (2016), 'Qualitative properties of ionic flows via Poisson-Nernst-Planck systems with Bikerman's local hard-sphere potential: Ion size effects', *Discrete and Continuous Dynamical Systemsseries B* 21, 1775–1802.
- Jung, Y. W., Lu, B. Z. & Mascagni, M. (2009), 'A computational study of ion conductance in the KcsA K+ channel using a Nernst-Planck model with explicit resident ions', *J. Chem. Phys.* **131**(215101).
- Karniadakis, G., Beskök, A. & Aluru, N. R. (2005), *Microflows and nanoflows : fundamentals and simulation*, Springer.
- Kilic, M. S., Bazant, M. Z. & Ajdari, A. (2007), 'Steric effects in the dynamics of electrolytes at large applied voltages. II. modified poisson-nernst-planck equations', *Phys. Rev. E* **75**(021503).
- Kirkwood, J. G. (1934), 'Theory of solution of molecules containing widely separated charges with special application to zwitterions', J. Comput. Phys. 7, 351 361.

- Kiselev, Y. V., Leda, M., Lobanov, A. I., Marenduzzo, D. & Goryachev, A. B. (2011), 'Lateral dynamics of charged lipids and peripheral proteins in spatially heterogeneous membranes: Comparison of continuous and monte carlo approaches', J. Chem. Phys. 135(155103).
- Klapper, I., Hagstrom, R., Fine, R., Sharp, K. & Honig, B. (1986), 'Focussing of electric fields in the active site of cu-zn superoxide dismutase: Effects of ionic strength and amino acid modification', *Protein* 1, 47 – 59.
- Kolomeisky, A. (2007), 'Channel-Facilitated Molecular Transport across Membranes: Attraction, Repulsion, and Asymmetry', *Phys. Rev. Lett.* **98**, 048105.
- Kurnikova, M. G., Coalson, R. D., Graf, P. & Nitzan, A. (1999), 'A lattice relaxation algorithm for Three-Dimensional Poisson-Nernst-Planck theory with application to ion transport through the Gramicidin A channel', *Biophysical Journal* 76, 642–656.
- Kuyucak, S., Andersen, O. S. & Chung, S.-H. (2001), 'Models of permeation in ion channels', *Rep. Prog. Phys* **64**, 1427–1472.
- Labro, A., Priest, M., Lacroix, J., Snyders, D. & Bezanilla, F. (2015), ' $K_v 3.1$ uses a timely resurgent K⁺ current to secure action potential repolarization', *Nature communications* **6**, 10173.
- Lacroix, J. J., Campos, F., Frezza, L. & Bezanilla, F. (2013), 'Molecular Bases for the Asynchronous Activation of Sodium and Potassium Channels Required for Nerve Impulse Generation', *Neuron* **79**, 651–657.
- Läuger, P. (1985), 'Ionic channels with conformational substates', Biophys J. 47, 581-590.
- Levitt, D. G. (1991a), 'General continuum theory for multiion channel. I. Theory', Biophys J. 59, 271-277.
- Levitt, D. G. (1991b), 'General continuum theory for multiion channel. II. Application to acetylcholine channel', Biophys J. 59, 278–288.
- Levitt, D. G. (1999), 'Modeling of ion channels', J. Gen. Physiol. 113(6), 789-794.
- Li, B., Lu, B. Z., Wang, Z. M. & McCammon, J. A. (2010), 'Solutions to a reduced Poisson-Nernst-Planck system and determination of reaction rates', *Physica A* **389**(7), 1329–1345.
- Li, H. & Lu, B. (2014), 'An ionic concentration and size dependent dielectric permittivity Poisson-Boltzmann model for biomolecular solvation studies', *J. Chem. Phys.* **141**, 024115.
- Lin, H., Cho, M.-H., Tang, H. & Cai, W. (2013), 'icBIE: A Boundary Integral Equation Program for an Ion Channel in Layered Membrane/Electrolyte Media.', *Computer Physics Communications* p. To appear.
- Lin, T.-C. & Eisenberg, B. (2014), 'A new approach to the Lennard-Jones potential and a new model: PNPsteric equations', *Communications in Mathematical Sciences* **12**, 149–173.
- Lin, T.-C. & Eisenberg, B. (2015), 'Multiple solutions of steady-state Poisson-Nernst-Planck equations with steric effects', Nonlinearity 28, 2053–2080.
- Liu, H. & Wang, Z. (2014), ' A free energy satisfying finite difference method for Poisson-Nernst-Planck equations', *Journal of Computational Physics* **268**, 363–376.
- Liu, H. & Wang, Z. (2017), 'A free energy satisfying discontinuous Galerkin method for one-dimensional Poisson-Nernst-Planck systems', *Journal of Computational Physics* **328**, 413–437.
- Liu, J.-L. & Eisenberg, B. (2014), 'Poisson-Nernst-Planck-Fermi Theory for Modeling Biological Ion Channels', Journal of Chemical Physics 141, 22D532.
- Liu, W. (2009), 'One-dimensional steady-state poisson-nernst-planck systems for ion channels with multiple ion species', *Journal of Differential Equations* **246**(1), 428–451.

- Liu, W. S. (2005), 'Geometric singular perturbation approach to steady-state Poisson-Nernst-Planck systems', SIAM J. Appl. Math. 65(3), 754–766.
- Long, S. B., Campbell, E. B. & Mackinnon, R. (2005), 'Crystal structure of a mammalian voltage-dependent Shaker family K+ channel', *Science* **309**, 897–903.
- Lu, B., Holst, M., McCammon, J. A. & Zhou, Y. C. (2010), 'Poisson-Nernst-Planck Equations for Simulating Biomolecular Diffusion-Reaction Processes I: Finite element solutions', *Journal of Computational Physics* 229, 6979–6994.
- Lu, B. & Zhou, Y. C. (2011), 'Poisson-Nernst-Planck Equations for Simulating Biomolecular Diffusion-Reaction Processes II: Size Effects on Ionic Distributions and Diffusion-Reaction Rates', *Biophysical Journal* 100, 2475–2485.
- MacKerell, A. D., J., Bashford, D., Bellot, M., Dunbrack, R. L., J., Evanseck, J. D., Field, M. J., Fischer, S., Gao, J., Guo, H., Ha, S., Joseph-McCarthy, D., Kuchnir, L., Kuczera, K., Lau, F. T. K., Mattos, C., Michnick, S., Ngo, T., Nguyen, D. T., Prodhom, B., Reiher, W. E., I., Roux, B., Schlenkrich, M., Smith, J. C., Stote, R., Straub, J., Watanabe, M., Wiorkiewicz-Kuczera, J., Yin, D. & Karplus, M. (1998), 'All-atom empirical potential for molecular modeling and dynamics studies of proteins', *Journal of Physical Chemistry B* 102(18), 3586–3616.
 URL: http://dx.doi.org/10.1021/jp973084f
- Mackoy, T., Harris, R. C., Johnson, J., Mascagni, M. & Fenley, M. O. (2013), 'Numerical Optimization of a Walk-on-Spheres Solver for the Linear Poisson-Boltzmann Equation', *Communications in Computational Physics* 13, 195–206.
- Madura, J. D., Briggs, J. M., Wade, R. C., Davis, M. E., Luty, B. A., Ilin, A., Antosiewicz, J., Gilson, M. K., Bagheri, B., Scott, L. R. & McCammon, J. A. (1995), 'Electrostatics and diffusion of molecules in solution - simulations with the University of Houston Brownian Dynamics program', *Computer Physics Communications* 91(1-3), 57–95. URL: http://dx.doi.org/10.1016/0010-4655(95)00043-F
- Mamonov, A. B., Kurnikova, M. G. & Coalson, R. D. (2006), 'Diffusion constant of K⁺ inside Gramicidin A: a comparative study of four computational methods', *Biophys Chem* **124**, 268–78.
- Marx, D. & Hutter, J. (2000), 'Ab initio molecular dynamics: Theory and implementation', Modern Methods and Algorithms of Quantum Chemistry, J. Grotendorst (Ed.), John von Neumann Institute for Computing, Jülich, NIC Series 3, 329–477.
- McCusker, E. C., Bagneris, C., Naylor, C. E., Cole, A. R., D'Avanzo, N., Nichols, C. G. & Wallace, B. A. (2012), 'Structure of a bacterial voltage-gated sodium channel pore reveals mechanisms of opening and closing', *Nature Communications* **3**, 1102.
- Meng, D., Zheng, B., Lin, G. & Sushko, M. L. (2014), 'Numerical Solution of 3D Poisson-Nernst-Planck Equations Coupled with Classical Density Functional Theory for Modeling Ion and Electron Transport in a Confined Environment', *Communications in Computational Physics* 16, 1298–1322.
- Metti, M. S., Xu, J. & Liu, C. (2016), 'Energetically stable discretizations for charge transport and electrokinetic models', *Journal of Computational Physics* **306**, 1–18.
- Miao, Y., Fu, R., Zhou, H.-X. & Cross, T. A. (2015), 'Dynamic short hydrogen bonds in histidine tetrad of full length M2 proton channel reveal tetrameric structural heterogeneity and functional mechanism', *Structure* 23, 2300–2308.
- Mitchell, P. (1976), 'Vectorial chemistry and the molecular mechanics of chemiosmotic coupling: power transmission by proticity', *Biochem Soc Trans* **4**, 399 430.

- Mukhopadhyay, A., Fenley, A. T., Tolokh, I. S. & Onufriev, A. V. (2012), 'Charge hydration asymmetry: The basic principle and how to use it to test and improve waster models', *J. Phys. Chem. B* **116**, 9776–9783.
- Nagle, J. F. & Morowitz, H. J. (1978), 'Molecular mechanisms for proton transport in membranes', Proc. Natl. Acad. Sci. U.S.A 1458(72), 298–302.
- Noskov, S., Berneche, S. & Roux, B. (2004), 'Control of ion selectivity in potassium channels by electrostatic and dynamic properties of carbonyl ligands', *Nature* **431**, 830–832.
- Onufriev, A., Case, D. A. & Bashford, D. (2002), 'Effective Born radii in the generalized Born approximation: the importance of being perfect', *Journal of Computational Chemistry* 23(14), 1297–304. URL: http://dx.doi.org/10.1002/jcc.10126
- Park, J. & Jerome, J. W. (1997), 'Qualitative properties of steady-state Poisson-Nernst-Planck systems: Mathematical study', *SIAM Journal on Applied Mathematics* **57**(3), 609-630.
- Park, J. K. & Wei, G. W. (2014), 'A molecular level prototype for mechanoelectrical transducers in mammalian hair cells', *Journal of Computational Neuroscience* 35, 231–241.
- Park, J., Xia, K. L. & Wei, G. W. (2015), 'Atomic scale design and three-dimensional simulations of nanofluidic systems', *Microfluidics and Nanofluidics* 19, 665–692.
- Payandeh, J., Gamal El-Din, T. M., Scheuer, T., Zheng, N. & Catterall, W. A. (2012), 'Crystal structure of a voltage-gated sodium channel in two potentially inactivated states', *Nature* **486**, 135–139.
- Perlman, D. A., Case, D. A., Caldwell, J. W., Ross, W. S., Cheatham, T. E., Debolt, S., Ferguson, D., Seibel, G. & Kollman, P. (1995), 'Amber, a package of computer programs for applying molecular mechanics, normal mode analysis, molecular dynamics and free energy calculations to simulate the structural and energetic properties of molecules', *Comp. Phys. Commun.* **91**, 1–41.
- Pomes, R. & Roux, B. (1996), 'Theoretical study of H⁺ translocation along a model proton wire', *J. Phys. Chem* **100**, 2519–2527.
- Pomes, R. & Roux, B. (2002*a*), 'Molecular mechanism of H⁺ conduction in the single-file water chain of the Gramicidin channel', *Biophysical Journal* **82**, 2304–2316.
- Pomes, R. & Roux, B. (2002b), 'Structure and Dynamics of a Proton Wire: A Theoretical Study of H⁺ Translocation along the Single-File Water Chain in the Gramicidin A Channel', *Biophysical Journal* **71**, 19– 39.
- Prabhu, N. V., Panda, M., Yang, Q. Y. & Sharp, K. A. (2008), 'Explicit ion, implicit water solvation for molecular dynamics of nucleic acids and highly charged molecules', J. Comput. Chem. 29, 1113–1130.
- Qiao, Y., Liu, X., Chen, M. & Lu, B. (2016), 'A Local Approximation of Fundamental Measure Theory Incorporated into Three Dimensional Poisson–Nernst–Planck Equations to Account for Hard Sphere Repulsion Among Ions', *Journal of Statistical Physics* 163(1), 156–174.
- Qiao, Y., Tu, B. & Lu, B. (2014), 'lonic size effects to molecular solvation energy and to ion current across a channel resulted from the nonuniform size-modified PNP equations', *Journal of Chemical Physics* **140**, 174102.
- Qin, S. & Zhou, H.-X. (2013), 'FFT-Based Method for Modeling Protein Folding and Binding under Crowding: Benchmarking on Ellipsoidal and All-Atom Crowders', J. Chem. Theory Comput. 9, 4633–4643.
- Qin, S. & Zhou, H.-X. (2014), 'Further Development of the FFT-based Method for Atomistic Modeling of Protein Folding and Binding under Crowding: Optimization of Accuracy and Speed', J. Chem. Theory Comput 10, 2824–2835.

- Ren, P., Chun, J., Thomas, D. G., Schnieders, M. J., Marucho, M., Zhang, J. & Baker, N. A. (2013), 'Biomolecular electrostatics and solvation: a computational perspective', *Quart. Rev. Biophys.* .
- Richards, F. M. (1977), 'Areas, volumes, packing, and protein structure', *Annual Review of Biophysics and Bioengineering* **6**(1), 151–176.
- Rocchia, W., Alexov, E. & Honig, B. (2001), 'Extending the applicability of the nonlinear poisson-boltzmann equation: Multiple dielectric constants and multivalent ions', *J. Phys. Chem.* **105**, 6507–6514.
- Rocchia, W., Sridharan, S., Nicholls, A., Alexov, E., Chiabrera, A. & Honig, B. (2002), 'Rapid grid-based construction of the molecular surface and the use of induced surface charge to calculate reaction field energies: Applications to the molecular systems and geometric objects', *Journal of Computational Chemistry* 23, 128 137.
- Roux, B. (2002), 'Computational studies of the Gramicidin channel', Acc. Chem. Res. 35, 366-375.
- Roux, B., Allen, T., Berneche, S. & Im, W. (2004), 'Theoretical and computational models of biological ionchannels', *Quarterly Reviews of Biophysics* **7**(1), 1–103.
- Sanner, M. F., Olson, A. J. & Spehner, J. C. (1996), 'Reduced surface: An efficient way to compute molecular surfaces', *Biopolymers* 38, 305–320.
- Schmitt, U. W. & Voth, G. A. (1999), 'The computer simulation of proton transport in water', Journal of Chemical Physics 111, 9361–9381.
- Schmuck, M. & Bazant, M. Z. (2015), 'Homogenization of the Poisson-Nernst-Planck equations for Ion Transport in Charged Porous Media', *SIAM Journal on Applied Mathematics* **75**(3), 1369–1401.
- Schnell, J. R. & Chou, J. J. (2008), 'Structure and mechanism of the M2 proton channel of influenza A virus', *Nature* 451, 591–596.
- Schumaker, M. F., Pomes, R. & Roux, B. (2000), 'A combined molecular dynamics and diffusion model of single proton conduction through Gramicidin', *Biophysical Journal* 79, 2840–2857.
- Sharma, M., Yi, M., Dong, H., Qin, H., Peterson, E., Busath, D. D., Zhou, H. & Cross, T. A. (2010), 'Insight into the mechanism of the influenza A proton channel from a structure in a lipid bilayer.', *Science* **330**, 509–12.
- Sharp, K. A. & Honig, B. (1990), 'Calculating total electrostatic energies with the nonlinear Poisson-Boltzmann equation', *Journal of Physical Chemistry* **94**, 7684–7692.
- Shaya, D., Findeisen, F., Abderemane-Ali, F., Arrigoni, C., Wong, S., Nurva, S. R., Loussouarn, G. & Minor, D. L. (2014), 'Structure of a prokaryotic sodium channel pore reveals essential gating elements and an outer ion binding site common to eukaryotic channels', *Journal of Molecular Biology* **426**, 467–483.
- Shepherd, L. M. S. & Morrison, C. A. (2010), 'Simulating Proton Transport through a Simplified Model for Trans-Membrane Proteins', Journal of Physical Chemistry 114, 7047–7055.
- Simakov, N. A. & Kurnikova, M. G. (2010), 'Soft wall ion channel in continuum representation with application to modeling ion currents in α -hemolysin', J. Phys. Chem. B **114**(46), 15180C15190.
- Simonov, N., Mascagni, M. & O, F. M. (2007), 'Monte Carlo-based linear Poisson-Boltzmann approach makes accurate salt-dependent solvation free energy predictions possible', *Journal of Chemical Physics* 127, 185105.
- Singer, A., Gillespie, D., Norbury, J. & Eisenberg, R. S. (2008), 'Singular perturbation analysis of the steady state Poisson-Nernst-Planck system: Applications to ion channels', *European Journal of Applied Mathematics* 19, 541–560.
- Singer, A. & Norbury, J. (2009), 'A Poisson-Nernst-Planck model for biological ion channels?an asymptotic analysis in a three-dimensional narrow funnel', SIAM Journal on Applied Mathematics 70(3), 949–968.

- Spolar, R. S. & Record Jr., M. T. (1994), 'Coupling of local folding to site-specific binding of proteins to dna', Science 263, 777–784.
- Sun, Y., Sun, P., Zheng, B. & Lin, G. (2016a), 'Error analysis of finite element method for Poisson-Nernst-Planck equations', *Journal of Computational and Applied Mathematics* **301**, 28–43.
- Sun, Y., Sun, P., Zheng, B. & Lin, G. (2016b), 'Error analysis of finite element method for Poisson-Nernst-Planck equations', *Journal of Computational and Applied Mathematics* **301**, 28-43.
- Tian, W. F. & Zhao, S. (2014), 'A fast ADI algorithm for geometric flow equations in biomolecular surface generations', *International Journal for Numerical Methods in Biomedical Engineering* **30**, 490–516.
- Tu, B., Chen, M. X., Xie, Y., Zhang, L. B., Eisenberg, B. & Lu, B. Z. (2013), 'A parallel finite element simulator for ion transport through three-dimensional ion channel systems', J. Comput. Chem. 34, 2065–2078.
- Vlachy, V. (1999), 'lonic effects beyond poisson-boltzmann theory', Annu. Rev. Phys. Chem. 50, 145-165.
- Wang, B. & Wei, G. W. (2015), 'Parameter optimization in differential geometry based solvation models', Journal Chemical Physics 143, 134119.
- Wang, B., Zhao, Z. & Wei, G. W. (submitted 2016), 'Feature functional theory binding predictor (fft-bp) for the blind prediction of binding free energy', *Journal of Chemical Theory and Computation*.
- Wang, J., Cai, Q., Xiang, Y. & Luo, R. (2012), 'Reducing Grid Dependence in Finite Difference Poisson Boltzmann Calculations', *Journal of Chemical Theory and Computation* **8**, 2741–2751.
- Wang, M. L., Lu, Z. Y. & Yang, W. T. (2006), 'Nuclear quantum effects on an enzyme-catalyzed reaction with reaction path potential: Proton transfer in triosephosphate isomerase', *Journal of Chemical Physics* 124(12), 124516. URL: *jGo to ISI*₂://000236464000053
- Wang, X.-S., He, D., Wylie, J. J. & Huang, H. (2014), 'Singular perturbation solutions of steady-state Poisson-Nernst-Planck systems', *Physical Review E* **89**(2), 022722.
- Wang, Y., Liu, C. & Tan, Z. (2016), 'A Generalized Poisson–Nernst–Planck–Navier–Stokes Model on the Fluid with the Crowded Charged Particles: Derivation and Its Well-Posedness', SIAM Journal on Mathematical Analysis 48(5), 3191–3235.
- Weeks, J. D., Chandler, D. & Andersen, H. C. (1971), 'Role of repulsive forces in determining the equilibrium structure of simple liquids', *Journal of Chemical Physics* 54(12), 5237–47. URL: http://dx.doi.org/10.1063/1.1674820
- Wei, G. W. (2010), 'Differential geometry based multiscale models', *Bulletin of Mathematical Biology* **72**, 1562 1622.
- Wei, G. W. (2013), 'Multiscale, multiphysics and multidomain models I: Basic theory', *Journal of Theoretical* and Computational Chemistry **12**(8), 1341006.
- Wei, G. W. (2016), 'Mathematical molecular bioscience and biophysics', SIAM News 49(7).
- Wei, G. W., Sun, Y. H., Zhou, Y. C. & Feig, M. (2005), 'Molecular multiresolution surfaces', arXiv:mathph/0511001v1 pp. 1 – 11.
- Wei, G.-W., Zheng, Q., Chen, Z. & Xia, K. (2012), 'Variational multiscale models for charge transport', SIAM Review 54(4), 699 – 754.
- Xia, K. L. & Wei, G. W. (2014), 'A Galerkin formulation of the MIB method for three dimensional elliptic interface problems', *Computers and Mathematics with Applications* **68**, 719–745.

- Xia, K. L. & Wei, G. W. (submitted 2016), 'A review of geometric, topological and graph theory apparatus for the modeling and analysis of biomolecular data'.
- Xia, K. L., Zhan, M., Wan, D. C. & Wei, G. W. (2012), 'Adaptively deformed mesh based matched interface and boundary (MIB) method for elliptic interface problems', *Journal of Computational Physics* **231**, 1440 –1461.
- Xia, K. L., Zhan, M. & Wei, G.-W. (2011), 'The matched interface and boundary (MIB) method for multidomain elliptic interface problems', *Journal of Computational Physics* 230, 8231–8258.
- Xia, K. L., Zhan, M. & Wei, G. W. (2014), 'MIB Galerkin method for elliptic interface problems', *Journal of Computational and Applied Mathematics* **272**, 195–220.
- Xie, D., Jiang, Y. & Scott, L. R. (2013), 'Efficient algorithms for solving a nonlocal dielectric model for protein in ionic solvent', *SIAM Journal on Scientific Computing* **38**, B1267–1284.
- Xie, D. & Zhou, S. Z. (2007), 'A new minimization protocol for solving nonlinear Poisson-Boltzmann mortar finite element equation', *BIT NUMERICAL MATHEMATICS* **47**, 853–871.
- Xu, S., Sheng, P. & Liu, C. (2014), 'An energetic variational approach for ion transport', *Communications in Mathematical Sciences* **12**, 779–789.
- Xu, Z., Ma, M. & Liu, P. (2014), 'Self-energy-modified Poisson-Nernst-Planck equations: WKB approximation and finite-difference approaches', *Physical Review E* **90**, 013307.
- Yan, S. H., Liang Zhang, L., Cukier, R. I. & Bu, Y. X. (2007), 'Exploration on regulating factors for proton transfer along hydrogen-bonded water chains', *Chem. Phys. Chem.* 8, 944–954.
- Yang, Y. & Lu, B. (2013), 'An error analysis for the finite element approximation to the steady-state Poisson-Nernst-Planck equations', Advances in Applied Mathematics and Mechanics 5(01), 113–130.
- Yu, S. N. & Wei, G. W. (2007), 'Three-dimensional matched interface and boundary (MIB) method for treating geometric singularities', J. Comput. Phys. 227, 602–632.
 URL: http://www.sciencedirect.com/science/article/B6WHY-4PF1WN4-1/2/a142bb0596d6f227a0537b4790d00bcf
- Yu, S. N., Zhou, Y. C. & Wei, G. W. (2007), 'Matched interface and boundary (MIB) method for elliptic problems with sharp-edged interfaces', J. Comput. Phys. 224(2), 729–756. URL: http://dx.doi.org/10.1016/j.jcp.2006.10.030
- Zhang, M. (2015), 'Asymptotic expansions and numerical simulations of I-V relations via a steady state Poisson-Nernst-Planck system', *Rocky Mountain Journal of Mathematics* **45**, 1681–1708.
- Zhao, S. (2011), 'Pseudo-time-coupled nonlinear models for biomolecular surface representation and solvation analysis', *International Journal for Numerical Methods in Biomedical Engineering* 27, 1964–1981.
- Zhao, S. (2012), 'Pseudo-time-coupled nonlinear models for biomolecular surface representation and solvation analysis', *International Journal for Numerical Methods in Biomedical Engineering* **27**, 1964–1981.
- Zhao, S. (2014), 'Operator splitting ADI schemes for pseudo-time coupled nonlinear solvation simulations', Journal of Computational Physics 257, 1000 – 1021.
- Zhao, S. & Wei, G. W. (2004), 'High-order FDTD methods via derivative matching for Maxwell's equations with material interfaces', J. Comput. Phys. 200(1), 60–103. URL: http://dx.doi.org/10.1016/j.jcp.2004.03.008
- Zheng, Q., Chen, D. & Wei, G. W. (2011), 'Second-order Poisson-Nernst-Planck solver for ion transport', Journal of Comput. Phys. 230, 5239 – 5262.

- Zheng, Q. & Wei, G. W. (2011), 'Poisson-Boltzmann-Nernst-Planck model', *Journal of Chemical Physics* 134, 194101.
- Zhou, H.-X. (2011*a*), 'A theory for the proton transport of the influenza virus M2 protein: extensive test against conductance data', *Biophys.* **100**, 912–921.
- Zhou, H.-X. (2011*b*), 'A Theory for the Proton Transport of the Influenza Virus M2 Protein: Extensive Test against Conductance Data', *Biophys J.* **100**, 912–921.
- Zhou, H.-X. (2011*c*), 'Mechanistic Insight into the H₂O/D₂O Isotope Effect in the Proton Transport of the Influenza Virus M2 Protein', *J. Membr Biol* **244**, 93–96.
- Zhou, S., Wang, Z. & Li, B. (2011), 'Mean-field description of ionic size effects with non-uniform ionic sizes: A numerical approach', *Phys. Rev. E* **84**, 021901.
- Zhou, Y. C., Feig, M. & Wei, G. W. (2008), 'Highly accurate biomolecular electrostatics in continuum dielectric environments', *Journal of Computational Chemistry* 29, 87–97.
- Zhou, Y. C. & Wei, G. W. (2006), 'On the fictitious-domain and interpolation formulations of the matched interface and boundary (MIB) method', J. Comput. Phys. 219(1), 228–246. URL: http://dx.doi.org/10.1016/j.jcp.2006.03.027
- Zhou, Y. C., Zhao, S., Feig, M. & Wei, G. W. (2006), 'High order matched interface and boundary method for elliptic equations with discontinuous coefficients and singular sources', J. Comput. Phys. 213(1), 1–30. URL: http://dx.doi.org/10.1016/j.jcp.2005.07.022
- Zhou, Y., Lu, B., Huber, G. A., Holst, M. J. & McCammon, J. A. (2008), 'Continuum simulations of acetylcholine consumption by acetylcholinesterase: A Poisson- Nernst- Planck approach', *The Journal of Physical Chemistry B* **112**(2), 270–275.
- Ziner, B. & Cai, W. (2016), 'A well-conditioned hypersingular boundary element method for electrostatic potentials in the presence of inhomogeneities within layered media', *Communication in Computational Physics* **19**, 970–997.
- Zinsl, J. (2015), 'Exponential convergence to equilibrium in a Poisson-Nernst-Planck-type system with nonlinear diffusion', arXiv preprint arXiv:1503.04029.