

## THE ROUND SPHERE MINIMIZES ENTROPY AMONG CLOSED SELF-SHRINKERS

TOBIAS HOLCK COLDING, TOM ILMANEN,  
WILLIAM P. MINICOZZI II & BRIAN WHITE

### Abstract

The entropy of a hypersurface is a geometric invariant that measures complexity and is invariant under rigid motions and dilations. It is given by the supremum over all Gaussian integrals with varying centers and scales. It is monotone under mean curvature flow, thus giving a Lyapunov functional. Therefore, the entropy of the initial hypersurface bounds the entropy at all future singularities. We show here that not only does the round sphere have the lowest entropy of any closed singularity, but there is a gap to the second lowest.

### 0. Introduction

The  $F$ -functional of a hypersurface  $\Sigma$  of Euclidean space  $\mathbf{R}^{n+1}$  is the Gaussian surface area

$$(0.1) \quad F(\Sigma) = (4\pi)^{-\frac{n}{2}} \int_{\Sigma} e^{-\frac{|x|^2}{4}} ,$$

whereas the Gaussian entropy is the supremum over all Gaussian surface areas given by

$$(0.2) \quad \lambda(\Sigma) = \sup (4\pi t_0)^{-\frac{n}{2}} \int_{\Sigma} e^{-\frac{|x-x_0|^2}{4t_0}} .$$

(Gaussian surface area has also been studied in convex geometry (see [B] and [Na]) and in theoretical computer science (see [K] and [KDS]).) Here the supremum is taken over all  $t_0 > 0$  and  $x_0 \in \mathbf{R}^{n+1}$ . Entropy is invariant under rigid motions and dilations.

Mean curvature flow (“MCF”) is an evolution equation where a one-parameter family of hypersurfaces  $M_t \subset \mathbf{R}^{n+1}$  evolves over time to minimize volume, satisfying the equation

$$(0.3) \quad (\partial_t x)^\perp = -H \mathbf{n} .$$

---

Received 5/25/2012.

Here  $H = \operatorname{div}_\Sigma \mathbf{n}$  is the mean curvature,  $\mathbf{n}$  is the outward pointing unit normal and  $x$  is the position vector. As a consequence of Huisken's monotonicity, entropy is monotone nonincreasing under MCF.

A hypersurface  $\Sigma \subset \mathbf{R}^{n+1}$  is a *self-shrinker* if it is the  $t = -1$  time-slice of a MCF moving by rescalings, i.e., where  $\Sigma_t \equiv \sqrt{-t}\Sigma$  is a MCF; see [A], [Ch], [KKM], [M], and [N] for examples. This is easily seen to be equivalent to the equation

$$(0.4) \quad H = \frac{\langle x, \mathbf{n} \rangle}{2}.$$

Since (0.4) is the Euler-Lagrange equation for  $F$ , self-shrinkers are critical points for  $F$ .

Under MCF, every closed hypersurface becomes singular and the main problem is to understand the singularities. By Huisken's monotonicity and an argument of [I1], [W2], blow ups of singularities of a MCF are self-shrinkers.

By section 7 in [CM1], the entropy of a self-shrinker is equal to the value of  $F$  and, thus, no supremum is needed. In [St], Stone computed the  $F$  functional, and therefore the entropy, for generalized cylinders  $\mathbf{S}^k \times \mathbf{R}^{n-k}$ . He showed that  $\lambda(\mathbf{S}^n)$  is decreasing in  $n$  and

$$(0.5) \quad \lambda(\mathbf{S}^1) = \sqrt{\frac{2\pi}{e}} \approx 1.5203 > \lambda(\mathbf{S}^2) = \frac{4}{e} \approx 1.4715 > \lambda(\mathbf{S}^3) > \dots > 1$$

$$(0.6) \quad = \lambda(\mathbf{R}^n).$$

Moreover, a simple computation shows that  $\lambda(\Sigma \times \mathbf{R}) = \lambda(\Sigma)$ .

It follows from Brakke's theorem, [Br], that  $\mathbf{R}^n$  has the least entropy of any self-shrinker and, in fact, there is a gap to the next lowest. Our main result is that the round sphere has the least entropy of any *closed* self-shrinker.

**Theorem 0.7.** *Given  $n$ , there exists  $\epsilon = \epsilon(n) > 0$  so that if  $\Sigma \subset \mathbf{R}^{n+1}$  is a closed self-shrinker not equal to the round sphere, then*

$$(0.8) \quad \lambda(\Sigma) \geq \lambda(\mathbf{S}^n) + \epsilon.$$

*Moreover, if  $\lambda(\Sigma) \leq \min\{\lambda(\mathbf{S}^{n-1}), \frac{3}{2}\}$ , then  $\Sigma$  is diffeomorphic to  $\mathbf{S}^n$ . (If  $n > 2$ , then  $\lambda(\mathbf{S}^{n-1}) < \frac{3}{2}$  and the minimum is unnecessary.)*

Theorem 0.7 is suggested by the dynamical approach to MCF of [CM1] and [CM2]. The idea is that MCF starting at a closed  $M$  becomes singular, the corresponding self-shrinker has lower entropy, and, by [CM1], the only self-shrinkers that cannot be perturbed away are  $\mathbf{S}^{n-k} \times \mathbf{R}^k$  and  $\lambda(\mathbf{S}^{n-k} \times \mathbf{R}^k) \geq \lambda(\mathbf{S}^n)$ . However, we are not able to make this approach rigorous. One of the difficulties is that if the self-shrinker is non-compact and we perturb it, then *a priori* it may flow smoothly

without ever becoming singular. (Theorem 0.10 in [CM1] is proven using an argument along these lines.) To overcome this, we combine results from [CM1] and [IW].

The dynamical picture also suggests two closely related conjectures; the first is for any closed hypersurface and the second is for self-shrinkers:

**Conjecture 0.9.** *Theorem 0.7 holds with  $\epsilon = 0$  for any closed hypersurface  $M^n$  with  $n \leq 6$ .*

**Conjecture 0.10.** *Theorem 0.7 holds for any non-flat self-shrinker  $\Sigma^n \subset \mathbf{R}^{n+1}$  with  $n \leq 6$ .*

Both conjectures are true for curves, i.e., when  $n = 1$ . The first conjecture follows for curves by combining Grayson's theorem, [G] (cf. [GaHa]), and the monotonicity of  $\lambda$  under curve shortening flow. The second conjecture follows for curves from the classification of self-shrinking curves by Abresch and Langer, [AbL].

Conjecture 0.9 would allow us to carry out the outline above to show that any closed hypersurface has entropy at least that of the sphere, proving Conjecture 0.10.

Furthermore, one could ask which self-shrinker has the third least entropy, etc. It is easy to see that the entropy of the ‘‘Simons cone’’ over  $\mathbf{S}^k \times \mathbf{S}^k$  in  $\mathbf{R}^{2k+2}$  is asymptotic to  $\sqrt{2}$  as  $k \rightarrow \infty$ , which is also the limit of  $\lambda(\mathbf{S}^{2k+1})$ . Thus, as the dimension increases, the Simons cones have lower entropy than some of the generalized cylinders. For example, the cone over  $\mathbf{S}^2 \times \mathbf{S}^2$  has entropy  $\frac{3}{2} < \lambda(\mathbf{S}^1 \times \mathbf{R}^4)$ . In other words, already for  $n = 5$ ,  $\mathbf{S}^k \times \mathbf{R}^{n-k}$  is not a complete list of the lowest entropy self-shrinkers.

It is easy to see that if an immersed hypersurface has entropy strictly less than two, then it is embedded; hence, we will always assume embeddedness below.

**Acknowledgments.** The first and third authors were partially supported by NSF Grants DMS 11040934, DMS 0906233, and NSF FRG grants DMS 0854774 and DMS 0853501. The fourth author was partially supported by NSF grant DMS 1105330.

## 1. Perturbing a self-shrinker

Throughout this paper, unless otherwise mentioned,  $L$  will be the second variation operator for the  $F$  functional from section 4 of [CM1], that is,

$$(1.1) \quad L = \mathcal{L} + |A|^2 + \frac{1}{2} = \Delta - \left\langle \frac{x}{2}, \nabla \cdot \right\rangle + |A|^2 + \frac{1}{2},$$

where  $A$  is the second fundamental form and the second equality defines the operator  $\mathcal{L}$ .

The first step is to perturb a closed self-shrinker inside itself, while reducing the entropy and making the self-shrinker version of mean curvature positive. This uses the classification of stable self-shrinkers from [CM1].

**Lemma 1.2.** *If  $\Sigma^n \subset \mathbf{R}^{n+1}$  is a closed self-shrinker (for any  $n$ ) and  $\Sigma$  is not round, then there is a nearby hypersurface  $\Gamma$  with the following properties:*

- 1)  $\lambda(\Gamma) < \lambda(\Sigma)$ .
- 2)  $\Gamma$  is inside of  $\Sigma$ , i.e., the compact region of  $\mathbf{R}^{n+1}$  bounded by  $\Sigma$  contains  $\Gamma$ .
- 3)  $(H - \frac{1}{2}\langle x, \mathbf{n} \rangle) > 0$  on  $\Gamma$ .

*Proof.* Let  $\Sigma_s$  be a one-parameter family of normal graphs given by

$$(1.3) \quad \Sigma_s = \{x + s u(x) \mathbf{n}(x) \mid x \in \Sigma\},$$

where  $\mathbf{n}(x)$  is the outward unit normal to  $\Sigma$ . Equation (4.10) in [CM1] computes that

$$(1.4) \quad \frac{d}{ds} \Big|_{s=0} \left( H - \frac{\langle x, \mathbf{n} \rangle}{2} \right) = -L u.$$

The following three facts are proved in [CM1]:

- The lowest eigenfunction  $u$  for  $L$  is positive and has  $L u = \mu u$  for some  $\mu > 1$ .
- There exists  $\epsilon > 0$ , so that for every  $s$  with  $0 < |s| < \epsilon$  we have  $\lambda(\Sigma_s) < \lambda(\Sigma)$ . Here  $\Sigma_s$  is the graph of  $s u$ .
- With  $s$  as above,  $\Sigma_s$  has

$$(1.5) \quad \left( H - \frac{1}{2} \langle x, \mathbf{n} \rangle \right) < 0 \text{ if } s > 0,$$

$$(1.6) \quad \left( H - \frac{1}{2} \langle x, \mathbf{n} \rangle \right) > 0 \text{ if } s < 0.$$

(See [CM1] corollary 5.15 and theorem 4.30, theorem 0.15, and equation (4.10), respectively.) Thus, we see that for  $s \in (-\epsilon, 0)$ ,  $\Gamma = \Sigma_s$  has the three properties. q.e.d.

## 2. Rescaled MCF

A one-parameter family of hypersurfaces  $M_t$  evolves by rescaled MCF if it satisfies

$$(2.1) \quad \partial_t x = - \left( H - \frac{1}{2} \langle x, \mathbf{n} \rangle \right) \mathbf{n}.$$

The rescaled MCF is equivalent to MCF, up to rescalings in space and a reparameterization of time. Namely, if  $\Sigma_t$  is a MCF, then  $t \rightarrow \Sigma_{-e^{-t}}/\sqrt{e^{-t}}$  is a solution to the rescaled MCF equation and vice versa.

Moreover, rescaled MCF is the negative gradient flow for the  $F$  functional and self-shrinkers are the fixed points for this flow.

**2.1. Mean convex rescaled MCF.** The next ingredient is a result from [IW] about “mean convex rescaled MCF” where

$$(2.2) \quad \left( H - \frac{1}{2} \langle x, \mathbf{n} \rangle \right) \geq 0.$$

We will often refer to this quantity as the rescaled mean curvature.

This result is the rescaled analog of that mean convexity is preserved for MCF.

**Lemma 2.3.** *Let  $M_t \subset \mathbf{R}^{n+1}$  be a smooth rescaled MCF of closed hypersurfaces for  $t \in [0, T]$ .*

- 1) *If (2.2) holds on  $M_0$ , then it also holds on  $M_t$  for every  $t \in [0, T]$ .*
- 2) *If in addition  $(H - \frac{1}{2} \langle x, \mathbf{n} \rangle) > 0$  at least at one point of  $M_0$ , then  $(H - \frac{1}{2} \langle x, \mathbf{n} \rangle) > 0$  on  $M_t$  for all  $t > 0$ .*

*In particular, the flow is monotone in that  $M_t$  is inside  $M_s$  for  $t > s$ .*

The key to proving this is a Simons type equation for rescaled MCF

$$(2.4) \quad (\partial_t - L) \left( H - \frac{\langle x, \mathbf{n} \rangle}{2} \right) = 0,$$

which follows from (1.4) with  $u = - \left( H - \frac{\langle x, \mathbf{n} \rangle}{2} \right)$ .

*Proof of Lemma 2.3.* Set  $\phi = H - \frac{1}{2} \langle x, \mathbf{n} \rangle$ . Equation (2.4) is equivalent to

$$(2.5) \quad (\partial_t - \Delta) \phi = \left( |A|^2 + \frac{1}{2} \right) \phi - \left\langle \frac{x}{2}, \nabla \phi \right\rangle.$$

Since  $\phi(x, 0) \geq 0$  by assumption, the parabolic maximum principle gives  $\phi(x, t) \geq 0$  for all  $t \geq 0$ , giving (1). The parabolic strong maximum principle then gives (2). The last claim follows immediately since  $\mathbf{n}$  is the outward pointing normal. q.e.d.

### 3. Simons type identity for the rescaled $A$

Using covariant derivatives, the operators  $\partial_t$  and  $L$  can be extended to tensors (see Hamilton, [Ha]; cf. lemma 10.8 in [CM1]). We will next show that the Simons type identity (2.4) is in reality the trace of a tensor equation:

**Lemma 3.1.** *If  $M_t$  is a rescaled MCF and  $L = \Delta - \langle \frac{x}{2}, \nabla \cdot \rangle + |A|^2 + \frac{1}{2}$ , then*

$$(3.2) \quad (\partial_t - L) A = -A,$$

$$(3.3) \quad (\partial_t - L) H = -H,$$

$$(3.4) \quad (\partial_t - L) \langle x, \mathbf{n} \rangle = -2H,$$

$$(3.5) \quad (\partial_t - L) e^t \left( A - \frac{\langle x, \mathbf{n} \rangle}{2n} g \right) = \frac{e^t}{n} \left( H - \frac{\langle x, \mathbf{n} \rangle}{2} \right) g.$$

The first equation is equivalent to  $(\partial_t - L) e^t A = 0$ .

*Proof.* Given a hypersurface and an orthonormal frame  $\{e_i\}$ , we follow the convention in [CM1] by setting

$$(3.6) \quad a_{ij} = A(e_i, e_j) = \langle \nabla_{e_i} e_j, \mathbf{n} \rangle,$$

so that  $H = -\sum_{i=1}^n a_{ii}$ . For a general hypersurface, the Laplacian of  $A$  is (see, e.g., lemma B.8 in [Si] where the sign convention for the  $a_{ij}$ 's is reversed)

$$(3.7) \quad (\Delta A)_{ij} = -|A|^2 a_{ij} - H a_{ik} a_{jk} - H_{ij}.$$

Here  $H_{ij}$  is the  $ij$  component of the Hessian of  $H$ .

If a hypersurface evolves by  $\partial_t x = f \mathbf{n}$  for a function  $f$ , then we have the following standard formulas for the variations of the components of the metric  $g_{ij}$ , the components of the second fundamental form  $a_{ij}$ , and the unit normal  $\mathbf{n}$  (see, e.g., lemmas 7.4 and 7.6 in [HP]):

$$(3.8) \quad \partial_t g_{ij} = -2f a_{ij},$$

$$(3.9) \quad \partial_t \mathbf{n} = -\nabla f,$$

$$(3.10) \quad \partial_t a_{ij} = f_{ij} - f a_{ik} a_{kj}.$$

In the rescaled MCF, we have  $f = \langle x, \mathbf{n} \rangle / 2 - H$ .

We will extend  $\partial_t$  to tensors by using the covariant derivative  $\nabla_{\partial_t}$  in the metric  $g_{ij} \times dt^2$  (cf. [Ha] or the appendix in [W4]). If we let  $e_i$  be the evolving frame on  $M_t$  coming from pushing forward the frame on  $M_0$ , then the formula for the Christoffel symbols gives

$$(3.11) \quad \partial_t e_i = \frac{1}{2} g^{jk} (\partial_t g_{ij}) e_k = -f g^{jk} a_{ij} e_k.$$

Using the Leibniz rule, we have

$$(3.12) \quad \partial_t g_{ij} = \partial_t (g(e_i, e_j)) = (\partial_t g)(e_i, e_j) + g(\partial_t e_i, e_j) + g(\partial_t e_j, e_i),$$

so we see that this choice makes  $\partial_t g = 0$ .

Similarly, working at a point where the  $e_i$ 's are orthonormal, the Leibniz rule gives that

$$(3.13) \quad \begin{aligned} (\partial_t A)(e_i, e_j) &= \partial_t a_{ij} - A(\partial_t e_i, e_j) - A(e_i, \partial_t e_j) \\ &= (f_{ij} - f a_{ik} a_{kj}) - 2(-f a_{ik} a_{kj}) = f_{ij} + f a_{ik} a_{kj}. \end{aligned}$$

Combining this with (3.7), we compute the heat operator on  $A$ :

$$\begin{aligned}
(\partial_t A - \Delta A)(e_i, e_j) &= f_{ij} + f a_{ik} a_{kj} + |A|^2 a_{ij} + H a_{ik} a_{kj} + H_{ij} \\
(3.14) \qquad \qquad \qquad &= (f + H)_{ij} + (f + H) a_{ik} a_{kj} + |A|^2 a_{ij} \\
&= \frac{1}{2} \langle x, \mathbf{n} \rangle_{ij} + \frac{1}{2} \langle x, \mathbf{n} \rangle a_{ik} a_{kj} + |A|^2 a_{ij}.
\end{aligned}$$

Rewriting this in terms of the  $L$  operator (using  $x^T$  for the tangential part of  $x$ ) gives

$$\begin{aligned}
(\partial_t A - LA)(e_i, e_j) &= (\partial_t A - \Delta A)(e_i, e_j) \\
&\quad + \left( \frac{1}{2} \nabla_{x^T} A - \frac{1}{2} A - |A|^2 A \right)(e_i, e_j) \\
(3.15) \qquad \qquad \qquad &= \frac{1}{2} \{ \langle x, \mathbf{n} \rangle_{ij} + \langle x, \mathbf{n} \rangle a_{ik} a_{kj} + (\nabla_{x^T} A)(e_i, e_j) - a_{ij} \}.
\end{aligned}$$

The next step is to compute the operators on  $\langle x, \mathbf{n} \rangle$ . To simplify the calculation, let  $e_i$  be an orthonormal frame and work at a point where its tangential covariant derivatives vanish. In particular, at this point we have  $\nabla_{e_i} e_j = a_{ij} \mathbf{n}$  and  $\nabla_{e_i} \mathbf{n} = -a_{ij} e_j$ . Using this and the general fact that  $\nabla_{e_i} x = e_i$  (cf. the proof of lemma 5.5 in [CM1]), we get at this point

$$(3.16) \qquad \langle x, \mathbf{n} \rangle_i = \nabla_{e_i} \langle x, \mathbf{n} \rangle = \langle e_i, \mathbf{n} \rangle + \langle x, \nabla_{e_i} \mathbf{n} \rangle = -A(x^T, e_i).$$

To compute the Hessian, we will differentiate this. Using first the Leibniz rule, then the fact that  $\nabla A$  is fully symmetric in all three inputs by the Codazzi equations and the fact that

$$(3.17) \qquad \langle \nabla_{e_j} x^T, e_k \rangle = \langle \nabla_{e_j} x, e_k \rangle - \langle \nabla_{e_j} x^\perp, e_k \rangle = \delta_{jk} + a_{jk} \langle x, \mathbf{n} \rangle,$$

we get that

$$\begin{aligned}
\langle x, \mathbf{n} \rangle_{ij} &= -(\nabla_{e_j} A)(x^T, e_i) - A((\nabla_{e_j} x^T)^T, e_i) - A(x^T, \nabla_{e_j} e_i) \\
(3.18) \qquad \qquad &= -(\nabla_{x^T} A)(e_i, e_j) - A(e_i, e_j) - a_{ij} a_{kj} \langle x, \mathbf{n} \rangle.
\end{aligned}$$

Using (3.18) in (3.15) gives the first claim. The second claim follows from the first since traces commute with covariant derivatives ( $g$  is parallel, even with respect to time). The third claim follows from the second claim and the Simons equation (2.4) for  $H - \frac{1}{2} \langle x, \mathbf{n} \rangle$ .

For the last claim, we use that  $(\partial_t - L) e^t A = 0$  by the first claim and, since the metric is time-parallel, the third claim gives that

$$\begin{aligned}
e^{-t} (\partial_t - L) (e^t \langle x, \mathbf{n} \rangle g) &= \langle x, \mathbf{n} \rangle g + ((\partial_t - L) \langle x, \mathbf{n} \rangle) g \\
(3.19) \qquad \qquad \qquad &= (\langle x, \mathbf{n} \rangle - 2H) g.
\end{aligned}$$

q.e.d.

**3.1. Bounding  $A$ .** In this subsection, we use the matrix equation in combination with the parabolic maximum principle to bound  $A$  for the rescaled MCF by the rescaled mean curvature when the latter is non-negative.

To achieve this bound, we will need some simple computations. First we will need the quotient rule. Let  $f$  be a symmetric 2-tensor and  $h$  a positive function

$$(3.20) \quad (\partial_t - \Delta) \frac{f}{h} = \frac{(\partial_t - \Delta) f}{h} - \frac{f(\partial_t - \Delta) h}{h^2} + 2 \frac{\nabla_{\nabla h} f}{h^2} - 2 \frac{f |\nabla h|^2}{h^3} \\ = \frac{(\partial_t - \Delta) f}{h} - \frac{f(\partial_t - \Delta) h}{h^2} + \frac{2}{h} \nabla_{\nabla h} \left( \frac{f}{h} \right).$$

Next let  $V$  be a vector field, and set  $\mathcal{L} f = \Delta f + \nabla_V f$ ; we will also use that

$$(3.21) \quad (\partial_t - \mathcal{L}) |f|^2 = 2 \langle f, (\partial_t - \mathcal{L}) f \rangle - 2 |\nabla f|^2 \leq 2 \langle f, (\partial_t - \mathcal{L}) f \rangle,$$

$$(3.22) \quad (\partial_t - \mathcal{L}) \left| \frac{f}{h} \right|^2 \leq 2 \left\langle \frac{f}{h}, (\partial_t - \mathcal{L}) \frac{f}{h} \right\rangle,$$

$$(3.23) \quad (\partial_t - \mathcal{L}) \frac{f}{h} = \frac{(\partial_t - \mathcal{L}) f}{h} - \frac{f(\partial_t - \mathcal{L}) h}{h^2} + \frac{2}{h} \nabla_{\nabla h} \left( \frac{f}{h} \right).$$

In particular, if  $L f = \mathcal{L} f + K f$  for some function  $K$ ,  $(\partial_t - L) f = 0$ , and  $(\partial_t - L) h = 0$ , then the two previous equations give

$$(3.24) \quad (\partial_t - \mathcal{L}) \left| \frac{f}{h} \right|^2 \leq 2 \left\langle \frac{f}{h}, (\partial_t - \mathcal{L}) \frac{f}{h} \right\rangle = \frac{4}{h} \left\langle \nabla_{\nabla h} \left( \frac{f}{h} \right), \frac{f}{h} \right\rangle \\ = \frac{2}{h} \left\langle \nabla \left| \frac{f}{h} \right|^2, \nabla h \right\rangle = 2 \left\langle \nabla \left| \frac{f}{h} \right|^2, \nabla \log h \right\rangle.$$

Applying the above to  $f = e^t A$  and  $h = H - \frac{\langle x, \mathbf{n} \rangle}{2}$  and using Lemma 3.1 yields the following differential inequality for the ratio

$$(3.25) \quad B = \frac{f}{h} = \frac{e^t A}{H - \frac{\langle x, \mathbf{n} \rangle}{2}}.$$

**Lemma 3.26.** *If  $M_t \subset \mathbf{R}^{n+1}$  are hypersurfaces flowing by the rescaled MCF,  $H - \frac{\langle x, \mathbf{n} \rangle}{2} > 0$  on the initial hypersurface, and  $B$  is as above, then*

$$(3.27) \quad (\partial_t - \mathcal{L}) |B|^2 \leq 2 \left\langle \nabla |B|^2, \nabla \log \left( H - \frac{\langle x, \mathbf{n} \rangle}{2} \right) \right\rangle.$$

The parabolic maximum principle now implies the following:

**Corollary 3.28.** *If  $M_t \subset \mathbf{R}^{n+1}$  are closed hypersurfaces flowing by the rescaled MCF with  $H - \frac{\langle x, \mathbf{n} \rangle}{2} > 0$  on the initial hypersurface, then*

$$(3.29) \quad |A|^2 \leq C e^{-2t} \left| H - \frac{\langle x, \mathbf{n} \rangle}{2} \right|^2,$$



for some constant  $C$  depending on the initial hypersurface.

*Proof.* This follows from the parabolic maximum principle and Lemma 3.26. q.e.d.

**3.2. Curvature bounds for blow ups.** The next lemma establishes that  $H$  bounds  $|A|$  on the regular part of any multiplicity one tangent flow when the initial hypersurface is closed and has positive rescaled mean curvature. In particular, any such tangent flow has  $H \geq 0$ .

**Lemma 3.30.** *Let  $M_t \subset \mathbf{R}^{n+1}$  be closed hypersurfaces flowing by the rescaled MCF with  $H - \frac{\langle x, \mathbf{n} \rangle}{2} \geq 0$  on the initial hypersurface  $M_0$ . There exists  $C > 0$  so that if  $T_t$  is a tangent flow at the first singular time and  $T_{-1}$  is multiplicity one, then:*

- $|A| \leq C H$  on the regular set  $\text{Reg}(T_{-1})$ .

*Proof.* Let  $\tau$  be the first singular time and  $y \in M_\tau$  the singular point.

By Lemma 2.3, every  $M_t$  has  $H - \frac{1}{2} \langle x, \mathbf{n} \rangle \geq 0$ , so the flow is nested and

$$(3.31) \quad \frac{1}{2} |\langle x, \mathbf{n} \rangle| \leq \frac{|x|}{2} \leq C_1 \equiv \frac{1}{2} \max_{M_0} |x|.$$

Corollary 3.28 gives a constant  $C_0 > 0$  depending on the initial hypersurface  $M_0$  so that

$$(3.32) \quad |A| \leq C_0 e^{-t} \left( H - \frac{\langle x, \mathbf{n} \rangle}{2} \right) \leq C_0 \left( H - \frac{\langle x, \mathbf{n} \rangle}{2} \right) \leq C_0 H + C_0 C_1.$$

Fix a compact subset  $\Omega \subset \text{Reg}(T_{-1})$ . Since  $\Omega$  is smooth and multiplicity one, White's version of Brakke's regularity theorem, [W2] (cf. Allard's theorem; see [AI] or [Si]), gives a sequence  $h_i \rightarrow 0$  so that (a subset of)

$$(3.33) \quad \Sigma_i \equiv \frac{1}{h_i} \left( M_{\tau-h_i^2} - y \right)$$

smoothly converges to  $\Omega$ . Let  $A_i$  and  $H_i$  be the second fundamental form and mean curvature, respectively, of  $\Sigma_i$ . It follows from (3.32) that

$$(3.34) \quad |A_i| = h_i |A| \leq h_i (C_0 H + C_0 C_1) = C_0 H_i + h_i C_0 C_1.$$

Since we have smooth convergence to  $\Omega$  and  $h_i \rightarrow 0$ , we can pass to limits to get that

$$(3.35) \quad |A_\Omega| \leq C_0 H_\Omega.$$

The lemma follows since  $\Omega$  is arbitrary. q.e.d.

**Remark 3.36.** A similar statement holds for blow ups at the first singular time.

#### 4. Finite time blow up for rescaled MCF

The next result shows that there is finite time blow up for solutions of the rescaled MCF when the initial closed hypersurface has a strict positive lower bound for  $H - \frac{1}{2} \langle x, \mathbf{n} \rangle$ .

**Lemma 4.1.** *Given  $c > 0$ , there exists  $T_c$  so that if  $M_0$  is a closed hypersurface in  $\mathbf{R}^{n+1}$  with  $H - \frac{1}{2} \langle x, \mathbf{n} \rangle \geq c$ , then the rescaled MCF  $M_t$  hits a singularity for  $t \leq T_c$ .*

*Proof.* Set  $\phi = H - \frac{1}{2} \langle x, \mathbf{n} \rangle$ . Let  $m(t)$  be the minimum of  $\phi$  at time  $t$ . By (2.5), we have

$$(4.2) \quad (\partial_t - \Delta) e^{-\frac{t}{2}} \phi \geq -\left\langle \frac{x}{2}, \nabla e^{-\frac{t}{2}} \phi \right\rangle,$$

so the parabolic maximum principle gives that the minimum of  $e^{-\frac{t}{2}} \phi$  is non-decreasing in time. This gives exponential growth of  $m(t)$ , i.e.,

$$(4.3) \quad m(t) \geq e^{\frac{t}{2}} m(0) \geq e^{\frac{t}{2}} c > 0.$$

Since  $M_t$  lies in a bounded set by Lemma 2.3 and  $|\mathbf{n}| = 1$ , we have a constant  $C$  so that  $|\frac{1}{2} \langle x, \mathbf{n} \rangle| \leq |x|/2 \leq C$  for all  $t \geq 0$ . Combining this with (4.3), we can choose  $T_1 > 0$  so that if  $t \geq T_1$ , then  $H \geq \frac{\phi}{2}$ . In particular, when  $t \geq T_1$ , this means that

$$(4.4) \quad n |A|^2 \geq H^2 \geq \frac{\phi^2}{4}.$$

Using this in (2.5), we get for  $t \geq T_1$  that

$$(4.5) \quad (\partial_t - \Delta) \phi = \left( |A|^2 + \frac{1}{2} \right) \phi - \frac{1}{2} \langle x, \nabla \phi \rangle \geq \frac{1}{4n} \phi^3 - \frac{1}{2} \langle x, \nabla \phi \rangle.$$

Using this, the parabolic maximum principle gives the differential inequality (for  $t \geq T_1$ )

$$(4.6) \quad m'(t) \geq \frac{1}{4n} m^3(t).$$

Comparing this with the ODE  $f'(t) \geq \frac{1}{4n} f^3(t)$ , this gives finite time blow up for  $\phi$  which implies the finite time blow up for  $|A|$ , giving the desired singularity. q.e.d.

#### 5. Tangent cones to self-shrinkers

In this section, we will study two types of  $n$ -dimensional rectifiable integral varifolds. The first are weak solutions of the self-shrinker equation (0.4); these are critical points for the  $F$  functional and are called  $F$ -stationary. The second are stationary with respect to the Euclidean volume and will simply be called stationary; these arise as blow ups of the first.

The singular set and regular set of a rectifiable varifold  $\Sigma$  are denoted by  $\text{Sing}(\Sigma)$  and  $\text{Reg}(\Sigma)$ , respectively.

The main result of this section is the following:

**Proposition 5.1.** *If  $\Sigma \subset \mathbf{R}^{n+1}$  is an  $F$ -stationary rectifiable varifold,  $\lambda(\Sigma) < \frac{3}{2}$ , and there is a constant  $C > 0$  so that*

$$(5.2) \quad |A| \leq C H \text{ on the regular set } \text{Reg}(\Sigma),$$

*then  $\Sigma$  is smooth.*

We will use a blow up analysis, where we analyze the tangent cones, to prove the proposition. It follows from [A1] (cf. section 42 in [Si]) that an integral  $n$ -rectifiable varifold has stationary integral rectifiable tangent cones as long as the generalized mean curvature  $H$  is locally in  $L^p$  for some  $p > n$ . This is trivially satisfied for stationary varifolds where  $H = 0$  and for  $F$ -stationary varifolds where  $H$  is locally in  $L_\infty$ .

**Lemma 5.3.** *Suppose that  $\Sigma \subset \mathbf{R}^{n+1}$  is an  $F$ -stationary rectifiable varifold satisfying (5.2). If  $\Gamma$  is any multiplicity one blow up of  $\Sigma$ , then  $\Gamma$  is stationary and  $|A| \equiv 0$  on  $\text{Reg}(\Gamma)$ .*

*Proof.* By definition, there are sequences  $s_i \rightarrow 0$  and  $y_i \rightarrow y \in \mathbf{R}^{n+1}$  so that

$$(5.4) \quad \Sigma_i \equiv \frac{1}{s_i} (\Sigma - y_i)$$

converges with multiplicity one to  $\Gamma$ . Hence, the regular part of each  $\Sigma_i$  satisfies  $|A_i| \leq C H_i$  with the constant  $C$  from (5.2) since this inequality is scale invariant. Since  $H = \frac{1}{2} \langle x, \mathbf{n} \rangle$  is locally bounded on  $\Sigma$ , the limit  $\Gamma$  is stationary.

Since  $\Gamma$  is multiplicity one, Allard's theorem (see [A1] or [Si]) implies that the  $\Sigma_i$ 's converge smoothly on the regular set  $\text{Reg}(\Gamma)$  and, thus,

$$(5.5) \quad |A_\Gamma| \leq C H_\Gamma \text{ on } \text{Reg}(\Gamma).$$

(Since the sequence of rescalings has locally bounded  $H$ , Allard gives uniform local  $C^{1,\alpha}$  estimate graphical estimates. Elliptic theory then gives uniform estimates on higher derivatives and, thus, smooth convergence.) Since  $\Gamma$  is stationary, we have  $H = 0$  on  $\text{Reg}(\Gamma)$  and the lemma follows. q.e.d.

Tangent cones are limits of rescalings about a fixed point. An iterated tangent cone is a tangent cone to a tangent cone to a tangent cone ... (with finitely many iterations).

**Corollary 5.6.** *Let  $\Sigma \subset \mathbf{R}^{n+1}$  be an  $F$ -stationary rectifiable varifold satisfying (5.2). If  $\lambda(\Sigma) < 2$ , then every iterated tangent cone  $\Gamma$  is stationary, has  $\lambda(\Gamma) \leq \lambda(\Sigma)$ , has multiplicity one, and satisfies  $|A| \equiv 0$  on  $\text{Reg}(\Gamma)$ .*

*Proof.* This follows from Lemma 5.3 since iterated tangent cones can be realized as blow ups by taking a diagonal sequence and any limit has multiplicity one since  $\lambda(\Sigma) < 2$ . q.e.d.

We will need two elementary lemmas from blow up analysis:

**Lemma 5.7.** *Suppose that  $\Gamma \subset \mathbf{R}^{n+1}$  is an  $n$ -dimensional stationary rectifiable varifold with  $|A| \equiv 0$  on  $\text{Reg}(\Gamma)$ . If  $\Gamma$  is a cone,  $\text{Sing}(\Gamma) \subset \{0\}$ , and  $n > 1$ , then  $\text{Sing}(\Gamma) = \emptyset$ .*

*Proof.* Since  $|A| \equiv 0$  on the regular set, the closure of each connected component of  $\text{Reg}(\Gamma)$  is contained in a hyperplane. Since 0 is the only singular point and  $\Gamma$  is a cone, the closure of each connected component of  $\text{Reg}(\Gamma)$  must in fact be a hyperplane through 0. However, since two distinct hyperplanes in  $\mathbf{R}^{n+1}$  with 0 in their closure always intersect away from 0 (since  $n > 1$ ),  $\Gamma$  consists of a single hyperplane through the origin and thus is, in particular, smooth, though of course it could have multiplicity. q.e.d.

The above lemma does not extend to  $n = 1$ ; in particular, three half-lines meeting at 0 with angles  $\frac{2\pi}{3}$  is a stationary configuration. Thus something else, like an entropy bound, is needed to rule out such a singularity. Also, as noted, it could have multiplicity.

The second elementary lemma that we will need is often implicitly used in Federer type dimension reduction arguments. It is the following:

**Lemma 5.8.** *Suppose that  $\Gamma \subset \mathbf{R}^{n+1}$  is a stationary rectifiable varifold and  $\Gamma$  is a cone. If  $y \in \text{Sing}(\Gamma) \setminus \{0\}$ , then every tangent cone to  $\Gamma$  at  $y$  is of the form  $\Gamma' \times \mathbf{R}_y$  where  $\mathbf{R}_y$  is the line in the direction  $y$  and  $\Gamma'$  is a conical stationary varifold in  $\mathbf{R}^n$ .*

*Proof of Proposition 5.1.* By Allard's regularity theorem, it is enough to show that every tangent cone to  $\Sigma$  is a multiplicity one hyperplane. We will prove this by contradiction. Suppose, therefore, that  $x \in \text{Sing}(\Sigma)$ ,  $\Gamma_1$  is a tangent cone to  $\Sigma$  at  $x$ , and  $\Gamma_1$  is not a multiplicity one hyperplane.

By assumption,  $\lambda(\Sigma) < 3/2$  and  $\Sigma$  satisfies (5.2); therefore, Corollary 5.6 implies that  $\Gamma_1$  is stationary, satisfies  $|A| \equiv 0$  on  $\text{Reg}(\Gamma_1)$ , and

$$(5.9) \quad \lambda(\Gamma_1) \leq \lambda(\Sigma) < 3/2.$$

Since  $\Gamma_1$  is not a hyperplane ( $x$  is singular), Lemma 5.7 gives a singular point  $y_1 \in \text{Sing}(\Gamma_1) \setminus \{0\}$ . Thus, Lemma 5.8 gives a multiplicity one tangent cone  $\Gamma_2$  to  $\Gamma_1$  at  $y_1$  with

$$(5.10) \quad \Gamma_2 = \Gamma'_2 \times \mathbf{R}_{y_1},$$

where  $\Gamma'_2$  is a stationary cone in  $\mathbf{R}^n$ . Since  $y_1$  was a singular point,  $\Gamma'_2$  is not a hyperplane. Furthermore, since Corollary 5.6 applies to all iterated tangent cones, we know that  $\Gamma_2$ , and thus also  $\Gamma'_2$  have  $|A| \equiv 0$  on their regular parts.

We can now repeat the argument to get a stationary cone  $\Gamma'_3 \subset \mathbf{R}^{n-1}$  that is multiplicity one, is not a hyperplane, and also has  $|A| \equiv 0$  on its regular part.

Repeating this  $n-1$  times eventually gives a stationary cone  $\Gamma'_n \subset \mathbf{R}^2$  that is not a line and where  $\Gamma'_n \times \mathbf{R}^{n-1}$  is an iterated tangent cone for  $\Gamma$ . In particular, since entropy is preserved under products with  $\mathbf{R}$ , we must have that

$$(5.11) \quad \lambda(\Gamma'_n) < 3/2.$$

This implies that  $\Gamma'_n$  consists of at most two rays through the origin. However, the only such configuration that is stationary is when there are exactly two rays meeting at angle  $\pi$  to form a line. This contradiction completes the proof. q.e.d.

## 6. Low entropy singularities for mean convex rescaled MCF

An important tool in this paper is a classification of singularities for rescaled MCF starting from a closed rescaled mean convex hypersurface with low entropy. Here, “rescaled mean convex” means that (2.2) holds. This classification will be given in Corollary 6.4 below. The first step in this direction is the following regularity theorem and partial classification:

**Proposition 6.1.** *Let  $M_t$  be a rescaled MCF of closed hypersurfaces in  $\mathbf{R}^{n+1}$ , satisfying (2.2) on  $[0, T)$  and  $\lambda(M_0) < 3/2$ . If there is a singularity at  $(y, T)$ , then there is a tangent flow  $T_t$  with:*

- $T_{-1}$  is a smooth, embedded, self-shrinker with  $\lambda(T_{-1}) < 3/2$  and multiplicity one.
- $T_{-1}$  has  $H \geq 0$  and is not flat.

*Proof.* The blow up argument given in lemma 8 of [I1] (cf. [W1], [W3]) gives a tangent flow  $T_t$  so that  $T_{-1}$  is an  $F$ -stationary rectifiable varifold. Huisken’s monotonicity and the properties of the entropy in [CM1] (see lemma 1.11 and section 7 there) give that

$$(6.2) \quad \lambda(T_{-1}) \leq \lambda(M_0) < 3/2,$$

so that  $T_{-1}$  has multiplicity one and is embedded. Hence, by Lemma 3.30,

- $|A| \leq C H$  on the regular set  $\text{Reg}(T_{-1})$ .

Finally, Proposition 5.1 gives that  $M_{-1}$  is smooth. q.e.d.

We will combine Proposition 6.1 with the following classification of smooth, embedded mean convex self-shrinkers in arbitrary dimension from theorem 0.17 in [CM1]:

**Theorem 6.3** ([CM1]).  $\mathbf{S}^k \times \mathbf{R}^{n-k}$  are the only smooth complete embedded self-shrinkers without boundary, with polynomial volume growth, and  $H \geq 0$  in  $\mathbf{R}^{n+1}$ .

The  $\mathbf{S}^k$  factor in Theorem 6.3 is round and has radius  $\sqrt{2k}$ ; we allow the possibilities of a sphere ( $n - k = 0$ ) or a hyperplane (i.e.,  $k = 0$ ), although Brakke's theorem rules out the multiplicity one hyperplane as a tangent flow at a singular point.

The classification of smooth embedded self-shrinkers with  $H \geq 0$  began with [H1], where Huisken showed that round spheres are the only closed ones. In [H2], Huisken showed that generalized cylinders  $\mathbf{S}^k \times \mathbf{R}^{n-k} \subset \mathbf{R}^{n+1}$  are the only open ones with polynomial volume growth and  $|A|$  bounded. Theorem 0.17 in [CM1] completed the classification by removing the  $|A|$  bound.

Combining Proposition 6.1 with Theorem 6.3 gives:

**Corollary 6.4.** *Let  $M_t$  be a rescaled MCF of closed hypersurfaces in  $\mathbf{R}^{n+1}$ , satisfying (2.2) on  $[0, T)$  and  $\lambda(M_0) < 3/2$ . If there is a singularity at  $(y, T)$ , then there is a multiplicity one tangent flow  $T_t$  of the form  $\mathbf{S}^k \times \mathbf{R}^{n-k}$  for some  $k > 0$ .*

Combining this with the monotonicity of the entropy will give the following lower bound for entropy and topological rigidity for rescaled mean convex hypersurfaces:

**Proposition 6.5.** *Let  $M^n$  be a closed hypersurface satisfying (2.2). If  $\lambda(M) < \min\{\lambda(\mathbf{S}^{n-1}), \frac{3}{2}\}$ , then  $M$  is diffeomorphic to  $\mathbf{S}^n$  and  $\lambda(M) \geq \lambda(\mathbf{S}^n)$ .*

*Proof.* We can assume that  $H - \frac{1}{2}\langle x, \mathbf{n} \rangle > 0$  at least at one point. To see this, suppose instead that  $H - \frac{1}{2}\langle x, \mathbf{n} \rangle \equiv 0$ , so that  $M$  is a self-shrinker. If  $M = \mathbf{S}^n$ , then we are done. Otherwise, Lemma 1.2 gives a nearby graph  $\tilde{M}$  over  $M$  with  $H - \frac{1}{2}\langle x, \mathbf{n} \rangle > 0$  and  $\lambda(\tilde{M}) < \lambda(M)$ .

Let  $M_t$  be the rescaled MCF with  $M_0 = M$ , so that Lemma 2.3 gives

$$(6.6) \quad H - \frac{1}{2}\langle x, \mathbf{n} \rangle > 0 \text{ for all } t > 0.$$

Lemmas 4.1 and 2.3 give a singularity in finite time inside of  $M$ . Thus, Corollary 6.4 gives a multiplicity one tangent flow  $T_t$  at this point of the form  $\mathbf{S}^k \times \mathbf{R}^{n-k}$  for some  $k > 0$ . By the monotonicity of the entropy under MCF (lemma 1.11 in [CM1]), its invariance under dilation, and its lower semi-continuity under limits, we have

$$(6.7) \quad \lambda(T_{-1}) \leq \lambda(M) < \min\{\lambda(\mathbf{S}^{n-1}), 3/2\}.$$

By [St], we have  $\lambda(\mathbf{S}^n) < \lambda(\mathbf{S}^{n-1} \times \mathbf{R}) < \dots$ , so we conclude that  $k = n$  and  $\lambda(\mathbf{S}^n) \leq \lambda(M)$ . Finally, White's version, [W2], of Brakke's theorem implies that  $T_{-1}$  is the smooth limit of rescalings of the  $M_t$ s. In particular,  $M$  itself is diffeomorphic to  $T_{-1} = \mathbf{S}^n$ . q.e.d.

## 7. Proof of the main theorem

*Proof of Theorem 0.7.* Let  $\Sigma$  be a closed self-shrinker in  $\mathbf{R}^{n+1}$  with  $\lambda(\Sigma) < \min\{\lambda(\mathbf{S}^{n-1}), 3/2\}$ . By Proposition 6.5,  $\tilde{\Sigma}$  is diffeomorphic to  $\mathbf{S}^n$  and  $\lambda(\tilde{\Sigma}) \geq \lambda(\mathbf{S}^n)$ .

We will show that there is a gap, i.e., some  $\epsilon > 0$  so that  $\lambda(\Sigma) \geq \lambda(\mathbf{S}^n) + \epsilon$  if  $\Sigma$  is not round. Suppose instead that there is a sequence of closed self-shrinkers  $\Sigma_i \neq \mathbf{S}^n$  with

$$(7.1) \quad \lambda(\mathbf{S}^n) \leq \lambda(\Sigma_i) < \lambda(\mathbf{S}^n) + 2^{-i}.$$

By Huisken, [H1], each  $\Sigma_i$  has negative mean curvature at some point since they are not round.

Perturbing the  $\Sigma_i$ 's with Lemma 1.2 and then applying rescaled MCF to the perturbations gives a sequence of rescaled MCFs  $\tilde{M}_{i,t}$  so that

- Each initial hypersurface  $\tilde{M}_{i,0}$  has  $\lambda(\tilde{M}_{i,0}) < \lambda(\Sigma_i)$  and has negative mean curvature at some point.
- Each  $\tilde{M}_{i,t}$  has a multiplicity one spherical singularity in finite time.

The second claim follows from Lemma 4.1.

We can now create a new sequence of rescaled MCFs  $M_{i,t}$  by rescaling the  $\tilde{M}_{i,t}$ s about the spherical singularity so that the new flows satisfy:

- 1) Each  $M_{i,t}$  converges smoothly to the round sphere as  $t \rightarrow \infty$ .
- 2) Each initial hypersurface  $M_{i,0}$  is a  $C^2$  graph over  $\mathbf{S}^n$  with  $C^2$  norm exactly  $\epsilon > 0$ .
- 3) Every hypersurface  $M_{i,t}$  for  $t \geq 0$  is a  $C^2$  graph over  $\mathbf{S}^n$  with  $C^2$  norm at most  $\epsilon$ .

By (3), we can choose a subsequence that converges smoothly to a limiting rescaled MCF  $M_t$ . Now consider the  $F$  functional along  $M_t$ . By (1), we have

$$(7.2) \quad \lim_{t \rightarrow \infty} F(M_t) = \lambda(\mathbf{S}^n).$$

On the other hand, we know that

$$(7.3) \quad F(M_0) \leq \lambda(M_0) \leq \liminf \lambda(\Sigma_i) = \lambda(\mathbf{S}^n).$$

Since the rescaled MCF is the gradient flow for  $F$ , we see that the flow must be constant and, thus, every  $M_t$  is round. This contradicts (2) which says that the initial hypersurfaces are strictly away from the round sphere, completing the proof. q.e.d.

## References

- [AbL] U. Abresch & J. Langer, *The normalized curve shortening flow and homothetic solutions*, JDG **23** (1986), 175–196, MR 0845704, Zbl 0592.53002.
- [Al] W.K Allard, *On the first variation of a varifold*. Ann. of Math. (2) **95** (1972), 417–491, MR 0307015, Zbl 0252.49028.

- [A] S.B. Angenent, *Shrinking doughnuts*, in: *Nonlinear diffusion equations and their equilibrium states*, Birkhäuser, Boston-Basel-Berlin, 3, 21–38, 1992, MR 1167827, Zbl 0762.53028.
- [B] K. Ball, *The reverse isoperimetric problem for Gaussian measure*, *Discrete Comput. Geom.* **10** (1993), 411–420, MR 1243336, Zbl 0788.52010.
- [Br] K. Brakke, *The motion of a surface by its mean curvature*, *Mathematical Notes*, **20**, Princeton University Press, Princeton, N.J., 1978, MR 0485012, Zbl 0386.53047.
- [Ch] D. Chopp, *Computation of self-similar solutions for mean curvature flow*, *Experiment. Math.* **3** (1994), no. 1, 1–15, MR 1302814, Zbl 0811.53011.
- [CM1] T.H. Colding & W.P. Minicozzi II, *Generic mean curvature flow I; generic singularities*, *Annals of Math.*, **175** (2012), 755–833, MR 2993752, Zbl 1239.53084.
- [CM2] ———, *Generic mean curvature flow II; dynamics of a closed smooth singularity*, in preparation.
- [CM3] ———, *Smooth compactness of self-shrinkers*, *Comm. Math. Helv.* **87** (2012), 463–487, MR 2914856, Zbl 1258.53069.
- [GaHa] M. Gage & R. Hamilton, *The heat equation shrinking convex plane curves*, *JDG* **23** (1986), 69–96, MR 0840401, Zbl 0621.53001.
- [G] M. Grayson, *The heat equation shrinks embedded plane curves to round points*, *JDG* **26** (1987), 285–314, MR 0906392, Zbl 0667.53001.
- [Ha] R.S. Hamilton, *Harnack estimate for the mean curvature flow*. *JDG* **41** (1995), 215–226, MR 1316556, Zbl 0827.53006.
- [H1] G. Huisken, *Asymptotic behavior for singularities of the mean curvature flow*. *JDG* **31** (1990), 285–299, MR 1030675, Zbl 0694.53005.
- [H2] ———, *Local and global behaviour of hypersurfaces moving by mean curvature*. *Differential geometry: partial differential equations on manifolds* (Los Angeles, CA, 1990), 175–191, *Proc. Sympos. Pure Math.*, **54**, Part 1, Amer. Math. Soc., Providence, RI, 1993, MR 1216584, Zbl 0791.58090.
- [HP] G. Huisken & A. Polden, *Geometric evolution equations for hypersurfaces*. *Calculus of variations and geometric evolution problems*, 45–84, *Lecture Notes in Math.*, 1713, Springer, Berlin, 1999, MR 1731639, Zbl 0942.35047.
- [HS1] G. Huisken & C. Sinestrari, *Mean curvature flow singularities for mean convex surfaces*, *Calc. Var. Partial Differ. Equ.* **8** (1999) 1–14, MR 1666878, Zbl 0992.53052.
- [HS2] ———, *Convexity estimates for mean curvature flow and singularities of mean convex surfaces*, *Acta Math.* **183** (1999), no. 1, 45–70, MR 1719551, Zbl 0992.53051.
- [I1] T. Ilmanen, *Singularities of Mean Curvature Flow of Surfaces*, preprint, 1995, <http://www.math.ethz.ch/~papers/pub.html>.
- [I2] ———, *Lectures on Mean Curvature Flow and Related Equations*, (Trieste Notes), 1995.
- [IW] T. Ilmanen & B. White, in preparation.
- [K] D.M. Kane. *The Gaussian Surface Area and Noise Sensitivity of Degree-d Polynomial Threshold Functions*, *Computational Complexity* **20** (2011) 389–412, MR 2822876, Zbl 1230.68169.



- [KDS] A. Klivans, R. O'Donnell & R. Servedio. *Learning geometric concepts via Gaussian surface area*, in Proc. 49th IEEE Symposium on Foundations of Computer Science FOCS '08.
- [KKM] N. Kapouleas, S. Kleene & N.M. Möller, *Mean curvature self-shrinkers of high genus: non-compact examples*, preprint, <http://arxiv.org/pdf/1106.5454>.
- [M] N.M. Möller, *Closed self-shrinking surfaces in  $\mathbf{R}^3$  via the torus*, <http://arxiv.org/pdf/1111.7318v1.pdf>.
- [Na] F. Nazarov, *On the maximal perimeter of a convex set in  $\mathbf{R}^n$  with respect to a Gaussian measure*. In Geometric aspects of functional analysis (2001-2002), pages 169–187. Lecture Notes in Math., Vol. 1807, Springer, 2003, MR 2083397, Zbl 1036.52014.
- [N] X.H. Nguyen, *Construction of Complete Embedded Self-Similar Surfaces under Mean Curvature Flow. Part III*, preprint, <http://arxiv.org/pdf/1106.5272v1>.
- [Si] L.M. Simon, *Lectures on Geometric Measure Theory*, Proc. of the CMA, ANU No. 3, Canberra, 1983, MR 0756417, Zbl 0546.49019.
- [St] A. Stone, *A density function and the structure of singularities of the mean curvature flow*, Calc. Var. **2** (1994), 443–480, MR 1383918, Zbl 0833.35062.
- [W1] B. White, *Stratification of minimal surfaces, mean curvature flows, and harmonic maps*. J. Reine Angew. Math. **488** (1997), 1–35, MR 1465365, Zbl 0874.58007.
- [W2] ———, *A local regularity theorem for mean curvature flow*, Ann. of Math. (2) **161** (2005), 1487–1519, MR 2180405, Zbl 1091.53045.
- [W3] ———, *The size of the singular set in mean curvature flow of mean-convex sets*. J. Amer. Math. Soc. **13** (2000), 665–695, MR 1758759, Zbl 0961.53039.
- [W4] ———, *The nature of singularities in mean curvature flow of mean-convex sets*, J. Amer. Math. Soc. **16** (2003), 123–138, MR 1937202, Zbl 1027.53078.

MIT DEPT. OF MATHEMATICS  
 77 MASSACHUSETTS AVENUE  
 CAMBRIDGE, MA 02139-4307, USA  
*E-mail address:* colding@math.mit.edu

DEPARTEMENT MATHEMATIK  
 ETH ZENTRUM  
 8092 ZÜRICH, SWITZERLAND  
*E-mail address:* tom.ilmanen@math.ethz.ch

MIT DEPT. OF MATHEMATICS  
 77 MASSACHUSETTS AVENUE  
 CAMBRIDGE, MA 02139-4307, USA  
*E-mail address:* minicozz@math.mit.edu

STANFORD UNIVERSITY  
 DEPT. OF MATHEMATICS  
 BLDG 380  
 STANFORD, CA 94305, USA  
*E-mail address:* white@math.stanford.edu