

quatrième série - tome 44 fascicule 1 janvier-février 2011

*ANNALES
SCIENTIFIQUES
de
L'ÉCOLE
NORMALE
SUPÉRIEURE*

Peng SHAN

*Crystals of Fock spaces and cyclotomic
rational double affine Hecke algebras*

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

CRYSTALS OF FOCK SPACES AND CYCLOTOMIC RATIONAL DOUBLE AFFINE HECKE ALGEBRAS

BY PENG SHAN

ABSTRACT. – We define the i -restriction and i -induction functors on the category \mathcal{O} of the cyclotomic rational double affine Hecke algebras. This yields a crystal on the set of isomorphism classes of simple modules, which is isomorphic to the crystal of a Fock space.

RÉSUMÉ. – On définit les foncteurs de i -restriction et i -induction sur la catégorie \mathcal{O} des algèbres de Hecke doublement affines rationnelles cyclotomiques. Ceci donne lieu à un cristal sur l'ensemble des classes d'isomorphismes de modules simples, qui est isomorphe au cristal d'un espace de Fock.

Introduction

In [1], S. Ariki defined the i -restriction and i -induction functors for cyclotomic Hecke algebras. He showed that the Grothendieck group of the category of finitely generated projective modules of these algebras admits a module structure over the affine Lie algebra of type $A^{(1)}$, with the action of Chevalley generators given by the i -restriction and i -induction functors.

The restriction and induction functors for rational DAHA's (= double affine Hecke algebras) were recently defined by R. Bezrukavnikov and P. Etingof. With these functors, we give an analogue of Ariki's construction for the category \mathcal{O} of cyclotomic rational DAHA's: we show that as a module over the type $A^{(1)}$ affine Lie algebra, the Grothendieck group of this category is isomorphic to a Fock space. We also construct a crystal on the set of isomorphism classes of simple modules in the category \mathcal{O} . It is isomorphic to the crystal of the Fock space. Recall that this Fock space also enters in some conjectural description of the decomposition numbers for the category \mathcal{O} considered here. See [16], [17], [14] for related works.

Notation

For A an algebra, we will write $A\text{-mod}$ for the category of finitely generated A -modules. For $f : A \rightarrow B$ an algebra homomorphism from A to another algebra B such that B is finitely generated over A , we will write

$$f_* : B\text{-mod} \rightarrow A\text{-mod}$$

for the restriction functor and we write

$$f^* : A\text{-mod} \rightarrow B\text{-mod}, \quad M \mapsto B \otimes_A M.$$

A \mathbb{C} -linear category \mathcal{C} is called artinian if the Hom sets are finite dimensional \mathbb{C} -vector spaces and every object has a finite length. Given an object M in \mathcal{C} , we denote by $\text{soc}(M)$ (resp. $\text{head}(M)$) the socle (resp. the head) of M , which is the largest semi-simple subobject (quotient) of M .

Let \mathcal{C} be an abelian category. The Grothendieck group of \mathcal{C} is the quotient of the free abelian group generated by objects in \mathcal{C} modulo the relations $M = M' + M''$ for all objects M, M', M'' in \mathcal{C} such that there is an exact sequence $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$. Let $K(\mathcal{C})$ denote the complexified Grothendieck group, a \mathbb{C} -vector space. For each object M in \mathcal{C} , let $[M]$ be its class in $K(\mathcal{C})$. Any exact functor $F : \mathcal{C} \rightarrow \mathcal{C}'$ between two abelian categories induces a vector space homomorphism $K(\mathcal{C}) \rightarrow K(\mathcal{C}')$, which we will denote by F again. Given an algebra A we will abbreviate $K(A) = K(A\text{-mod})$.

Denote by $\text{Fct}(\mathcal{C}, \mathcal{C}')$ the category of functors from a category \mathcal{C} to a category \mathcal{C}' . For $F \in \text{Fct}(\mathcal{C}, \mathcal{C}')$ write $\text{End}(F)$ for the ring of endomorphisms of the functor F . We denote by $1_F : F \rightarrow F$ the identity element in $\text{End}(F)$. Let $G \in \text{Fct}(\mathcal{C}', \mathcal{C}'')$ be a functor from \mathcal{C}' to another category \mathcal{C}'' . For any $X \in \text{End}(F)$ and any $X' \in \text{End}(G)$ we write $X'X : G \circ F \rightarrow G \circ F$ for the morphism of functors given by $X'X(M) = X'(F(M)) \circ G(X(M))$ for any $M \in \mathcal{C}$.

Let $e \geq 2$ be an integer and z be a formal parameter. Denote by \mathfrak{sl}_e the Lie algebra of traceless $e \times e$ complex matrices. The type $A^{(1)}$ affine Lie algebra is

$$\tilde{\mathfrak{sl}}_e = \mathfrak{sl}_e \otimes \mathbb{C}[z, z^{-1}] \oplus \mathbb{C}c \oplus \mathbb{C}\partial$$

equipped with the Lie bracket

$$[\xi \otimes z^m + ac + b\partial, \xi' \otimes z^n + a'c + b'\partial] = [\xi, \xi'] \otimes z^{m+n} + m\delta_{m,-n} \text{tr}(\xi\xi')c + nb\xi' \otimes z^n - mb'\xi \otimes z^m,$$

for $\xi, \xi' \in \mathfrak{sl}_e, a, a', b, b' \in \mathbb{C}$. Here $\text{tr} : \mathfrak{sl}_e \rightarrow \mathbb{C}$ is the trace map. Let

$$\widehat{\mathfrak{sl}}_e = \mathfrak{sl}_e \otimes \mathbb{C}[z, z^{-1}] \oplus \mathbb{C}c.$$

It is the Lie subalgebra of $\tilde{\mathfrak{sl}}_e$ generated by the Chevalley generators

$$\begin{aligned} e_i &= E_{i,i+1} \otimes 1, & f_i &= E_{i+1,i} \otimes 1, & 1 \leq i \leq e-1 \\ e_0 &= E_{e1} \otimes z, & f_0 &= E_{1e} \otimes z^{-1}. \end{aligned}$$

Here E_{ij} is the elementary matrix with 1 in the position (i, j) and 0 elsewhere. Let $h_i = [e_i, f_i]$ for $0 \leq i \leq e-1$. We consider the Cartan subalgebra

$$\mathfrak{t} = \bigoplus_{i \in \mathbb{Z}/e\mathbb{Z}} \mathbb{C}h_i \oplus \mathbb{C}\partial,$$

and its dual \mathfrak{t}^* . For $i \in \mathbb{Z}/e\mathbb{Z}$ let $\alpha_i \in \mathfrak{t}^*$ (resp. $\alpha_i^\vee \in \mathfrak{t}$) be the simple root (resp. coroot) corresponding to e_i . The fundamental weights are $\{\Lambda_i \in \mathfrak{t}^* : i \in \mathbb{Z}/e\mathbb{Z}\}$ such that $\Lambda_i(\alpha_j^\vee) = \delta_{ij}$ and $\Lambda_i(\partial) = 0$ for any $i, j \in \mathbb{Z}/e\mathbb{Z}$. Let $\delta \in \mathfrak{t}^*$ be the element given by $\delta(h_i) = 0$ for all i and $\delta(\partial) = 1$. We will write P for the weight lattice of $\widetilde{\mathfrak{sl}}_e$. It is the free abelian group generated by the fundamental weights and δ .

1. Reminders on Hecke algebras, rational DAHA's and restriction functors

1.1. Hecke algebras

Let \mathfrak{h} be a finite dimensional vector space over \mathbb{C} . Recall that a pseudo-reflection is a non trivial element s of $GL(\mathfrak{h})$ which acts trivially on a hyperplane, called the reflecting hyperplane of s . Let $W \subset GL(\mathfrak{h})$ be a finite subgroup generated by pseudo-reflections. Let \mathcal{J} be the set of pseudo-reflections in W and \mathcal{A} be the set of reflecting hyperplanes. We set $\mathfrak{h}_{\text{reg}} = \mathfrak{h} - \bigcup_{H \in \mathcal{A}} H$, it is stable under the action of W . Fix $x_0 \in \mathfrak{h}_{\text{reg}}$ and identify it with its image in $\mathfrak{h}_{\text{reg}}/W$. By definition the braid group attached to (W, \mathfrak{h}) , denoted by $B(W, \mathfrak{h})$, is the fundamental group $\pi_1(\mathfrak{h}_{\text{reg}}/W, x_0)$.

For any $H \in \mathcal{A}$, let W_H be the pointwise stabilizer of H . This is a cyclic group. Write e_H for the order of W_H . Let s_H be the unique element in W_H whose determinant is $\exp(\frac{2\pi\sqrt{-1}}{e_H})$. Let q be a map from \mathcal{J} to \mathbb{C}^* that is constant on the W -conjugacy classes. Following [6, Definition 4.21] the Hecke algebra $\mathcal{H}_q(W, \mathfrak{h})$ attached to (W, \mathfrak{h}) with parameter q is the quotient of the group algebra $\mathbb{C}B(W, \mathfrak{h})$ by the relations:

$$(1.1) \quad (T_{s_H} - 1) \prod_{t \in W_H \cap \mathcal{J}} (T_{s_H} - q(t)) = 0, \quad H \in \mathcal{A}.$$

Here T_{s_H} is a generator of the monodromy around H in $\mathfrak{h}_{\text{reg}}/W$ such that the lift of T_{s_H} in $\pi_1(W, \mathfrak{h}_{\text{reg}})$ via the map $\mathfrak{h}_{\text{reg}} \rightarrow \mathfrak{h}_{\text{reg}}/W$ is represented by a path from x_0 to $s_H(x_0)$. See [6, Section 2B] for a precise definition. When the subspace \mathfrak{h}^W of fixed points of W in \mathfrak{h} is trivial, we abbreviate

$$B_W = B(W, \mathfrak{h}), \quad \mathcal{H}_q(W) = \mathcal{H}_q(W, \mathfrak{h}).$$

1.2. Parabolic restriction and induction for Hecke algebras

In this section we will assume that $\mathfrak{h}^W = 1$. A parabolic subgroup W' of W is by definition the stabilizer of a point $b \in \mathfrak{h}$. By a theorem of Steinberg, the group W' is also generated by pseudo-reflections. Let q' be the restriction of q to $\mathcal{J}' = W' \cap \mathcal{J}$. There is an explicit inclusion $\iota_q : \mathcal{H}_{q'}(W') \hookrightarrow \mathcal{H}_q(W)$ given by [6, Section 2D]. The restriction functor

$${}^{\mathcal{H}}\text{Res}_{W'}^W : \mathcal{H}_q(W)\text{-mod} \rightarrow \mathcal{H}_{q'}(W')\text{-mod}$$

is the functor $(\iota_q)_*$. The induction functor

$${}^{\mathcal{H}}\text{Ind}_{W'}^W = \mathcal{H}_q(W) \otimes_{\mathcal{H}_{q'}(W')} -$$

is left adjoint to ${}^{\mathcal{H}}\text{Res}_{W'}^W$. The coinduction functor

$${}^{\mathcal{H}}\text{coInd}_{W'}^W = \text{Hom}_{\mathcal{H}_{q'}(W')}(\mathcal{H}_q(W), -)$$

is right adjoint to ${}^{\mathcal{H}}\text{Res}_{W'}^W$. The three functors above are all exact.

Let us recall the definition of ι_q . It is induced from an inclusion $\iota : B_{W'} \hookrightarrow B_W$, which is in turn the composition of three morphisms ℓ, κ, j defined as follows. First, let $\mathcal{O}' \subset \mathcal{O}$ be the set of reflecting hyperplanes of W' . Write

$$\bar{\mathfrak{h}} = \mathfrak{h}/\mathfrak{h}^{W'}, \quad \bar{\mathcal{O}} = \{\bar{H} = H/\mathfrak{h}^{W'} : H \in \mathcal{O}'\}, \quad \bar{\mathfrak{h}}_{\text{reg}} = \bar{\mathfrak{h}} - \bigcup_{\bar{H} \in \bar{\mathcal{O}}} \bar{H}, \quad \mathfrak{h}'_{\text{reg}} = \mathfrak{h} - \bigcup_{H \in \mathcal{O}'} H.$$

The canonical epimorphism $p : \mathfrak{h} \rightarrow \bar{\mathfrak{h}}$ induces a trivial W' -equivariant fibration $p : \mathfrak{h}'_{\text{reg}} \rightarrow \bar{\mathfrak{h}}_{\text{reg}}$, which yields an isomorphism

$$(1.2) \quad \ell : B_{W'} = \pi_1(\bar{\mathfrak{h}}_{\text{reg}}/W', p(x_0)) \xrightarrow{\sim} \pi_1(\mathfrak{h}'_{\text{reg}}/W', x_0).$$

Endow \mathfrak{h} with a W -invariant hermitian scalar product. Let $\|\cdot\|$ be the associated norm. Set

$$(1.3) \quad \Omega = \{x \in \mathfrak{h} : \|x - b\| < \varepsilon\},$$

where ε is a positive real number such that the closure of Ω does not intersect any hyperplane that is in the complement of \mathcal{O}' in \mathcal{O} . Let $\gamma : [0, 1] \rightarrow \mathfrak{h}$ be a path such that $\gamma(0) = x_0$, $\gamma(1) = b$ and $\gamma(t) \in \mathfrak{h}_{\text{reg}}$ for $0 < t < 1$. Let $u \in [0, 1[$ such that $x_1 = \gamma(u)$ belongs to Ω , write γ_u for the restriction of γ to $[0, u]$. Consider the homomorphism

$$\sigma : \pi_1(\Omega \cap \mathfrak{h}_{\text{reg}}, x_1) \rightarrow \pi_1(\mathfrak{h}_{\text{reg}}, x_0), \quad \lambda \mapsto \gamma_u^{-1} \cdot \lambda \cdot \gamma_u.$$

The canonical inclusion $\mathfrak{h}_{\text{reg}} \hookrightarrow \mathfrak{h}'_{\text{reg}}$ induces a homomorphism $\pi_1(\mathfrak{h}_{\text{reg}}, x_0) \rightarrow \pi_1(\mathfrak{h}'_{\text{reg}}, x_0)$. Composing it with σ gives an invertible homomorphism

$$\pi_1(\Omega \cap \mathfrak{h}_{\text{reg}}, x_1) \rightarrow \pi_1(\mathfrak{h}'_{\text{reg}}, x_0).$$

Since Ω is W' -invariant, its inverse gives an isomorphism

$$(1.4) \quad \kappa : \pi_1(\mathfrak{h}'_{\text{reg}}/W', x_0) \xrightarrow{\sim} \pi_1((\Omega \cap \mathfrak{h}_{\text{reg}})/W', x_1).$$

Finally, we see from above that σ is injective. So it induces an inclusion

$$\pi_1((\Omega \cap \mathfrak{h}_{\text{reg}})/W', x_1) \hookrightarrow \pi_1(\mathfrak{h}_{\text{reg}}/W', x_0).$$

Composing it with the canonical inclusion $\pi_1(\mathfrak{h}_{\text{reg}}/W', x_0) \hookrightarrow \pi_1(\mathfrak{h}_{\text{reg}}/W, x_0)$ gives an injective homomorphism

$$(1.5) \quad j : \pi_1((\Omega \cap \mathfrak{h}_{\text{reg}})/W', x_1) \hookrightarrow \pi_1(\mathfrak{h}_{\text{reg}}/W, x_0) = B_W.$$

By composing ℓ, κ, j we get the inclusion

$$(1.6) \quad \iota = j \circ \kappa \circ \ell : B_{W'} \hookrightarrow B_W.$$

It is proved in [6, Section 4C] that ι preserves the relations in (1.1). So it induces an inclusion of Hecke algebras which is the desired inclusion

$$\iota_q : \mathcal{H}_{q'}(W') \hookrightarrow \mathcal{H}_q(W).$$

For $\iota, \iota' : B_{W'} \hookrightarrow B_W$ two inclusions defined as above via different choices of the path γ , there exists an element $\rho \in P_W = \pi_1(\mathfrak{h}_{\text{reg}}, x_0)$ such that for any $a \in B_{W'}$ we have $\iota(a) = \rho \iota'(a) \rho^{-1}$. In particular, the functors ι_* and $(\iota')_*$ from $B_{W'}$ -mod to B_W -mod are isomorphic. Also, we have $(\iota_q)_* \cong (\iota'_q)_*$. So there is a unique restriction functor ${}^{\mathcal{H}_q} \text{Res}_{B_{W'}}^W$ up to isomorphisms.

1.3. Rational DAHA's

Let c be a map from \mathcal{J} to \mathbb{C} that is constant on the W -conjugacy classes. The rational DAHA attached to W with parameter c is the quotient $H_c(W, \mathfrak{h})$ of the smash product of $\mathbb{C}W$ and the tensor algebra of $\mathfrak{h} \oplus \mathfrak{h}^*$ by the relations

$$[x, x'] = 0, \quad [y, y'] = 0, \quad [y, x] = \langle x, y \rangle - \sum_{s \in \mathcal{J}} c_s \langle \alpha_s, y \rangle \langle x, \alpha_s^\vee \rangle s,$$

for all $x, x' \in \mathfrak{h}^*$, $y, y' \in \mathfrak{h}$. Here $\langle \cdot, \cdot \rangle$ is the canonical pairing between \mathfrak{h}^* and \mathfrak{h} , the element α_s is a generator of $\text{Im}(s|_{\mathfrak{h}^*} - 1)$ and α_s^\vee is the generator of $\text{Im}(s|_{\mathfrak{h}} - 1)$ such that $\langle \alpha_s, \alpha_s^\vee \rangle = 2$.

For $s \in \mathcal{J}$ write λ_s for the non trivial eigenvalue of s in \mathfrak{h}^* . Let $\{x_i\}$ be a basis of \mathfrak{h}^* and let $\{y_i\}$ be the dual basis. Let

$$(1.7) \quad \mathbf{eu} = \sum_i x_i y_i + \frac{\dim(\mathfrak{h})}{2} - \sum_{s \in \mathcal{J}} \frac{2c_s}{1 - \lambda_s} s$$

be the Euler element in $H_c(W, \mathfrak{h})$. Its definition is independent of the choice of the basis $\{x_i\}$. We have

$$(1.8) \quad [\mathbf{eu}, x_i] = x_i, \quad [\mathbf{eu}, y_i] = -y_i, \quad [\mathbf{eu}, s] = 0.$$

1.4. The category \mathcal{O}

The category \mathcal{O} of $H_c(W, \mathfrak{h})$ is the full subcategory $\mathcal{O}_c(W, \mathfrak{h})$ of the category of $H_c(W, \mathfrak{h})$ -modules consisting of objects that are finitely generated as $\mathbb{C}[\mathfrak{h}]$ -modules and \mathfrak{h} -locally nilpotent. We recall from [10, Section 3] the following properties of $\mathcal{O}_c(W, \mathfrak{h})$.

The action of the Euler element \mathbf{eu} on a module in $\mathcal{O}_c(W, \mathfrak{h})$ is locally finite. The category $\mathcal{O}_c(W, \mathfrak{h})$ is a highest weight category. In particular, it is artinian. Write $\text{Irr}(W)$ for the set of isomorphism classes of irreducible representations of W . The poset of standard modules in $\mathcal{O}_c(W, \mathfrak{h})$ is indexed by $\text{Irr}(W)$ with the partial order given by [10, Theorem 2.19]. More precisely, for $\xi \in \text{Irr}(W)$, equip it with a $\mathbb{C}W \rtimes \mathbb{C}[\mathfrak{h}^*]$ -module structure by letting the elements in $\mathfrak{h} \subset \mathbb{C}[\mathfrak{h}^*]$ act by zero, the standard module corresponding to ξ is

$$\Delta(\xi) = H_c(W, \mathfrak{h}) \otimes_{\mathbb{C}W \rtimes \mathbb{C}[\mathfrak{h}^*]} \xi.$$

It is an indecomposable module with a simple head $L(\xi)$. The set of isomorphism classes of simple modules in $\mathcal{O}_c(W, \mathfrak{h})$ is

$$\{[L(\xi)] : \xi \in \text{Irr}(W)\}.$$

It is a basis of the \mathbb{C} -vector space $K(\mathcal{O}_c(W, \mathfrak{h}))$. The set $\{[\Delta(\xi)] : \xi \in \text{Irr}(W)\}$ gives another basis of $K(\mathcal{O}_c(W, \mathfrak{h}))$.

We say a module N in $\mathcal{O}_c(W, \mathfrak{h})$ has a standard filtration if it admits a filtration

$$0 = N_0 \subset N_1 \subset \cdots \subset N_n = N$$

such that each quotient N_i/N_{i-1} is isomorphic to a standard module. We denote by $\mathcal{O}_c^\Delta(W, \mathfrak{h})$ the full subcategory of $\mathcal{O}_c(W, \mathfrak{h})$ consisting of such modules.

LEMMA 1.1. – (1) Any projective object in $\mathcal{O}_c(W, \mathfrak{h})$ has a standard filtration.

(2) A module in $\mathcal{O}_c(W, \mathfrak{h})$ has a standard filtration if and only if it is free as a $\mathbb{C}[\mathfrak{h}]$ -module.

Both (1) and (2) are given by [10, Proposition 2.21].

The category $\mathcal{O}_c(W, \mathfrak{h})$ has enough projective objects and has finite homological dimension [10, Section 4.3.1]. In particular, any module in $\mathcal{O}_c(W, \mathfrak{h})$ has a finite projective resolution. Write $\text{Proj}_c(W, \mathfrak{h})$ for the full subcategory of projective modules in $\mathcal{O}_c(W, \mathfrak{h})$. Let

$$I : \text{Proj}_c(W, \mathfrak{h}) \rightarrow \mathcal{O}_c(W, \mathfrak{h})$$

be the canonical embedding functor. We have the following lemma.

LEMMA 1.2. – *For any abelian category \mathcal{A} and any right exact functors F_1, F_2 from $\mathcal{O}_c(W, \mathfrak{h})$ to \mathcal{A} , the homomorphism of vector spaces*

$$r_I : \text{Hom}(F_1, F_2) \rightarrow \text{Hom}(F_1 \circ I, F_2 \circ I), \quad \gamma \mapsto \gamma 1_I$$

is an isomorphism.

In particular, if the functor $F_1 \circ I$ is isomorphic to $F_2 \circ I$, then we have $F_1 \cong F_2$.

Proof. – We need to show that for any morphism of functors $\nu : F_1 \circ I \rightarrow F_2 \circ I$ there is a unique morphism $\tilde{\nu} : F_1 \rightarrow F_2$ such that $\tilde{\nu} 1_I = \nu$. Since $\mathcal{O}_c(W, \mathfrak{h})$ has enough projectives, for any $M \in \mathcal{O}_c(W, \mathfrak{h})$ there exist P_0, P_1 in $\text{Proj}_c(W, \mathfrak{h})$ and an exact sequence in $\mathcal{O}_c(W, \mathfrak{h})$

$$(1.9) \quad P_1 \xrightarrow{d_1} P_0 \xrightarrow{d_0} M \longrightarrow 0.$$

Applying the right exact functors F_1, F_2 to this sequence we get the two exact sequences in the diagram below. The morphism of functors $\nu : F_1 \circ I \rightarrow F_2 \circ I$ yields well defined morphisms $\nu(P_1), \nu(P_0)$ such that the square commutes

$$\begin{array}{ccccc} F_1(P_1) & \xrightarrow{F_1(d_1)} & F_1(P_0) & \xrightarrow{F_1(d_0)} & F_1(M) & \longrightarrow & 0 \\ \downarrow \nu(P_1) & & \downarrow \nu(P_0) & & & & \\ F_2(P_1) & \xrightarrow{F_2(d_1)} & F_2(P_0) & \xrightarrow{F_2(d_0)} & F_2(M) & \longrightarrow & 0. \end{array}$$

Define $\tilde{\nu}(M)$ to be the unique morphism $F_1(M) \rightarrow F_2(M)$ that makes the diagram commute. Its definition is independent of the choice of P_0, P_1 , and it is independent of the choice of the exact sequence (1.9). The assignment $M \mapsto \tilde{\nu}(M)$ gives a morphism of functor $\tilde{\nu} : F_1 \rightarrow F_2$ such that $\tilde{\nu} 1_I = \nu$. It is unique by the uniqueness of the morphism $\tilde{\nu}(M)$. \square

1.5. The Knizhnik-Zamolodchikov functor

The Knizhnik-Zamolodchikov functor is an exact functor from the category $\mathcal{O}_c(W, \mathfrak{h})$ to the category $\mathcal{H}_q(W, \mathfrak{h})\text{-mod}$, where q is a certain parameter associated with c . Let us recall its definition from [10, Section 5.3].

Let $\mathcal{D}(\mathfrak{h}_{\text{reg}})$ be the algebra of differential operators on $\mathfrak{h}_{\text{reg}}$. Write

$$H_c(W, \mathfrak{h}_{\text{reg}}) = H_c(W, \mathfrak{h}) \otimes_{\mathbb{C}[\mathfrak{h}]} \mathbb{C}[\mathfrak{h}_{\text{reg}}].$$

We consider the Dunkl isomorphism, which is an isomorphism of algebras

$$H_c(W, \mathfrak{h}_{\text{reg}}) \xrightarrow{\sim} \mathcal{D}(\mathfrak{h}_{\text{reg}}) \rtimes \mathbb{C}W$$

given by $x \mapsto x, w \mapsto w$ for $x \in \mathfrak{h}^*, w \in W$, and

$$y \mapsto \partial_y + \sum_{s \in \phi} \frac{2c_s}{1 - \lambda_s} \frac{\alpha_s(y)}{\alpha_s} (s - 1), \quad \text{for } y \in \mathfrak{h}.$$

For any $M \in \mathcal{O}_c(W, \mathfrak{h})$, write

$$M_{\mathfrak{h}_{\text{reg}}} = M \otimes_{\mathbb{C}[\mathfrak{h}]} \mathbb{C}[\mathfrak{h}_{\text{reg}}].$$

It identifies via the Dunkl isomorphism with a $\mathcal{D}(\mathfrak{h}_{\text{reg}}) \rtimes W$ -module which is finitely generated over $\mathbb{C}[\mathfrak{h}_{\text{reg}}]$. Hence $M_{\mathfrak{h}_{\text{reg}}}$ is a W -equivariant vector bundle on $\mathfrak{h}_{\text{reg}}$ with an integrable connection ∇ given by $\nabla_y(m) = \partial_y m$ for $m \in M, y \in \mathfrak{h}$. It is proved in [10, Proposition 5.7] that the connection ∇ has regular singularities. Now, regard $\mathfrak{h}_{\text{reg}}$ as a complex manifold endowed with the transcendental topology. Denote by $\mathcal{O}_{\mathfrak{h}_{\text{reg}}}^{\text{an}}$ the sheaf of holomorphic functions on $\mathfrak{h}_{\text{reg}}$. For any free $\mathbb{C}[\mathfrak{h}_{\text{reg}}]$ -module N of finite rank, we consider

$$N^{\text{an}} = N \otimes_{\mathbb{C}[\mathfrak{h}_{\text{reg}}]} \mathcal{O}_{\mathfrak{h}_{\text{reg}}}^{\text{an}}.$$

It is an analytic locally free sheaf on $\mathfrak{h}_{\text{reg}}$. For ∇ an integrable connection on N , the sheaf of holomorphic horizontal sections

$$N^{\nabla} = \{n \in N^{\text{an}} : \nabla_y(n) = 0 \text{ for all } y \in \mathfrak{h}\}$$

is a W -equivariant local system on $\mathfrak{h}_{\text{reg}}$. Hence it identifies with a local system on $\mathfrak{h}_{\text{reg}}/W$. So it yields a finite dimensional representation of $\mathbb{C}B(W, \mathfrak{h})$. For $M \in \mathcal{O}_c(W, \mathfrak{h})$ it is proved in [10, Theorem 5.13] that the action of $\mathbb{C}B(W, \mathfrak{h})$ on $(M_{\mathfrak{h}_{\text{reg}}})^{\nabla}$ factors through the Hecke algebra $\mathcal{H}_q(W, \mathfrak{h})$. The formula for the parameter q is given in [10, Section 5.2].

The Knizhnik-Zamolodchikov functor is the functor

$$\text{KZ}(W, \mathfrak{h}) : \mathcal{O}_c(W, \mathfrak{h}) \rightarrow \mathcal{H}_q(W, \mathfrak{h})\text{-mod}, \quad M \mapsto (M_{\mathfrak{h}_{\text{reg}}})^{\nabla}.$$

By definition it is exact. Let us recall some of its properties following [10]. Assume in the rest of this subsection that *the algebras $\mathcal{H}_q(W, \mathfrak{h})$ and $\mathbb{C}W$ have the same dimension over \mathbb{C}* . We abbreviate $\text{KZ} = \text{KZ}(W, \mathfrak{h})$. The functor KZ is represented by a projective object P_{KZ} in $\mathcal{O}_c(W, \mathfrak{h})$. More precisely, there is an algebra homomorphism

$$\rho : \mathcal{H}_q(W, \mathfrak{h}) \rightarrow \text{End}_{\mathcal{O}_c(W, \mathfrak{h})}(P_{\text{KZ}})^{\text{op}}$$

such that KZ is isomorphic to the functor $\text{Hom}_{\mathcal{O}_c(W, \mathfrak{h})}(P_{\text{KZ}}, -)$. By [10, Theorem 5.15] the homomorphism ρ is an isomorphism. In particular $\text{KZ}(P_{\text{KZ}})$ is isomorphic to $\mathcal{H}_q(W, \mathfrak{h})$ as $\mathcal{H}_q(W, \mathfrak{h})$ -modules.

Now, recall that the center of a category \mathcal{C} is the algebra $Z(\mathcal{C})$ of endomorphisms of the identity functor $Id_{\mathcal{C}}$. So there is a canonical map

$$Z(\mathcal{O}_c(W, \mathfrak{h})) \rightarrow \text{End}_{\mathcal{O}_c(W, \mathfrak{h})}(P_{\text{KZ}}).$$

The composition of this map with ρ^{-1} yields an algebra homomorphism

$$\gamma : Z(\mathcal{O}_c(W, \mathfrak{h})) \rightarrow Z(\mathcal{H}_q(W, \mathfrak{h})),$$

where $Z(\mathcal{H}_q(W, \mathfrak{h}))$ denotes the center of $\mathcal{H}_q(W, \mathfrak{h})$.

LEMMA 1.3. – (1) *The homomorphism γ is an isomorphism.*

(2) *For a module M in $\mathcal{O}_c(W, \mathfrak{h})$ and an element f in $Z(\mathcal{O}_c(W, \mathfrak{h}))$ the morphism*

$$\mathrm{KZ}(f(M)) : \mathrm{KZ}(M) \rightarrow \mathrm{KZ}(M)$$

is the multiplication by $\gamma(f)$.

See [10, Corollary 5.18] for (1). Part (2) follows from the construction of γ .

The functor KZ is a quotient functor, see [10, Theorem 5.14]. Therefore it has a right adjoint $S : \mathcal{H}_q(W, \mathfrak{h}) \rightarrow \mathcal{O}_c(W, \mathfrak{h})$ such that the canonical adjunction map $\mathrm{KZ} \circ S \rightarrow \mathrm{Id}_{\mathcal{H}_q(W, \mathfrak{h})}$ is an isomorphism of functors. We have the following proposition.

PROPOSITION 1.4. – *Let Q be a projective object in $\mathcal{O}_c(W, \mathfrak{h})$.*

(1) *For any object $M \in \mathcal{O}_c(W, \mathfrak{h})$, the following morphism of \mathbb{C} -vector spaces is an isomorphism*

$$\mathrm{Hom}_{\mathcal{O}_c(W, \mathfrak{h})}(M, Q) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{H}_q(W)}(\mathrm{KZ}(M), \mathrm{KZ}(Q)), \quad f \mapsto \mathrm{KZ}(f).$$

In particular, the functor KZ is fully faithful over $\mathrm{Proj}_c(W, \mathfrak{h})$.

(2) *The canonical adjunction map gives an isomorphism $Q \xrightarrow{\sim} S \circ \mathrm{KZ}(Q)$.*

See [10, Theorems 5.3, 5.16].

1.6. Parabolic restriction and induction for rational DAHA's

From now on we will always assume that $\mathfrak{h}^W = 1$. Recall from Section 1.2 that $W' \subset W$ is the stabilizer of a point $b \in \mathfrak{h}$ and that $\bar{\mathfrak{h}} = \mathfrak{h}/\mathfrak{h}^{W'}$. Let us recall from [4] the definition of the parabolic restriction and induction functors

$$\mathrm{Res}_b : \mathcal{O}_c(W, \mathfrak{h}) \rightarrow \mathcal{O}_{c'}(W', \bar{\mathfrak{h}}), \quad \mathrm{Ind}_b : \mathcal{O}_{c'}(W', \bar{\mathfrak{h}}) \rightarrow \mathcal{O}_c(W, \mathfrak{h}).$$

First we need some notation. For any point $p \in \mathfrak{h}$ we write $\mathbb{C}[[\mathfrak{h}]]_p$ for the completion of $\mathbb{C}[\mathfrak{h}]$ at p , and we write $\widehat{\mathbb{C}[\mathfrak{h}]}_p$ for the completion of $\mathbb{C}[\mathfrak{h}]$ at the W -orbit of p in \mathfrak{h} . Note that we have $\mathbb{C}[[\mathfrak{h}]]_0 = \widehat{\mathbb{C}[\mathfrak{h}]}_0$. For any $\mathbb{C}[\mathfrak{h}]$ -module M let

$$\widehat{M}_p = \widehat{\mathbb{C}[\mathfrak{h}]}_p \otimes_{\mathbb{C}[\mathfrak{h}]} M.$$

The completions $\widehat{H}_c(W, \mathfrak{h})_b, \widehat{H}_{c'}(W', \mathfrak{h})_0$ are well defined algebras. We denote by $\widehat{\mathcal{O}}_c(W, \mathfrak{h})_b$ the category of $\widehat{H}_c(W, \mathfrak{h})_b$ -modules that are finitely generated over $\widehat{\mathbb{C}[\mathfrak{h}]}_b$, and we denote by $\widehat{\mathcal{O}}_{c'}(W', \mathfrak{h})_0$ the category of $\widehat{H}_{c'}(W', \mathfrak{h})_0$ -modules that are finitely generated over $\widehat{\mathbb{C}[\mathfrak{h}]}_0$. Let $P = \mathrm{Fun}_{W'}(W, \widehat{H}_c(W', \mathfrak{h})_0)$ be the set of W' -invariant maps from W to $\widehat{H}_c(W', \mathfrak{h})_0$. Let $Z(W, W', \widehat{H}_c(W', \mathfrak{h})_0)$ be the ring of endomorphisms of the right $\widehat{H}_c(W', \mathfrak{h})_0$ -module P . We have the following proposition given by [4, Theorem 3.2].

PROPOSITION 1.5. – *There is an isomorphism of algebras*

$$\Theta : \widehat{H}_c(W, \mathfrak{h})_b \longrightarrow Z(W, W', \widehat{H}_{c'}(W', \mathfrak{h})_0)$$

defined as follows: for $f \in P$, $\alpha \in \mathfrak{h}^*$, $a \in \mathfrak{h}$, $u \in W$,

$$\begin{aligned} (\Theta(u)f)(w) &= f(wu), \\ (\Theta(x_\alpha)f)(w) &= (x_{w\alpha}^{(b)} + \alpha(w^{-1}b))f(w), \\ (\Theta(y_a)f)(w) &= y_{wa}^{(b)}f(w) + \sum_{s \in \phi, s \notin W'} \frac{2c_s}{1 - \lambda_s} \frac{\alpha_s(wa)}{x_{\alpha_s}^{(b)} + \alpha_s(b)} (f(sw) - f(w)), \end{aligned}$$

where $x_\alpha \in \mathfrak{h}^* \subset H_c(W, \mathfrak{h})$, $x_\alpha^{(b)} \in \mathfrak{h}^* \subset H_{c'}(W', \mathfrak{h})$, $y_a \in \mathfrak{h} \subset H_c(W, \mathfrak{h})$, $y_a^{(b)} \in \mathfrak{h} \subset H_{c'}(W', \mathfrak{h})$.

Using Θ we will identify $\widehat{H}_c(W, \mathfrak{h})_b$ -modules with $Z(W, W', \widehat{H}_{c'}(W', \mathfrak{h})_0)$ -modules. So the module $P = \text{Fun}_{W'}(W, \widehat{H}_c(W, \mathfrak{h})_0)$ becomes an $(\widehat{H}_c(W, \mathfrak{h})_b, \widehat{H}_{c'}(W', \mathfrak{h})_0)$ -bimodule. Hence for any $N \in \widehat{\mathcal{O}}_{c'}(W', \mathfrak{h})_0$ the module $P \otimes_{\widehat{H}_{c'}(W', \mathfrak{h})_0} N$ lives in $\widehat{\mathcal{O}}_c(W, \mathfrak{h})_b$. It is naturally identified with $\text{Fun}_{W'}(W, N)$, the set of W' -invariant maps from W to N . For any $\mathbb{C}[\mathfrak{h}^*]$ -module M write $E(M) \subset M$ for the locally nilpotent part of M under the action of \mathfrak{h} .

The ingredients for defining the functors Res_b and Ind_b consist of:

- the adjoint pair of functors $(\widehat{}_b, E^b)$ with

$$\widehat{}_b : \mathcal{O}_c(W, \mathfrak{h}) \rightarrow \widehat{\mathcal{O}}_c(W, \mathfrak{h})_b, \quad M \mapsto \widehat{M}_b,$$

$$E^b : \widehat{\mathcal{O}}_c(W, \mathfrak{h})_b \rightarrow \mathcal{O}_c(W, \mathfrak{h}), \quad N \rightarrow E(N),$$

- the Morita equivalence

$$J : \widehat{\mathcal{O}}_{c'}(W', \mathfrak{h})_0 \rightarrow \widehat{\mathcal{O}}_c(W, \mathfrak{h})_b, \quad N \mapsto \text{Fun}_{W'}(W, N),$$

and its quasi-inverse R given in Section 1.7 below,

- the equivalence of categories

$$E : \widehat{\mathcal{O}}_{c'}(W', \mathfrak{h})_0 \rightarrow \mathcal{O}_{c'}(W', \mathfrak{h}), \quad M \mapsto E(M)$$

and its quasi-inverse given by $N \mapsto \widehat{N}_0$,

- the equivalence of categories

$$(1.10) \quad \zeta : \mathcal{O}_{c'}(W', \mathfrak{h}) \rightarrow \mathcal{O}_{c'}(W', \bar{\mathfrak{h}}), \quad M \mapsto \{v \in M : yv = 0, \text{ for all } y \in \mathfrak{h}^{W'}\}$$

and its quasi-inverse ζ^{-1} given in Section 1.8 below.

For $M \in \mathcal{O}_c(W, \mathfrak{h})$ and $N \in \mathcal{O}_{c'}(W', \bar{\mathfrak{h}})$ the functors Res_b and Ind_b are defined by

$$(1.11) \quad \begin{aligned} \text{Res}_b(M) &= \zeta \circ E \circ R(\widehat{M}_b), \\ \text{Ind}_b(N) &= E^b \circ J(\widehat{\zeta^{-1}(N)}_0). \end{aligned}$$

We refer to [4, Section 2,3] for details.

1.7. The idempotent x_{pr} and the functor R

We give some details on the isomorphism Θ for a future use. Fix elements $1 = u_1, u_2, \dots, u_r$ in W such that $W = \bigsqcup_{i=1}^r W' u_i$. Let $\text{Mat}_r(\widehat{H}_{c'}(W', \mathfrak{h})_0)$ be the algebra of $r \times r$ matrices with coefficients in $\widehat{H}_{c'}(W', \mathfrak{h})_0$. We have an algebra isomorphism

$$(1.12) \quad \begin{aligned} \Phi : Z(W, W', \widehat{H}_{c'}(W', \mathfrak{h})_0) &\rightarrow \text{Mat}_r(\widehat{H}_{c'}(W', \mathfrak{h})_0), \\ A &\mapsto (\Phi(A)_{ij})_{1 \leq i, j \leq r} \end{aligned}$$

such that

$$(Af)(u_i) = \sum_{j=1}^r \Phi(A)_{ij} f(u_j), \quad \text{for all } f \in P, 1 \leq i \leq r.$$

Denote by E_{ij} , $1 \leq i, j \leq r$, the elementary matrix in $\text{Mat}_r(\widehat{H}_{c'}(W', \mathfrak{h})_0)$ with coefficient 1 in the position (i, j) and zero elsewhere. Note that the algebra isomorphism

$$\Phi \circ \Theta : \widehat{H}_c(W, \mathfrak{h})_b \xrightarrow{\sim} \text{Mat}_r(\widehat{H}_{c'}(W', \mathfrak{h})_0)$$

restricts to an isomorphism of subalgebras

$$(1.13) \quad \widehat{\mathbb{C}[\mathfrak{h}]_b} \cong \bigoplus_{i=1}^r \mathbb{C}[[\mathfrak{h}]]_0 E_{ii}.$$

Indeed, there is a unique isomorphism of algebras

$$(1.14) \quad \varpi : \widehat{\mathbb{C}[\mathfrak{h}]_b} \cong \bigoplus_{i=1}^r \mathbb{C}[[\mathfrak{h}]]_{u_i^{-1}b},$$

extending the algebra homomorphism

$$\mathbb{C}[\mathfrak{h}] \rightarrow \bigoplus_{i=1}^r \mathbb{C}[\mathfrak{h}], \quad x \mapsto (x, x, \dots, x), \quad \forall x \in \mathfrak{h}^*.$$

For each i consider the isomorphism of algebras

$$\phi_i : \mathbb{C}[[\mathfrak{h}]]_{u_i^{-1}b} \rightarrow \mathbb{C}[[\mathfrak{h}]]_0, \quad x \mapsto u_i x + x(u_i^{-1}b), \quad \forall x \in \mathfrak{h}^*.$$

The isomorphism (1.13) is exactly the composition of ϖ with the direct sum $\bigoplus_{i=1}^r \phi_i$. Here E_{ii} is the image of the idempotent in $\widehat{\mathbb{C}[\mathfrak{h}]_b}$ corresponding to the component $\mathbb{C}[[\mathfrak{h}]]_{u_i^{-1}b}$. We will denote by x_{pr} the idempotent in $\widehat{\mathbb{C}[\mathfrak{h}]_b}$ corresponding to $\mathbb{C}[[\mathfrak{h}]]_b$, i.e., $\Phi \circ \Theta(x_{\text{pr}}) = E_{11}$. Then the functor

$$R : \widehat{\mathcal{O}}_c(W, \mathfrak{h})_b \rightarrow \widehat{\mathcal{O}}_{c'}(W', \mathfrak{h})_0, \quad M \mapsto x_{\text{pr}} M$$

is a quasi-inverse of J . Here, the action of $\widehat{H}_{c'}(W', \mathfrak{h})_0$ on $R(M) = x_{\text{pr}} M$ is given by the following formulas deduced from Proposition 1.5. For any $\alpha \in \mathfrak{h}^*$, $w \in W'$, $a \in \mathfrak{h}^*$, $m \in M$ we have

$$(1.15) \quad x_{\alpha}^{(b)} x_{\text{pr}}(m) = x_{\text{pr}}((x_{\alpha} - \alpha(b))m),$$

$$(1.16) \quad w x_{\text{pr}}(m) = x_{\text{pr}}(wm),$$

$$(1.17) \quad y_a^{(b)} x_{\text{pr}}(m) = x_{\text{pr}} \left(\left(y_a + \sum_{s \in \mathcal{J}, s \notin W'} \frac{2c_s}{1 - \lambda_s} \frac{\alpha_s(a)}{\alpha_s} \right) m \right).$$

In particular, we have

$$(1.18) \quad R(M) = \phi_1^*(x_{\text{pr}}(M))$$

as $\mathbb{C}[[\mathfrak{h}]]_0 \rtimes W'$ -modules. Finally, note that the following equality holds in $\widehat{H}_c(W, \mathfrak{h})_b$

$$(1.19) \quad x_{\text{pr}} u x_{\text{pr}} = 0, \quad \forall u \in W - W'.$$

1.8. A quasi-inverse of ζ

Let us recall from [4, Section 2.3] the following facts. Let $\mathfrak{h}^{*W'}$ be the subspace of \mathfrak{h}^* consisting of fixed points of W' . Set

$$(\mathfrak{h}^{*W'})^\perp = \{v \in \mathfrak{h} : f(v) = 0 \text{ for all } f \in \mathfrak{h}^{*W'}\}.$$

We have a W' -invariant decomposition

$$\mathfrak{h} = (\mathfrak{h}^{*W'})^\perp \oplus \mathfrak{h}^{W'}.$$

The W' -space $(\mathfrak{h}^{*W'})^\perp$ is canonically identified with $\bar{\mathfrak{h}}$. Since the action of W' on $\mathfrak{h}^{W'}$ is trivial, we have an obvious algebra isomorphism

$$(1.20) \quad H_{c'}(W', \mathfrak{h}) \cong H_{c'}(W', \bar{\mathfrak{h}}) \otimes \mathcal{D}(\mathfrak{h}^{W'}).$$

It maps an element y in the subset $\mathfrak{h}^{W'}$ of $H_{c'}(W', \mathfrak{h})$ to the operator ∂_y in $\mathcal{D}(\mathfrak{h}^{W'})$. Write $\mathcal{O}(1, \mathfrak{h}^{W'})$ for the category of finitely generated $\mathcal{D}(\mathfrak{h}^{W'})$ -modules that are ∂_y -locally nilpotent for all $y \in \mathfrak{h}^{W'}$. The algebra isomorphism above yields an equivalence of categories

$$\mathcal{O}_{c'}(W', \mathfrak{h}) \cong \mathcal{O}_{c'}(W', \bar{\mathfrak{h}}) \otimes \mathcal{O}(1, \mathfrak{h}^{W'}).$$

The functor ζ in (1.10) is an equivalence, because it is induced by the functor

$$\mathcal{O}(1, \mathfrak{h}^{W'}) \xrightarrow{\sim} \mathbb{C}\text{-mod}, \quad M \rightarrow \{m \in M, \partial_y(m) = 0 \text{ for all } y \in \mathfrak{h}^{W'}\},$$

which is an equivalence by Kashiwara's lemma upon taking Fourier transforms. In particular, a quasi-inverse of ζ is given by

$$(1.21) \quad \zeta^{-1} : \mathcal{O}_{c'}(W', \bar{\mathfrak{h}}) \rightarrow \mathcal{O}_{c'}(W', \mathfrak{h}), \quad N \mapsto N \otimes \mathbb{C}[\mathfrak{h}^{W'}],$$

where $\mathbb{C}[\mathfrak{h}^{W'}] \in \mathcal{O}(1, \mathfrak{h}^{W'})$ is the polynomial representation of $\mathcal{D}(\mathfrak{h}^{W'})$.

Moreover, the functor ζ maps a standard module in $\mathcal{O}_{c'}(W', \mathfrak{h})$ to a standard module in $\mathcal{O}_{c'}(W', \bar{\mathfrak{h}})$. Indeed, for any $\xi \in \text{Irr}(W')$, we have an isomorphism of $H_{c'}(W', \mathfrak{h})$ -modules

$$H_{c'}(W', \mathfrak{h}) \otimes_{\mathbb{C}[\mathfrak{h}^*] \rtimes W'} \xi = (H_{c'}(W', \bar{\mathfrak{h}}) \otimes_{\mathbb{C}[(\bar{\mathfrak{h}})^*] \rtimes W'} \xi) \otimes (\mathcal{D}(\mathfrak{h}^{W'}) \otimes_{\mathbb{C}[(\mathfrak{h}^{W'})^*]} \mathbb{C}).$$

On the right hand side \mathbb{C} denotes the trivial module of $\mathbb{C}[(\mathfrak{h}^{W'})^*]$, and the latter is identified with the subalgebra of $\mathcal{D}(\mathfrak{h}^{W'})$ generated by ∂_y for all $y \in \mathfrak{h}^{W'}$. We have

$$\mathcal{D}(\mathfrak{h}^{W'}) \otimes_{\mathbb{C}[(\mathfrak{h}^{W'})^*]} \mathbb{C} \cong \mathbb{C}[\mathfrak{h}^{W'}]$$

as $\mathcal{D}(\mathfrak{h}^{W'})$ -modules. So ζ maps the standard module $\Delta(\xi)$ for $H_{c'}(W', \mathfrak{h})$ to the standard module $\Delta(\xi)$ for $H_{c'}(W', \bar{\mathfrak{h}})$.

1.9. – Here are some properties of Res_b and Ind_b .

PROPOSITION 1.6. – (1) *Both functors Res_b and Ind_b are exact. The functor Res_b is left adjoint to Ind_b . In particular the functor Res_b preserves projective objects and Ind_b preserves injective objects.*

(2) *Let $\text{Res}_{W'}^W$ and $\text{Ind}_{W'}^W$ be respectively the restriction and induction functors of groups. We have the following commutative diagram*

$$\begin{array}{ccc} K(\mathcal{O}_c(W, \mathfrak{h})) & \xrightarrow[\sim]{\omega} & K(\mathbb{C}W) \\ \text{Ind}_b \uparrow \downarrow \text{Res}_b & & \text{Ind}_{W'}^W \uparrow \downarrow \text{Res}_{W'}^W \\ K(\mathcal{O}_{c'}(W', \bar{\mathfrak{h}})) & \xrightarrow[\sim]{\omega'} & K(\mathbb{C}W') \end{array}$$

Here the isomorphism ω (resp. ω') is given by mapping $[\Delta(\xi)]$ to $[\xi]$ for any $\xi \in \text{Irr}(W)$ (resp. $\xi \in \text{Irr}(W')$).

See [4, Proposition 3.9, Theorem 3.10] for (1), [4, Proposition 3.14] for (2).

1.10. Restriction of modules having a standard filtration

In the rest of Section 1, we study the actions of the restriction functors on modules having a standard filtration in $\mathcal{O}_c(W, \mathfrak{h})$ (Proposition 1.9). We will need the following lemmas.

LEMMA 1.7. – *Let M be an object in $\mathcal{O}_c^\Delta(W, \mathfrak{h})$.*

(1) *There is a finite dimensional subspace V of M such that V is stable under the action of $\mathbb{C}W$ and the map*

$$\mathbb{C}[\mathfrak{h}] \otimes V \rightarrow M, \quad p \otimes v \mapsto pv$$

is an isomorphism of $\mathbb{C}[\mathfrak{h}] \rtimes W$ -modules.

(2) *The map $\omega : K(\mathcal{O}_c(W, \mathfrak{h})) \rightarrow K(\mathbb{C}W)$ in Proposition 1.6(2) satisfies*

$$(1.22) \quad \omega([M]) = [V].$$

Proof. – Let

$$0 = M_0 \subset M_1 \subset \cdots \subset M_l = M$$

be a filtration of M such that for any $1 \leq i \leq l$ we have $M_i/M_{i-1} \cong \Delta(\xi_i)$ for some $\xi_i \in \text{Irr}(W)$. We prove (1) and (2) by recurrence on l . If $l = 1$, then M is a standard module. Both (1) and (2) hold by definition. For $l > 1$, by induction we may suppose that there is a subspace V' of M_{l-1} such that the properties in (1) and (2) are satisfied for M_{l-1} and V' . Now, consider the exact sequence

$$0 \longrightarrow M_{l-1} \longrightarrow M \xrightarrow{j} \Delta(\xi_l) \longrightarrow 0.$$

From the isomorphism of $\mathbb{C}[\mathfrak{h}] \rtimes W$ -modules $\Delta(\xi_l) \cong \mathbb{C}[\mathfrak{h}] \otimes \xi$ we see that $\Delta(\xi_l)$ is a projective $\mathbb{C}[\mathfrak{h}] \rtimes W$ -module. Hence there exists a morphism of $\mathbb{C}[\mathfrak{h}] \rtimes W$ -modules $s : \Delta(\xi_l) \rightarrow M$ that provides a section of j . Let $V = V' \oplus s(\xi_l) \subset M$. It is stable under the action of $\mathbb{C}W$. The map $\mathbb{C}[\mathfrak{h}] \otimes V \rightarrow M$ in (1) is an injective morphism of $\mathbb{C}[\mathfrak{h}] \rtimes W$ -modules. Its image is $M_{l-1} \oplus s(\Delta(\xi_l))$, which is equal to M . So it is an isomorphism. We have

$$\omega([M]) = \omega([M_{l-1}]) + \omega([\Delta(\xi_l)]),$$

by assumption $\omega([M_{l-1}]) = [V']$, so $\omega([M]) = [V'] + [\xi_l] = [V]$. \square

LEMMA 1.8. – (1) Let M be an $\widehat{H}_c(W, \mathfrak{h})_0$ -module free over $\mathbb{C}[[\mathfrak{h}]]_0$. If there exist generalized eigenvectors v_1, \dots, v_n of \mathbf{eu} which form a basis of M over $\mathbb{C}[[\mathfrak{h}]]_0$, then for $f_1, \dots, f_n \in \mathbb{C}[[\mathfrak{h}]]_0$ the element $m = \sum_{i=1}^n f_i v_i$ is \mathbf{eu} -finite if and only if f_1, \dots, f_n all belong to $\mathbb{C}[\mathfrak{h}]$.

(2) Let N be an object in $\mathcal{O}_c(W, \mathfrak{h})$. If \widehat{N}_0 is a free $\mathbb{C}[[\mathfrak{h}]]_0$ -module, then N is a free $\mathbb{C}[\mathfrak{h}]$ -module. It admits a basis consisting of generalized eigenvectors v_1, \dots, v_n of \mathbf{eu} .

Proof. – (1) It follows from the proof of [4, Theorem 2.3].

(2) Since N belongs to $\mathcal{O}_c(W, \mathfrak{h})$, it is finitely generated over $\mathbb{C}[\mathfrak{h}]$. Denote by \mathfrak{m} the maximal ideal of $\mathbb{C}[[\mathfrak{h}]]_0$. The canonical map $N \rightarrow \widehat{N}_0/\mathfrak{m}\widehat{N}_0$ is surjective. So there exist v_1, \dots, v_n in N such that their images form a basis of $\widehat{N}_0/\mathfrak{m}\widehat{N}_0$ over \mathbb{C} . Moreover, we may choose v_1, \dots, v_n to be generalized eigenvectors of \mathbf{eu} , because the \mathbf{eu} -action on N is locally finite. Since \widehat{N}_0 is free over $\mathbb{C}[[\mathfrak{h}]]_0$, Nakayama's lemma yields that v_1, \dots, v_n form a basis of \widehat{N}_0 over $\mathbb{C}[[\mathfrak{h}]]_0$. By part (1) the set N' of \mathbf{eu} -finite elements in \widehat{N}_0 is the free $\mathbb{C}[\mathfrak{h}]$ -submodule generated by v_1, \dots, v_n . On the other hand, since \widehat{N}_0 belongs to $\widehat{\mathcal{O}}_c(W, \mathfrak{h})_0$, by [4, Proposition 2.4] an element in \widehat{N}_0 is \mathfrak{h} -nilpotent if and only if it is \mathbf{eu} -finite. So $N' = E(\widehat{N}_0)$. On the other hand, the canonical inclusion $N \subset E(\widehat{N}_0)$ is an equality by [4, Theorem 3.2]. Hence $N = N'$. This implies that N is free over $\mathbb{C}[\mathfrak{h}]$, with a basis given by v_1, \dots, v_n , which are generalized eigenvectors of \mathbf{eu} . \square

PROPOSITION 1.9. – Let M be an object in $\mathcal{O}_c^\Delta(W, \mathfrak{h})$.

(1) The object $\text{Res}_b(M)$ has a standard filtration.

(2) Let V be a subspace of M that has the properties of Lemma 1.7(1). Then there is an isomorphism of $\mathbb{C}[\mathfrak{h}] \rtimes W'$ -modules

$$\text{Res}_b(M) \cong \mathbb{C}[\mathfrak{h}] \otimes \text{Res}_{W'}^W(V).$$

Proof. – (1) By the end of Section 1.8 the equivalence ζ maps a standard module in $\mathcal{O}_{c'}(W', \mathfrak{h})$ to a standard one in $\mathcal{O}_{c'}(W', \widehat{\mathfrak{h}})$. Hence to prove that $\text{Res}_b(M) = \zeta \circ E \circ R(\widehat{M}_b)$ has a standard filtration, it is enough to show that $N = E \circ R(\widehat{M}_b)$ has one. We claim that the module N is free over $\mathbb{C}[\mathfrak{h}]$. So the result follows from Lemma 1.1(2).

Let us prove the claim. Recall from (1.18) that we have $R(\widehat{M}_b) = \phi_1^*(x_{\text{pr}} \widehat{M}_b)$ as $\mathbb{C}[[\mathfrak{h}]]_0 \rtimes W'$ -modules. Using the isomorphism of $\mathbb{C}[\mathfrak{h}] \rtimes W'$ -modules $M \cong \mathbb{C}[\mathfrak{h}] \otimes V$ given in Lemma 1.7(1), we deduce an isomorphism of $\mathbb{C}[[\mathfrak{h}]]_0 \rtimes W'$ -modules

$$\begin{aligned} R(\widehat{M}_b) &\cong \phi_1^*(x_{\text{pr}}(\widehat{\mathbb{C}[\mathfrak{h}]_b} \otimes V)) \\ &\cong \mathbb{C}[[\mathfrak{h}]]_0 \otimes V. \end{aligned}$$

So the module $R(\widehat{M}_b)$ is free over $\mathbb{C}[[\mathfrak{h}]]_0$. The completion of the module N at 0 is isomorphic to $R(\widehat{M}_b)$. By Lemma 1.8(2) the module N is free over $\mathbb{C}[\mathfrak{h}]$. The claim is proved.

(2) Since $\text{Res}_b(M)$ has a standard filtration, by Lemma 1.7 there exists a finite dimensional vector space $V' \subset \text{Res}_b(M)$ such that V' is stable under the action of $\mathbb{C}W'$ and we have an isomorphism of $\mathbb{C}[\mathfrak{h}] \rtimes W'$ -modules

$$\text{Res}_b(M) \cong \mathbb{C}[\mathfrak{h}] \otimes V'.$$

Moreover, we have $\omega'([\text{Res}_b(M)]) = [V']$ where ω' is the map in Proposition 1.6(2). The same proposition yields that $\text{Res}_{W'}^W(\omega[M]) = \omega'([\text{Res}_b(M)])$. Since $\omega([M]) = [V]$ by

(1.22), the $\mathbb{C}W'$ -module V' is isomorphic to $\text{Res}_{W'}^W(V)$. So we have an isomorphism of $\mathbb{C}[\bar{\mathfrak{h}}] \rtimes W'$ -modules

$$\text{Res}_b(M) \cong \mathbb{C}[\bar{\mathfrak{h}}] \otimes \text{Res}_{W'}^W(V). \quad \square$$

2. KZ commutes with restriction functors

In this section, we relate the restriction and induction functors for rational DAHA's to the corresponding functors for Hecke algebras via the functor KZ . We will always assume that the Hecke algebras have the same dimension as the corresponding group algebras. Thus the Knizhnik-Zamolodchikov functors admit the properties recalled in Section 1.5.

2.1. – Let W be a complex reflection group acting on \mathfrak{h} . Let b be a point in \mathfrak{h} and let W' be its stabilizer in W . We will abbreviate $\text{KZ} = \text{KZ}(W, \mathfrak{h})$, $\text{KZ}' = \text{KZ}(W', \bar{\mathfrak{h}})$.

THEOREM 2.1. – *There is an isomorphism of functors*

$$\text{KZ}' \circ \text{Res}_b \cong \mathcal{R} \text{Res}_{W'}^W \circ \text{KZ}.$$

Proof. – We will regard $\text{KZ} : \mathcal{O}_c(W, \mathfrak{h}) \rightarrow \mathcal{H}_q(W)\text{-mod}$ as a functor from $\mathcal{O}_c(W, \mathfrak{h})$ to $B_W\text{-mod}$ in the obvious way. Similarly we will regard KZ' as a functor to $B_{W'}\text{-mod}$. Recall the inclusion $\iota : B_{W'} \hookrightarrow B_W$ from (1.6). The theorem amounts to prove that for any $M \in \mathcal{O}_c(W, \mathfrak{h})$ there is a natural isomorphism of $B_{W'}$ -modules

$$(2.1) \quad \text{KZ}' \circ \text{Res}_b(M) \cong \iota_* \circ \text{KZ}(M).$$

Step 1. Recall the functor $\zeta : \mathcal{O}_{c'}(W', \mathfrak{h}) \rightarrow \mathcal{O}_{c'}(W', \bar{\mathfrak{h}})$ from (1.10) and its quasi-inverse ζ^{-1} in (1.21). Let

$$N = \zeta^{-1}(\text{Res}_b(M)).$$

We have $N \cong \text{Res}_b(M) \otimes \mathbb{C}[\mathfrak{h}^{W'}]$. Since the canonical epimorphism $\mathfrak{h} \rightarrow \bar{\mathfrak{h}}$ induces a fibration $\mathfrak{h}'_{\text{reg}} \rightarrow \bar{\mathfrak{h}}_{\text{reg}}$, see Section 1.2, we have

$$(2.2) \quad N_{\mathfrak{h}'_{\text{reg}}} \cong \text{Res}_b(M)_{\bar{\mathfrak{h}}_{\text{reg}}} \otimes \mathbb{C}[\mathfrak{h}^{W'}].$$

By Dunkl isomorphisms, the left hand side is a $\mathcal{D}(\mathfrak{h}'_{\text{reg}}) \rtimes W'$ -module while the right hand side is a $(\mathcal{D}(\bar{\mathfrak{h}}_{\text{reg}}) \rtimes W') \otimes \mathcal{D}(\mathfrak{h}^{W'})$ -module. Identify these two algebras in the obvious way. The isomorphism (2.2) is compatible with the W' -equivariant \mathcal{D} -module structures. Hence we have

$$(N_{\mathfrak{h}'_{\text{reg}}})^\nabla \cong (\text{Res}_b(M)_{\bar{\mathfrak{h}}_{\text{reg}}})^\nabla \otimes \mathbb{C}[\mathfrak{h}^{W'}]^\nabla.$$

Since $\mathbb{C}[\mathfrak{h}^{W'}]^\nabla = \mathbb{C}$, this yields a natural isomorphism

$$\ell_* \circ \text{KZ}(W', \mathfrak{h})(N) \cong \text{KZ}' \circ \text{Res}_b(M),$$

where ℓ is the homomorphism defined in (1.2).

Step 2. Consider the W' -equivariant algebra isomorphism

$$\phi : \mathbb{C}[\mathfrak{h}] \rightarrow \mathbb{C}[\mathfrak{h}], \quad x \mapsto x + x(b) \text{ for } x \in \mathfrak{h}^*.$$

It induces an isomorphism $\hat{\phi} : \mathbb{C}[[\mathfrak{h}]]_b \xrightarrow{\sim} \mathbb{C}[[\mathfrak{h}]]_0$. The latter yields an algebra isomorphism

$$\mathbb{C}[[\mathfrak{h}]]_b \otimes_{\mathbb{C}[\mathfrak{h}]} \mathbb{C}[\mathfrak{h}_{\text{reg}}] \simeq \mathbb{C}[[\mathfrak{h}]]_0 \otimes_{\mathbb{C}[\mathfrak{h}]} \mathbb{C}[\mathfrak{h}'_{\text{reg}}].$$

To see this note first that by definition, the left hand side is $\mathbb{C}[[\mathfrak{h}]]_b[\alpha_s^{-1}, s \in \mathcal{J}]$. For $s \in \mathcal{J}$, $s \notin W'$ the element α_s is invertible in $\mathbb{C}[[\mathfrak{h}]]_b$, so we have

$$\mathbb{C}[[\mathfrak{h}]]_b \otimes_{\mathbb{C}[\mathfrak{h}]} \mathbb{C}[\mathfrak{h}_{\text{reg}}] = \mathbb{C}[[\mathfrak{h}]]_b[\alpha_s^{-1}, s \in \mathcal{J} \cap W'].$$

For $s \in \mathcal{J} \cap W'$ we have $\alpha_s(b) = 0$, so $\hat{\phi}(\alpha_s) = \alpha_s$. Hence

$$\begin{aligned} \hat{\phi}(\mathbb{C}[[\mathfrak{h}]]_b)[\hat{\phi}(\alpha_s)^{-1}, s \in \mathcal{J} \cap W'] &= \mathbb{C}[[\mathfrak{h}]]_0[\alpha_s^{-1}, s \in \mathcal{J} \cap W'] \\ &= \mathbb{C}[[\mathfrak{h}]]_0 \otimes_{\mathbb{C}[\mathfrak{h}]} \mathbb{C}[\mathfrak{h}'_{\text{reg}}]. \end{aligned}$$

Step 3. We will assume in Steps 3, 4, 5 that M is a module in $\mathcal{O}_c^\Delta(W, \mathfrak{h})$. In this step we prove that N is isomorphic to $\phi^*(M)$ as $\mathbb{C}[\mathfrak{h}] \rtimes W'$ -modules. Let V be a subspace of M as in Lemma 1.7(1). So we have an isomorphism of $\mathbb{C}[\mathfrak{h}] \rtimes W$ -modules

$$(2.3) \quad M \cong \mathbb{C}[\mathfrak{h}] \otimes V.$$

Also, by Proposition 1.9(2) there is an isomorphism of $\mathbb{C}[\mathfrak{h}] \rtimes W'$ -modules

$$N \cong \mathbb{C}[\mathfrak{h}] \otimes \text{Res}_{W'}^W(V).$$

So N is isomorphic to $\phi^*(M)$ as $\mathbb{C}[\mathfrak{h}] \rtimes W'$ -modules.

Step 4. In this step we compare $(\widehat{(\phi^*(M))}_0)_{\mathfrak{h}'_{\text{reg}}}$ and $(\widehat{N}_0)_{\mathfrak{h}'_{\text{reg}}}$ as $\widehat{\mathcal{D}(\mathfrak{h}'_{\text{reg}})}_0$ -modules. The definition of these $\widehat{\mathcal{D}(\mathfrak{h}'_{\text{reg}})}_0$ -module structures will be given below in terms of connections. By (1.11) we have $N = E \circ R(\widehat{M}_b)$, so we have $\widehat{N}_0 \cong R(\widehat{M}_b)$. Next, by (1.18) we have an isomorphism of $\mathbb{C}[[\mathfrak{h}]]_0 \rtimes W'$ -modules

$$\begin{aligned} R(\widehat{M}_b) &= \hat{\phi}^*(x_{\text{pr}}(\widehat{M}_b)) \\ &= \widehat{(\phi^*(M))}_0. \end{aligned}$$

So we get an isomorphism of $\mathbb{C}[[\mathfrak{h}]]_0 \rtimes W'$ -modules

$$\hat{\Psi} : \widehat{(\phi^*(M))}_0 \rightarrow \widehat{N}_0.$$

Now, let us consider connections on these modules. Note that by Step 2 we have

$$\widehat{(\phi^*(M))}_0_{\mathfrak{h}'_{\text{reg}}} = \hat{\phi}^*(x_{\text{pr}}(\widehat{M}_b)_{\mathfrak{h}_{\text{reg}}}).$$

Write ∇ for the connection on $M_{\mathfrak{h}_{\text{reg}}}$ given by the Dunkl isomorphism for $H_c(W, \mathfrak{h}_{\text{reg}})$. We equip $\widehat{(\phi^*(M))}_0_{\mathfrak{h}'_{\text{reg}}}$ with the connection $\tilde{\nabla}$ given by

$$\tilde{\nabla}_a(x_{\text{pr}}m) = x_{\text{pr}}(\nabla_a(m)), \quad \forall m \in (\widehat{M}_b)_{\mathfrak{h}_{\text{reg}}}, \quad a \in \mathfrak{h}.$$

Let $\nabla^{(b)}$ be the connection on $N_{\mathfrak{h}'_{\text{reg}}}$ given by the Dunkl isomorphism for $H_{c'}(W', \mathfrak{h}'_{\text{reg}})$. This restricts to a connection on $(\widehat{N}_0)_{\mathfrak{h}'_{\text{reg}}}$. We claim that Ψ is compatible with these connections, i.e., we have

$$(2.4) \quad \nabla_a^{(b)}(x_{\text{pr}}m) = x_{\text{pr}}\nabla_a(m), \quad \forall m \in (\widehat{M}_b)_{\mathfrak{h}_{\text{reg}}}.$$

Recall the subspace V of M from Step 3. By Lemma 1.7(1) the map

$$(\widehat{\mathbb{C}[\mathfrak{h}]}_b \otimes_{\mathbb{C}[\mathfrak{h}]} \mathbb{C}[\mathfrak{h}_{\text{reg}}]) \otimes V \rightarrow (\widehat{M}_b)_{\mathfrak{h}_{\text{reg}}}, \quad p \otimes v \mapsto pv$$

is a bijection. So it is enough to prove (2.4) for $m = pv$ with $p \in \widehat{\mathbb{C}[\mathfrak{h}]_b} \otimes_{\mathbb{C}[\mathfrak{h}]} \mathbb{C}[\mathfrak{h}_{\text{reg}}]$, $v \in V$. We have

$$\begin{aligned}
 \nabla_a^{(b)}(x_{\text{pr}}pv) &= (y_a^{(b)} - \sum_{s \in \mathcal{J} \cap W'} \frac{2c_s}{1 - \lambda_s} \frac{\alpha_s(a)}{x_{\alpha_s}^{(b)}}(s - 1))(x_{\text{pr}}pv) \\
 &= x_{\text{pr}}(y_a + \sum_{s \in \mathcal{J}, s \notin W'} \frac{2c_s}{1 - \lambda_s} \frac{\alpha_s(a)}{x_{\alpha_s}} \\
 &\quad - \sum_{s \in \mathcal{J} \cap W'} \frac{2c_s}{1 - \lambda_s} \frac{\alpha_s(a)}{x_{\alpha_s}}(s - 1))(x_{\text{pr}}pv) \\
 &= x_{\text{pr}}(\nabla_a + \sum_{s \in \mathcal{J}, s \notin W'} \frac{2c_s}{1 - \lambda_s} \frac{\alpha_s(a)}{x_{\alpha_s}}s)(x_{\text{pr}}pv) \\
 (2.5) \qquad \qquad \qquad &= x_{\text{pr}}\nabla_a(x_{\text{pr}}pv).
 \end{aligned}$$

Here the first equality is by the Dunkl isomorphism for $H_{c'}(W', \mathfrak{h}'_{\text{reg}})$. The second is by (1.15), (1.16), (1.17) and the fact that $x_{\text{pr}}^2 = x_{\text{pr}}$. The third is by the Dunkl isomorphism for $H_c(W, \mathfrak{h}_{\text{reg}})$. The last is by (1.19). Next, since x_{pr} is the idempotent in $\widehat{\mathbb{C}[\mathfrak{h}]_b}$ corresponding to the component $\mathbb{C}[[\mathfrak{h}]]_b$ in the decomposition (1.14), we have

$$\begin{aligned}
 \nabla_a(x_{\text{pr}}pv) &= (\partial_a(x_{\text{pr}}p))v + x_{\text{pr}}p(\nabla_a v) \\
 &= x_{\text{pr}}(\partial_a(p))v + x_{\text{pr}}p(\nabla_a v) \\
 &= x_{\text{pr}}\nabla_a(pv).
 \end{aligned}$$

Together with (2.5) this implies that

$$\nabla_a^{(b)}(x_{\text{pr}}pv) = x_{\text{pr}}\nabla_a(pv).$$

So (2.4) is proved.

Step 5. In this step we prove the isomorphism (2.1) for $M \in \mathcal{O}_c^\Delta(W, \mathfrak{h})$. Here we need some more notation. For $X = \mathfrak{h}$ or $\mathfrak{h}'_{\text{reg}}$, let U be an open analytic subvariety of X , write $i : U \hookrightarrow X$ for the canonical embedding. For F an analytic coherent sheaf on X we write $i^*(F)$ for the restriction of F to U . If U contains 0 , for an analytic locally free sheaf E over U , we write \widehat{E} for the restriction of E to the formal disc at 0 .

Let $\Omega \subset \mathfrak{h}$ be the open ball defined in (1.3). Let $f : \mathfrak{h} \rightarrow \mathfrak{h}$ be the morphism defined by ϕ . The preimage of Ω via f is an open ball Ω_0 in \mathfrak{h} centered at 0 . We have

$$f(\Omega_0 \cap \mathfrak{h}'_{\text{reg}}) = \Omega \cap \mathfrak{h}_{\text{reg}}.$$

Let $u : \Omega_0 \cap \mathfrak{h}'_{\text{reg}} \hookrightarrow \mathfrak{h}$ and $v : \Omega \cap \mathfrak{h}_{\text{reg}} \hookrightarrow \mathfrak{h}$ be the canonical embeddings. By Step 3 there is an isomorphism of W' -equivariant analytic locally free sheaves over $\Omega_0 \cap \mathfrak{h}'_{\text{reg}}$

$$u^*(N^{\text{an}}) \cong \phi^*(v^*(M^{\text{an}})).$$

By Step 4 there is an isomorphism

$$\widehat{u^*(N^{\text{an}})} \xrightarrow{\sim} \widehat{\phi^*(v^*(M^{\text{an}}))}$$

which is compatible with their connections. It follows from Lemma 2.2 below that there is an isomorphism

$$(u^*(N^{\text{an}}))^{\nabla^{(b)}} \cong \phi^*((v^*(M^{\text{an}}))^{\nabla}).$$

Since $\Omega_0 \cap \widehat{\mathfrak{h}}'_{\text{reg}}$ is homotopy equivalent to $\widehat{\mathfrak{h}}'_{\text{reg}}$ via u , the left hand side is isomorphic to $(N_{\widehat{\mathfrak{h}}'_{\text{reg}}})^{\nabla^{(b)}}$. So we have

$$\kappa_* \circ j_* \circ \text{KZ}(M) \cong \text{KZ}(W', \mathfrak{h})(N),$$

where κ, j are as in (1.4), (1.5). Combined with Step 1 we have the following isomorphisms

$$\begin{aligned} \text{KZ}' \circ \text{Res}_b(M) &\cong \ell_* \circ \text{KZ}(W', \mathfrak{h})(N) \\ (2.6) \quad &\cong \ell_* \circ \kappa_* \circ j_* \circ \text{KZ}(M) \\ &= \iota_* \circ \text{KZ}(M). \end{aligned}$$

They are functorial on M .

LEMMA 2.2. – *Let E be an analytic locally free sheaf over the complex manifold $\widehat{\mathfrak{h}}'_{\text{reg}}$. Let ∇_1, ∇_2 be two integrable connections on E with regular singularities. If there exists an isomorphism $\widehat{\psi} : (\widehat{E}, \nabla_1) \rightarrow (\widehat{E}, \nabla_2)$, then the local systems E^{∇_1} and E^{∇_2} are isomorphic.*

Proof. – Write $\text{End}(E)$ for the sheaf of endomorphisms of E . Then $\text{End}(E)$ is a locally free sheaf over $\widehat{\mathfrak{h}}'_{\text{reg}}$. The connections ∇_1, ∇_2 define a connection ∇ on $\text{End}(E)$ as follows,

$$\nabla : \text{End}(E) \rightarrow \text{End}(E), \quad f \mapsto \nabla_2 \circ f - f \circ \nabla_1.$$

So the isomorphism $\widehat{\psi}$ is a horizontal section of $(\widehat{\text{End}(E)}, \nabla)$. Let $(\text{End}(E)^{\nabla})_0$ be the set of germs of horizontal sections of $(\text{End}(E), \nabla)$ on zero. By the Comparison theorem [12, Theorem 6.3.1] the canonical map $(\text{End}(E)^{\nabla})_0 \rightarrow (\widehat{\text{End}(E)})^{\nabla}$ is bijective. Hence there exists a holomorphic isomorphism $\psi : (E, \nabla_1) \rightarrow (E, \nabla_2)$ which maps to $\widehat{\psi}$. Now, let U be an open ball in $\widehat{\mathfrak{h}}'_{\text{reg}}$ centered at 0 with radius ε small enough such that the holomorphic isomorphism ψ converges in U . Write E_U for the restriction of E to U . Then ψ induces an isomorphism of local systems $(E_U)^{\nabla_1} \cong (E_U)^{\nabla_2}$. Since $\widehat{\mathfrak{h}}'_{\text{reg}}$ is homotopy equivalent to U , we have

$$E^{\nabla_1} \cong E^{\nabla_2}. \quad \square$$

Step 6. Finally, write I for the inclusion of $\text{Proj}_c(W, \mathfrak{h})$ into $\mathcal{O}_c(W, \mathfrak{h})$. By Lemma 1.1(1) any projective object in $\mathcal{O}_c(W, \mathfrak{h})$ has a standard filtration, so (2.6) yields an isomorphism of functors

$$\text{KZ}' \circ \text{Res}_b \circ I \rightarrow \iota_* \circ \text{KZ} \circ I.$$

Applying Lemma 1.2 to the exact functors $\text{KZ}' \circ \text{Res}_b$ and $\iota_* \circ \text{KZ}$ yields that there is an isomorphism of functors

$$\text{KZ}' \circ \text{Res}_b \cong \iota_* \circ \text{KZ}. \quad \square$$

2.2. – We give some corollaries of Theorem 2.1.

COROLLARY 2.3. – *There is an isomorphism of functors*

$$\text{KZ} \circ \text{Ind}_b \cong \mathscr{L} \text{coInd}_{W'}^W \circ \text{KZ}'.$$

Proof. – To simplify notation let us write

$$\mathcal{O} = \mathcal{O}_c(W, \mathfrak{h}), \quad \mathcal{O}' = \mathcal{O}_{c'}(W', \bar{\mathfrak{h}}), \quad \mathcal{H} = \mathcal{H}_q(W), \quad \mathcal{H}' = \mathcal{H}_{q'}(W').$$

Recall that the functor KZ is represented by a projective object P_{KZ} in \mathcal{O} . So for any $N \in \mathcal{O}'$ we have a morphism of \mathcal{H} -modules

$$\begin{aligned} \text{KZ} \circ \text{Ind}_b(N) &\cong \text{Hom}_{\mathcal{O}}(P_{\text{KZ}}, \text{Ind}_b(N)) \\ &\cong \text{Hom}_{\mathcal{O}'}(\text{Res}_b(P_{\text{KZ}}), N) \\ (2.7) \quad &\rightarrow \text{Hom}_{\mathcal{H}'}(\text{KZ}'(\text{Res}_b(P_{\text{KZ}})), \text{KZ}'(N)). \end{aligned}$$

By Theorem 2.1 we have

$$\text{KZ}' \circ \text{Res}_b(P_{\text{KZ}}) \cong {}^{\mathcal{H}}\text{Res}_{W'}^W \circ \text{KZ}(P_{\text{KZ}}).$$

Recall from Section 1.5 that the \mathcal{H} -module $\text{KZ}(P_{\text{KZ}})$ is isomorphic to \mathcal{H} . So as \mathcal{H}' -modules $\text{KZ}'(\text{Res}_b(P_{\text{KZ}}))$ is also isomorphic to \mathcal{H} . Therefore the morphism (2.7) rewrites as

$$(2.8) \quad \chi(N) : \text{KZ} \circ \text{Ind}_b(N) \rightarrow \text{Hom}_{\mathcal{H}'}(\mathcal{H}, \text{KZ}'(N)).$$

It yields a morphism of functors

$$\chi : \text{KZ} \circ \text{Ind}_b \rightarrow {}^{\mathcal{H}}\text{coInd}_{W'}^W \circ \text{KZ}'.$$

Note that if N is a projective object in \mathcal{O}' , then $\chi(N)$ is an isomorphism by Proposition 1.4(1). So Lemma 1.2 implies that χ is an isomorphism of functors, because both functors $\text{KZ} \circ \text{Ind}_b$ and ${}^{\mathcal{H}}\text{coInd}_{W'}^W \circ \text{KZ}'$ are exact. \square

2.3. – The following lemma will be useful to us.

LEMMA 2.4. – *Let K, L be two right exact functors from \mathcal{O}_1 to \mathcal{O}_2 , where \mathcal{O}_1 and \mathcal{O}_2 can be either $\mathcal{O}_c(W, \mathfrak{h})$ or $\mathcal{O}_{c'}(W', \bar{\mathfrak{h}})$. Let KZ_2 denote the KZ -functor on \mathcal{O}_2 . Suppose that K, L map projective objects to projective ones. Then the vector space homomorphism*

$$(2.9) \quad \text{Hom}(K, L) \rightarrow \text{Hom}(\text{KZ}_2 \circ K, \text{KZ}_2 \circ L), \quad f \mapsto 1_{\text{KZ}_2} f,$$

is an isomorphism.

Notice that if $K = L$, this is even an isomorphism of rings.

Proof. – Let $\text{Proj}_1, \text{Proj}_2$ be respectively the subcategory of projective objects in $\mathcal{O}_1, \mathcal{O}_2$. Write \tilde{K}, \tilde{L} for the functors from Proj_1 to Proj_2 given by the restrictions of K, L , respectively. Let \mathcal{H}_2 be the Hecke algebra corresponding to \mathcal{O}_2 . Since the functor KZ_2 is fully faithful over Proj_2 by Proposition 1.4(1), the following functor

$$\text{Fct}(\text{Proj}_1, \text{Proj}_2) \rightarrow \text{Fct}(\text{Proj}_1, \mathcal{H}_2\text{-mod}), \quad G \mapsto \text{KZ}_2 \circ G$$

is also fully faithful. This yields an isomorphism

$$\text{Hom}(\tilde{K}, \tilde{L}) \xrightarrow{\sim} \text{Hom}(\text{KZ}_2 \circ \tilde{K}, \text{KZ}_2 \circ \tilde{L}), \quad f \mapsto 1_{\text{KZ}_2} f.$$

Next, by Lemma 1.2 the canonical morphisms

$$\text{Hom}(K, L) \rightarrow \text{Hom}(\tilde{K}, \tilde{L}), \quad \text{Hom}(\text{KZ}_2 \circ K, \text{KZ}_2 \circ L) \rightarrow \text{Hom}(\text{KZ}_2 \circ \tilde{K}, \text{KZ}_2 \circ \tilde{L})$$

are isomorphisms. So the map (2.9) is also an isomorphism. \square

Let $b(W, W'')$ be a point in \mathfrak{h} whose stabilizer is W'' . Let $b(W', W'')$ be its image in $\bar{\mathfrak{h}} = \mathfrak{h}/\mathfrak{h}^{W'}$ via the canonical projection. Write $b(W, W') = b$.

COROLLARY 2.5. – *There are isomorphisms of functors*

$$\begin{aligned} \text{Res}_{b(W', W'')} \circ \text{Res}_{b(W, W')} &\cong \text{Res}_{b(W, W'')}, \\ \text{Ind}_{b(W, W')} \circ \text{Ind}_{b(W', W'')} &\cong \text{Ind}_{b(W, W'')}. \end{aligned}$$

Proof. – Since the restriction functors map projective objects to projective ones by Proposition 1.6(1), Lemma 2.4 applied to the categories $\mathcal{O}_1 = \mathcal{O}_c(W, \mathfrak{h})$, $\mathcal{O}_2 = \mathcal{O}_{c''}(W'', \mathfrak{h}/\mathfrak{h}^{W''})$ yields an isomorphism

$$\begin{aligned} \text{Hom}(\text{Res}_{b(W', W'')} \circ \text{Res}_{b(W, W')}, \text{Res}_{b(W, W'')}) \\ \cong \text{Hom}(\text{KZ}'' \circ \text{Res}_{b(W', W'')} \circ \text{Res}_{b(W, W')}, \text{KZ}'' \circ \text{Res}_{b(W, W'')}). \end{aligned}$$

By Theorem 2.1 the set on the second row is

$$(2.10) \quad \text{Hom}({}^{\mathcal{H}}\text{Res}_{W''}^{W'} \circ {}^{\mathcal{H}}\text{Res}_{W'}^W \circ \text{KZ}, {}^{\mathcal{H}}\text{Res}_{W''}^W \circ \text{KZ}).$$

By the presentations of Hecke algebras in [6, Proposition 4.22], there is an isomorphism

$$\sigma : {}^{\mathcal{H}}\text{Res}_{W''}^{W'} \circ {}^{\mathcal{H}}\text{Res}_{W'}^W \xrightarrow{\sim} {}^{\mathcal{H}}\text{Res}_{W''}^W.$$

Hence the element $\sigma 1_{\text{KZ}}$ in the set (2.10) maps to an isomorphism

$$\text{Res}_{b(W', W'')} \circ \text{Res}_{b(W, W')} \cong \text{Res}_{b(W, W'')}.$$

This proves the first isomorphism in the corollary. The second one follows from the uniqueness of right adjoint functor. \square

2.4. Biadjointness of Res_b and Ind_b

Recall that a finite dimensional \mathbb{C} -algebra A is symmetric if A is isomorphic to $A^* = \text{Hom}_{\mathbb{C}}(A, \mathbb{C})$ as (A, A) -bimodules.

LEMMA 2.6. – *Assume that $\mathcal{H}_q(W)$ and $\mathcal{H}_{q'}(W')$ are symmetric algebras. Then the functors ${}^{\mathcal{H}}\text{Ind}_{W'}^W$ and ${}^{\mathcal{H}}\text{coInd}_{W'}^W$ are isomorphic, i.e., the functor ${}^{\mathcal{H}}\text{Ind}_{W'}^W$ is biadjoint to ${}^{\mathcal{H}}\text{Res}_{W'}^W$.*

Proof. – We abbreviate $\mathcal{H} = \mathcal{H}_q(W)$ and $\mathcal{H}' = \mathcal{H}_{q'}(W')$. Since \mathcal{H} is free as a left \mathcal{H}' -module, for any \mathcal{H}' -module M the map

$$(2.11) \quad \text{Hom}_{\mathcal{H}'}(\mathcal{H}, \mathcal{H}') \otimes_{\mathcal{H}'} M \rightarrow \text{Hom}_{\mathcal{H}'}(\mathcal{H}, M)$$

given by multiplication is an isomorphism of \mathcal{H} -modules. By assumption \mathcal{H}' is isomorphic to $(\mathcal{H}')^*$ as $(\mathcal{H}', \mathcal{H}')$ -bimodules. Thus we have the following $(\mathcal{H}, \mathcal{H}')$ -bimodule isomorphisms

$$\begin{aligned} \text{Hom}_{\mathcal{H}'}(\mathcal{H}, \mathcal{H}') &\cong \text{Hom}_{\mathcal{H}'}(\mathcal{H}, (\mathcal{H}')^*) \\ &\cong \text{Hom}_{\mathbb{C}}(\mathcal{H}' \otimes_{\mathcal{H}'} \mathcal{H}, \mathbb{C}) \\ &\cong \mathcal{H}^* \\ &\cong \mathcal{H}. \end{aligned}$$

The last isomorphism follows from the fact the \mathcal{H} is symmetric. Thus, by (2.11) the functors ${}^{\mathcal{H}}\text{Ind}_W^W$, and ${}^{\mathcal{H}}\text{coInd}_W^W$, are isomorphic. \square

REMARK 2.7. – It is proved that $\mathcal{H}_q(W)$ is a symmetric algebra for all irreducible complex reflection group W except for some of the 34 exceptional groups in the Shephard-Todd classification. See [5, Section 2A] for details.

The biadjointness of Res_b and Ind_b was conjectured in [4, Remark 3.18] and was announced by I. Gordon and M. Martino. We give a proof in Proposition 2.9 since it seems not yet to be available in the literature. Let us first consider the following lemma.

LEMMA 2.8. – (1) Let A, B be noetherian algebras and T be a functor

$$T : A\text{-mod} \rightarrow B\text{-mod}.$$

If T is right exact and commutes with direct sums, then it has a right adjoint.

(2) The functor

$$\text{Res}_b : \mathcal{O}_c(W, \mathfrak{h}) \rightarrow \mathcal{O}_{c'}(W', \bar{\mathfrak{h}})$$

has a left adjoint.

Proof. – (1) Consider the (B, A) -bimodule $M = T(A)$. We claim that the functor T is isomorphic to the functor $M \otimes_A -$. Indeed, by definition we have $T(A) \cong M \otimes_A A$ as B -modules. Now, for any $N \in A\text{-mod}$, since N is finitely generated and A is noetherian there exist $m, n \in \mathbb{N}$ and an exact sequence

$$A^{\oplus n} \rightarrow A^{\oplus m} \rightarrow N \rightarrow 0.$$

Since both T and $M \otimes_A -$ are right exact and they commute with direct sums, the fact that $T(A) \cong M \otimes_A A$ implies that $T(N) \cong M \otimes_A N$ as B -modules. This proved the claim. Now, the functor $M \otimes_A -$ has a right adjoint $\text{Hom}_B(M, -)$, so T also has a right adjoint.

(2) Recall that for any complex reflection group W , a contravariant duality functor

$$(-)^\vee : \mathcal{O}_c(W, \mathfrak{h}) \rightarrow \mathcal{O}_{c^\dagger}(W, \mathfrak{h}^*)$$

was defined in [10, Section 4.2], here $c^\dagger : \mathcal{J} \rightarrow \mathbb{C}$ is another parameter explicitly determined by c . Consider the functor

$$\text{Res}_b^\vee = (-)^\vee \circ \text{Res}_b \circ (-)^\vee : \mathcal{O}_{c^\dagger}(W, \mathfrak{h}^*) \rightarrow \mathcal{O}_{c'^\dagger}(W', (\bar{\mathfrak{h}})^*).$$

The category $\mathcal{O}_{c^\dagger}(W, \mathfrak{h}^*)$ has a projective generator P . The algebra $\text{End}_{\mathcal{O}_{c^\dagger}(W, \mathfrak{h}^*)}(P)^{\text{op}}$ is finite dimensional over \mathbb{C} and by Morita theory we have an equivalence of categories

$$\mathcal{O}_{c^\dagger}(W, \mathfrak{h}^*) \cong \text{End}_{\mathcal{O}_{c^\dagger}(W, \mathfrak{h}^*)}(P)^{\text{op}}\text{-mod}.$$

Since the functor Res_b^\vee is exact and obviously commutes with direct sums, by part (1) it has a right adjoint Ψ . Then it follows that $(-)^{\vee} \circ \Psi \circ (-)^{\vee}$ is left adjoint to Res_b . The lemma is proved. \square

PROPOSITION 2.9. – Assume that $\mathcal{H}_q(W)$ and $\mathcal{H}_{q'}(W')$ are symmetric algebras. Then the functor Ind_b is left adjoint to Res_b .

Proof. – Step 1. We abbreviate $\mathcal{O} = \mathcal{O}_c(W, \mathfrak{h})$, $\mathcal{O}' = \mathcal{O}_{c'}(W', \bar{\mathfrak{h}})$, $\mathcal{H} = \mathcal{H}_q(W)$, $\mathcal{H}' = \mathcal{H}_{q'}(W')$, and write $\text{Id}_{\mathcal{O}}, \text{Id}_{\mathcal{O}'}, \text{Id}_{\mathcal{H}}, \text{Id}_{\mathcal{H}'}$ for the identity functor on the corresponding categories. We also abbreviate $E^{\mathcal{H}} = {}^{\mathcal{H}}\text{Res}_{W'}^W$, $F^{\mathcal{H}} = {}^{\mathcal{H}}\text{Ind}_{W'}^W$, and $E = \text{Res}_b$. By Lemma 2.8 the functor E has a left adjoint. We denote it by $F : \mathcal{O}' \rightarrow \mathcal{O}$. Recall the functors

$$\text{KZ} : \mathcal{O} \rightarrow \mathcal{H}\text{-mod}, \quad \text{KZ}' : \mathcal{O}' \rightarrow \mathcal{H}'\text{-mod}.$$

The goal of this step is to show that there exists an isomorphism of functors

$$\text{KZ} \circ F \cong F^{\mathcal{H}} \circ \text{KZ}'.$$

To this end, let S, S' be respectively the right adjoints of KZ, KZ' , see Section 1.5. We will first give an isomorphism of functors

$$F^{\mathcal{H}} \cong \text{KZ} \circ F \circ S'.$$

Let $M \in \mathcal{H}'\text{-mod}$ and $N \in \mathcal{H}\text{-mod}$. Consider the following equalities given by adjunctions

$$\begin{aligned} \text{Hom}_{\mathcal{H}}(\text{KZ} \circ F \circ S'(M), N) &= \text{Hom}_{\mathcal{O}}(F \circ S'(M), S(N)) \\ &= \text{Hom}_{\mathcal{O}'}(S'(M), E \circ S(N)). \end{aligned}$$

The functor KZ' yields a map

$$(2.12) \quad a(M, N) : \text{Hom}_{\mathcal{O}'}(S'(M), E \circ S(N)) \rightarrow \text{Hom}_{\mathcal{H}'}(\text{KZ}' \circ S'(M), \text{KZ}' \circ E \circ S(N)).$$

Since the canonical adjunction maps $\text{KZ}' \circ S' \rightarrow \text{Id}_{\mathcal{H}'}, \text{KZ} \circ S \rightarrow \text{Id}_{\mathcal{H}}$ are isomorphisms (see Section 1.5) and since we have an isomorphism of functors $\text{KZ}' \circ E \cong E^{\mathcal{H}} \circ \text{KZ}$ by Theorem 2.1, we get the following equalities

$$\begin{aligned} \text{Hom}_{\mathcal{H}'}(\text{KZ}' \circ S'(M), \text{KZ}' \circ E \circ S(N)) &= \text{Hom}_{\mathcal{H}'}(M, E^{\mathcal{H}} \circ \text{KZ} \circ S(N)) \\ &= \text{Hom}_{\mathcal{H}'}(M, E^{\mathcal{H}}(N)) \\ &= \text{Hom}_{\mathcal{H}}(F^{\mathcal{H}}(M), N). \end{aligned}$$

In the last equality we used that $F^{\mathcal{H}}$ is left adjoint to $E^{\mathcal{H}}$. So the map (2.12) can be rewritten into the following form

$$a(M, N) : \text{Hom}_{\mathcal{H}}(\text{KZ} \circ F \circ S'(M), N) \rightarrow \text{Hom}_{\mathcal{H}}(F^{\mathcal{H}}(M), N).$$

Now, take $N = \mathcal{H}$. Recall that \mathcal{H} is isomorphic to $\text{KZ}(P_{\text{KZ}})$ as \mathcal{H} -modules. Since P_{KZ} is projective, by Proposition 1.4(2) we have a canonical isomorphism in \mathcal{O}

$$P_{\text{KZ}} \cong S(\text{KZ}(P_{\text{KZ}})) = S(\mathcal{H}).$$

Further E maps projectives to projectives by Proposition 1.6(1), so $E \circ S(\mathcal{H})$ is also projective. Hence Proposition 1.4(1) implies that in this case (2.12) is an isomorphism for any M , i.e., we get an isomorphism

$$a(M, \mathcal{H}) : \text{Hom}_{\mathcal{H}}(\text{KZ} \circ F \circ S'(M), \mathcal{H}) \xrightarrow{\sim} \text{Hom}_{\mathcal{H}}(F^{\mathcal{H}}(M), \mathcal{H}).$$

Further this is an isomorphism of right \mathcal{H} -modules with respect to the \mathcal{H} -actions induced by the right action of \mathcal{H} on itself. Now, the fact that \mathcal{H} is a symmetric algebra yields that

for any finite dimensional \mathcal{H} -module N we have isomorphisms of right \mathcal{H} -modules

$$\begin{aligned}\mathrm{Hom}_{\mathcal{H}}(N, \mathcal{H}) &\cong \mathrm{Hom}_{\mathcal{H}}(N, \mathrm{Hom}_{\mathbb{C}}(\mathcal{H}, \mathbb{C})) \\ &\cong \mathrm{Hom}_{\mathbb{C}}(N, \mathbb{C}).\end{aligned}$$

Therefore $a(M, \mathcal{H})$ yields an isomorphism of right \mathcal{H} -modules

$$\mathrm{Hom}_{\mathbb{C}}(\mathrm{KZ} \circ F \circ S'(M), \mathbb{C}) \rightarrow \mathrm{Hom}_{\mathbb{C}}(F^{\mathcal{H}}(M), \mathbb{C}).$$

We deduce a natural isomorphism of left \mathcal{H} -modules

$$\mathrm{KZ} \circ F \circ S'(M) \cong F^{\mathcal{H}}(M)$$

for any \mathcal{H}' -module M . This gives an isomorphism of functors

$$\psi : \mathrm{KZ} \circ F \circ S' \xrightarrow{\sim} F^{\mathcal{H}}.$$

Finally, consider the canonical adjunction map $\eta : \mathrm{Id}_{\mathcal{O}' } \rightarrow S' \circ \mathrm{KZ}'$. We have a morphism of functors

$$\phi = (1_{\mathrm{KZ} \circ F} \eta) \circ (\psi 1_{\mathrm{KZ}'}) : \mathrm{KZ} \circ F \rightarrow F^{\mathcal{H}} \circ \mathrm{KZ}'.$$

Note that $\psi 1_{\mathrm{KZ}'}$ is an isomorphism of functors. If Q is a projective object in \mathcal{O}' , then by Proposition 1.4(2) the morphism $\eta(Q) : Q \rightarrow S' \circ \mathrm{KZ}'(Q)$ is also an isomorphism, so $\phi(Q)$ is an isomorphism. This implies that ϕ is an isomorphism of functors by Lemma 1.2, because both $\mathrm{KZ} \circ F$ and $F^{\mathcal{H}} \circ \mathrm{KZ}'$ are right exact functors. Here the right exactness of F follows from that it is left adjoint to E . So we get the desired isomorphism of functors

$$\mathrm{KZ} \circ F \cong F^{\mathcal{H}} \circ \mathrm{KZ}'.$$

Step 2. Let us now prove that F is right adjoint to E . By uniqueness of adjoint functors, this will imply that F is isomorphic to Ind_b . First, by Lemma 2.6 the functor $F^{\mathcal{H}}$ is isomorphic to ${}^{\mathcal{H}}\mathrm{coInd}_W^W$. So $F^{\mathcal{H}}$ is right adjoint to $E^{\mathcal{H}}$, i.e., we have morphisms of functors

$$\varepsilon^{\mathcal{H}} : E^{\mathcal{H}} \circ F^{\mathcal{H}} \rightarrow \mathrm{Id}_{\mathcal{H}'}, \quad \eta^{\mathcal{H}} : \mathrm{Id}_{\mathcal{H}} \rightarrow F^{\mathcal{H}} \circ E^{\mathcal{H}}$$

such that

$$(\varepsilon^{\mathcal{H}} 1_{E^{\mathcal{H}}}) \circ (1_{E^{\mathcal{H}}} \eta^{\mathcal{H}}) = 1_{E^{\mathcal{H}}}, \quad (1_{F^{\mathcal{H}}} \varepsilon^{\mathcal{H}}) \circ (\eta^{\mathcal{H}} 1_{F^{\mathcal{H}}}) = 1_{F^{\mathcal{H}}}.$$

Next, both F and E have exact right adjoints, given respectively by E and Ind_b . Therefore F and E map projective objects to projective ones. Applying Lemma 2.4 to $\mathcal{O}_1 = \mathcal{O}_2 = \mathcal{O}'$, $K = E \circ F$, $L = \mathrm{Id}_{\mathcal{O}'}$ yields that the following map is bijective

$$(2.13) \quad \mathrm{Hom}(E \circ F, \mathrm{Id}_{\mathcal{O}'}) \rightarrow \mathrm{Hom}(\mathrm{KZ}' \circ E \circ F, \mathrm{KZ}' \circ \mathrm{Id}_{\mathcal{O}}), \quad f \mapsto 1_{\mathrm{KZ}'} f.$$

By Theorem 2.1 and Step 1 there exist isomorphisms of functors

$$\phi_E : E^{\mathcal{H}} \circ \mathrm{KZ} \xrightarrow{\sim} \mathrm{KZ}' \circ E, \quad \phi_F : F^{\mathcal{H}} \circ \mathrm{KZ}' \xrightarrow{\sim} \mathrm{KZ} \circ F.$$

Let

$$\begin{aligned}\phi_{EF} &= (\phi_E 1_F) \circ (1_{E^{\mathcal{H}}} \phi_F) : E^{\mathcal{H}} \circ F^{\mathcal{H}} \circ \mathrm{KZ}' \xrightarrow{\sim} \mathrm{KZ}' \circ E \circ F, \\ \phi_{FE} &= (\phi_F 1_E) \circ (1_{F^{\mathcal{H}}} \phi_E) : F^{\mathcal{H}} \circ E^{\mathcal{H}} \circ \mathrm{KZ} \xrightarrow{\sim} \mathrm{KZ} \circ F \circ E.\end{aligned}$$

Identify

$$\mathrm{KZ} \circ \mathrm{Id}_{\mathcal{O}} = \mathrm{Id}_{\mathcal{H}} \circ \mathrm{KZ}, \quad \mathrm{KZ}' \circ \mathrm{Id}_{\mathcal{O}'} = \mathrm{Id}_{\mathcal{H}'} \circ \mathrm{KZ}'.$$

We have a bijective map

$$\mathrm{Hom}(\mathrm{KZ}' \circ E \circ F, \mathrm{KZ}' \circ \mathrm{Id}_{\mathcal{O}'}) \xrightarrow{\sim} \mathrm{Hom}(E^{\mathcal{H}} \circ F^{\mathcal{H}} \circ \mathrm{KZ}', \mathrm{Id}_{\mathcal{H}'} \circ \mathrm{KZ}'), \quad g \mapsto g \circ \phi_{EF}.$$

Together with (2.13), it implies that there exists a unique morphism $\varepsilon : E \circ F \rightarrow \mathrm{Id}_{\mathcal{O}'}$ such that

$$(\mathbf{1}_{\mathrm{KZ}'} \varepsilon) \circ \phi_{EF} = \varepsilon^{\mathcal{H}} \mathbf{1}_{\mathrm{KZ}'}.$$

Similarly, there exists a unique morphism $\eta : \mathrm{Id}_{\mathcal{O}} \rightarrow F \circ E$ such that

$$(\phi_{FE})^{-1} \circ (\mathbf{1}_{\mathrm{KZ}} \eta) = \eta^{\mathcal{H}} \mathbf{1}_{\mathrm{KZ}}.$$

Now, we have the following commutative diagram

$$\begin{array}{ccccc} E^{\mathcal{H}} \circ \mathrm{KZ} & \xlongequal{\quad} & E^{\mathcal{H}} \circ \mathrm{KZ} & \xrightarrow{\phi_E} & \mathrm{KZ}' \circ E \\ \downarrow \mathbf{1}_{E^{\mathcal{H}}} \eta^{\mathcal{H}} \mathbf{1}_{\mathrm{KZ}} & & \downarrow \mathbf{1}_{E^{\mathcal{H}}} \mathbf{1}_{\mathrm{KZ}} \eta & & \downarrow \mathbf{1}_{\mathrm{KZ}'} \mathbf{1}_E \eta \\ E^{\mathcal{H}} \circ F^{\mathcal{H}} \circ E^{\mathcal{H}} \circ \mathrm{KZ} & \xrightarrow{\mathbf{1}_{E^{\mathcal{H}}} \phi_{FE}} & E^{\mathcal{H}} \circ \mathrm{KZ} \circ F \circ E & \xrightarrow{\phi_E \mathbf{1}_F \mathbf{1}_E} & \mathrm{KZ}' \circ E \circ F \circ E \\ \parallel & & \uparrow \mathbf{1}_{E^{\mathcal{H}}} \phi_F \mathbf{1}_E & & \parallel \\ E^{\mathcal{H}} \circ F^{\mathcal{H}} \circ E^{\mathcal{H}} \circ \mathrm{KZ} & \xrightarrow{\mathbf{1}_{E^{\mathcal{H}}} \mathbf{1}_F \phi_E} & E^{\mathcal{H}} \circ F^{\mathcal{H}} \circ \mathrm{KZ}' \circ E & \xrightarrow{\phi_{EF} \mathbf{1}_E} & \mathrm{KZ}' \circ E \circ F \circ E \\ \downarrow \varepsilon^{\mathcal{H}} \mathbf{1}_{E^{\mathcal{H}}} \mathbf{1}_{\mathrm{KZ}} & & \downarrow \varepsilon^{\mathcal{H}} \mathbf{1}_{\mathrm{KZ}'} \mathbf{1}_E & & \downarrow \mathbf{1}_{\mathrm{KZ}'} \varepsilon \mathbf{1}_E \\ E^{\mathcal{H}} \circ \mathrm{KZ} & \xrightarrow{\phi_E} & \mathrm{KZ}' \circ E & \xlongequal{\quad} & \mathrm{KZ}' \circ E. \end{array}$$

It yields that

$$(\mathbf{1}_{\mathrm{KZ}'} \varepsilon \mathbf{1}_E) \circ (\mathbf{1}_{\mathrm{KZ}'} \mathbf{1}_E \eta) = \phi_E \circ (\varepsilon^{\mathcal{H}} \mathbf{1}_{E^{\mathcal{H}}} \mathbf{1}_{\mathrm{KZ}}) \circ (\mathbf{1}_{E^{\mathcal{H}}} \eta^{\mathcal{H}} \mathbf{1}_{\mathrm{KZ}}) \circ (\phi_E)^{-1}.$$

We deduce that

$$(2.14) \quad \begin{aligned} \mathbf{1}_{\mathrm{KZ}'}((\varepsilon \mathbf{1}_E) \circ (\mathbf{1}_E \eta)) &= \phi_E \circ (\mathbf{1}_{E^{\mathcal{H}}} \mathbf{1}_{\mathrm{KZ}}) \circ (\phi_E)^{-1} \\ &= \mathbf{1}_{\mathrm{KZ}'} \mathbf{1}_E. \end{aligned}$$

By applying Lemma 2.4 to $\mathcal{O}_1 = \mathcal{O}$, $\mathcal{O}_2 = \mathcal{O}'$, $K = L = E$, we deduce that the following map is bijective

$$\mathrm{End}(E) \rightarrow \mathrm{End}(\mathrm{KZ}' \circ E), \quad f \mapsto \mathbf{1}_{\mathrm{KZ}'} f.$$

Hence (2.14) implies that

$$(\varepsilon \mathbf{1}_E) \circ (\mathbf{1}_E \eta) = \mathbf{1}_E.$$

Similarly, we have $(\mathbf{1}_F \varepsilon) \circ (\eta \mathbf{1}_F) = \mathbf{1}_F$. So E is left adjoint to F . By uniqueness of adjoint functors this implies that F is isomorphic to Ind_b . Therefore Ind_b is biadjoint to Res_b . \square

3. Reminders on the cyclotomic case

From now on we will concentrate on the cyclotomic rational DAHA's. We fix some notation in this section.

3.1. – Let l, n be positive integers. Write $\varepsilon = \exp(\frac{2\pi\sqrt{-1}}{l})$. Let $\mathfrak{h} = \mathbb{C}^n$, write $\{y_1, \dots, y_n\}$ for its standard basis. For $1 \leq i, j, k \leq n$ with i, j, k distinct, let ε_k, s_{ij} be the following elements of $GL(\mathfrak{h})$:

$$\varepsilon_k(y_k) = \varepsilon y_k, \quad \varepsilon_k(y_j) = y_j, \quad s_{ij}(y_i) = y_j, \quad s_{ij}(y_k) = y_k.$$

Let $B_n(l)$ be the subgroup of $GL(\mathfrak{h})$ generated by ε_k and s_{ij} for $1 \leq k \leq n$ and $1 \leq i < j \leq n$. It is a complex reflection group with the set of reflections

$$\phi_n = \{\varepsilon_i^p : 1 \leq i \leq n, 1 \leq p \leq l-1\} \sqcup \{s_{ij}^{(p)} = s_{ij}\varepsilon_i^p\varepsilon_j^{-p} : 1 \leq i < j \leq n, 1 \leq p \leq l\}.$$

Note that there is an obvious inclusion $\phi_{n-1} \hookrightarrow \phi_n$. It yields an embedding

$$(3.1) \quad B_{n-1}(l) \hookrightarrow B_n(l).$$

This embedding identifies $B_{n-1}(l)$ with the parabolic subgroup of $B_n(l)$ given by the stabilizer of the point $b_n = (0, \dots, 0, 1) \in \mathbb{C}^n$.

The cyclotomic rational DAHA is the algebra $H_c(B_n(l), \mathfrak{h})$. We will use another presentation in which we replace the parameter c by an l -tuple $\mathbf{h} = (h, h_1, \dots, h_{l-1})$ such that

$$c_{s_{ij}^{(p)}} = -h, \quad c_{\varepsilon_p} = \frac{-1}{2} \sum_{p'=1}^{l-1} (\varepsilon^{-pp'} - 1)h_{p'}.$$

We will denote $H_c(B_n(l), \mathfrak{h})$ by $H_{\mathbf{h},n}$. The corresponding category \mathcal{O} will be denoted by $\mathcal{O}_{\mathbf{h},n}$. In the rest of the paper, we will fix the positive integer l . We will also fix a positive integer $e \geq 2$ and an l -tuple of integers $\mathbf{s} = (s_1, \dots, s_l)$. We will always assume that the parameter \mathbf{h} is given by the following formulas,

$$(3.2) \quad h = \frac{-1}{e}, \quad h_p = \frac{s_{p+1} - s_p}{e} - \frac{1}{l}, \quad 1 \leq p \leq l-1.$$

The functor $\text{KZ}(B_n(l), \mathbb{C}^n)$ goes from $\mathcal{O}_{\mathbf{h},n}$ to the category of finite dimensional modules of a certain Hecke algebra $\mathcal{H}_{\mathbf{q},n}$ attached to the group $B_n(l)$. Here the parameter is $\mathbf{q} = (q, q_1, \dots, q_l)$ with

$$q = \exp(2\pi\sqrt{-1}/e), \quad q_p = q^{s_p}, \quad 1 \leq p \leq l.$$

The algebra $\mathcal{H}_{\mathbf{q},n}$ has the following presentation:

- Generators: T_0, T_1, \dots, T_{n-1} ,
- Relations:

$$\begin{aligned} (T_0 - q_1) \cdots (T_0 - q_l) &= (T_i + 1)(T_i - q) = 0, \quad 1 \leq i \leq n-1, \\ T_0 T_1 T_0 T_1 &= T_1 T_0 T_1 T_0, \\ T_i T_j &= T_j T_i, \quad \text{if } |i - j| > 1, \\ T_i T_{i+1} T_i &= T_{i+1} T_i T_{i+1}, \quad 1 \leq i \leq n-2. \end{aligned}$$

The algebra $\mathcal{H}_{\mathbf{q},n}$ satisfies the assumption of Section 2, i.e., it has the same dimension as $\mathbb{C}B_n(l)$.

3.2. – For each positive integer n , the embedding (3.1) of $B_{n-1}(l)$ into $B_n(l)$ yields an embedding of Hecke algebras

$$\iota_{\mathbf{q}} : \mathcal{H}_{\mathbf{q},n-1} \hookrightarrow \mathcal{H}_{\mathbf{q},n},$$

see Section 1.2. Under the presentation above this embedding is given by

$$\iota_{\mathbf{q}}(T_i) = T_i, \quad \forall 0 \leq i \leq n-2,$$

see [6, Proposition 2.29].

We will consider the following restriction and induction functors:

$$\begin{aligned} E(n) &= \text{Res}_{b_n}, & E(n)^{\mathcal{H}} &= {}^{\mathcal{H}}\text{Res}_{B_{n-1}(l)}^{B_n(l)}, \\ F(n) &= \text{Ind}_{b_n}, & F(n)^{\mathcal{H}} &= {}^{\mathcal{H}}\text{Ind}_{B_{n-1}(l)}^{B_n(l)}. \end{aligned}$$

The algebra $\mathcal{H}_{\mathbf{q},n}$ is symmetric (see Remark 2.7). Hence by Lemma 2.6 we have

$$F(n)^{\mathcal{H}} \cong {}^{\mathcal{H}}\text{coInd}_{B_{n-1}(l)}^{B_n(l)}.$$

We will abbreviate

$$\mathcal{O}_{\mathbf{h},\mathbb{N}} = \bigoplus_{n \in \mathbb{N}} \mathcal{O}_{\mathbf{h},n}, \quad \text{KZ} = \bigoplus_{n \in \mathbb{N}} \text{KZ}(B_n(l), \mathbb{C}^n), \quad \mathcal{H}_{\mathbf{q},\mathbb{N}}\text{-mod} = \bigoplus_{n \in \mathbb{N}} \mathcal{H}_{\mathbf{q},n}\text{-mod}.$$

So KZ is the Knizhnik-Zamolodchikov functor from $\mathcal{O}_{\mathbf{h},\mathbb{N}}$ to $\mathcal{H}_{\mathbf{q},\mathbb{N}}\text{-mod}$. Let

$$\begin{aligned} E &= \bigoplus_{n \geq 1} E(n), & E^{\mathcal{H}} &= \bigoplus_{n \geq 1} E(n)^{\mathcal{H}}, \\ F &= \bigoplus_{n \geq 1} F(n), & F^{\mathcal{H}} &= \bigoplus_{n \geq 1} F(n)^{\mathcal{H}}. \end{aligned}$$

So $(E^{\mathcal{H}}, F^{\mathcal{H}})$ is a pair of biadjoint endo-functors of $\mathcal{H}_{\mathbf{q},\mathbb{N}}\text{-mod}$, and (E, F) is a pair of biadjoint endo-functors of $\mathcal{O}_{\mathbf{h},\mathbb{N}}$ by Proposition 2.9.

3.3. Fock spaces

Recall that an l -partition is an l -tuple $\lambda = (\lambda^1, \dots, \lambda^l)$ with each λ^j a partition, that is a sequence of integers $(\lambda^j)_1 \geq \dots \geq (\lambda^j)_k > 0$. To any l -partition $\lambda = (\lambda^1, \dots, \lambda^l)$ we attach the set

$$\Upsilon_{\lambda} = \{(a, b, j) \in \mathbb{N} \times \mathbb{N} \times (\mathbb{Z}/l\mathbb{Z}) : 0 < b \leq (\lambda^j)_a\}.$$

Write $|\lambda|$ for the number of elements in this set, we say that λ is an l -partition of $|\lambda|$. For $n \in \mathbb{N}$ we denote by $\mathcal{P}_{n,l}$ the set of l -partitions of n . For any l -partition μ such that Υ_{μ} contains Υ_{λ} , we write μ/λ for the complement of Υ_{λ} in Υ_{μ} . Let $|\mu/\lambda|$ be the number of elements in this set. To each element (a, b, j) in Υ_{λ} we attach an element

$$\text{res}((a, b, j)) = b - a + s_j \in \mathbb{Z}/e\mathbb{Z},$$

called the residue of (a, b, j) . Here s_j is the j -th component of our fixed l -tuple \mathbf{s} .

The Fock space with multi-charge \mathbf{s} is the \mathbb{C} -vector space $\mathcal{F}_{\mathbf{s}}$ spanned by the l -partitions, i.e.,

$$\mathcal{F}_{\mathbf{s}} = \bigoplus_{n \in \mathbb{N}} \bigoplus_{\lambda \in \mathcal{P}_{n,l}} \mathbb{C}\lambda.$$

It admits an integrable $\tilde{\mathfrak{sl}}_e$ -module structure such that the Chevalley generators act as follows (cf. [11]): for any $i \in \mathbb{Z}/e\mathbb{Z}$,

$$(3.3) \quad e_i(\lambda) = \sum_{|\lambda/\mu|=1, \text{res}(\lambda/\mu)=i} \mu, \quad f_i(\lambda) = \sum_{|\mu/\lambda|=1, \text{res}(\mu/\lambda)=i} \mu.$$

Let n_i be the number of elements in the set $\{(a, b, j) \in \Upsilon_\lambda : \text{res}((a, b, j)) = i\}$. The element $\partial \in \tilde{\mathfrak{sl}}_e$ acts on \mathcal{F}_s by

$$\partial(\lambda) = -n_0\lambda.$$

For each $n \in \mathbb{Z}$ set $\Lambda_n = \Lambda_{\underline{n}}$, where \underline{n} is the image of n in $\mathbb{Z}/e\mathbb{Z}$ and $\Lambda_{\underline{n}}$ is the corresponding fundamental weight of $\tilde{\mathfrak{sl}}_e$. Set

$$\Lambda_s = \Lambda_{s_1} + \cdots + \Lambda_{s_l}.$$

Each l -partition λ is a weight vector of \mathcal{F}_s with weight

$$(3.4) \quad \text{wt}(\lambda) = \Lambda_s - \sum_{i \in \mathbb{Z}/e\mathbb{Z}} n_i \alpha_i.$$

We will call $\text{wt}(\lambda)$ the weight of λ .

In [14, Section 6.1.1] an explicit bijection was given between the sets $\text{Irr}(B_n(l))$ and $\mathcal{P}_{n,l}$. Using this bijection we identify these two sets and index the standard and simple modules in $\mathcal{O}_{\mathbf{h},\mathbb{N}}$ by l -partitions. In particular, we have an isomorphism of \mathbb{C} -vector spaces

$$(3.5) \quad \theta : K(\mathcal{O}_{\mathbf{h},\mathbb{N}}) \xrightarrow{\sim} \mathcal{F}_s, \quad [\Delta(\lambda)] \mapsto \lambda.$$

3.4. – We end this section by the following lemma. Recall that the functor KZ gives a map $K(\mathcal{O}_{\mathbf{h},n}) \rightarrow K(\mathcal{H}_{\mathbf{q},n})$. For any l -partition λ of n let S_λ be the corresponding Specht module in $\mathcal{H}_{\mathbf{q},n}$ -mod, see [2, Definition 13.22] for its definition.

LEMMA 3.1. – *In $K(\mathcal{H}_{\mathbf{q},n})$, we have $\text{KZ}([\Delta(\lambda)]) = [S_\lambda]$.*

Proof. – Let R be any commutative ring over \mathbb{C} . For any l -tuple $\mathbf{z} = (z, z_1, \dots, z_{l-1})$ of elements in R one defines the rational DAHA over R attached to $B_n(l)$ with parameter \mathbf{z} in the same way as before. Denote it by $H_{R,\mathbf{z},n}$. The standard modules $\Delta_R(\lambda)$ are also defined as before. For any $(l+1)$ -tuple $\mathbf{u} = (u, u_1, \dots, u_l)$ of invertible elements in R the Hecke algebra $\mathcal{H}_{R,\mathbf{u},n}$ over R attached to $B_n(l)$ with parameter \mathbf{u} is defined by the same presentation as in Section 3.1. The Specht modules $S_{R,\lambda}$ are also well-defined (see [2]). If R is a field, we will write $\text{Irr}(\mathcal{H}_{R,\mathbf{u},n})$ for the set of isomorphism classes of simple $\mathcal{H}_{R,\mathbf{u},n}$ -modules.

Now, fix R to be the ring of holomorphic functions of one variable ϖ . We choose $\mathbf{z} = (z, z_1, \dots, z_{l-1})$ to be given by

$$z = l\varpi, \quad z_p = (s_{p+1} - s_p)l\varpi + e\varpi, \quad 1 \leq p \leq l-1.$$

Write $x = \exp(-2\pi\sqrt{-1}\varpi)$. Let $\mathbf{u} = (u, u_1, \dots, u_l)$ be given by

$$u = x^l, \quad u_p = \varepsilon^{p-1} x^{s_p l - (p-1)e}, \quad 1 \leq p \leq l.$$

By [6, Theorem 4.12] the same definition as in Section 1.5 yields a well defined $\mathcal{H}_{R,\mathbf{u},n}$ -module

$$T_R(\lambda) = \text{KZ}_R(\Delta_R(\lambda)).$$

It is a free R -module of finite rank and it commutes with the base change functor by the existence and unicity theorem for linear differential equations, i.e., for any ring homomorphism $R \rightarrow R'$ over \mathbb{C} , we have a canonical isomorphism of $\mathcal{H}_{R',\mathbf{u},n}$ -modules

$$(3.6) \quad T_{R'}(\lambda) = \text{KZ}_{R'}(\Delta_{R'}(\lambda)) \cong T_R(\lambda) \otimes_R R'.$$

In particular, for any ring homomorphism $a : R \rightarrow \mathbb{C}$. Write \mathbb{C}_a for the vector space \mathbb{C} equipped with the R -module structure given by a . Let $a(\mathbf{z}), a(\mathbf{u})$ denote the images of \mathbf{z}, \mathbf{u} by a . Note that we have $H_{a(\mathbf{z}),n} = H_{R,\mathbf{z},n} \otimes_R \mathbb{C}_a$ and $\mathcal{H}_{a(\mathbf{u}),n} = \mathcal{H}_{R,\mathbf{u},n} \otimes_R \mathbb{C}_a$. Denote the Knizhnik-Zamolodchikov functor of $H_{a(\mathbf{z}),n}$ by $\text{KZ}_{a(\mathbf{z})}$ and the standard module corresponding to λ by $\Delta_{a(\mathbf{z})}(\lambda)$. Then we have an isomorphism of $\mathcal{H}_{a(\mathbf{u}),n}$ -modules

$$T_R(\lambda) \otimes_R \mathbb{C}_a \cong \text{KZ}_{a(\mathbf{z})}(\Delta_{a(\mathbf{z})}(\lambda)).$$

Let K be the fraction field of R . By [10, Theorem 2.19] the category $\mathcal{O}_{K,\mathbf{z},n}$ is split semisimple. In particular, the standard modules are simple. We have

$$\{T_K(\lambda), \lambda \in \mathcal{P}_{n,l}\} = \text{Irr}(\mathcal{H}_{K,\mathbf{u},n}).$$

The Hecke algebra $\mathcal{H}_{K,\mathbf{u},n}$ is also split semisimple and we have

$$\{S_{K,\lambda}, \lambda \in \mathcal{P}_{n,l}\} = \text{Irr}(\mathcal{H}_{K,\mathbf{u},n}),$$

see for example [2, Corollary 13.9]. Thus there is a bijection $\varphi : \mathcal{P}_{n,l} \rightarrow \mathcal{P}_{n,l}$ such that $T_K(\lambda)$ is isomorphic to $S_{K,\varphi(\lambda)}$ for all λ . We claim that φ is identity. To see this, consider the algebra homomorphism $a_0 : R \rightarrow \mathbb{C}$ given by $\varpi \mapsto 0$. Then $\mathcal{H}_{a_0(\mathbf{u}),n}$ is canonically isomorphic to the group algebra $\mathbb{C}B_n(l)$, thus it is semi-simple. Let \overline{K} be the algebraic closure of K . Let \overline{R} be the integral closure of R in \overline{K} and fix an extension \overline{a}_0 of a_0 to \overline{R} . By Tit's deformation theorem (see for example [9, Section 68A]), there is a bijection

$$\psi : \text{Irr}(\mathcal{H}_{\overline{K},\mathbf{u},n}) \xrightarrow{\sim} \text{Irr}(\mathcal{H}_{\overline{a}_0(\mathbf{u}),n})$$

such that

$$\psi(T_{\overline{K}}(\lambda)) = T_{\overline{R}}(\lambda) \otimes_{\overline{R}} \mathbb{C}_{\overline{a}_0}, \quad \psi(S_{\overline{K},\lambda}) = S_{\overline{R},\lambda} \otimes_{\overline{R}} \mathbb{C}_{\overline{a}_0}.$$

By the definition of Specht modules we have $S_{\overline{R},\lambda} \otimes_{\overline{R}} \mathbb{C}_{\overline{a}_0} \cong \lambda$ as $\mathbb{C}B_n(l)$ -modules. On the other hand, since $a_0(\mathbf{z}) = 0$, by (3.6) we have the following isomorphisms

$$\begin{aligned} T_{\overline{R}}(\lambda) \otimes_{\overline{R}} \mathbb{C}_{\overline{a}_0} &\cong T_R(\lambda) \otimes_R \mathbb{C}_{a_0} \\ &\cong \text{KZ}_0(\Delta_0(\lambda)) \\ &= \lambda. \end{aligned}$$

So $\psi(T_{\overline{K}}(\lambda)) = \psi(S_{\overline{K},\lambda})$. Hence we have $T_{\overline{K}}(\lambda) \cong S_{\overline{K},\lambda}$. Since $T_{\overline{K}}(\lambda) = T_K(\lambda) \otimes_K \overline{K}$ is isomorphic to $S_{\overline{K},\varphi(\lambda)} = S_{K,\varphi(\lambda)} \otimes_K \overline{K}$, we deduce that $\varphi(\lambda) = \lambda$. The claim is proved.

Finally, let \mathfrak{m} be the maximal ideal of R consisting of the functions vanishing at $\varpi = -1/el$. Let \widehat{R} be the completion of R at \mathfrak{m} . It is a discrete valuation ring with residue field \mathbb{C} . Let $a_1 : \widehat{R} \rightarrow \widehat{R}/\mathfrak{m}\widehat{R} = \mathbb{C}$ be the quotient map. We have $a_1(\mathbf{z}) = \mathbf{h}$ and $a_1(\mathbf{u}) = \mathbf{q}$. Let \widehat{K} be the fraction field of \widehat{R} . Recall that the decomposition map is given by

$$d : K(\mathcal{H}_{\widehat{K},\mathbf{u},n}) \rightarrow K(\mathcal{H}_{\mathbf{q},n}), \quad [M] \mapsto [L \otimes_{\widehat{R}} \mathbb{C}_{a_1}].$$

Here L is any free \widehat{R} -submodule of M such that $L \otimes_{\widehat{R}} \widehat{K} = M$. The choice of L does not affect the class $[L \otimes_{\widehat{R}} \mathbb{C}_{a_1}]$ in $K(\mathcal{H}_{\mathbf{q},n})$. See [2, Section 13.3] for details on this map. Now, observe that we have

$$\begin{aligned} d([S_{\widehat{K},\lambda}]) &= [S_{\widehat{R},\lambda} \otimes_{\widehat{R}} \mathbb{C}_{a_1}] = [S_\lambda], \\ d([T_{\widehat{K}}(\lambda)]) &= [T_{\widehat{R}}(\lambda) \otimes_{\widehat{R}} \mathbb{C}_{a_1}] = [\text{KZ}(\Delta(\lambda))]. \end{aligned}$$

Since \widehat{K} is an extension of K , by the last paragraph we have $[S_{\widehat{K},\lambda}] = [T_{\widehat{K}}(\lambda)]$. We deduce that $[\text{KZ}(\Delta(\lambda))] = [S_\lambda]$. □

4. i -restriction and i -induction

We define in this section the i -restriction and i -induction functors for the cyclotomic rational DAHA's. This is done in parallel with the Hecke algebra case.

4.1. – Let us recall the definition of the i -restriction and i -induction functors for $\mathcal{H}_{\mathbf{q},n}$. First define the Jucy-Murphy elements J_0, \dots, J_{n-1} in $\mathcal{H}_{\mathbf{q},n}$ by

$$J_0 = T_0, \quad J_i = q^{-1}T_i J_{i-1} T_i \quad \text{for } 1 \leq i \leq n-1.$$

Write $Z(\mathcal{H}_{\mathbf{q},n})$ for the center of $\mathcal{H}_{\mathbf{q},n}$. For any symmetric polynomial σ of n variables the element $\sigma(J_0, \dots, J_{n-1})$ belongs to $Z(\mathcal{H}_{\mathbf{q},n})$ (cf. [2, Section 13.1]). In particular, if z is a formal variable the polynomial $C_n(z) = \prod_{i=0}^{n-1} (z - J_i)$ in $\mathcal{H}_{\mathbf{q},n}[z]$ has coefficients in $Z(\mathcal{H}_{\mathbf{q},n})$.

Now, for any $a(z) \in \mathbb{C}(z)$ let $P_{n,a(z)}$ be the exact endo-functor of the category $\mathcal{H}_{\mathbf{q},n}$ -mod that maps an object M to the generalized eigenspace of $C_n(z)$ in M with the eigenvalue $a(z)$.

For any $i \in \mathbb{Z}/e\mathbb{Z}$ the i -restriction functor and i -induction functor

$$E_i(n)^{\mathscr{C}} : \mathcal{H}_{\mathbf{q},n}\text{-mod} \rightarrow \mathcal{H}_{\mathbf{q},n-1}\text{-mod}, \quad F_i(n)^{\mathscr{C}} : \mathcal{H}_{\mathbf{q},n-1}\text{-mod} \rightarrow \mathcal{H}_{\mathbf{q},n}\text{-mod}$$

are defined as follows (cf. [2, Definition 13.33]):

$$(4.1) \quad E_i(n)^{\mathscr{C}} = \bigoplus_{a(z) \in \mathbb{C}(z)} P_{n-1,a(z)/(z-q^i)} \circ E(n)^{\mathscr{C}} \circ P_{n,a(z)},$$

$$(4.2) \quad F_i(n)^{\mathscr{C}} = \bigoplus_{a(z) \in \mathbb{C}(z)} P_{n,a(z)(z-q^i)} \circ F(n)^{\mathscr{C}} \circ P_{n-1,a(z)}.$$

We will write

$$(4.3) \quad E_i^{\mathscr{C}} = \bigoplus_{n \geq 1} E_i(n)^{\mathscr{C}}, \quad F_i^{\mathscr{C}} = \bigoplus_{n \geq 1} F_i(n)^{\mathscr{C}}.$$

They are endo-functors of $\mathcal{H}_{\mathbf{q},\mathbb{N}}$ -mod. For each $\lambda \in \mathscr{P}_{n,l}$ set

$$a_\lambda(z) = \prod_{v \in \Upsilon_\lambda} (z - q^{\text{res}(v)}).$$

We recall some properties of these functors in the following proposition.

PROPOSITION 4.1. – (1) *The functors $E_i(n)^{\mathscr{E}}, F_i(n)^{\mathscr{E}}$ are exact. The functor $E_i(n)^{\mathscr{E}}$ is biadjoint to $F_i(n)^{\mathscr{E}}$.*

(2) *For any $\lambda \in \mathscr{P}_{n,l}$ the element $C_n(z)$ has a unique eigenvalue on the Specht module S_λ . It is equal to $a_\lambda(z)$.*

(3) *We have*

$$E_i(n)^{\mathscr{E}}([S_\lambda]) = \sum_{\text{res}(\lambda/\mu)=i} [S_\mu], \quad F_i(n)^{\mathscr{E}}([S_\lambda]) = \sum_{\text{res}(\mu/\lambda)=i} [S_\mu].$$

(4) *We have*

$$E(n)^{\mathscr{E}} = \bigoplus_{i \in \mathbb{Z}/e\mathbb{Z}} E_i(n)^{\mathscr{E}}, \quad F(n)^{\mathscr{E}} = \bigoplus_{i \in \mathbb{Z}/e\mathbb{Z}} F_i(n)^{\mathscr{E}}.$$

Proof. – Part (1) is obvious. See [2, Theorem 13.21(2)] for (2) and [2, Lemma 13.37] for (3). Part (4) follows from (3) and [2, Lemma 13.32]. \square

4.2. – By Lemma 1.3(1) we have an algebra isomorphism

$$\gamma : Z(\mathscr{O}_{\mathbf{h},n}) \xrightarrow{\sim} Z(\mathscr{H}_{\mathbf{q},n}).$$

So there are unique elements $K_1, \dots, K_n \in Z(\mathscr{O}_{\mathbf{h},n})$ such that the polynomial

$$D_n(z) = z^n + K_1 z^{n-1} + \dots + K_n$$

maps to $C_n(z)$ by γ . Since the elements K_1, \dots, K_n act on simple modules by scalars and the category $\mathscr{O}_{\mathbf{h},n}$ is artinian, every module M in $\mathscr{O}_{\mathbf{h},n}$ is a direct sum of generalized eigenspaces of $D_n(z)$. For $a(z) \in \mathbb{C}(z)$ let $Q_{n,a(z)}$ be the exact endo-functor of $\mathscr{O}_{\mathbf{h},n}$ which maps an object M to the generalized eigenspace of $D_n(z)$ in M with the eigenvalue $a(z)$.

DEFINITION 4.2. – The *i*-restriction functor and the *i*-induction functor

$$E_i(n) : \mathscr{O}_{\mathbf{h},n} \rightarrow \mathscr{O}_{\mathbf{h},n-1}, \quad F_i(n) : \mathscr{O}_{\mathbf{h},n-1} \rightarrow \mathscr{O}_{\mathbf{h},n}$$

are given by

$$E_i(n) = \bigoplus_{a(z) \in \mathbb{C}(z)} Q_{n-1,a(z)/(z-q^i)} \circ E(n) \circ Q_{n,a(z)},$$

$$F_i(n) = \bigoplus_{a(z) \in \mathbb{C}(z)} Q_{n,a(z)(z-q^i)} \circ F(n) \circ Q_{n-1,a(z)}.$$

We will write

$$(4.4) \quad E_i = \bigoplus_{n \geq 1} E_i(n), \quad F_i = \bigoplus_{n \geq 1} F_i(n).$$

We have the following proposition.

PROPOSITION 4.3. – *For any $i \in \mathbb{Z}/e\mathbb{Z}$ there are isomorphisms of functors*

$$\text{KZ} \circ E_i(n) \cong E_i(n)^{\mathscr{E}} \circ \text{KZ}, \quad \text{KZ} \circ F_i(n) \cong F_i(n)^{\mathscr{E}} \circ \text{KZ}.$$

Proof. – Since $\gamma(D_n(z)) = C_n(z)$, by Lemma 1.3(2) for any $a(z) \in \mathbb{C}(z)$ we have

$$\text{KZ} \circ Q_{n,a(z)} \cong P_{n,a(z)} \circ \text{KZ}.$$

So the proposition follows from Theorem 2.1 and Corollary 2.3. \square

The next proposition is the DAHA version of Proposition 4.1.

PROPOSITION 4.4. – (1) *The functors $E_i(n)$, $F_i(n)$ are exact. The functor $E_i(n)$ is biadjoint to $F_i(n)$.*

(2) *For any $\lambda \in \mathcal{P}_{n,l}$ the unique eigenvalue of $D_n(z)$ on the standard module $\Delta(\lambda)$ is $a_\lambda(z)$.*

(3) *We have the following equalities*

$$(4.5) \quad E_i(n)([\Delta(\lambda)]) = \sum_{\text{res}(\lambda/\mu)=i} [\Delta(\mu)], \quad F_i(n)([\Delta(\lambda)]) = \sum_{\text{res}(\mu/\lambda)=i} [\Delta(\mu)].$$

(4) *We have*

$$E(n) = \bigoplus_{i \in \mathbb{Z}/e\mathbb{Z}} E_i(n), \quad F(n) = \bigoplus_{i \in \mathbb{Z}/e\mathbb{Z}} F_i(n).$$

Proof. – (1) This is by construction and by Proposition 2.9.

(2) Since a standard module is indecomposable, the element $D_n(z)$ has a unique eigenvalue on $\Delta(\lambda)$. By Lemma 3.1 this eigenvalue is the same as the eigenvalue of $C_n(z)$ on S_λ .

(3) Let us prove the equality for $E_i(n)$. The Pieri rule for the group $B_n(l)$ together with Proposition 1.6(2) yields

$$(4.6) \quad E(n)([\Delta(\lambda)]) = \sum_{|\lambda/\mu|=1} [\Delta(\mu)], \quad F(n)([\Delta(\lambda)]) = \sum_{|\mu/\lambda|=1} [\Delta(\mu)].$$

So we have

$$\begin{aligned} E_i(n)([\Delta(\lambda)]) &= \bigoplus_{a(z) \in \mathbb{C}[z]} Q_{n-1, a(z)/(z-q^i)}(E(n)(Q_{n, a(z)}([\Delta(\lambda)]))) \\ &= Q_{n-1, a_\lambda(z)/(z-q^i)}(E(n)(Q_{n, a_\lambda(z)}([\Delta(\lambda)]))) \\ &= Q_{n-1, a_\lambda(z)/(z-q^i)}(E(n)([\Delta(\lambda)])) \\ &= Q_{n-1, a_\lambda(z)/(z-q^i)}\left(\sum_{|\lambda/\mu|=1} [\Delta(\mu)]\right) \\ &= \sum_{\text{res}(\lambda/\mu)=i} [\Delta(\mu)]. \end{aligned}$$

The last equality follows from the fact that for any l -partition μ such that $|\lambda/\mu| = 1$ we have $a_\lambda(z) = a_\mu(z)(z - q^{\text{res}(\lambda/\mu)})$. The proof for $F_i(n)$ is similar.

(4) It follows from part (3) and (4.6). □

COROLLARY 4.5. – *Under the isomorphism θ in (3.5) the operators E_i and F_i on $K(\mathcal{O}_{\mathbf{h}, \mathbb{N}})$ go respectively to the operators e_i and f_i on $\mathcal{F}_{\mathbf{s}}$. When i runs over $\mathbb{Z}/e\mathbb{Z}$ they yield an action of $\widehat{\mathfrak{sl}}_e$ on $K(\mathcal{O}_{\mathbf{h}, \mathbb{N}})$ such that θ is an isomorphism of $\widehat{\mathfrak{sl}}_e$ -modules.*

Proof. – This is clear from Proposition 4.4(3) and from (3.3). □

5. $\widehat{\mathfrak{sl}}_e$ -categorification

In this section, we construct an $\widehat{\mathfrak{sl}}_e$ -categorification on the category $\mathcal{O}_{\mathbf{h}, \mathbb{N}}$ (Theorem 5.1).

5.1. – Recall that we put $q = \exp(\frac{2\pi\sqrt{-1}}{e})$ and P denotes the weight lattice of $\widetilde{\mathfrak{sl}}_e$. Let \mathcal{C} be a \mathbb{C} -linear artinian abelian category. For any functor $F : \mathcal{C} \rightarrow \mathcal{C}$ and any $X \in \text{End}(F)$, the generalized eigenspace of X acting on F with eigenvalue $a \in \mathbb{C}$ will be called the a -eigenspace of X in F . By [15, Definition 5.29] an $\widetilde{\mathfrak{sl}}_e$ -categorification on \mathcal{C} is the data of

- (a) an adjoint pair (U, V) of exact functors $\mathcal{C} \rightarrow \mathcal{C}$,
- (b) $X \in \text{End}(U)$ and $T \in \text{End}(U^2)$,
- (c) a decomposition $\mathcal{C} = \bigoplus_{\tau \in P} \mathcal{C}_\tau$,

such that, set U_i (resp. V_i) to be the q^i -eigenspace of X in U (resp. in V) ⁽¹⁾ for $i \in \mathbb{Z}/e\mathbb{Z}$, we have

- (1) $U = \bigoplus_{i \in \mathbb{Z}/e\mathbb{Z}} U_i$,
- (2) the endomorphisms X and T satisfy

$$(5.1) \quad \begin{aligned} (1_U T) \circ (T 1_U) \circ (1_U T) &= (T 1_U) \circ (1_U T) \circ (T 1_U), \\ (T + 1_{U^2}) \circ (T - q 1_{U^2}) &= 0, \\ T \circ (1_U X) \circ T &= q X 1_U, \end{aligned}$$

- (3) the action of $e_i = U_i$, $f_i = V_i$ on $K(\mathcal{C})$ with i running over $\mathbb{Z}/e\mathbb{Z}$ gives an integrable representation of $\widehat{\mathfrak{sl}}_e$.
- (4) $U_i(\mathcal{C}_\tau) \subset \mathcal{C}_{\tau+\alpha_i}$ and $V_i(\mathcal{C}_\tau) \subset \mathcal{C}_{\tau-\alpha_i}$,
- (5) V is isomorphic to a left adjoint of U .

5.2. – We construct an $\widetilde{\mathfrak{sl}}_e$ -categorification on $\mathcal{O}_{\mathbf{h}, \mathbb{N}}$ in the following way. The adjoint pair will be given by (E, F) . To construct the part (b) of the data we need to go back to Hecke algebras. Following [7, Section 7.2.2] let $X^{\mathcal{H}}$ be the endomorphism of $E^{\mathcal{H}}$ given on $E(n)^{\mathcal{H}}$ as the multiplication by the Jucy-Murphy element J_{n-1} . Let $T^{\mathcal{H}}$ be the endomorphism of $(E^{\mathcal{H}})^2$ given on $E(n)^{\mathcal{H}} \circ E(n-1)^{\mathcal{H}}$ as the multiplication by the element T_{n-1} in $\mathcal{H}_{\mathbf{q}, n}$. The endomorphisms $X^{\mathcal{H}}$ and $T^{\mathcal{H}}$ satisfy the relations (5.1). Moreover the q^i -eigenspace of $X^{\mathcal{H}}$ in $E^{\mathcal{H}}$ and $F^{\mathcal{H}}$ gives respectively the i -restriction functor $E_i^{\mathcal{H}}$ and the i -induction functor $F_i^{\mathcal{H}}$ for any $i \in \mathbb{Z}/e\mathbb{Z}$.

By Theorem 2.1 we have an isomorphism $\text{KZ} \circ E \cong E^{\mathcal{H}} \circ \text{KZ}$. This yields an isomorphism

$$\text{End}(\text{KZ} \circ E) \cong \text{End}(E^{\mathcal{H}} \circ \text{KZ}).$$

By Proposition 1.6(1) the functor E maps projective objects to projective ones, so Lemma 2.4 applied to $\mathcal{O}_1 = \mathcal{O}_2 = \mathcal{O}_{\mathbf{h}, \mathbb{N}}$ and $K = L = E$ yields an isomorphism

$$\text{End}(E) \cong \text{End}(\text{KZ} \circ E).$$

Composing it with the isomorphism above gives a ring isomorphism

$$(5.2) \quad \sigma_E : \text{End}(E) \xrightarrow{\sim} \text{End}(E^{\mathcal{H}} \circ \text{KZ}).$$

Replacing E by E^2 we get another isomorphism

$$\sigma_{E^2} : \text{End}(E^2) \xrightarrow{\sim} \text{End}((E^{\mathcal{H}})^2 \circ \text{KZ}).$$

⁽¹⁾ Here X acts on V via the isomorphism $\text{End}(U) \cong \text{End}(V)^{op}$ given by adjunction, see [7, Section 4.1.2] for the precise definition.

The data of $X \in \text{End}(E)$ and $T \in \text{End}(E^2)$ in our $\widetilde{\mathfrak{sl}}_e$ -categorification on $\mathcal{O}_{\mathbf{h},\mathbb{N}}$ will be provided by

$$X = \sigma_E^{-1}(X^{\mathcal{H}} 1_{\text{KZ}}), \quad T = \sigma_{E^2}^{-1}(T^{\mathcal{H}} 1_{\text{KZ}}).$$

Finally, the part (c) of the data will be given by the block decomposition of the category $\mathcal{O}_{\mathbf{h},\mathbb{N}}$. Recall from [13, Theorem 2.11] that the block decomposition of the category $\mathcal{H}_{\mathbf{q},\mathbb{N}}\text{-mod}$ is

$$\mathcal{H}_{\mathbf{q},\mathbb{N}}\text{-mod} = \bigoplus_{\tau \in P} (\mathcal{H}_{\mathbf{q},\mathbb{N}}\text{-mod})_{\tau},$$

where $(\mathcal{H}_{\mathbf{q},\mathbb{N}}\text{-mod})_{\tau}$ is the subcategory generated by the composition factors of the Specht modules S_{λ} with λ running over l -partitions of weight τ . By convention $(\mathcal{H}_{\mathbf{q},\mathbb{N}}\text{-mod})_{\tau}$ is zero if such λ does not exist. By Lemma 1.3 the functor KZ induces a bijection between the blocks of the category $\mathcal{O}_{\mathbf{h},\mathbb{N}}$ and the blocks of $\mathcal{H}_{\mathbf{q},\mathbb{N}}\text{-mod}$. So the block decomposition of $\mathcal{O}_{\mathbf{h},\mathbb{N}}$ is

$$\mathcal{O}_{\mathbf{h},\mathbb{N}} = \bigoplus_{\tau \in P} (\mathcal{O}_{\mathbf{h},\mathbb{N}})_{\tau},$$

where $(\mathcal{O}_{\mathbf{h},\mathbb{N}})_{\tau}$ is the block corresponding to $(\mathcal{H}_{\mathbf{q},\mathbb{N}}\text{-mod})_{\tau}$ via KZ .

5.3. – Now we prove the following theorem.

THEOREM 5.1. – *The data of*

- (a) *the adjoint pair (E, F) ,*
- (b) *the endomorphisms $X \in \text{End}(E)$, $T \in \text{End}(E^2)$,*
- (c) *the decomposition $\mathcal{O}_{\mathbf{h},\mathbb{N}} = \bigoplus_{\tau \in P} (\mathcal{O}_{\mathbf{h},\mathbb{N}})_{\tau}$*

is an $\widetilde{\mathfrak{sl}}_e$ -categorification on $\mathcal{O}_{\mathbf{h},\mathbb{N}}$.

Proof. – First, we prove that for any $i \in \mathbb{Z}/e\mathbb{Z}$ the q^i -generalized eigenspaces of X in E and F are respectively the i -restriction functor E_i and the i -induction functor F_i as defined in (4.4). Recall from Proposition 4.1(4) and Proposition 4.4(4) that we have

$$E = \bigoplus_{i \in \mathbb{Z}/e\mathbb{Z}} E_i \quad \text{and} \quad E^{\mathcal{H}} = \bigoplus_{i \in \mathbb{Z}/e\mathbb{Z}} E_i^{\mathcal{H}}.$$

By the proof of Proposition 4.3 we see that any isomorphism

$$\text{KZ} \circ E \cong E^{\mathcal{H}} \circ \text{KZ}$$

restricts to an isomorphism $\text{KZ} \circ E_i \cong E_i^{\mathcal{H}} \circ \text{KZ}$ for each $i \in \mathbb{Z}/e\mathbb{Z}$. So the isomorphism σ_E in (5.2) maps $\text{Hom}(E_i, E_j)$ to $\text{Hom}(E_i^{\mathcal{H}} \circ \text{KZ}, E_j^{\mathcal{H}} \circ \text{KZ})$. Write

$$X = \sum_{i,j \in \mathbb{Z}/e\mathbb{Z}} X_{ij}, \quad X^{\mathcal{H}} 1_{\text{KZ}} = \sum_{i,j \in \mathbb{Z}/e\mathbb{Z}} (X^{\mathcal{H}} 1_{\text{KZ}})_{ij}$$

with $X_{ij} \in \text{Hom}(E_i, E_j)$ and $(X^{\mathcal{H}} 1_{\text{KZ}})_{ij} \in \text{Hom}(E_i^{\mathcal{H}} \circ \text{KZ}, E_j^{\mathcal{H}} \circ \text{KZ})$. We have

$$\sigma_E(X_{ij}) = (X^{\mathcal{H}} 1_{\text{KZ}})_{ij}.$$

Since $E_i^{\mathcal{H}}$ is the q^i -eigenspace of $X^{\mathcal{H}}$ in $E^{\mathcal{H}}$, we have $(X^{\mathcal{H}} 1_{\text{KZ}})_{ij} = 0$ for $i \neq j$ and $(X^{\mathcal{H}} 1_{\text{KZ}})_{ii} - q^i$ is nilpotent for any $i \in \mathbb{Z}/e\mathbb{Z}$. Since σ_E is an isomorphism of rings, this implies that $X_{ij} = 0$ and $X_{ii} - q^i$ is nilpotent in $\text{End}(E)$. So E_i is the q^i -eigenspace of X in E . The fact that F_i is the q^i -eigenspace of X in F follows from adjunction.

Now, let us check the conditions (1)–(5):

(1) It is given by Proposition 4.4(4).

(2) Since $X^{\mathcal{H}}$ and $T^{\mathcal{H}}$ satisfy relations in (5.1), the endomorphisms X and T also satisfy them. Because these relations are preserved by ring homomorphisms.

(3) It follows from Corollary 4.5.

(4) By the definition of $(\mathcal{O}_{\mathbf{h},\mathbb{N}})_{\tau}$ and Lemma 3.1, the standard modules in $(\mathcal{O}_{\mathbf{h},\mathbb{N}})_{\tau}$ are all the $\Delta(\lambda)$ such that $\text{wt}(\lambda) = \tau$. By (3.4) if μ is an l -partition such that $\text{res}(\lambda/\mu) = i$ then $\text{wt}(\mu) = \text{wt}(\lambda) + \alpha_i$. Now, the result follows from (4.5).

(5) This is Proposition 2.9. \square

6. Crystals

Using the $\tilde{\mathfrak{sl}}_e$ -categorification in Theorem 5.1 we construct a crystal on the classes of simple objects in $\mathcal{O}_{\mathbf{h},\mathbb{N}}$ and prove that it coincides with the crystal of the Fock space $\mathcal{F}_{\mathbf{s}}$ (Theorem 6.3).

6.1. – A *crystal* (or more precisely, an $\tilde{\mathfrak{sl}}_e$ -crystal) is a set B together with maps

$$\text{wt} : B \rightarrow P, \quad \tilde{e}_i, \tilde{f}_i : B \rightarrow B \sqcup \{0\}, \quad \epsilon_i, \varphi_i : B \rightarrow \mathbb{Z} \sqcup \{-\infty\},$$

such that

- we have $\varphi_i(b) = \epsilon_i(b) + \langle \alpha_i^{\vee}, \text{wt}(b) \rangle$,
- if $\tilde{e}_i b \in B$, then $\text{wt}(\tilde{e}_i b) = \text{wt}(b) + \alpha_i$, $\epsilon_i(\tilde{e}_i b) = \epsilon_i(b) - 1$, $\varphi_i(\tilde{e}_i b) = \varphi_i(b) + 1$,
- if $\tilde{f}_i b \in B$, then $\text{wt}(\tilde{f}_i b) = \text{wt}(b) - \alpha_i$, $\epsilon_i(\tilde{f}_i b) = \epsilon_i(b) + 1$, $\varphi_i(\tilde{f}_i b) = \varphi_i(b) - 1$,
- let $b, b' \in B$, then $\tilde{f}_i b = b'$ if and only if $\tilde{e}_i b' = b$,
- if $\varphi_i(b) = -\infty$, then $\tilde{e}_i b = 0$ and $\tilde{f}_i b = 0$.

Let \mathfrak{b} be the Lie subalgebra of $\tilde{\mathfrak{sl}}_e$ generated by the elements $e_i, i \in \mathbb{Z}/e\mathbb{Z}$ and t . We say that an $\tilde{\mathfrak{sl}}_e$ -module V is \mathfrak{b} -locally finite if

- we have $V = \bigoplus_{\mu \in P} V_{\mu}$, where $V_{\mu} = \{v \in V : hv = \mu(h)v, \forall h \in t\}$,
- for any $v \in V$, the \mathfrak{b} -submodule of V generated by v is finite dimensional.

Let V be a \mathfrak{b} -locally finite $\tilde{\mathfrak{sl}}_e$ -module. For any nonzero vector $v \in V$ and any $i \in \mathbb{Z}/e\mathbb{Z}$ we set

$$l_i(v) = \max\{l \in \mathbb{N} : e_i^l(v) \neq 0\}.$$

Write $l_i(0) = -\infty$. For $l \geq 0$ let

$$V_i^{<l} = \{v \in V : l_i(v) < l\}.$$

A weight basis of V is a basis B of V such that each element of B is a weight vector. Following A. Berenstein and D. Kazhdan (cf. [3, Definition 5.30]), a *perfect basis* of V is a weight basis B together with maps $\tilde{e}_i, \tilde{f}_i : B \rightarrow B \sqcup \{0\}$ for $i \in \mathbb{Z}/e\mathbb{Z}$ such that

- for $b, b' \in B$ we have $\tilde{f}_i b = b'$ if and only if $\tilde{e}_i b' = b$,
- we have $\tilde{e}_i(b) \neq 0$ if and only if $e_i(b) \neq 0$,
- if $e_i(b) \neq 0$ then we have

$$(6.1) \quad e_i(b) \in \mathbb{C}^* \tilde{e}_i(b) + V_i^{<l_i(b)-1}.$$

We denote it by $(B, \tilde{e}_i, \tilde{f}_i)$. For such a basis let $\text{wt}(b)$ be the weight of b , let $\epsilon_i(b) = l_i(b)$ and let

$$\varphi_i(b) = \epsilon_i(b) + \langle \alpha_i^\vee, \text{wt}(b) \rangle$$

for all $b \in B$. The data

$$(6.2) \quad (B, \text{wt}, \tilde{e}_i, \tilde{f}_i, \epsilon_i, \varphi_i)$$

is a crystal. We will always attach this crystal structure to $(B, \tilde{e}_i, \tilde{f}_i)$. We call $b \in B$ a primitive element if $e_i(b) = 0$ for all $i \in \mathbb{Z}/e\mathbb{Z}$. Let B^+ be the set of primitive elements in B . Let V^+ be the vector space spanned by all the primitive vectors in V . The following lemma is [3, Claim 5.32].

LEMMA 6.1. – *For any perfect basis $(B, \tilde{e}_i, \tilde{f}_i)$ the set B^+ is a basis of V^+ .*

Proof. – By definition we have $B^+ \subset V^+$. Given a vector $v \in V^+$, there exist $\zeta_1, \dots, \zeta_r \in \mathbb{C}^*$ and distinct elements $b_1, \dots, b_r \in B$ such that $v = \sum_{j=1}^r \zeta_j b_j$. For any $i \in \mathbb{Z}/e\mathbb{Z}$ let $l_i = \max\{l_i(b_j) : 1 \leq j \leq r\}$ and $J = \{j : l_i(b_j) = l_i, 1 \leq j \leq r\}$. Then by the third property of perfect basis there exist $\eta_j \in \mathbb{C}^*$ for $j \in J$ and a vector $w \in V^{<l_i-1}$ such that $0 = e_i(v) = \sum_{j \in J} \zeta_j \eta_j \tilde{e}_i(b_j) + w$. For distinct $j, j' \in J$, we have $b_j \neq b_{j'}$, so $\tilde{e}_i(b_j)$ and $\tilde{e}_i(b_{j'})$ are different unless they are zero. Moreover, since $l_i(\tilde{e}_i(b_j)) = l_i - 1$, the equality yields that $\tilde{e}_i(b_j) = 0$ for all $j \in J$. So $l_i = 0$. Hence $b_j \in B^+$ for $j = 1, \dots, r$. \square

6.2. – Given an $\tilde{\mathfrak{sl}}_e$ -categorification on a \mathbb{C} -linear artinian abelian category \mathcal{C} with the adjoint pair of endo-functors (U, V) , $X \in \text{End}(U)$ and $T \in \text{End}(U^2)$, assume that the $\tilde{\mathfrak{sl}}_e$ -module $K(\mathcal{C})$ is \mathfrak{b} -locally finite, then one can construct a perfect basis of $K(\mathcal{C})$ as follows. For $i \in \mathbb{Z}/e\mathbb{Z}$ let U_i, V_i be the q^i -eigenspaces of X in U and V . By definition, the action of X restricts to each U_i . One can prove that T also restricts to endomorphism of $(U_i)^2$, see for example the beginning of Section 7 in [7]. It follows that the data (U_i, V_i, X, T) gives an \mathfrak{sl}_2 -categorification on \mathcal{C} in the sense of [7, Section 5.21]. By [7, Proposition 5.20] this implies that for any simple object L in \mathcal{C} , the object $\text{head}(U_i(L))$ (resp. $\text{soc}(V_i(L))$) is simple unless it is zero.

Let $B_{\mathcal{C}}$ be the set of isomorphism classes of simple objects in \mathcal{C} . As part of the data of the $\tilde{\mathfrak{sl}}_e$ -categorification, we have a decomposition $\mathcal{C} = \bigoplus_{\tau \in P} \mathcal{C}_{\tau}$. For a simple module $L \in \mathcal{C}_{\tau}$, the weight of $[L]$ in $K(\mathcal{C})$ is τ . Hence $B_{\mathcal{C}}$ is a weight basis of $K(\mathcal{C})$. Now for $i \in \mathbb{Z}/e\mathbb{Z}$ define the maps

$$\begin{aligned} \tilde{e}_i : B_{\mathcal{C}} &\rightarrow B_{\mathcal{C}} \sqcup \{0\}, & [L] &\mapsto [\text{head}(U_i L)], \\ \tilde{f}_i : B_{\mathcal{C}} &\rightarrow B_{\mathcal{C}} \sqcup \{0\}, & [L] &\mapsto [\text{soc}(V_i L)]. \end{aligned}$$

PROPOSITION 6.2. – *The data $(B_{\mathcal{C}}, \tilde{e}_i, \tilde{f}_i)$ is a perfect basis of $K(\mathcal{C})$.*

Proof. – Fix $i \in \mathbb{Z}/e\mathbb{Z}$. Let us check the conditions in order. First, for two simple modules $L, L' \in \mathcal{C}$, we have $\tilde{e}_i([L]) = [L']$, if and only if $0 \neq \text{Hom}(U_i L, L') = \text{Hom}(L, V_i L')$, if and only if $\tilde{f}_i([L']) = [L]$. The second condition follows from the fact that any non trivial module has a non trivial head. The third condition is [7, Proposition 5.20(d)]. \square

6.3. – Let $B_{\mathcal{F}_s}$ be the set of l -partitions. In [11] this set is given a crystal structure. We will call it the crystal of the Fock space \mathcal{F}_s .

THEOREM 6.3. – (1) *The set*

$$B_{\mathcal{O}_{\mathbf{h},\mathbb{N}}} = \{[L(\lambda)] \in K(\mathcal{O}_{\mathbf{h},\mathbb{N}}) : \lambda \in \mathcal{P}_{n,l}, n \in \mathbb{N}\}$$

and the maps

$$\begin{aligned} \tilde{e}_i : B_{\mathcal{O}_{\mathbf{h},\mathbb{N}}} &\rightarrow B_{\mathcal{O}_{\mathbf{h},\mathbb{N}}} \sqcup \{0\}, & [L] &\mapsto [\text{head}(E_i L)], \\ \tilde{f}_i : B_{\mathcal{O}_{\mathbf{h},\mathbb{N}}} &\rightarrow B_{\mathcal{O}_{\mathbf{h},\mathbb{N}}} \sqcup \{0\}, & [L] &\mapsto [\text{soc}(F_i L)]. \end{aligned}$$

define a crystal structure on $B_{\mathcal{O}_{\mathbf{h},\mathbb{N}}}$.

(2) *The crystal $B_{\mathcal{O}_{\mathbf{h},\mathbb{N}}}$ given by (1) is isomorphic to the crystal $B_{\mathcal{F}_s}$.*

Proof. – (1) The Fock space \mathcal{F}_s is a locally finite \mathfrak{b} -module. So applying Proposition 6.2 to the $\tilde{\mathfrak{sl}}_e$ -categorification in Theorem 5.1 yields that $(B_{\mathcal{O}_{\mathbf{h},\mathbb{N}}}, \tilde{e}_i, \tilde{f}_i)$ is a perfect basis. Therefore it defines a crystal structure on $B_{\mathcal{O}_{\mathbf{h},\mathbb{N}}}$ by (6.2).

(2) It is known that $B_{\mathcal{F}_s}$ is a perfect basis of \mathcal{F}_s . Identify the $\tilde{\mathfrak{sl}}_e$ -modules \mathcal{F}_s and $K(\mathcal{O}_{\mathbf{h},\mathbb{N}})$. By Lemma 6.1 the set $B_{\mathcal{F}_s}^+$ and $B_{\mathcal{O}_{\mathbf{h},\mathbb{N}}}^+$ are two weight bases of \mathcal{F}_s^+ . So there is a bijection $\psi : B_{\mathcal{F}_s}^+ \rightarrow B_{\mathcal{O}_{\mathbf{h},\mathbb{N}}}^+$ such that $\text{wt}(b) = \text{wt}(\psi(b))$. Since \mathcal{F}_s is a direct sum of highest weight simple $\tilde{\mathfrak{sl}}_e$ -modules, this bijection extends to an automorphism ψ of the $\tilde{\mathfrak{sl}}_e$ -module \mathcal{F}_s . By [3, Main Theorem 5.37] any automorphism of \mathcal{F}_s which maps $B_{\mathcal{F}_s}^+$ to $B_{\mathcal{O}_{\mathbf{h},\mathbb{N}}}^+$ induces an isomorphism of crystals $B_{\mathcal{F}_s} \cong B_{\mathcal{O}_{\mathbf{h},\mathbb{N}}}$. \square

REMARK 6.4. – One can prove that if $n < e$ then a simple module $L \in \mathcal{O}_{\mathbf{h},n}$ has finite dimension over \mathbb{C} if and only if the class $[L]$ is a primitive element in $B_{\mathcal{O}_{\mathbf{h},\mathbb{N}}}$. In the case $n = 1$, we have $B_n(l) = \mu_l$, the cyclic group, and the primitive elements in the crystal $B_{\mathcal{F}_s}$ have explicit combinatorial descriptions. This yields another proof of the classification of finite dimensional simple modules of $H_{\mathbf{h}}(\mu_l)$, which was first given by W. Crawley-Boevey and M. P. Holland. See type A case of [8, Theorem 7.4].

Acknowledgements

I would like to thank my advisor E. Vasserot for suggesting me this problem, for his guidance and his patience. I would also like to thank I. Gordon and M. Martino for enlightening discussions. Independently, they also obtained the result on crystals in their recent joint work. I am grateful to D. Yang and I. Losev for pointing out some errors in the preliminary version of this paper.

REFERENCES

- [1] S. ARIKI, On the decomposition numbers of the Hecke algebra of $G(m, 1, n)$, *J. Math. Kyoto Univ.* **36** (1996), 789–808.
- [2] S. ARIKI, *Representations of quantum algebras and combinatorics of Young tableaux*, University Lecture Series **26**, Amer. Math. Soc., 2002.

- [3] A. BERENSTEIN, D. KAZHDAN, Geometric and unipotent crystals. II. From unipotent bicrystals to crystal bases, in *Quantum groups*, Contemp. Math. **433**, Amer. Math. Soc., 2007, 13–88.
- [4] R. BEZRUKAVNIKOV, P. ETINGOF, Parabolic induction and restriction functors for rational Cherednik algebras, *Selecta Math. (N.S.)* **14** (2009), 397–425.
- [5] M. BROUÉ, G. MALLE, J. MICHEL, Towards spetses. I, *Transform. Groups* **4** (1999), 157–218.
- [6] M. BROUÉ, G. MALLE, R. ROUQUIER, Complex reflection groups, braid groups, Hecke algebras, *J. reine angew. Math.* **500** (1998), 127–190.
- [7] J. CHUANG, R. ROUQUIER, Derived equivalences for symmetric groups and \mathfrak{sl}_2 -categorification, *Ann. of Math.* **167** (2008), 245–298.
- [8] W. CRAWLEY-BOEVEY, M. P. HOLLAND, Noncommutative deformations of Kleinian singularities, *Duke Math. J.* **92** (1998), 605–635.
- [9] C. W. CURTIS, I. REINER, *Methods of representation theory. Vol. II, with applications to finite groups and orders*, Pure and Applied Mathematics, John Wiley & Sons Inc., 1987.
- [10] V. GINZBURG, N. GUAY, E. OPDAM, R. ROUQUIER, On the category \mathcal{O} for rational Cherednik algebras, *Invent. Math.* **154** (2003), 617–651.
- [11] M. JIMBO, K. C. MISRA, T. MIWA, M. OKADO, Combinatorics of representations of $U_q(\widehat{\mathfrak{sl}}(n))$ at $q = 0$, *Comm. Math. Phys.* **136** (1991), 543–566.
- [12] M. KASHIWARA, T. KAWAI, On holonomic systems of microdifferential equations. III. Systems with regular singularities, *Publ. Res. Inst. Math. Sci.* **17** (1981), 813–979.
- [13] S. LYLE, A. MATHAS, Blocks of cyclotomic Hecke algebras, *Adv. Math.* **216** (2007), 854–878.
- [14] R. ROUQUIER, q -Schur algebras and complex reflection groups, *Mosc. Math. J.* **8** (2008), 119–158.
- [15] R. ROUQUIER, 2-Kac-Moody algebras, preprint [arXiv:0812.5023](https://arxiv.org/abs/0812.5023).
- [16] D. UGLOV, Canonical bases of higher-level q -deformed Fock spaces and Kazhdan-Lusztig polynomials, in *Physical combinatorics (Kyoto, 1999)*, Progr. Math. **191**, Birkhäuser, 2000, 249–299.
- [17] X. YVONNE, A conjecture for q -decomposition matrices of cyclotomic v -Schur algebras, *J. Algebra* **304** (2006), 419–456.

(Manuscrit reçu le 20 mai 2009 ;
 accepté, après révision, le 16 février 2010.)

Peng SHAN
 Université Paris 7
 Institut de Mathématiques de Jussieu
 Théorie des groupes et des représentations
 Case 7012
 2 place Jussieu
 75251 Paris Cedex 05, France
 E-mail: shan@math.jussieu.fr