

ALTERNATING ARM EXPONENTS FOR THE CRITICAL PLANAR ISING MODEL

BY HAO WU*

Yau Mathematical Sciences Center, Tsinghua University, China

We derive the alternating arm exponents of the critical Ising model. We obtain six different patterns of alternating boundary arm exponents which correspond to the boundary conditions $(\ominus\oplus)$, $(\ominus\text{free})$ and (free free) , and the alternating interior arm exponents.

CONTENTS

1	Introduction	1
2	Preliminaries on SLE	5
	2.1 \mathbb{H} -hull and Loewner chain	5
	2.2 SLE processes	8
3	SLE Boundary Arm Exponents	11
	3.1 Definitions and Statements	11
	3.2 Estimates on the Derivatives	14
	3.3 Proof of Propositions 3.1 to 3.3	20
4	SLE Interior Arm Exponents	26
5	Ising Model	32
	5.1 Definitions	32
	5.2 Quasi-Multiplicativity	34
	5.3 Proof of Theorems 1.1 and 1.2	37
6	Appendix: One-Point Estimate of the Intersection of SLE with the Boundary	44
	References	48

1. Introduction. The Lenz-Ising model is one of the simplest models in statistical physics. It is a model on the spin configurations. Each vertex x has a spin σ_x which is \oplus or \ominus . Each configuration of spins $\sigma = (\sigma_x, x \in V)$ has an intrinsic energy—the Hamiltonian: $H(\sigma) = -\sum_{x \sim y} \sigma_x \sigma_y$. A natural way to sample the random configuration is the Boltzman measure:

$$\mu[\sigma] \propto \exp(-\beta H(\sigma)),$$

*This work is supported by NCCR/SwissMAP, ERC AG COMPASP, the Swiss NSF.

MSC 2010 subject classifications: Primary 60J67; secondary 60K35

Keywords and phrases: Schramm Loewner Evolution, Critical Planar Ising Model, Arm Exponent

where $\beta > 0$ is the inverse-temperature. This measure favors configurations with low energy. Due to recent celebrated work of Chelkak and Smirnov [CS12, CDCH⁺14], it is proved that at the critical temperature, the interface of the Ising model is conformally invariant and converges to a random curve—Schramm Loewner Evolution (SLE₃). In this paper, we derive the alternating arm exponents of the critical Ising model.

An arm is a simple path of \oplus or of \ominus . We are interested in the decay of the probability that there are a certain number of arms of certain pattern in the semi-annulus $A^+(n, N)$ or annulus $A(n, N)$ connecting the inner boundary to the outer boundary. This probability should decay like a power in N as $N \rightarrow \infty$, and the exponent in the power is called the critical arm exponents.

In [LSW01, LSW02, SW01], the authors derived the value of the arm exponents for the critical percolation. As explained in [SW01], to derive the arm exponents, one needs three inputs: (1) the convergence of the interface to SLE; (2) the arm exponents of SLE; and (3) the quasi-multiplicativity. This strategy also works for the critical Ising model. In this paper, we derive the boundary arm exponents and the interior arm exponents of SLE _{κ} and its variant SLE _{κ} (ρ), and then explain how to apply these formulae to get the alternating arm exponents of the critical Ising model.

THEOREM 1.1. *For the critical planar Ising model on the square lattice, we have the following six different patterns of the boundary arm exponents (the arm patterns are in clockwise order). Fix $j \geq 1$. We write b.c. for “boundary conditions”.*

- Consider the b.c. ($\oplus\oplus$) and the pattern ($\ominus \oplus \ominus \cdots \oplus \ominus$) with length $2j - 1$. The corresponding boundary arm exponents are given by

$$(1.1) \quad \alpha_{2j-1}^+ = j(4j + 1)/3.$$

- Consider the b.c. ($\ominus\oplus$) and the pattern ($\oplus \ominus \cdots \oplus \ominus$) with length $2j$. The corresponding boundary arm exponents are given by

$$(1.2) \quad \alpha_{2j}^+ = j(4j + 5)/3.$$

- Consider the b.c. (\ominus free) and the pattern ($\oplus \ominus \oplus \cdots \ominus \oplus$) with length $2j - 1$. The corresponding boundary arm exponents are given by

$$(1.3) \quad \beta_{2j-1}^+ = 2j(2j - 1)/3.$$

- Consider the b.c. (\ominus free) and the pattern ($\oplus \ominus \oplus \cdots \oplus \ominus$) with length $2j$. The corresponding boundary arm exponents are given by

$$(1.4) \quad \beta_{2j}^+ = 2j(2j + 1)/3.$$

- Consider the b.c. (free free) and the pattern $(\ominus \oplus \ominus \cdots \oplus \ominus)$ with length $2j - 1$. The corresponding boundary arm exponents are given by

$$(1.5) \quad \gamma_{2j-1}^+ = (2j - 1)(4j - 3)/6.$$

- Consider the b.c. (free free) and the pattern $(\ominus \oplus \cdots \ominus \oplus)$ with length $2j$. The corresponding boundary arm exponents are given by

$$(1.6) \quad \gamma_{2j}^+ = j(4j - 1)/3.$$




Fig 1.1: The six different patterns of boundary arm exponents in Theorem 1.1.

THEOREM 1.2. *For the critical planar Ising model on the square lattice, the alternating interior arm exponents with length $2j$ for $j \geq 1$ are given by*

$$(1.7) \quad \alpha_{2j} = (16j^2 - 1)/24.$$

REMARK 1.3. *In Theorem 1.1, the arm exponent $\gamma_2^+ = 1$ is a universal arm exponent of the critical Ising model. In other words, the fact that $\gamma_2^+ = 1$ can be obtained by standard proof of universal arm exponents using RSW.*

REMARK 1.4. *For the critical planar Ising model (on the square lattice) in a topological rectangle $(\Omega; a, b, c, d)$ with free boundary conditions, consider the probability that there exists a path of \oplus connecting the boundary arc (ab)*

to the boundary arc (cd) . It is proved in [BDCH16] that, as the mesh-size goes to zero, this probability converges to a function f which maps topological rectangles to $[0, 1]$ and it is conformally invariant. Therefore, the limit of this probability only depends on the extremal distance of the rectangle. Whereas, the exact formula for f is unknown. As a consequence of Theorem 1.1, we can give the asymptotics of this function f . Consider the rectangle $[0, \pi L] \times [0, 1]$ and let $f(L)$ be the limit of the probability that the Ising model with free boundary conditions has a \oplus horizontal crossing of the rectangle. Then

$$f(L) = \exp(-L(1/6 + o(1))).$$

Relation to previous works. In this paper, we derive the arm exponents for $\text{SLE}_\kappa(\rho)$ with $\kappa \in (0, 4)$ and $\rho \in (-2, 0]$. In [Wu17+], we derive the arm exponents of SLE_κ for $\kappa \in (4, 8)$. The boundary 1-arm exponent γ_1^+ is related to the Hausdorff dimension of the intersection of $\text{SLE}_\kappa(\rho)$ with the boundary which is $1 - \gamma_1^+$. This dimension was obtained in [WW13, MW17]. The formulae (1.1) and (1.2) are also obtained in [WZ16]. The formulae (1.1) and (1.7) were predicted by KPZ in [Dup03, Eq. (11.42), Eq. (11.43)] and our work justifies those predictions.

The techniques developed in this paper are more complicated than those in [SW01, MW17, WZ16, Wu17+]. One difficulty is that, when we estimate the arm events of $\text{SLE}_\kappa(\rho)$ with $\kappa \in (0, 4)$, we have two more variables to take care of than the cases in [SW01, WZ16, Wu17+], see the informal discussion at the end of Section 3.1. Another difficulty is that, when we derive the arm exponents by iteration, we do not allow error terms in the exponents; whereas, when one derives the one-point estimate for the intersection probability as in many other papers calculating the Hausdorff dimension of SLE curves, one is allowed to have error terms in the exponents (e.g. the conclusion as in Theorem 6.1 is sufficient to derive the upper bound for the Hausdorff dimension). We treat these two difficulties in Section 3. There we obtain up-to-constant one-point estimates which guarantee the iteration.

Outline. We give preliminaries on SLE in Section 2. We derive the boundary arm exponents of $\text{SLE}_\kappa(\rho)$ with $\kappa \in (0, 4)$ and $\rho \in (-2, 0]$ in Section 3. We derive the interior arm exponents of SLE_κ with $\kappa \in (0, 4)$ in Section 4. Finally, we explain how to apply these formulae to obtain the alternating arm exponents of the critical Ising in Section 5 and complete the proof of Theorems 1.1 and 1.2.

Acknowledgment. The author acknowledges Hugo Duminil-Copin, Matan Harel, Aran Raoufi, Stanislav Smirnov, and Vincent Tassion for helpful discussion on Ising and FK-Ising model. The author thanks Gregory Lawler,

David Wilson and Dapeng Zhan for helpful discussion on SLE estimates. The author appreciates the advice from two anonymous referees.

2. Preliminaries on SLE.

Notations. We denote by $f \lesssim g$ if f/g is bounded from above by universal finite constant, by $f \gtrsim g$ if f/g is bounded from below by universal positive constant, and by $f \asymp g$ if $f \lesssim g$ and $f \gtrsim g$. We denote by

$$f(\epsilon) = g(\epsilon)^{1+o(1)} \quad \text{if} \quad \lim_{\epsilon \rightarrow 0} \frac{\log f(\epsilon)}{\log g(\epsilon)} = 1.$$

For $z \in \mathbb{C}, r > 0$, we denote $B(z, r) = \{w \in \mathbb{C} : |w - z| < r\}$. We denote the unit disc $B(0, 1)$ by \mathbb{U} .

For two subsets $A, B \subset \mathbb{C}$, we denote $\text{dist}(A, B) = \inf\{|x - y| : x \in A, y \in B\}$. We assume that $\text{dist}(A, \emptyset) = \infty$.

Let Ω be an open set and let V_1, V_2 be two sets such that $V_1 \cap \overline{\Omega} \neq \emptyset$ and $V_2 \cap \overline{\Omega} \neq \emptyset$. We denote the extremal distance between V_1 and V_2 in Ω by $d_\Omega(V_1, V_2)$, see [Ahl10, Section 4] for the definition.

2.1. \mathbb{H} -hull and Loewner chain. We call a compact subset K of $\overline{\mathbb{H}}$ an \mathbb{H} -hull if $\mathbb{H} \setminus K$ is simply connected. Riemann's Mapping Theorem and Schwarz Reflection Principle assert that (see e.g. [Law05, Proposition 3.34]) there exists a unique conformal map g_K from $\mathbb{H} \setminus K$ onto \mathbb{H} such that $\lim_{|z| \rightarrow \infty} |g_K(z) - z| = 0$. We call such g_K the conformal map from $\mathbb{H} \setminus K$ onto \mathbb{H} normalized at ∞ .

The following Lemmas 2.1 to 2.3 are technical and they study the image of balls under conformal maps. They are crucial in the iteration when we derive the arm exponents in Section 3.

LEMMA 2.1. *Let K be an \mathbb{H} -hull and let g_K be the conformal map from $\mathbb{H} \setminus K$ onto \mathbb{H} normalized at ∞ . Fix $x > 0$ and $\epsilon > 0$ and assume that $x > \max(K \cap \mathbb{R})$. Denote by γ the connected component of $\mathbb{H} \cap (\partial B(x, \epsilon) \setminus K)$ whose closure contains $x + \epsilon$. Then $g_K(\gamma)$ is contained in the ball with center $g_K(x + \epsilon)$ and radius $3(g_K(x + 3\epsilon) - g_K(x + \epsilon))$, hence it is also contained in the ball with center $g_K(x + 3\epsilon)$ and radius $8\epsilon g'_K(x + 3\epsilon)$.*

PROOF. Define $r^* = \sup\{|z - g_K(x + \epsilon)| : z \in g_K(\gamma)\}$. It suffices to show

$$(2.1) \quad r^* \leq 3(g_K(x + 3\epsilon) - g_K(x + \epsilon)).$$

We will prove (2.1) by estimating the extremal distance: $d_{\mathbb{H}}(g_K(\gamma), [g_K(x + 3\epsilon), \infty))$. By the conformal invariance and the comparison principle [Ahl10, Section 4.3], we can obtain the lower bound:

$$\begin{aligned} d_{\mathbb{H}}(g_K(\gamma), [g_K(x + 3\epsilon), \infty)) &= d_{\mathbb{H} \setminus K}(\gamma, [x + 3\epsilon, \infty)) \\ &\geq d_{\mathbb{H} \setminus B(x, \epsilon)}(B(x, \epsilon), [x + 3\epsilon, \infty)) \\ &= d_{\mathbb{H} \setminus \mathbb{U}}(\mathbb{U}, [3, \infty)) = d_{\mathbb{H}}([-1, 0], [1/3, \infty)). \end{aligned}$$

In the last equality, we use the conformal map $\varphi(z) := (z + 1/z)/4 - 1/2$ which sends $\mathbb{H} \setminus \mathbb{U}$ onto \mathbb{H} . Under this conformal map, we see that \mathbb{U} is mapped to the interval $[-1, 0]$ and $[3, \infty)$ is mapped to the interval $[1/3, \infty)$.

On the other hand, we will give an upper bound. Recall a fact for extremal distance: for $x < y$ and $r > 0$, the extremal distance in \mathbb{H} between $[y, \infty)$ and a connected set $S \subset \overline{\mathbb{H}}$ with $x \in \overline{S} \subset \overline{B(x, r)}$ is maximized when $S = [x - r, x]$, see [Ahl06, Chapter I-E, Chapter III-A]. Since $g_K(\gamma)$ is connected and $g_K(x + \epsilon) \in \mathbb{R} \cap g_K(\gamma)$, by the above fact, we have the upper bound:

$$\begin{aligned} d_{\mathbb{H}}(g_K(\gamma), [g_K(x + 3\epsilon), \infty)) &\leq d_{\mathbb{H}}([g_K(x + \epsilon) - r^*, g_K(x + \epsilon)], [g_K(x + 3\epsilon), \infty)) \\ &= d_{\mathbb{H}}([-r^*, 0], [g_K(x + 3\epsilon) - g_K(x + \epsilon), \infty)). \end{aligned}$$

Combining the lower bound with the upper bound, we have

$$d_{\mathbb{H}}([-1, 0], [1/3, \infty)) \leq d_{\mathbb{H}}([-r^*, 0], [g_K(x + 3\epsilon) - g_K(x + \epsilon), \infty)).$$

This implies (2.1) and completes the proof. \square

The following lemma is a standard estimate using the Koebe 1/4 theorem.

LEMMA 2.2. *Fix $z \in \overline{\mathbb{H}}$ and $\epsilon > 0$. Let K be an \mathbb{H} -hull and let g_K be the conformal map from $\mathbb{H} \setminus K$ onto \mathbb{H} normalized at ∞ . Assume that $\text{dist}(K, z) \geq 16\epsilon$. Then $g_K(B(z, \epsilon))$ is contained in the ball with center $g_K(z)$ and radius $4\epsilon|g'_K(z)|$.*

Loewner chain is a collection of \mathbb{H} -hulls $(K_t, t \geq 0)$ associated with the family of conformal maps $(g_t, t \geq 0)$ obtained by solving the Loewner equation: for each $z \in \mathbb{H}$,

$$(2.2) \quad \partial_t g_t(z) = \frac{2}{g_t(z) - W_t}, \quad g_0(z) = z,$$

where $(W_t, t \geq 0)$ is a one-dimensional real-valued continuous function which we call the driving function. Let T_z be the swallowing time of z defined as $\sup\{t \geq 0 : \min_{s \in [0, t]} |g_s(z) - W_s| > 0\}$. Let $K_t := \overline{\{z \in \mathbb{H} : T_z \leq t\}}$. Then g_t is the unique conformal map from $H_t := \mathbb{H} \setminus K_t$ onto \mathbb{H} normalized at ∞ .

LEMMA 2.3. *Suppose that $(K_t, t \geq 0)$ is a Loewner chain which is generated by a continuous curve $(\eta(t), t \geq 0)$. Fix $y \leq -4r < 0 < x$. Let σ be the first time that η hits $B(y, r)$ and assume that x is not swallowed by $\eta[0, \sigma]$ and that $y - r$ is not swallowed by $\eta[0, \sigma]$. Then we have*

$$g_\sigma(x) - W_\sigma \geq (x - y - 2r)/2.$$

PROOF. Let γ be the right side of $\eta[0, \sigma]$. We prove the conclusion by estimating the extremal distance $d_{\mathbb{H} \setminus \eta[0, \sigma]}((-\infty, y - r), \gamma \cup [0, x])$. Denote $g_\sigma - W_\sigma$ by f . On the one hand, by the conformal invariance of the extremal distance, we have

$$\begin{aligned} d_{\mathbb{H} \setminus \eta[0, \sigma]}((-\infty, y - r), \gamma \cup [0, x]) &= d_{\mathbb{H}}((-\infty, f(y - r)), (0, f(x))) \\ &= d_{\mathbb{H}}\left((-\infty, 0), \left(1, \frac{f(x) - f(y - r)}{-f(y - r)}\right)\right). \end{aligned}$$

On the other hand, we give the following upper bound. Since any rectifiable arc in $\mathbb{H} \setminus B(y, r)$ connecting $(-\infty, y - r)$ and $(y + r, x)$ contains a rectifiable arc in $\mathbb{H} \setminus \eta[0, \sigma]$ connecting $(-\infty, y - r)$ and $\gamma \cup [0, x]$ (recall that σ is the first time that η hits $B(y, r)$). By the comparison principle of the extremal distance [Ahl10, Section 4.3], we have

$$\begin{aligned} d_{\mathbb{H} \setminus \eta[0, \sigma]}((-\infty, y - r), \gamma \cup [0, x]) &\leq d_{\mathbb{H} \setminus B(y, r)}((-\infty, y - r), (y + r, x)) \\ &= d_{\mathbb{H}}\left((-\infty, 0), \left(1, \frac{1}{2} + \frac{x - y}{4r} + \frac{r}{4(x - y)}\right)\right). \end{aligned}$$

Comparing these two parts, we have

$$\frac{f(x) - f(y - r)}{-f(y - r)} \geq \frac{1}{2} + \frac{x - y}{4r} + \frac{r}{4(x - y)} \geq \frac{1}{2} + \frac{x - y}{4r}.$$

Thus

$$\frac{g_\sigma(x) - W_\sigma}{W_\sigma - g_\sigma(y - r)} \geq \frac{x - y}{4r} - \frac{1}{2}.$$

Since the quantity $g_t(x) - g_t(y - r)$ is increasing in t , we have

$$g_\sigma(x) - g_\sigma(y - r) \geq x - y + r.$$

We denote $g_\sigma(x) - W_\sigma$ by A and $W_\sigma - g_\sigma(y - r)$ by B . Then the above two estimates can be written as

$$A \geq \frac{x - y - 2r}{4r} B; \quad A + B \geq x - y + r.$$

As a consequence, we obtain

$$A \geq \frac{x - y - 2r}{x - y + 2r} (A + B) \geq \frac{x - y - 2r}{x - y + 2r} (x - y + r) \geq (x - y - 2r)/2.$$

This completes the proof. \square

2.2. *SLE processes.* In this section, we introduce standard SLE_κ process, the process with one extra marked point $\text{SLE}_\kappa(\rho)$, and the process with multiple marked points $\text{SLE}_\kappa(\underline{\rho})$. The main statements in Sections 3 and 4 only concern $\text{SLE}_\kappa(\rho)$ process; but, to derive those conclusions, one needs estimates for $\text{SLE}_\kappa(\underline{\rho})$ process.

SLE_κ and $\text{SLE}_\kappa(\rho)$ processes. An SLE_κ is the random Loewner chain $(K_t, t \geq 0)$ driven by $W_t = \sqrt{\kappa}B_t$ where $(B_t, t \geq 0)$ is a standard one-dimensional Brownian motion. In [RS05], the authors prove that $(K_t, t \geq 0)$ is almost surely generated by a continuous transient curve, i.e. there almost surely exists a continuous curve η such that for each $t \geq 0$, H_t is the unbounded connected component of $\mathbb{H} \setminus \eta[0, t]$ and that $\lim_{t \rightarrow \infty} |\eta(t)| = \infty$.

For $\kappa > 0$ and $\rho \in \mathbb{R}$, an $\text{SLE}_\kappa(\rho)$ process is a Loewner chain with one marked point $x \geq 0$. It is the Loewner chain driven by W_t which is the solution to the system of SDEs:

$$dW_t = \sqrt{\kappa}dB_t + \frac{\rho dt}{W_t - V_t}, \quad W_0 = 0; \quad dV_t = \frac{2dt}{V_t - W_t}, \quad V_0 = x.$$

When $\rho > -2$, the process is well-defined for all time and it is almost surely generated by a continuous transient curve. When $\rho \leq -2$, the process is well-defined up to T_x —the swallowing time of x . Moreover, the process is almost surely generated by a continuous curve up to and including T_x .

We summarize the behaviors of SLE for different ρ 's in the following, see [Dub09, Lemma 15]. Fix $\kappa > 0, \rho \in \mathbb{R}$ and $x > 0$. Suppose that η is an $\text{SLE}_\kappa(\rho)$ process with force point x . The following facts hold almost surely.

- When $\rho \geq \kappa/2 - 2$, the curve η never hits the interval (x, ∞) .
- When $\kappa/2 - 2 > \rho > \kappa/2 - 4$, the curve η accumulates at a point in (x, ∞) at finite time.
- When $\rho \leq \kappa/2 - 4$, the curve η converges to the point x at finite time.

$\text{SLE}_\kappa(\underline{\rho})$ processes. Next, we define an $\text{SLE}_\kappa(\underline{\rho}^L; \underline{\rho}^R)$ process with multiple force points $(\underline{x}^L; \underline{x}^R)$ where

$$\underline{\rho}^L = (\rho^{1,L}, \dots, \rho^{1,L}), \quad \underline{\rho}^R = (\rho^{1,R}, \dots, \rho^{r,R}) \quad \text{with } \rho^{i,q} \in \mathbb{R};$$

$$\underline{x}^L = (x^{1,L} < \dots < x^{1,L} \leq 0), \quad \underline{x}^R = (0 \leq x^{1,R} < \dots < x^{r,R}).$$

It is the Loewner chain driven by W_t which is the solution to the following systems of SDEs:

$$dW_t = \sqrt{\kappa}dB_t + \sum_i \frac{\rho^{i,L} dt}{W_t - V_t^{i,L}} + \sum_i \frac{\rho^{i,R} dt}{W_t - V_t^{i,R}}, \quad W_0 = 0;$$

$$dV_t^{i,L} = \frac{2dt}{V_t^{i,L} - W_t}, \quad V_0^{i,L} = x^{i,L}; \quad dV_t^{i,R} = \frac{2dt}{V_t^{i,R} - W_t}, \quad V_0^{i,R} = x^{i,R}.$$

The solution exists and is unique up to the continuation threshold is hit—the first time t that $W_t = V_t^{j,q}$ where $\sum_1^j \rho^{i,q} \leq -2$ for some $q \in \{L, R\}$. Moreover, the corresponding Loewner chain is almost surely generated by a continuous curve, see [MS16, Section 2 and Theorem 1.3].

In fact, in this paper, we only need the definitions and properties of SLE with three force points: $\text{SLE}_\kappa(\rho^L; \rho^{1,R}, \rho^{2,R})$ with force points $(x^L; x^{1,R}, x^{2,R})$. To simplify notation, we will focus on these SLE processes in this section. From the Girsanov Theorem, it follows that the law of an $\text{SLE}_\kappa(\rho)$ process can be constructed by reweighting the law of an ordinary SLE_κ up to the first time that the Loewner chain swallows any force point [SW05, Theorem 6].

LEMMA 2.4. *Suppose $x^L < 0 < x^{1,R} < x^{2,R}$ and $\rho^L, \rho^{1,R}, \rho^{2,R} \in \mathbb{R}$. The following process is a local martingale for SLE_κ .*

$$\begin{aligned} M_t &= g_t'(x^L)^{\rho^L(\rho^L+4-\kappa)/(4\kappa)} (g_t(x^L) - W_t)^{\rho^L/\kappa} \\ &\quad \times g_t'(x^{1,R})^{\rho^{1,R}(\rho^{1,R}+4-\kappa)/(4\kappa)} (g_t(x^{1,R}) - W_t)^{\rho^{1,R}/\kappa} \\ &\quad \times g_t'(x^{2,R})^{\rho^{2,R}(\rho^{2,R}+4-\kappa)/(4\kappa)} (g_t(x^{2,R}) - W_t)^{\rho^{2,R}/\kappa} \\ &\quad \times (g_t(x^{1,R}) - g_t(x^L))^{\rho^L \rho^{1,R}/(2\kappa)} (g_t(x^{2,R}) - g_t(x^L))^{\rho^L \rho^{2,R}/(2\kappa)} \\ &\quad \times (g_t(x^{2,R}) - g_t(x^{1,R}))^{\rho^{1,R} \rho^{2,R}/(2\kappa)}. \end{aligned}$$

The law of SLE_κ weighted by M (up to the first time that W hits one of the force points) is equal to the law of $\text{SLE}_\kappa(\rho^L; \rho^{1,R}, \rho^{2,R})$ with force points $(x^L; x^{1,R}, x^{2,R})$.

The following two lemmas are technical estimates for SLE process with two marked points on the right. These two lemmas estimate the probability for SLE curves to have nice behavior and give lower bound for the probability uniform over the location of force points. We will use them in Section 3.

LEMMA 2.5. *Fix $\kappa \in (0, 4)$, $\rho > -2$, $\nu \in \mathbb{R}$ that $\rho + \nu < \kappa/2 - 4$. Suppose that η is an $\text{SLE}_\kappa(\rho, \nu)$ process with force points (v, x) where $0 \leq v < x$. For $\epsilon > 0$, let τ be the first time that η hits $B(x, \epsilon)$. For $C \geq 4$, $1/4 \geq c > 0$, define*

$$\mathcal{F} = \{\eta[0, \tau] \subset B(0, Cx), \text{dist}(\eta[0, \tau], [x - \epsilon, Cx]) \geq c\epsilon\}.$$

Then, there exist constants $c, C, u_0 > 0$ which are uniform over v, x, ϵ such that $\mathbb{P}[\mathcal{F}] \geq u_0$.

PROOF. By the scale invariance of SLE, we may assume $x = 1$. Let $\varphi(z) = \epsilon z / (1 - z)$. Then φ is the Möbius transformation of \mathbb{H} that sends the triplet $(0, 1, \infty)$ to $(0, \infty, -\epsilon)$. Let us check the images of η , $B(0, C)$, and the $c\epsilon$ -neighborhood of $[x - \epsilon, C\epsilon]$ under φ respectively.

Denote the image of η under φ by $\tilde{\eta}$, and denote its law by $\tilde{\mathbb{P}}$. Note that $\tilde{\eta}$ is an $\text{SLE}_\kappa(\rho^L; \rho^R)$ with force points $(-\epsilon; \epsilon v / (1 - v))$ where

$$\rho^L = \kappa - 6 - \rho - \nu > \kappa/2 - 2, \quad \rho^R = \rho > -2.$$

For $r \in (0, 1/4)$ and $y \in (-1, 0)$, let $\tilde{T} = \inf\{t : \tilde{\eta}(t) \in \partial B(y, r|y|)\}$ and $\tilde{S} = \inf\{t : \tilde{\eta}(t) \in \partial B(0, 1)\}$. Since $\rho^L > \kappa/2 - 2$, by [MW17, Corollary 3.3] or Lemma 6.3, there exists $A > 1$ depending only on κ, ρ^L, ρ^R such that,

$$(2.3) \quad \tilde{\mathbb{P}} \left[\tilde{T} < \tilde{S}, \text{Im } \tilde{\eta}(\tilde{T}) \geq r|y|/4 \right] \leq r^A.$$

Consider the image of $\mathbb{H} \setminus B(0, C)$ under φ . It is contained in the ball $B(-\epsilon, 2\epsilon/C)$. By Lemma 6.5, there exists a function $q(C)$, which depends on C and is uniform over ϵ , such that the probability for $\tilde{\eta}$ to hit $B(-\epsilon, 2\epsilon/C)$ is bounded by $q(C)$ and $q(C) \rightarrow 0$ as $C \rightarrow \infty$.

Consider the image of $c\epsilon$ -neighborhood of $[1 + \epsilon, C]$ under φ . Since $c\epsilon$ -neighborhood of $[1 + \epsilon, C]$ is contained in the union of the balls $B(1 + k\epsilon/4, 4c\epsilon)$ for $4/c \leq k \leq C/\epsilon$, its image under φ is contained in the union of the following balls

$$B(-4/(ck) - \epsilon, 256/(ck^2)), \quad [4/c] \leq k \leq \lceil C/\epsilon \rceil.$$

Define $\tilde{\mathcal{F}}$ to be the event that $\tilde{\eta}$ exits the unit disc without touching the union of $B(-\epsilon, 2\epsilon/C)$ and the image of $c\epsilon$ -neighborhood of $[1 + \epsilon, C]$ under φ . Then, by (2.3), we have

$$1 - \mathbb{P}[\mathcal{F}] \leq 1 - \tilde{\mathbb{P}}[\tilde{\mathcal{F}}] \lesssim q(C) + \sum_{k=4/c}^{C/\epsilon} \left(\frac{1}{4k + \epsilon ck^2} \right)^A \lesssim q(C) + c^{A-1}.$$

This implies the conclusion. In this proof, it is important that $A > 1$ in (2.3). This explains the requirement $\rho + \nu < \kappa/2 - 4$. \square

LEMMA 2.6. *Fix $\kappa \in (0, 4), \rho > -2, \nu \in \mathbb{R}$ that $\rho + \nu > -2$. Suppose that η is an $\text{SLE}_\kappa(\rho, \nu)$ process with force points (v, x) where $0 \leq v \leq x$. For $r > 0 > y$ and $M > 1$, assume $r < |y| \leq Mr$. Let σ be the first time that η hits $B(y, r)$. For $C \geq 4, 1/4 \geq c > 0$, define*

$$\mathcal{F} = \{\sigma < \infty, \text{dist}(\eta[0, \sigma], x) \geq cx, \eta[0, \sigma] \subset B(0, C|y|), \text{dist}(\eta[0, \sigma], [Cy, y]) \geq cr\}.$$

Then there exist constants $c, C, u_0 > 0$ which may depend on M but are uniform over v, x, y, r such that $\mathbb{P}[\mathcal{F}] \geq u_0$.

PROOF. From Lemma 6.5, there exists a function $p(\delta)$ which is uniform over v, x such that $p(\delta) \downarrow 0$ as $\delta \downarrow 0$ and that

$$(2.4) \quad \mathbb{P}[\text{dist}(\eta, x) \geq \delta x] \geq 1 - p(\delta).$$

By scale invariance, we may assume $y = -1$ and $r \in [1/M, 1)$. Next, we estimate the probability for the following event:

$$\mathcal{G} = \{\sigma < \infty, \eta[0, \sigma] \subset B(0, 4), \text{dist}(\eta[0, \sigma], [-4, -1]) \geq r/4\}.$$

Denote by $f(v, x, r) := \mathbb{P}[\mathcal{G}]$. By a similar argument as in the proof of Lemma 6.5, we see that f is continuous and $q(M) := \inf f(v, x, r) > 0$ where the infimum is over $0 \leq v \leq x$ and $r \in [1/M, 1)$. Thus, $\mathbb{P}[\mathcal{G}] \geq q(M)$. Combining with (2.4), we have $\mathbb{P}[\mathcal{F}] \geq q(M) - p(c)$. This implies the conclusion. \square

3. SLE Boundary Arm Exponents.

3.1. *Definitions and Statements.* Fix $\kappa \in (0, 4)$ and $\rho > -2$. Let η be an $\text{SLE}_\kappa(\rho)$ with force point $v \geq 0$. Assume $y \leq -4r < 0 < \epsilon \leq v \leq x$ and we consider the crossings of η between $B(x, \epsilon)$ and $B(y, r)$. Let T_x be the first time that η swallows x . We write c.c. for ‘‘connected component’’. We have four different types of crossing events.

Set $\tau_0 = \sigma_0 = 0$. Let τ_1 be the first time that η hits $B(x, \epsilon)$ and let σ_1 be the first time after τ_1 that η hits the c.c. of $\partial B(y, r) \setminus \eta[0, \tau_1]$ containing $y - r$. For $j \geq 1$, let τ_j be the first time after σ_{j-1} that η hits the c.c. of $\partial B(x, \epsilon) \setminus \eta[0, \sigma_{j-1}]$ containing $x + \epsilon$, and let σ_j be the first time after τ_j that η hits the c.c. of $\partial B(y, r) \setminus \eta[0, \tau_j]$ containing $y - r$. Define

$$(3.1) \quad \mathcal{H}_{2j-1}^\alpha(\epsilon, x, y, r; v) = \{\tau_j < T_x\}, \quad \mathcal{H}_{2j}^\beta(\epsilon, x, y, r; v) = \{\sigma_j < T_x\}.$$

In the definition of $\mathcal{H}_{2j-1}^\alpha$ and \mathcal{H}_{2j}^β , we are interested in the case when x, y, r are fixed and $\epsilon > 0$ small. Imagine η is the interface of the lattice model, then $\mathcal{H}_{2j-1}^\alpha$ means that there are $2j - 1$ arms connecting $B(x, \epsilon)$ to far away place; and \mathcal{H}_{2j}^β means that there are $2j$ arms connecting $B(x, \epsilon)$ to far away place.

Next, we define the other two types of crossing events. We emphasize that we will change the definition of the stopping times in the following. Set $\tau_0 = \sigma_0 = 0$. Let σ_1 be the first time that η hits $B(y, r)$ and τ_1 be the first time after σ_1 that η hits the c.c. of $\partial B(x, \epsilon) \setminus \eta[0, \sigma_1]$ containing $x + \epsilon$. For $j \geq 1$, let σ_j be the first time after τ_{j-1} that η hits the c.c. of

$\partial B(y, r) \setminus \eta[0, \tau_{j-1}]$ containing $y - r$ and let τ_j be the first time after σ_j that η hits the c.c. of $\partial B(x, \epsilon) \setminus \eta[0, \sigma_j]$ containing $x + \epsilon$. Define

$$(3.2) \quad \mathcal{H}_{2j}^\alpha(\epsilon, x, y, r; v) = \{\tau_j < T_x\}, \quad \mathcal{H}_{2j+1}^\beta(\epsilon, x, y, r; v) = \{\sigma_{j+1} < T_x\}.$$

In the definition of \mathcal{H}_{2j}^α and \mathcal{H}_{2j+1}^β , we are interested in the case when y, r are fixed and $x = \epsilon > 0$ small. Imagine η is the interface of the lattice model, then \mathcal{H}_{2j}^α means that there are $2j$ arms connecting $B(x, \epsilon)$ to far away place; and \mathcal{H}_{2j+1}^β means that there are $2j + 1$ arms connecting $B(x, \epsilon)$ to far away place.

The definition here might be confusing at first sight, but these definitions avoid confusions in the proof. We emphasize that we define \mathcal{H}_n^α for odd n in (3.1) and for even n in (3.2); and that we define \mathcal{H}_n^β for even n in (3.1) and for odd n in (3.2).

Propositions 3.1 and 3.2 study the probability of \mathcal{H}_n^α and \mathcal{H}_n^β when the force point v is close to x ; Proposition 3.3 studies the probability of \mathcal{H}_n^α and \mathcal{H}_n^β when the force point v is far from x . Set $\alpha_0^+ = 0$, $\beta_0^+ = 0$ and $\gamma_0^+ = 0$. Assume $j \geq 1$.

PROPOSITION 3.1. *Fix $\kappa \in (0, 4)$ and $\rho \in (-2, 0]$. Define*

$$\alpha_{2j-1}^+ = 2j(2j + \rho + 2 - \kappa/2)/\kappa, \quad \alpha_{2j}^+ = 2j(2j + \rho + 4 - \kappa/2)/\kappa.$$

Suppose $r \geq 1 \vee (200\epsilon)$. We have

$$(3.3) \quad \mathbb{P}[\mathcal{H}_{2j-1}^\alpha(\epsilon, x, y, r; v)] \lesssim x^{\alpha_{2j-2}^+ - \alpha_{2j-1}^+} \epsilon^{\alpha_{2j-1}^+},$$

provided $0 \leq x - v \lesssim \epsilon$, and $|y| \geq (40)^{2j-1}r$,

$$(3.4) \quad \mathbb{P}[\mathcal{H}_{2j}^\alpha(\epsilon, x, y, r; v)] \lesssim x^{\alpha_{2j}^+ - \alpha_{2j-1}^+} \epsilon^{\alpha_{2j-1}^+},$$

provided $0 \leq x - v \lesssim \epsilon$, and $|y| \geq (40)^{2j}r$,

where the constants in \lesssim are uniform over x and ϵ . We also have

$$(3.5) \quad \mathbb{P}[\mathcal{H}_{2j-1}^\alpha(\epsilon, x, y, r; v)] \gtrsim x^{\alpha_{2j-2}^+ - \alpha_{2j-1}^+} \epsilon^{\alpha_{2j-1}^+},$$

provided $0 \leq x - v \lesssim \epsilon$, and $x \asymp r \leq |y| \lesssim r$,

$$(3.6) \quad \mathbb{P}[\mathcal{H}_{2j}^\alpha(\epsilon, x, y, r; v)] \gtrsim x^{\alpha_{2j}^+ - \alpha_{2j-1}^+} \epsilon^{\alpha_{2j-1}^+},$$

provided $0 \leq x - v \lesssim \epsilon$, and $r \leq |y| \lesssim r$,

where the constants in \gtrsim are uniform over x and ϵ . In particular, we have

$$\begin{aligned} \mathbb{P} \left[\mathcal{H}_{2j-1}^\alpha(\epsilon, x, y, r; v) \right] &\asymp \epsilon^{\alpha_{2j-1}^+}, \\ &\text{provided } 0 \leq x - v \lesssim \epsilon, \text{ and } x \asymp r \leq (40)^{2j-1} r \leq |y| \lesssim r, \\ \mathbb{P} \left[\mathcal{H}_{2j}^\alpha(\epsilon, x, y, r; v) \right] &\asymp \epsilon^{\alpha_{2j}^+}, \\ &\text{provided } x \asymp v \asymp \epsilon, \text{ and } (40)^{2j} r \leq |y| \lesssim r, \end{aligned}$$

where the constants in \asymp are uniform over ϵ .

PROPOSITION 3.2. Fix $\kappa \in (0, 4)$ and $\rho \in (-2, \kappa/2 - 2)$. Define

$$\beta_{2j-1}^+ = 2j(2j + \kappa/2 - 4 - \rho)/\kappa, \quad \beta_{2j}^+ = 2j(2j + \kappa/2 - 2 - \rho)/\kappa.$$

Suppose $r \geq 1 \vee (200\epsilon)$. We have

$$(3.7) \quad \mathbb{P} \left[\mathcal{H}_{2j}^\beta(\epsilon, x, y, r; v) \right] \lesssim x^{\beta_{2j-1}^+ - \beta_{2j}^+} \epsilon^{\beta_{2j}^+},$$

provided $0 \leq x - v \lesssim \epsilon$, and $|y| \geq (40)^{2j} r$,

$$(3.8) \quad \mathbb{P} \left[\mathcal{H}_{2j-1}^\beta(\epsilon, x, y, r; v) \right] \lesssim x^{\beta_{2j-1}^+ - \beta_{2j-2}^+} \epsilon^{\beta_{2j-2}^+},$$

provided $0 \leq x - v \lesssim \epsilon$, and $|y| \geq (40)^{2j-1} r$,

where the constants in \lesssim are uniform over x and ϵ . We also have

$$(3.9) \quad \mathbb{P} \left[\mathcal{H}_{2j}^\beta(\epsilon, x, y, r; v) \right] \gtrsim x^{\beta_{2j-1}^+ - \beta_{2j}^+} \epsilon^{\beta_{2j}^+},$$

provided $0 \leq x - v \lesssim \epsilon$, and $x \asymp r \leq |y| \lesssim r$,

$$(3.10) \quad \mathbb{P} \left[\mathcal{H}_{2j-1}^\beta(\epsilon, x, y, r; v) \right] \gtrsim x^{\beta_{2j-1}^+ - \beta_{2j-2}^+} \epsilon^{\beta_{2j-2}^+},$$

provided $0 \leq x - v \lesssim \epsilon$, and $r \leq |y| \lesssim r$,

where the constants in \gtrsim are uniform over x and ϵ . In particular, we have

$$\begin{aligned} \mathbb{P} \left[\mathcal{H}_{2j}^\beta(\epsilon, x, y, r; v) \right] &\asymp \epsilon^{\beta_{2j}^+}, \\ &\text{provided } 0 \leq x - v \lesssim \epsilon, \text{ and } x \asymp r \leq (40)^{2j} r \leq |y| \lesssim r, \\ \mathbb{P} \left[\mathcal{H}_{2j-1}^\beta(\epsilon, x, y, r; v) \right] &\asymp \epsilon^{\beta_{2j-1}^+}, \\ &\text{provided } x \asymp v \asymp \epsilon, \text{ and } (40)^{2j-1} r \leq |y| \lesssim r, \end{aligned}$$

where the constants in \asymp are uniform over ϵ .

PROPOSITION 3.3. Fix $\kappa \in (0, 4)$ and $\rho \in (-2, \kappa/2 - 2)$. Define

$$\gamma_{2j-1}^+ = (2j + \rho)(2j + \rho + 2 - \kappa/2)/\kappa, \quad \gamma_{2j}^+ = 2j(2j + \kappa/2 - 2)/\kappa.$$

Define the event

$$\mathcal{F} = \{\tau_1 < T_x, \eta[0, \tau_1] \subset B(0, Cx), \text{dist}(\eta[0, \tau_1], [x - \epsilon, x + 3\epsilon]) \geq c\epsilon\},$$

where c, C are the constants from Lemma 2.5. We have

$$(3.11) \quad \mathbb{P}[\mathcal{H}_{2j-1}^\alpha(\epsilon, x, y, r; 0^+) \cap \mathcal{F}] \asymp \epsilon^{\gamma_{2j-1}^+},$$

provided $Cx \leq r \leq (40)^{2j-1}r \leq |y| \lesssim r$,

$$(3.12) \quad \mathbb{P}[\mathcal{H}_{2j}^\beta(\epsilon, x, y, r; 0^+) \cap \mathcal{F}] \asymp \epsilon^{\gamma_{2j}^+},$$

provided $Cx \leq r \leq (40)^{2j}r \leq |y| \lesssim r$,

where the constants in \asymp are uniform over ϵ .

We end this section by an informal discussion. Consider Proposition 3.1. Suppose we are allowed to ignore the evolution of the variables y, r and v , and eliminate them from the notation. Then we can prove the conclusion by iteration. Let τ be the first time that η hits the ball $B(x, \epsilon)$, and denote $g_\tau - W_\tau$ by f . The image of the ball $B(x, \epsilon)$ under f is roughly a ball centered at $f(x)$ with radius $g'_\tau(x)\epsilon$. Thus we can write

$$\mathbb{P}[\mathcal{H}_{2j}^\alpha(\epsilon, x)] \approx \mathbb{E}[(g'_\tau(x)\epsilon)^{\alpha_{2j-1}^+} \mathbb{1}_{\{\tau < \infty\}}],$$

and the conclusion can be deduced by an estimate on the expectation of $g'_\tau(x)^\lambda$ for $\lambda \geq 0$. This is a natural first trial. However, the SLE curves can behave badly with small positive chance, and the evolution of the variables y, r and v can be arbitrary. In order to fulfill the above iteration procedure, we need to take care of all the variables. This explains the hard and lengthy work in Sections 3.2 and 3.3. In Section 3.2, we derive a more general version of the estimate on $\mathbb{E}[g'_\tau(x)^\lambda]$. In Section 3.3, we prove Propositions 3.1 to 3.3 by iteration where the results in Section 3.2 play a crucial role.

3.2. *Estimates on the Derivatives.* Suppose η is an $\text{SLE}_\kappa(\rho)$ with force point $v \geq 0$. We use the following notations: g_t is the conformal map from the Loewner chain, W_t is the driving function, V_t is the evolution of the force point and O_t is the rightmost point of $\eta[0, t] \cap \mathbb{R}$ under g_t .

LEMMA 3.4. Fix $\kappa \in (0, 4)$ and $\rho > -2$. For $\lambda \geq 0$, define

$$u_1(\lambda) = \frac{1}{\kappa}(\rho + 4 - \kappa/2) + \frac{1}{\kappa}\sqrt{4\kappa\lambda + (\rho + 4 - \kappa/2)^2}.$$

For $b \in \mathbb{R}$, assume

$$(3.13) \quad 4b \geq (\lambda - b)(2\rho + \kappa(\lambda - b) + 4 - \kappa).$$

Suppose $x \geq v > \epsilon > 0$ and let η be an $\text{SLE}_\kappa(\rho)$ with force point v . Define τ to be the first time that η hits $B(x, \epsilon)$ and T to be the swallowing time of x . If $x = v$, we have

$$(3.14) \quad \mathbb{E} \left[g'_\tau(x)^b (g_\tau(x) - W_\tau)^{\lambda-b} \mathbb{1}_{\{\tau < T\}} \right] \asymp x^{-u_1(\lambda)} \epsilon^{u_1(\lambda) + \lambda - b},$$

where the constants in \asymp are uniform over x and ϵ . For $C \geq 4, 1/4 \geq c > 0$, define

$$\mathcal{F} = \{\tau < T, \text{Im } \eta(\tau) \geq c\epsilon, \eta[0, \tau] \subset B(0, Cx), \text{dist}(\eta[0, \tau], [-Cx, y+r]) \geq cr\}.$$

There exist constants C, c depending only on κ and ρ such that

$$(3.15) \quad \mathbb{E} \left[g'_\tau(x)^\lambda \mathbb{1}_{\mathcal{F}} \right] \asymp x^{-u_1(\lambda)} \epsilon^{u_1(\lambda)},$$

provided $0 \leq x - v \lesssim \epsilon$, and $x \asymp r \leq |y| \lesssim r$,

where the constants in \asymp are uniform over x and ϵ .

To prove Lemma 3.4, we only need to show the upper bound in (3.14) and the lower bound in (3.15). To show the upper bound in (3.14), we need the following Lemma 3.5 which is similar in the spirit of [VL12, Section 6.3]; to show the lower bound in (3.15), we use Lemma 2.5.

LEMMA 3.5. Fix $\kappa > 0$ and $\nu \leq \kappa/2 - 4$. Let η be an $\text{SLE}_\kappa(\nu)$ with force point 1. Set $\Upsilon_t = (g_t(1) - O_t)/g'_t(1)$, $\sigma(s) = \inf\{t : \Upsilon_t = e^{-2s}\}$, and $J_t = (V_t - O_t)/(V_t - W_t)$. Let T be the swallowing time of the point 1. We have, for $\beta \geq 0$,

$$(3.16) \quad \mathbb{E} \left[J_{\sigma(s)}^{-\beta} \mathbb{1}_{\{\sigma(s) < T\}} \right] \asymp 1, \quad \text{when } 8 + 2\nu + \kappa\beta < 2\kappa,$$

where the constants in \asymp depend only on κ, ν, β .

PROOF. Since $0 \leq J_t \leq 1$, we only need to show the upper bound. Set $X_t = V_t - W_t$. We know that

$$dW_t = \sqrt{\kappa}dB_t + \frac{\nu dt}{W_t - V_t}, \quad dV_t = \frac{2dt}{V_t - W_t}.$$

By Itô's formula, we have

$$dJ_t = \frac{J_t}{X_t^2} \left(\kappa - \nu - 2 - \frac{2}{1 - J_t} \right) dt + \frac{J_t}{X_t} \sqrt{\kappa}dB_t, \quad d\Upsilon_t = \Upsilon_t \frac{-2J_t dt}{X_t^2(1 - J_t)}.$$

Recall that $\sigma(s) = \inf\{t : \Upsilon_t = e^{-2s}\}$, and denote by $\hat{X}, \hat{J}, \hat{\Upsilon}$ the processes indexed by $\sigma(s)$. Then we have

$$d\sigma(s) = \hat{X}_s^2 \frac{1 - \hat{J}_s}{\hat{J}_s} ds, \quad d\hat{J}_s = \left(\kappa - \nu - 4 - (\kappa - \nu - 2)\hat{J}_s \right) ds + \sqrt{\kappa\hat{J}_s(1 - \hat{J}_s)} d\hat{B}_s,$$

where \hat{B} is a standard 1-dimensional Brownian motion. By [Law15, Eq. (56), (62)], we know that \hat{J} has an invariant density on $(0, 1)$, which is proportional to $y^{1-(8+2\nu)/\kappa}(1-y)^{4/\kappa-1}$. Moreover, since $\hat{J}_0 = 1$, by a standard coupling argument, we may couple (\hat{J}_s) with the stationary process (\tilde{J}_s) that satisfies the same equation as (\hat{J}_s) , such that $\hat{J}_s \geq \tilde{J}_s$ for all $s \geq 0$. Then we get $\mathbb{E}[\hat{J}_s^{-\beta}] \leq \mathbb{E}[\tilde{J}_s^{-\beta}]$, which is a finite constant if $8 + 2\nu + \kappa\beta < 2\kappa$. This gives the upper bound in (3.16) and completes the proof. \square

PROOF OF LEMMA 3.4–(3.14). Define $\Upsilon_t = (g_t(x) - O_t)/g'_t(x)$, $J_t = (g_t(x) - O_t)/(g_t(x) - W_t)$ and $\hat{\tau}_\epsilon = \inf\{t : \Upsilon_t = \epsilon\}$. Set

$$M_t = g'_t(x)^{(\nu-\rho)(\nu+\rho+4-\kappa)/(4\kappa)} (g_t(x) - W_t)^{(\nu-\rho)/\kappa},$$

where $\nu = \kappa/2 - 4 - \sqrt{4\kappa\lambda + (\rho + 4 - \kappa/2)^2}$. Then M is a local martingale and the law of η weighted by M becomes the law of $\text{SLE}_\kappa(\nu)$ with force point x (see Lemma 2.4). Set $\beta = u_1(\lambda) + \lambda - b$. By the choice of ν , we can rewrite

$$M_t = g'_t(x)^b (g_t(x) - W_t)^{\lambda-b} \Upsilon_t^{-\beta} J_t^\beta.$$

At time $\hat{\tau}_\epsilon < \infty$, we have $\Upsilon_{\hat{\tau}_\epsilon} = \epsilon$. Thus

$$\begin{aligned} & \mathbb{E} \left[g'_{\hat{\tau}_\epsilon}(x)^b (g_{\hat{\tau}_\epsilon}(x) - W_{\hat{\tau}_\epsilon})^{\lambda-b} \mathbb{1}_{\{\hat{\tau}_\epsilon < T\}} \right] \\ &= \epsilon^\beta M_0 \mathbb{E}^* \left[\left(J_{\hat{\tau}_\epsilon}^* \right)^{-\beta} \right] = \epsilon^\beta x^{-u_1(\lambda)} \mathbb{E}^* \left[\left(J_{\hat{\tau}_\epsilon}^* \right)^{-\beta} \right] \asymp \epsilon^\beta x^{-u_1(\lambda)}, \end{aligned}$$

where \mathbb{P}^* is the law of η weighted by M and $\hat{\tau}_\epsilon^*, J^*$ are defined accordingly. The last relation is due to Lemma 3.5. Thus we have

$$(3.17) \quad \mathbb{E} \left[g'_{\hat{\tau}_\epsilon}(x)^b (g_{\hat{\tau}_\epsilon}(x) - W_{\hat{\tau}_\epsilon})^{\lambda-b} \mathbb{1}_{\{\hat{\tau}_\epsilon < T\}} \right] \asymp x^{-u_1(\lambda)} \epsilon^{u_1(\lambda) + \lambda - b}.$$

Consider the process $(U_t := g'_t(x)^b (g_t(x) - W_t)^{\lambda-b})_{t \geq 0}$. We can check that it is a super martingale by Itô's formula when (3.13) holds. Combining with the fact $\hat{\tau}_{\epsilon/4} \geq \tau \geq \hat{\tau}_{4\epsilon}$, we have

$$\mathbb{E} \left[U_{\hat{\tau}_{\epsilon/4}} \mathbb{1}_{\{\hat{\tau}_{\epsilon/4} < T\}} \right] \leq \mathbb{E} \left[U_\tau \mathbb{1}_{\{\tau < T\}} \right] \leq \mathbb{E} \left[U_{\hat{\tau}_{4\epsilon}} \mathbb{1}_{\{\hat{\tau}_{4\epsilon} < T\}} \right].$$

Combining with (3.17), we obtain (3.14). \square

PROOF OF LEMMA 3.4–(3.15). We may assume $x > v$. Define

$$M_t = g'_t(x)^{\nu(\nu+4-\kappa)/(4\kappa)} (g_t(x) - W_t)^{\nu/\kappa} (g_t(x) - g_t(v))^{\nu\rho/(2\kappa)},$$

where $\nu = -\kappa u_1(\lambda)$. Then M is a local martingale for η and the law of η weighted by M is an $\text{SLE}_\kappa(\rho, \nu)$ with force points (v, x) (see Lemma 2.4). We argue that

$$(3.18) \quad g_\tau(x) - g_\tau(v) \asymp (x - v) g'_\tau(x).$$

There are two possibilities: v is swallowed by $\eta[0, \tau]$ or not. If v is not swallowed by $\eta[0, \tau]$, then by the Koebe 1/4 theorem, we know that $g_\tau(x) - g_\tau(v) \asymp (x - v) g'_\tau(x)$. If v is swallowed by $\eta[0, \tau]$, then we must have $x - v \geq \epsilon$. By the Koebe 1/4 theorem, we have $g_\tau(x) - g_\tau(v) \asymp \epsilon g'_\tau(x)$. Since $\epsilon \leq x - v \lesssim \epsilon$, we have $g_\tau(x) - g_\tau(v) \asymp (x - v) g'_\tau(x)$. These complete the proof of (3.18).

On $\{\text{Im } \eta(\tau) \geq c\epsilon\}$, we also have $g_\tau(x) - W_\tau \asymp \epsilon g'_\tau(x)$. Combining with (3.18) and the choice of ν , we have

$$M_\tau \asymp \epsilon^{\nu/\kappa} (x - v)^{\nu\rho/(2\kappa)} g'_\tau(x)^\lambda, \quad \text{on } \mathcal{F}.$$

Therefore,

$$\mathbb{E} \left[g'_\tau(x)^\lambda \mathbb{1}_{\mathcal{F}} \right] \asymp \epsilon^{-\nu/\kappa} (x - v)^{-\nu\rho/(2\kappa)} M_0 \mathbb{P}^*[\mathcal{F}^*] = \epsilon^{u_1(\lambda)} x^{-u_1(\lambda)} \mathbb{P}^*[\mathcal{F}^*],$$

where \mathbb{P}^* is the law of η weighted by M and \mathcal{F}^* is defined accordingly. Note that $\rho > -2, \rho + \nu < \kappa/2 - 4$. By a similar proof of Lemma 2.5, we know that there exists constants C, c such that $\mathbb{P}^*[\mathcal{F}^*] \asymp 1$. This completes the proof. \square

REMARK 3.6. Taking $\lambda = b = 0$ in Lemma 3.4, we see Proposition 3.1 holds for \mathcal{H}_1^α with $\alpha_1^+ = u_1(0) = 2(\rho + 4 - \kappa/2)/\kappa$. Precisely, taking $\lambda = 0$ in (3.15), we have $\mathbb{P}[\eta \text{ hits } B(x, \epsilon)] \gtrsim (\epsilon/x)^{u_1(0)}$. This gives the lower bound. Taking $\lambda = b = 0$ in (3.14), we have $\mathbb{P}[\eta \text{ hits } B(v, \epsilon)] \asymp (\epsilon/v)^{u_1(0)}$. Since $0 \leq x - v \lesssim \epsilon$, we know that $B(x, \epsilon)$ is contained in $B(v, \tilde{C}\epsilon)$ for some constant \tilde{C} , thus

$$\mathbb{P}[\eta \text{ hits } B(x, \epsilon)] \leq \mathbb{P}[\eta \text{ hits } B(v, \tilde{C}\epsilon)] \asymp (\epsilon/v)^{u_1(0)} \lesssim (\epsilon/x)^{u_1(0)}.$$

This gives the upper bound.

LEMMA 3.7. Fix $\kappa \in (0, 4)$ and $\rho \in (-2, 0]$. For $\lambda \geq 0$, define

$$u_2(\lambda) = \frac{1}{\kappa}(\kappa/2 - 2 - \rho) + \frac{1}{\kappa}\sqrt{4\kappa\lambda + (\kappa/2 - 2 - \rho)^2}.$$

Let η be an $\text{SLE}_\kappa(\rho)$ with force point $v > 0$. For $r > 0 > -r > y$, and $0 < v \leq x$, define σ to be the first time that η hits $B(y, r)$ and T to be the swallowing time of x . For $b \leq u_2(\lambda)$ and $x \geq v$, we have

$$(3.19) \quad \mathbb{E} \left[g'_\sigma(x)^\lambda (g_\sigma(x) - W_\sigma)^b \mathbb{1}_{\{\sigma < T\}} \right] \lesssim x^{u_2(\lambda)} (x - y - 2r)^{b - u_2(\lambda)},$$

where the constant in \lesssim is uniform over x, y, r . Assume $r < |y| \lesssim r$, define

$$\mathcal{F} = \{\sigma < T, \text{dist}(\eta[0, \sigma], x) \geq cx, \eta[0, \sigma] \subset B(0, C|y|), \text{dist}(\eta[0, \sigma], [Cy, y]) \geq cr\},$$

where the constants C, c are from Lemma 2.6. Then, for $b \leq u_2(\lambda)$ and $x \geq v \geq (1 - c)x$, we have

$$(3.20) \quad \mathbb{E} \left[g'_\sigma(x)^\lambda (g_\sigma(x) - W_\sigma)^b \mathbb{1}_{\mathcal{F}} \right] \gtrsim x^{u_2(\lambda)} |y|^{b - u_2(\lambda)},$$

where the constant in \gtrsim is uniform over x and y .

PROOF OF LEMMA 3.7–(3.19). We may assume $x > v$. Set

$$M_t = g'_t(x)^{\nu(\nu+4-\kappa)/(4\kappa)} (g_t(x) - W_t)^{\nu/\kappa} (g_t(x) - g_t(v))^{\nu\rho/(2\kappa)},$$

where $\nu = \kappa u_2(\lambda) \geq 0$. By Lemma 2.4, we know that M is a local martingale for η . Note that $g_t(x) - g_t(v) \leq (x - v)g'_t(x)$ and $\nu\rho \leq 0$. We have

$$M_t \geq g'_t(x)^\lambda (g_t(x) - W_t)^{u_2(\lambda)} (x - v)^{\nu\rho/(2\kappa)}.$$

Therefore,

$$\begin{aligned}
& \mathbb{E} \left[g'_\sigma(x)^\lambda (g_\sigma(x) - W_\sigma)^b \mathbb{1}_{\{\sigma < T\}} \right] \\
& \leq (x - v)^{-\nu\rho/(2\kappa)} M_0 \mathbb{E}^* \left[(g_{\sigma^*}^*(x) - W_{\sigma^*}^*)^{b-u_2(\lambda)} \mathbb{1}_{\{\sigma^* < T^*\}} \right] \\
& = x^{u_2(\lambda)} \mathbb{E}^* \left[(g_{\sigma^*}^*(x) - W_{\sigma^*}^*)^{b-u_2(\lambda)} \mathbb{1}_{\{\sigma^* < T^*\}} \right] \\
\text{(by Lemma 2.3)} & \lesssim x^{u_2(\lambda)} (x - y - 2r)^{b-u_2(\lambda)},
\end{aligned}$$

where \mathbb{P}^* is the law of η weighted by M and g^*, W^*, σ^*, T^* are defined accordingly. This implies the conclusion. \square

PROOF OF LEMMA 3.7–(3.20). Assume the same notations as in the proof of (3.19). On $\{\text{dist}(\eta[0, \sigma], x) \geq cx\}$, since $0 \leq x - v \leq cx$, by the Koebe 1/4 theorem, we have $g_t(x) - g_t(v) \geq (x - v)g'_t(x)/4$. Thus

$$M_t \lesssim g'_t(x)^\lambda (g_t(x) - W_t)^{u_2(\lambda)} (x - v)^{\nu\rho/(2\kappa)}.$$

On $\{\eta[0, \sigma] \subset B(0, C|y|)\}$, we have $g_\sigma(x) - W_\sigma \lesssim |y|$. Therefore,

$$\mathbb{E} \left[g'_\sigma(x)^\lambda (g_\sigma(x) - W_\sigma)^b \mathbb{1}_{\mathcal{F}} \right] \gtrsim x^{u_2(\lambda)} |y|^{b-u_2(\lambda)} \mathbb{P}^*[\mathcal{F}^*],$$

where \mathbb{P}^* is the law of η weighted by M and \mathcal{F}^* is defined accordingly. By Lemma 2.6, we have $\mathbb{P}^*[\mathcal{F}^*] \asymp 1$. This completes the proof. \square

REMARK 3.8. Taking $\lambda = b = 0$ in Lemma 3.7, we see Proposition 3.2 holds for \mathcal{H}_1^β with $\beta_1^+ = u_2(0) = 2(\kappa/2 - 2 - \rho)/\kappa$.

LEMMA 3.9. Fix $\kappa \in (0, 4)$ and $\rho > -2$. For $\lambda \geq 0$, define

$$u_3(\lambda) = \frac{(\rho + 2)}{2\kappa} \left(\rho + 4 - \kappa/2 + \sqrt{4\kappa\lambda + (\rho + 4 - \kappa/2)^2} \right).$$

Let η be an $\text{SLE}_\kappa(\rho)$ with force point 0^+ . For $x > \epsilon > 0$, define τ to be the first time that η hits $B(x, \epsilon)$ and T to be the swallowing time of x . Define

$$\mathcal{G} = \{\tau < T, \text{Im } \eta(\tau) \geq c\epsilon\},$$

$$\mathcal{F} = \mathcal{G} \cap \{\eta[0, \tau] \subset B(0, Cx), \text{dist}(\eta[0, \tau], [x - \epsilon, x + 3\epsilon]) \geq c\epsilon\}$$

where c, C are the constants from Lemma 2.5. Then we have

$$\mathbb{E} \left[g'_\tau(x)^\lambda \mathbb{1}_{\mathcal{F}} \right] \asymp \mathbb{E} \left[g'_\tau(x)^\lambda \mathbb{1}_{\mathcal{G}} \right] \asymp x^{-u_3(\lambda)} \epsilon^{u_3(\lambda)},$$

where the constants in \asymp are uniform over x and ϵ .

PROOF. Set $\nu = \kappa/2 - 4 - \rho - \sqrt{4\kappa\lambda + (\kappa/2 - 4 - \rho)^2}$ and

$$M_t = g'_t(x)^{\nu(\nu+4-\kappa)/(4\kappa)}(g_t(x) - W_t)^{\nu/\kappa}(g_t(x) - V_t)^{\nu\rho/(2\kappa)}.$$

Then M is a local martingale and the law of η weighted by M becomes $\text{SLE}_\kappa(\rho, \nu)$ with force points $(0^+, x)$ (see Lemma 2.4). On \mathcal{G} , we have $g_\tau(x) - W_\tau \asymp g_\tau(x) - V_\tau \asymp \epsilon g'_\tau(x)$. Combining with the choice of ν , we have

$$M_\tau \asymp g'_\tau(x)^\lambda \epsilon^{-u_3(\lambda)}, \quad \text{on } \mathcal{G}.$$

Therefore,

$$\mathbb{E} \left[g'_\tau(x)^\lambda \mathbb{1}_{\mathcal{G}} \right] \asymp \epsilon^{u_3(\lambda)} x^{-u_3(\lambda)} \mathbb{P}^*[\mathcal{G}^*], \quad \mathbb{E} \left[g'_\tau(x)^\lambda \mathbb{1}_{\mathcal{F}} \right] \asymp \epsilon^{u_3(\lambda)} x^{-u_3(\lambda)} \mathbb{P}^*[\mathcal{F}^*],$$

where η^* is an $\text{SLE}_\kappa(\rho, \nu)$ with force points $(0^+, x)$, and \mathbb{P}^* denotes its law and $\mathcal{G}^*, \mathcal{F}^*$ are defined accordingly. By Lemma 2.5, we have $\mathbb{P}^*[\mathcal{F}^*] \asymp 1$. This completes the proof. \square

REMARK 3.10. *Taking $\lambda = 0$ in Lemma 3.9, we see Proposition 3.3 holds for \mathcal{H}_1^α with $\gamma_1^+ = u_3(0) = (\rho + 2)(\rho + 4 - \kappa/2)/\kappa$.*

3.3. *Proof of Propositions 3.1 to 3.3.* Fix $\kappa \in (0, 4)$ and $\rho > -2$. Suppose η is an $\text{SLE}_\kappa(\rho)$ with force point $v \geq 0$. We keep the same notations as before: g_t is the conformal map from the Loewner chain, W_t is the driving function, V_t is the evolution of the force point and O_t is the rightmost point of $\eta[0, t] \cap \mathbb{R}$ under g_t , and T is the swallowing time of x . Assume $j \geq 1$.

LEMMA 3.11. *If (3.3) holds for $\mathcal{H}_{2j-1}^\alpha$, then (3.4) holds for \mathcal{H}_{2j}^α .*

PROOF. Let σ be the first time that η hits the ball $B(y, 16(40)^{2j-1}r)$. Denote $g_\sigma - W_\sigma$ by f . Let $\tilde{\eta}$ be the image of $\eta[\sigma, \infty)$ under f . We know that $\tilde{\eta}$ is an $\text{SLE}_\kappa(\rho)$ with force point $f(v)$. Define $\tilde{\mathcal{H}}_{2j-1}^\alpha$ for $\tilde{\eta}$. We have the following observations.

- Consider the image of $B(y, r)$ under f . By Lemma 2.2, we know that $f(B(y, r))$ is contained in the ball with center $f(y)$ and radius $4r f'(y)$. By the Koebe 1/4 theorem, we have

$$|f(y)| \geq 4(40)^{2j-1} r f'(y).$$

- Consider the image of the connected component of $\partial B(x, \epsilon) \setminus \eta[0, \sigma]$ containing $x + \epsilon$ under f . By Lemma 2.1, it is contained in the ball with center $f(x + 3\epsilon)$ and radius $8\epsilon f'(x + 3\epsilon)$. Moreover, we have

$$f(x + 3\epsilon) - f(v) \leq (x + 3\epsilon - v) f'(x + 3\epsilon) \lesssim \epsilon f'(x + 3\epsilon).$$

Combining these two facts with (3.3), we have

$$\begin{aligned} & \mathbb{P} \left[\mathcal{H}_{2j}^\alpha(\epsilon, x, y, r; v) \mid \eta[0, \sigma] \right] \\ & \leq \mathbb{P} \left[\tilde{\mathcal{H}}_{2j-1}^\alpha(8\epsilon f'(x+3\epsilon), f(x+3\epsilon), f(y), 4rf'(y); f(v)) \right] \\ & \lesssim (g_\sigma(x+3\epsilon) - W_\sigma)^{\alpha_{2j-2}^+ - \alpha_{2j-1}^+} (\epsilon g'_\sigma(x+3\epsilon))^{\alpha_{2j-1}^+}. \end{aligned}$$

By Lemma 3.7 and the fact that the swallowing time of $x+3\epsilon$ is greater than T , we have

$$\begin{aligned} & \mathbb{P} \left[\mathcal{H}_{2j}^\alpha(\epsilon, x, y, r; v) \right] \\ & \lesssim \mathbb{E} \left[(g_\sigma(x+3\epsilon) - W_\sigma)^{\alpha_{2j-2}^+ - \alpha_{2j-1}^+} (\epsilon g'_\sigma(x+3\epsilon))^{\alpha_{2j-1}^+} \mathbb{1}_{\{\sigma < T\}} \right] \\ & \lesssim \epsilon^{\alpha_{2j-1}^+} (x+3\epsilon)^{u_2(\alpha_{2j-1}^+)} (x-y-32(40)^{2j-1}r)^{\alpha_{2j-2}^+ - \alpha_{2j}^+} \\ & \lesssim x^{\alpha_{2j}^+ - \alpha_{2j-1}^+} \epsilon^{\alpha_{2j-1}^+}. \end{aligned}$$

The last line is because $x \geq \epsilon$ and $|y| \geq (40)^{2j}r$. \square

LEMMA 3.12. *If (3.4) holds for \mathcal{H}_{2j}^α , then (3.3) holds for $\mathcal{H}_{2j+1}^\alpha$.*

PROOF. If $x \leq 64\epsilon$, then $\mathbb{P} \left[\mathcal{H}_{2j+1}^\alpha(\epsilon, x, y, r; v) \right] \leq \mathbb{P} \left[\mathcal{H}_{2j}^\alpha(\epsilon, x, y, r; v) \right]$. This gives the conclusion. In the following, we may assume $x > 64\epsilon$. Let τ be the first time that η hits $B(x, 16\epsilon)$. Denote $g_\tau - W_\tau$ by f . Let $\tilde{\eta}$ be the image of $\eta[\tau, \infty)$ under f . We know that $\tilde{\eta}$ is an SLE $_\kappa(\rho)$ with force point $f(v)$. Define $\tilde{\mathcal{H}}_{2j}^\alpha$ for $\tilde{\eta}$. We have the following observations.

- Consider the image of the connected component of $\partial B(y, r) \setminus \eta[0, \tau]$ containing $y-r$ under f . By Lemma 2.1, we know that it is contained in the ball with center $f(y-3r)$ and radius $8rf'(y-3r)$. By Lemma 2.3, we have

$$|f(y-3r)| \geq (x-y+3r-32\epsilon)/2 \geq (40)^{2j}8r.$$

- Consider the image of $B(x, \epsilon)$ under f . By Lemma 2.2, we know that $B(x, \epsilon)$ is contained in the ball with center $f(x)$ and radius $4\epsilon f'(x)$. Moreover,

$$f(x) - f(v) \leq (x-v)f'(x) \lesssim \epsilon f'(x).$$

Combining these two facts with (3.4), we have

$$\begin{aligned} & \mathbb{P} \left[\mathcal{H}_{2j+1}^\alpha(\epsilon, x, y, r; v) \mid \eta[0, \tau] \right] \\ & \leq \mathbb{P} \left[\tilde{\mathcal{H}}_{2j}^\alpha(4\epsilon f'(x), f(x), f(y-3r), 8rf'(y-3r); f(v)) \right] \\ & \lesssim (g_\tau(x) - W_\tau)^{\alpha_{2j}^+ - \alpha_{2j-1}^+} (\epsilon g'_\tau(x))^{\alpha_{2j-1}^+}. \end{aligned}$$

If $x = v$, by Lemma 3.4, since α_{2j-1}^+ and α_{2j}^+ satisfy (3.13):

$$\kappa \left(\alpha_{2j}^+ - \alpha_{2j-1}^+ \right) \left(2\rho + 4 - \kappa + \kappa \left(\alpha_{2j}^+ - \alpha_{2j-1}^+ \right) \right) = 4\kappa\alpha_{2j-1}^+,$$

we have

$$\begin{aligned} & \mathbb{P} \left[\mathcal{H}_{2j+1}^\alpha(\epsilon, v, y, r; v) \right] \\ & \lesssim \mathbb{E} \left[\left(g_\tau(v) - W_\tau \right)^{\alpha_{2j}^+ - \alpha_{2j-1}^+} \left(\epsilon g'_\tau(v) \right)^{\alpha_{2j-1}^+} \mathbb{1}_{\{\hat{\tau} < T_v\}} \right] \\ & \lesssim v^{-u_1(\alpha_{2j}^+)} \epsilon^{\alpha_{2j+1}^+} = v^{\alpha_{2j}^+ - \alpha_{2j+1}^+} \epsilon^{\alpha_{2j+1}^+}. \end{aligned}$$

For $0 \leq x - v \lesssim \epsilon$, we know that $B(x, \epsilon)$ is contained in $B(v, \tilde{C}\epsilon)$ for some constant \tilde{C} , thus

$$\begin{aligned} & \mathbb{P} \left[\mathcal{H}_{2j+1}^\alpha(\epsilon, x, y, r; v) \right] \\ & \leq \mathbb{P} \left[\mathcal{H}_{2j+1}^\alpha(\tilde{C}\epsilon, v, y, r; v) \right] \lesssim v^{\alpha_{2j}^+ - \alpha_{2j+1}^+} \epsilon^{\alpha_{2j+1}^+} \lesssim x^{\alpha_{2j}^+ - \alpha_{2j+1}^+} \epsilon^{\alpha_{2j+1}^+}. \end{aligned}$$

This gives the conclusion. \square

LEMMA 3.13. *If (3.5) holds for $\mathcal{H}_{2j-1}^\alpha$, then (3.6) holds for \mathcal{H}_{2j}^α .*

PROOF. Let σ be the first time that η hits $B(y, r)$. Define

$$\mathcal{F} = \{ \sigma < T, \text{dist}(\eta[0, \sigma], x) \geq cx, \eta[0, \sigma] \subset B(0, C|y|), \text{dist}(\eta[0, \sigma], [Cy, y]) \geq cr \},$$

where c, C are the constants from Lemma 2.5. Denote $g_\sigma - W_\sigma$ by f . Let $\tilde{\eta}$ be the image of $\eta[\sigma, \infty)$ under f , then $\tilde{\eta}$ is an SLE $_\kappa(\rho)$ with force point $f(v)$. Given $\eta[0, \sigma]$ and on \mathcal{F} , we have the following observations.

- Consider the image of $B(y, r)$ under f . By the Koebe 1/4 theorem, it contains the ball with center $f(y)$ and radius $rf'(y)/4$. On $\{ \text{dist}(\eta[0, \sigma], [Cy, y]) \geq cr \}$, we have

$$rf'(y)/4 \leq |f(y)| \lesssim rf'(y).$$

- Consider the image of $B(x, \epsilon)$ under f . On $\{ \text{dist}(\eta[0, \sigma], x) \geq cx \}$, by the Koebe 1/4 theorem, it contains the ball with the center $f(x)$ and radius $c\epsilon f'(x)/4$. Since $x - v \lesssim \epsilon$, we have

$$f(x) - f(v) \leq (x - v)f'(x) \lesssim \epsilon f'(x).$$

- Compare $f(x)$ and $|f(y)| \asymp rf'(y)$. On $\{ \eta[0, \sigma] \subset B(0, C|y|) \}$, we have $f(x) \lesssim |y|$. On $\{ \text{dist}(\eta[0, \sigma], [Cy, y]) \geq cr \}$, we have $|f(y)| \gtrsim |y|$. Thus, on \mathcal{F} , we have

$$f(x) \lesssim |y| \lesssim |f(y)| \asymp rf'(y).$$

Combining these three facts with (3.5), we have

$$\begin{aligned} & \mathbb{P} [\mathcal{H}_{2j}^\alpha(\epsilon, x, y, r; v) \mid \eta[0, \sigma], \mathcal{F}] \\ & \geq \mathbb{P} [\tilde{\mathcal{H}}_{2j-1}^\alpha(\epsilon f'(x)/4, f(x), f(y), r f'(y)/4; f(v))] \\ & \gtrsim (g_\sigma(x) - W_\sigma)^{\alpha_{2j-2}^+ - \alpha_{2j-1}^+} (\epsilon g'_\sigma(x))^{\alpha_{2j-1}^+}. \end{aligned}$$

By Lemma 3.7, we have

$$\begin{aligned} & \mathbb{P} [\mathcal{H}_{2j}^\alpha(\epsilon, x, y, r; v) \cap \mathcal{F}] \\ & \gtrsim \mathbb{E} \left[(g_\sigma(x) - W_\sigma)^{\alpha_{2j-2}^+ - \alpha_{2j-1}^+} (\epsilon g'_\sigma(x))^{\alpha_{2j-1}^+} \mathbb{1}_{\mathcal{F}} \right] \\ & \gtrsim x^{u_2(\alpha_{2j-1}^+)} \epsilon^{\alpha_{2j-1}^+} = x^{\alpha_{2j}^+ - \alpha_{2j-1}^+} \epsilon^{\alpha_{2j-1}^+}. \end{aligned}$$

This gives the conclusion. \square

LEMMA 3.14. *If (3.6) holds for \mathcal{H}_{2j}^α , then (3.5) holds for $\mathcal{H}_{2j+1}^\alpha$.*

PROOF. Let τ be the first time that η hits $B(x, \epsilon)$. Define

$$\mathcal{F} = \{\tau < T, \text{Im } \eta(\tau) \geq c\epsilon, \eta[0, \tau] \subset B(0, Cx), \text{dist}(\eta[0, \tau], [-Cx, y+r]) \geq cr\},$$

where c, C are constants from Lemma 3.4. Denote $g_\tau - W_\tau$ by f . Let $\tilde{\eta}$ be the image of $\eta[\tau, \infty)$ under f , then $\tilde{\eta}$ is an SLE $_\kappa(\rho)$ with force point $f(v)$. Define $\tilde{\mathcal{H}}_{2j}^\alpha$ for $\tilde{\eta}$. Given $\eta[0, \tau]$ and on \mathcal{F} , we have the following observations.

- Consider the image of $B(y, r)$ under f . On \mathcal{F} , we know that $f(B(y, r))$ contains the ball with center $f(y)$ and radius $cr f'(y)/4$; moreover, we have

$$cr f'(y)/4 \leq |f(y)| \lesssim r f'(y).$$

- Consider the image of $B(x, \epsilon)$ under f . By the Koebe 1/4 theorem, it contains the ball with center $f(x)$ and radius $\epsilon f'(x)/4$. On $\{\text{Im } \eta(\tau) \geq c\epsilon\}$, we have

$$f(x) \asymp \epsilon f'(x).$$

Since $x - v \lesssim \epsilon$, we have

$$f(x) - f(v) \leq (x - v) f'(x) \lesssim \epsilon f'(x).$$

Combining these two facts with (3.6), we have

$$\begin{aligned} & \mathbb{P} [\mathcal{H}_{2j+1}^\alpha(\epsilon, x, y, r; v) \mid \eta[0, \tau], \mathcal{F}] \\ & \geq \mathbb{P} [\tilde{\mathcal{H}}_{2j}^\alpha(\epsilon f'(x)/4, f(x), f(y), r f'(y)/4; f(v))] \gtrsim (\epsilon g'_\tau(x))^{\alpha_{2j}^+}. \end{aligned}$$

By Lemma 3.4, we have

$$\begin{aligned} & \mathbb{P} \left[\mathcal{H}_{2j+1}^\alpha(\epsilon, x, y, r; v) \cap \mathcal{F} \right] \\ & \gtrsim \mathbb{E} \left[(\epsilon g'_\tau(x))^{\alpha_{2j}^+} \mathbb{1}_{\mathcal{F}} \right] \asymp x^{-u_1(\alpha_{2j}^+)} \epsilon^{u_1(\alpha_{2j}^+) + \alpha_{2j}^+} = x^{\alpha_{2j}^+ - \alpha_{2j+1}^+} \epsilon^{\alpha_{2j+1}^+}. \end{aligned}$$

□

PROOF OF PROPOSITION 3.1. Note that

$$\alpha_{2j+1}^+ = \alpha_{2j}^+ + u_1(\alpha_{2j}^+), \quad \alpha_{2j}^+ = \alpha_{2j-1}^+ + u_2(\alpha_{2j-1}^+).$$

Combining Remark 3.6 with Lemmas 3.11, 3.12, 3.13 and 3.14, we obtain the conclusion. □

PROOF OF PROPOSITION 3.2. By Remark 3.8, we know the conclusion is true for \mathcal{H}_1^β . Note that

$$\beta_{2j}^+ = \beta_{2j-1}^+ + u_1(\beta_{2j-1}^+), \quad \beta_{2j+1}^+ = \beta_{2j}^+ + u_2(\beta_{2j}^+).$$

Moreover, the exponents β_{2j-2}^+ and β_{2j-1}^+ satisfy (3.13):

$$\kappa \left(\beta_{2j-1}^+ - \beta_{2j-2}^+ \right) \left(2\rho + 4 - \kappa + \kappa \left(\beta_{2j-1}^+ - \beta_{2j-2}^+ \right) \right) = 4\kappa\beta_{2j-2}^+.$$

We can prove the following:

If (3.8) holds for \mathcal{H}_{2j-1}^β , then (3.7) holds for \mathcal{H}_{2j}^β (by the proof of Lemma 3.12).
 If (3.7) holds for \mathcal{H}_{2j}^β , then (3.8) holds for \mathcal{H}_{2j+1}^β (by the proof of Lemma 3.11).
 If (3.10) holds for \mathcal{H}_{2j-1}^β , then (3.9) holds for \mathcal{H}_{2j}^β (by the proof of Lemma 3.14).
 If (3.9) holds for \mathcal{H}_{2j}^β , then (3.10) holds for \mathcal{H}_{2j+1}^β (by the proof of Lemma 3.13).
 Combining all these, we complete the proof. □

PROOF OF PROPOSITION 3.3–(3.11)–UPPER BOUND. By Remark 3.10, we know that the conclusion is true for \mathcal{H}_1^α . We will prove the conclusion for $\mathcal{H}_{2j+1}^\alpha$ for $j \geq 1$. Recall that η is an SLE $_\kappa(\rho)$ with force point 0^+ . Let τ be the first time that η hits $B(x, \epsilon)$, and T be the first time that η swallows x . Recall that

$$\mathcal{F} = \{ \tau < T, \eta[0, \tau] \subset B(0, Cx), \text{dist}(\eta[0, \tau], [x - \epsilon, x + 3\epsilon]) \geq c\epsilon \}.$$

Given $\eta[0, \tau]$, denote $g_\tau - W_\tau$ by f . Let $\tilde{\eta}$ be the image of $\eta[\tau, \infty)$ under f , then $\tilde{\eta}$ is an SLE $_\kappa(\rho)$ with force point $f(0^+)$. Define \mathcal{H}_{2j}^α for $\tilde{\eta}$. We have the following observations.

- Consider the image of the connected component of $\partial B(x, \epsilon) \setminus \eta[0, \tau]$ containing $x + \epsilon$ under f . By Lemma 2.1, we know that it is contained in the ball with center $f(x + 3\epsilon)$ and radius $8\epsilon f'(x + 3\epsilon)$. On the event $\{\text{dist}(\eta[0, \tau], [x - \epsilon, x + 3\epsilon]) \geq c\epsilon\}$, by Koebe distortion theorem [Pom92, Chapter I, Theorem 1.3], we know that there exists some universal constant \tilde{C} such that the ball with center $f(x + 3\epsilon)$ and radius $8\epsilon f'(x + 3\epsilon)$ is contained in the ball with center $f(x)$ and radius $\tilde{C}\epsilon f'(x)$. Moreover, on the event $\{\text{dist}(\eta[0, \tau], [x - \epsilon, x + 3\epsilon]) \geq c\epsilon\}$, we have

$$f(x) \asymp f(x) - f(0^+) \asymp \epsilon f'(x).$$

- Consider the image of the connected component of $\partial B(y, r) \setminus \eta[0, \tau]$ containing $y - r$ under f . By Lemma 2.1, we know that it is contained in the ball with center $f(y - 3r)$ and radius $8r f'(y - 3r)$. By Lemma 2.3, we know that

$$|f(y - 3r)| \geq (x - y + 3r - 2\epsilon)/2 \geq |y|/2 \geq (40)^{2j} 8r.$$

Combining these two facts with (3.4), we have

$$\mathbb{P} [\mathcal{H}_{2j+1}^\alpha(\epsilon, x, y, r; 0^+) \mid \eta[0, \tau], \mathcal{F}] \lesssim (\epsilon g'_\tau(x))^{\alpha_{2j}^+}.$$

By Lemma 3.9, we have

$$\mathbb{P} [\mathcal{H}_{2j+1}^\alpha(\epsilon, x, y, r; 0^+) \cap \mathcal{F}] \lesssim \mathbb{E} \left[(\epsilon g'_\tau(x))^{\alpha_{2j}^+} \mathbb{1}_{\mathcal{F}} \right] \asymp \epsilon^{u_3(\alpha_{2j}^+) + \alpha_{2j}^+}.$$

Note that

$$\gamma_{2j+1}^+ = u_3(\alpha_{2j}^+) + \alpha_{2j}^+.$$

This completes the proof. \square

PROOF OF PROPOSITION 3.3–(3.11)–LOWER BOUND. Assume the same notations as in the proof of the upper bound. We have the following observations.

- Consider the image of $B(x, \epsilon)$ under f . By Koebe 1/4 theorem, it contains the ball with center $f(x)$ and radius $\epsilon f'(x)/4$. Moreover, on the event \mathcal{F} , we have

$$f(x) \asymp f(x) - f(0^+) \asymp \epsilon f'(x).$$

- Consider the image of $B(y, r)$ under f . Note that $r \geq Cx$ and $|y| \geq (40)^{2j+1}r$. Thus, on the event $\{\eta[0, \tau] \subset B(0, Cx)\}$, we know that $\eta[0, \tau]$ does not hit $B(y, r)$. Thus $f(B(y, r))$ contains the ball with

center $f(y)$ and radius $rf'(y)/4$. On the event $\{\eta[0, \tau] \subset B(0, Cx)\}$, we know that

$$rf'(y)/4 \leq |f(y)| \leq |y| + (Cx)^2/|y| \leq 2|y| \asymp r.$$

Combining these two facts with (3.6), we have

$$\mathbb{P} [\mathcal{H}_{2j+1}^\alpha(\epsilon, x, y, r; 0^+) \mid \eta[0, \tau], \mathcal{F}] \gtrsim (\epsilon g'_\tau(x))^{\alpha_{2j}^+}.$$

By Lemma 3.9, we have

$$\mathbb{P} [\mathcal{H}_{2j+1}^\alpha(\epsilon, x, y, r; 0^+) \cap \mathcal{F}] \gtrsim \mathbb{E} \left[(\epsilon g'_\tau(x))^{\alpha_{2j}^+} \mathbb{1}_{\mathcal{F}} \right] \asymp \epsilon^{u_3(\alpha_{2j}^+) + \alpha_{2j}^+}.$$

This completes the proof. \square

PROOF OF PROPOSITION 3.3–(3.12). By the same proof of (3.11), we can prove that

$$\mathbb{P} [\mathcal{H}_{2j}^\alpha(\epsilon, x, y, r; 0^+) \cap \mathcal{F}] \asymp \mathbb{E} \left[(\epsilon g'_\tau(x))^{\beta_{2j-1}^+} \right] \asymp \epsilon^{u_3(\beta_{2j-1}^+) + \beta_{2j-1}^+}.$$

Note that

$$\gamma_{2j}^+ = u_3(\beta_{2j-1}^+) + \beta_{2j-1}^+.$$

This completes the proof. \square

4. SLE Interior Arm Exponents. Fix $\kappa \in (0, 4)$ and let η be an SLE_κ in \mathbb{H} from 0 to ∞ . We keep the same notations as before: g_t is the family of conformal maps for the Loewner chain, W_t is the driving function. We write c.c. for “connected component”.

Fix $z \in \mathbb{H}$ with $|z| = 1$ and suppose $r > 0$ and $y \leq -4r$. Let τ_1 be the first time that η hits $B(z, \epsilon)$. Define

$$\mathcal{E}_2(\epsilon, z) = \{\tau_1 < \infty\}.$$

Let σ_1 be the first time after τ_1 that η hits the c.c. of $\partial B(y, r) \setminus \eta[0, \tau_1]$ containing $y - r$. Define \mathcal{E}^g to be the event that z is in the unbounded c.c. of $\mathbb{H} \setminus (\eta[0, \sigma_1] \cup B(y, r))$.

Given $\eta[0, \sigma_1]$, we know that $B(z, \epsilon) \setminus \eta[0, \sigma_1]$ has one c.c. that contains z , denoted by C_z . The boundary ∂C_z consists of pieces of $\eta[0, \sigma_1]$ and pieces of $\partial B(z, \epsilon)$. Consider $\partial C_z \cap \partial B(z, \epsilon)$. There may be several c.c.s, but there is only one which can be connected to ∞ in $\mathbb{H} \setminus (\eta[0, \sigma_1] \cup B(z, \epsilon))$. We denote this c.c. by C_z^b , oriented it clockwise and denote the end point as X_z^b . See Fig 4.1.




Fig 4.1: The gray part is the c.c. of $B(z, \epsilon) \setminus \eta[0, \sigma_1]$ that contains z , which is denoted by C_z . The bold part of ∂C_z is C_z^b . The point X_z^b is denoted in the figure.

Let τ_2 be the first time after σ_1 that η hits C_z^b , and let σ_2 be the first time after τ_2 that η hits the c.c. of $\partial B(y, r) \setminus \eta[0, \tau_2]$ containing $y - r$. For $j \geq 2$, let τ_j be the first time after σ_{j-1} such that η hits the c.c. of $C_z^b \setminus \eta[0, \sigma_{j-1}]$ containing X_z^b and let σ_j be the first time after τ_j that η hits the c.c. of $\partial B(y, r) \setminus \eta[0, \tau_j]$ containing $y - r$. For $j \geq 2$, define

$$\mathcal{E}_{2j}(\epsilon, z, y, r) = \mathcal{E}^g \cap \{\tau_j < T_z\}.$$

We will prove the following estimate on the probability of \mathcal{E}_{2j} .

PROPOSITION 4.1. *Fix $\kappa \in (0, 4)$ and $z \in \mathbb{H}$ with $|z| = 1$. For $j \geq 1$, define*

$$\alpha_{2j} = (16j^2 - (\kappa - 4)^2)/(8\kappa).$$

Define $\mathcal{F} = \{\eta[0, \tau_1] \subset B(0, R)\}$ where R is the constant from Lemma 4.2. Then we have, for $j \geq 1$,

$$(4.1) \quad \mathbb{P}[\mathcal{E}_{2j}(\epsilon, z, y, r) \cap \mathcal{F}] = \epsilon^{\alpha_{2j} + o(1)},$$

provided $R \leq r \leq (40)^{2j} r \leq |y| \lesssim r$.

We will first explain the choice of the constant R in Lemma 4.2, and then prove the lower bound and the upper bound of (4.1) separately. The lower bound is easier, and the upper bound requires the estimates in Lemmas 4.3 and 4.4.

LEMMA 4.2. Fix $\kappa \in (0, 8)$ and let η be an SLE_κ in \mathbb{H} from 0 to ∞ . For $\lambda \geq 0$, define

$$\rho = \kappa/2 - 4 - \sqrt{4\kappa\lambda + (\kappa/2 - 4)^2}, \quad v(\lambda) = \frac{1}{2} - \frac{\kappa}{16} - \frac{\lambda}{2} + \frac{1}{8}\sqrt{4\kappa\lambda + (\kappa/2 - 4)^2}.$$

Fix $z \in \mathbb{H}$ with $|z| = 1$. For $\epsilon > 0$, let τ be the first time that η hits $B(z, \epsilon)$. Define $\Theta_t = \arg(g_t(z) - W_t)$. For $\delta \in (0, 1/16)$, $R \geq 4$, define

$$\mathcal{G} = \{\tau < \infty, \Theta_\tau \in (\delta, \pi - \delta)\}, \quad \mathcal{F} = \{\eta[0, \tau] \subset B(0, R)\}.$$

There exists a constant R depending only on κ and z such that the following is true:

$$\epsilon^{v(\lambda)} \lesssim \mathbb{E} \left[|g'_\tau(z)|^\lambda \mathbb{1}_{\mathcal{F} \cap \mathcal{G}} \right] \leq \mathbb{E} \left[|g'_\tau(z)|^\lambda \mathbb{1}_{\mathcal{G}} \right] \lesssim \epsilon^{v(\lambda)} \delta^{-v(\lambda) - \rho^2/(2\kappa)},$$

where the constants in \lesssim are uniform over ϵ, δ .

PROOF. Similar results were proved in [VL12, Section 6.3] and [MW17, Lemmas 4.1, 4.2] with constants in \lesssim only uniform over ϵ . In our setting, we need the explicit dependence on δ . Set

$$M_t = |g'_t(z)|^{\rho(\rho+8-2\kappa)/(8\kappa)} (\text{Im } g_t(z))^{\rho^2/(8\kappa)} |g_t(z) - W_t|^{\rho/\kappa}.$$

From [SW05, Theorem 6], the process M is a local martingale and the law of η weighted by M becomes the law of $\text{SLE}_\kappa(\rho)$ with force point z . We introduce two other quantities:

$$\Upsilon_t = \frac{\text{Im } g_t(z)}{|g'_t(z)|}, \quad S_t = \sin \Theta_t = \frac{\text{Im } g_t(z)}{|g_t(z) - W_t|}.$$

Then we can rewrite M as follows:

$$M_t = |g'_t(z)|^\lambda \Upsilon_t^{-v(\lambda)} S_t^{v(\lambda) + \rho^2/(8\kappa)}.$$

By the Koebe 1/4 theorem, we know that $\Upsilon_\tau \asymp \epsilon$. On \mathcal{G} , we know that $S_\tau \geq \delta/2$ for $\delta < 1/16$. Thus

$$\epsilon^{v(\lambda)} \mathbb{P}^*[\mathcal{F}^* \cap \mathcal{G}^*] \lesssim \mathbb{E} \left[|g'_t(z)|^\lambda \mathbb{1}_{\mathcal{F} \cap \mathcal{G}} \right] \leq \mathbb{E} \left[|g'_t(z)|^\lambda \mathbb{1}_{\mathcal{G}} \right] \lesssim \epsilon^{v(\lambda)} \delta^{-v(\lambda) - \rho^2/(8\kappa)},$$

where η^* is an $\text{SLE}_\kappa(\rho)$ with force point z , \mathbb{P}^* denotes its law and $\tau^*, \Theta^*, \mathcal{F}^*, \mathcal{G}^*$ are defined accordingly. By [MW17, Eq. (4.7), (4.8)], we have

$$\mathbb{P}^*[\eta^*[0, \tau^*] \subset B(0, R)] \rightarrow 1, \quad \text{as } R \rightarrow \infty; \quad \text{and} \quad \mathbb{P}^*[\Theta_{\tau^*}^* \in (1/16, \pi - 1/16)] \asymp 1.$$

Therefore, there exists a constant R depending only on κ and z such that

$$\mathbb{P}^*[\mathcal{F}^* \cap \mathcal{G}^*] \geq \mathbb{P}^*[\eta^*[0, \tau^*] \subset B(0, R), \Theta_{\tau^*}^* \in (1/16, \pi - 1/16)] \asymp 1.$$

This completes the proof. \square

We will fix the constant R from Lemma 4.2 in the following of the section. The conclusion for \mathcal{E}_2 was proved in [Bef08, Proposition 4], we will prove the conclusion for \mathcal{E}_{2j+2} for $j \geq 1$. We will need the following conclusion from Section 3. For $j \geq 1$, taking $\rho = 0$ in Proposition 3.1, we have $\alpha_{2j}^+ = 2j(2j + 4 - \kappa/2)/\kappa$ and

$$(4.2) \quad \mathbb{P} [\mathcal{H}_{2j}^\alpha(\epsilon, x, y, r)] \asymp x^{\alpha_{2j}^+} \epsilon^{-\alpha_{2j}^+} \epsilon^{\alpha_{2j-1}^+}, \quad \text{provided } (40)^{2j} r \leq |y| \lesssim r.$$

Note that, since $\rho = 0$, we may assume $v = x$ and we eliminate the force point in the definition of \mathcal{H}_{2j}^α .

PROOF OF PROPOSITION 4.1–(4.1)–LOWER BOUND. We will prove the lower bound for the probability of \mathcal{E}_{2j+2} . Let η be an SLE $_\kappa$ in \mathbb{H} from 0 to ∞ . Let τ be the first time that η hits $B(z, \epsilon)$. Denote the centered conformal map $g_t - W_t$ by f_t for $t \geq 0$. Recall that $\mathcal{F} = \{\eta[0, \tau] \subset B(0, R)\}$. Fix some $\delta > 0$ and define $\mathcal{G} = \mathcal{F} \cap \{\Theta_\tau \in (\delta, \pi - \delta)\}$.

We run η until the time τ . On \mathcal{G} , by the Koebe 1/4 theorem, we know that $f_\tau(B(z, \epsilon))$ contains the ball with center $w := f_\tau(z)$ and radius $u := \epsilon |f'_\tau(z)|/4$ and

$$\arg(w) \in (\delta, \pi - \delta), \quad u \leq \text{Im } w \leq 16u.$$

We wish to apply (4.2), however this ball is centered at $w = f_\tau(z)$ which does not satisfy the conditions in (4.2). We will fix this problem by running η a little further and argue that there is a positive chance that η does the right thing.

Let $\tilde{\eta}$ be the image of $\eta[\tau, \infty)$ under f_τ . Let γ be the broken line from 0 to w and then to $-u + ui$ and let A_u be the $u/4$ -neighborhood of γ . Let S_1 be the first time that $\tilde{\eta}$ exits A_u and let S_2 be the first time that $\tilde{\eta}$ hits the ball with center $-u + ui$ and radius $u/4$. By [MW17, Lemma 2.5], we know that $\mathbb{P}[S_2 < S_1]$ is bounded from below by positive constant depending only on κ and δ . On $\{S_2 < S_1\}$, it is clear that there exist constants $x_\delta, c_\delta > 0$ depending only on δ such that $f_{S_2}(B(z, \epsilon))$ contains the ball with center $x_\delta u$ and radius $c_\delta u$.

Consider the image of $B(y, r)$ under f_{S_2} . On $\mathcal{F} \cap \{S_2 < S_1\}$, we know that the image of $B(y, r)$ under f_{S_2} contains the ball with center $f_{S_2}(y)$ and radius $r f'_{S_2}(y)/4$ where

$$2y \leq f_{S_2}(y) \leq y, \quad f'_{S_2}(y) \asymp 1.$$

Combining with (4.2), we have

$$\mathbb{P} [\mathcal{E}_{2j+2}(\epsilon, z, y, r) \mid \eta[0, S_2], \mathcal{G} \cap \{S_2 < S_1\}] \gtrsim (\epsilon |g'_\tau(z)|)^{\alpha_{2j}^+}.$$

Since $\{S_2 < S_1\}$ has a positive chance, we have

$$\mathbb{P}[\mathcal{E}_{2j+2}(\epsilon, z, y, r) \mid \eta[0, \tau], \mathcal{G}] \gtrsim (\epsilon |g'_\tau(z)|)^{\alpha_{2j}^+}.$$

Therefore, by Lemma 4.2, we have

$$\mathbb{P}[\mathcal{E}_{2j+2}(\epsilon, z, y, r)] \gtrsim \mathbb{E}\left[(\epsilon |g'_\tau(z)|)^{\alpha_{2j}^+} \mathbf{1}_{\mathcal{G}}\right] \asymp \epsilon^{v(\alpha_{2j}^+) + \alpha_{2j}^+} = \epsilon^{\alpha_{2j+2}},$$

where the constants in \gtrsim and \asymp are uniform over ϵ . This completes the proof. \square

LEMMA 4.3. *Fix $\kappa \in (0, 4)$ and let η be an SLE_κ in \mathbb{H} from 0 to ∞ . Fix $z \in \mathbb{H}$ with $|z| = 1$. Let $\Theta_t = \arg(g_t(z) - W_t)$. For $C \geq 16$, let ξ be the first time that η hits $\partial B(z, C\epsilon)$. For $\delta \in (0, 1/16)$, define*

$$\mathcal{F} = \{\xi < \infty, \Theta_\xi \in (\delta, \pi - \delta), \eta[0, \xi] \subset B(0, R)\}.$$

Then we have

$$\mathbb{P}[\mathcal{E}_{2j+2}(\epsilon, z, y, r) \cap \mathcal{F}] \lesssim C^A \delta^{-B} \epsilon^{\alpha_{2j+2}}, \quad \text{provided } y \leq -20r, r \geq R.$$

where A, B are some constants depending on κ and j , and the constant in \lesssim is uniform over δ, C, ϵ .

PROOF. We run the curve up to time ξ and let $f = g_\xi - W_\xi$. We have the following observations.

- Consider $f(B(z, \epsilon))$. By Lemma 2.2, we know that $f(B(z, \epsilon))$ is contained in the ball with center $f(z)$ and radius $u := 4\epsilon |f'(z)|$. Applying the Koebe 1/4 theorem to f , we have

$$(4.3) \quad C\epsilon |f'(z)|/4 \leq \text{Im } f(z) \leq 4C\epsilon |f'(z)|.$$

Next, we argue that $f(B(z, \epsilon))$ is contained in the ball with center $|f(z)| \in \mathbb{R}$ and radius $8Cu/\delta$. Since $f(B(z, \epsilon))$ is contained in the ball with center $f(z)$ and radius u , it is clear that $f(B(z, \epsilon))$ is contained in the ball with center $|f(z)|$ with radius $u + 2|f(z)|$. By (4.3), we have

$$Cu/16 \leq |f(z)| \sin \Theta_\xi \leq Cu.$$

Since $\Theta_\xi \in (\delta, \pi - \delta)$, we know that, for $\delta > 0$ small, we have $\sin \Theta_\xi \geq \delta/2$. Thus, $Cu/16 \leq |f(z)| \leq 2Cu/\delta$. Therefore, $f(B(z, \epsilon))$ is contained in the ball with center $|f(z)|$ with radius $8Cu/\delta$. In summary, we know that $f(B(z, \epsilon))$ is contained in the ball with center $|f(z)|$ and radius $32C\epsilon |f'(z)|/\delta$ where

$$C\epsilon |f'(z)|/4 \leq |f(z)| \leq 8C\epsilon |f'(z)|/\delta.$$

- Consider $f(B(y, r))$. Since $\{\eta[0, \xi] \subset B(0, R)\}$ and $y \leq -20r$ with $r \geq R$, we know that $f(B(y, r))$ is contained in the ball with center $f(y)$ and radius $4rf'(y)$ where

$$2y \leq f(y) \leq y, \quad f'(y) \asymp 1.$$

Combining these two facts with (4.2), we have

$$\mathbb{P}[\mathcal{E}_{2j+2}(\epsilon, z, y, r) \mid \eta[0, \xi], \mathcal{F}] \lesssim (C\epsilon|f'(z)|/\delta)^{\alpha_{2j}^+},$$

where the constant in \lesssim is uniform over C, ϵ, δ . Thus, by Lemma 4.2, we have

$$\mathbb{P}[\mathcal{E}_{2j+2}(\epsilon, z, y, r) \cap \mathcal{F}] \lesssim C^A \delta^{-B} \epsilon^{\alpha_{2j+2}},$$

where A, B are some constants depending on κ and j . This completes the proof. \square

LEMMA 4.4. *Fix $\kappa \in (0, 8)$ and let η be an SLE $_{\kappa}$ in \mathbb{H} from 0 to ∞ . Fix $z \in \mathbb{H}$ with $|z| = 1$. Let T_z be the first time that η swallows z and set $\Theta_t = \arg(g_t(z) - W_t)$. Take $n \in \mathbb{N}$ such that $B(z, 16\epsilon 2^n)$ is contained in \mathbb{H} . For $1 \leq m \leq n$, let ξ_m be the first time that η hits $B(z, 16\epsilon 2^{n-m+1})$. Note that ξ_1, \dots, ξ_n is an increasing sequence of stopping times and ξ_1 is the first time that η hits $B(z, 16\epsilon 2^n)$ and ξ_n is the first time that η hits $B(z, 32\epsilon)$. For $1 \leq m \leq n$, for $\delta > 0$, define*

$$\mathcal{F}_m = \{\xi_m < T_z, \Theta_{\xi_m} \notin (\delta, \pi - \delta)\}$$

There exists a function $p : (0, 1) \rightarrow [0, 1]$ with $p(\delta) \downarrow 0$ as $\delta \downarrow 0$ such that

$$\mathbb{P}[\cap_1^n \mathcal{F}_m] \leq p(\delta)^n.$$

PROOF. For $w \in \mathbb{H}$ with $\arg(w) \notin (\delta, \pi - \delta)$, by (6.3), we know that

$$(4.4) \quad \mathbb{P}[\eta \text{ hits } B(w, \text{Im } w)] \leq C\delta^{8/\kappa-1},$$

where C is some universal constant.

For $1 \leq m \leq n$, let $f_m = g_{\xi_m} - W_{\xi_m}$. Note that ξ_m is the first time that η hits $B(z, 16\epsilon 2^{n-m+1})$. We denote $\epsilon 2^{n-m+1}$ by u . By Lemma 2.2, we know that the ball $f_m(B(z, u))$ is contained in the ball with center $f_m(z)$ and radius $4u|f'_m(z)|$, moreover

$$4u|f'_m(z)| \leq \text{Im } f_m(z) \leq 64u|f'_m(z)|.$$

Therefore, by (4.4), we have $\mathbb{P}[\mathcal{F}_{m+4} | \eta[0, \xi_m]] \leq C\delta^{8/\kappa-1}$. Iterating this inequality, we have

$$\mathbb{P}[\cap_1^n \mathcal{F}_m] \leq \left(C\delta^{8/\kappa-1}\right)^{n/4},$$

where C is some universal constant. This implies the conclusion. \square

PROOF OF PROPOSITION 4.1–(4.1)–UPPER BOUND. Assume the same notations as in Lemma 4.4. Recall that $\mathcal{F} = \{\eta[0, \tau_1] \subset B(0, R)\}$. By Lemma 4.3, we have, for $1 \leq m \leq n$

$$\mathbb{P}[\mathcal{E}_{2j+2} \cap \mathcal{F} \cap \mathcal{F}_m^c] \lesssim 2^{nA} \delta^{-B} \epsilon^{\alpha_{2j+2}},$$

where A, B are some constants depending on κ and j . Combining with Lemma 4.4, we have, for any n and $\delta > 0$,

$$\mathbb{P}[\mathcal{E}_{2j+2}(\epsilon, z, y, r) \cap \mathcal{F}] \lesssim n 2^{nA} \delta^{-B} \epsilon^{\alpha_{2j+2}} + p(\delta)^n,$$

where $p(\delta) \downarrow 0$ as $\delta \downarrow 0$. This implies the conclusion. \square

5. Ising Model.

5.1. Definitions. We focus on the square lattice \mathbb{Z}^2 . Two vertices $x = (x_1, x_2)$ and $y = (y_1, y_2)$ are neighbors if $|x_1 - y_1| + |x_2 - y_2| = 1$, and we write $x \sim y$. We denote by $\Lambda_n(x)$ the box centered at x :

$$\Lambda_n(x) = x + [-n, n]^2, \quad \Lambda_n = \Lambda_n(0).$$

Let Ω be a finite subset of \mathbb{Z}^2 , and the edge-set of Ω consists of all edges of \mathbb{Z}^2 that link two vertices of Ω . The boundary of Ω is defined to be $\partial\Omega = \{e = (x, y) : x \sim y, x \in \Omega, y \notin \Omega\}$. We sometimes identify a boundary edge (x, y) with one of its endpoints. Two vertices $x = (x_1, x_2)$ and $y = (y_1, y_2)$ are \star -neighbors if $\max\{|x_1 - y_1|, |x_2 - y_2|\} = 1$. With this definition, each vertex has eight \star -neighbors instead of four.

The Ising model with free boundary conditions is a random assignment $\sigma \in \{\ominus, \oplus\}^\Omega$ of spins $\sigma_x \in \{\ominus, \oplus\}$, where σ_x denotes the spin at the vertex x . The Hamiltonian of the Ising model is defined by $H_\Omega^{\text{free}}(\sigma) = -\sum_{x \sim y} \sigma_x \sigma_y$. The Ising measure is the Boltzmann measure with Hamiltonian H_Ω^{free} and inverse-temperature $\beta > 0$:

$$\mu_{\beta, \Omega}^{\text{free}}[\sigma] = \frac{\exp(-\beta H_\Omega^{\text{free}}(\sigma))}{Z_{\beta, \Omega}^{\text{free}}}, \quad \text{where } Z_{\beta, \Omega}^{\text{free}} = \sum_{\sigma} \exp(-\beta H_\Omega^{\text{free}}(\sigma)).$$

For a graph Ω and $\tau \in \{\ominus, \oplus\}^{\mathbb{Z}^2}$, one may also define the Ising model with boundary conditions τ by the Hamiltonian

$$H_{\Omega}^{\tau}(\sigma) = - \sum_{x \sim y, \{x, y\} \cap \Omega \neq \emptyset} \sigma_x \sigma_y, \quad \text{if } \sigma_x = \tau_x, \forall x \notin \Omega.$$

The Ising model has the following Domain Markov Property: Suppose $\Omega \subset \Omega'$ are two finite subsets of \mathbb{Z}^2 . Let $\tau \in \{\ominus, \oplus\}^{\mathbb{Z}^2}$ and let $\beta > 0$. Suppose X is a random variable measurable with respect to $\{\sigma_x : x \in \Omega\}$. Then we have

$$\mu_{\beta, \Omega'}^{\tau}[X \mid \sigma_x = \tau_x, \forall x \in \Omega' \setminus \Omega] = \mu_{\beta, \Omega}^{\tau}[X].$$

Dobrushin domains are the discrete analogue of simply connected domains with two marked points on their boundary. Suppose $(\Omega; a, b)$ is a Dobrushin domain. Assume that $\partial\Omega$ can be divided into two \star -connected paths from a to b (counterclockwise) and from b to a . Several boundary conditions will be of particular interest in this paper.

- We denote by μ^{free} for free boundary conditions. We denote by μ^{\oplus} (resp. μ^{\ominus}) for the boundary conditions $\tau_x = \oplus$ for all x (resp. $\tau_x = \ominus$ for all x).
- $(\ominus\oplus)$ boundary conditions: \oplus along $\partial\Omega$ from a to b , and \ominus along $\partial\Omega$ from b to a . These boundary conditions are also called Dobrushin boundary conditions, or domain-wall boundary conditions.
- $(\ominus\text{free})$ boundary conditions: free along $\partial\Omega$ from a to b , and \ominus along $\partial\Omega$ from b to a .

The set $\{\ominus, \oplus\}^{\Omega}$ is equipped with a partial order: $\sigma \leq \sigma'$ if $\sigma_x \leq \sigma'_x$ for all $x \in \Omega$. A random variable X is increasing if $\sigma \leq \sigma'$ implies $X(\sigma) \leq X(\sigma')$. An event \mathcal{A} is increasing if $\mathbb{1}_{\mathcal{A}}$ is increasing. The following inequality is the FKG inequality for the Ising model: Let Ω be a finite subset, let τ be the boundary conditions, and let $\beta > 0$. For any two increasing events \mathcal{A} and \mathcal{B} , we have

$$\mu_{\beta, \Omega}^{\tau}[\mathcal{A} \cap \mathcal{B}] \geq \mu_{\beta, \Omega}^{\tau}[\mathcal{A}] \mu_{\beta, \Omega}^{\tau}[\mathcal{B}].$$

As a consequence of the FKG inequality, we have the following comparison between boundary conditions: For boundary conditions $\tau_1 \leq \tau_2$ and an increasing event \mathcal{A} , we have

$$(5.1) \quad \mu_{\beta, \Omega}^{\tau_1}[\mathcal{A}] \leq \mu_{\beta, \Omega}^{\tau_2}[\mathcal{A}].$$

The Ising model with inverse-temperature $\beta > 0$ is related to the random-cluster model with parameters $(p, 2)$ through the Edwards-Sokal coupling, thus the critical value $p_c(2)$ for the random-cluster model gives the critical

value of $\beta : \beta_c = (1/2) \log(1 + \sqrt{2})$. We focus on the critical Ising model on the square lattice and derive the arm exponents. To this end, we need three inputs:

- The convergence of the scaling limit of the interface in the critical Ising model. This is proved in [CDCH⁺14, HK13], see Theorems 5.7 and 5.8.
- The arm exponents of SLE₃. This is the topic of Sections 3 and 4.
- A stronger version of Russo-Seymour-Welsh inequality for the critical Ising model. This is proved in [CDCH16], see Proposition 5.1. We can deduce the so-called quasi-multiplicativity from the RSW inequality.

With these three inputs at hand, we can apply the same strategy as in [SW01] where the authors derived the arm exponents of the critical percolation. The content in this section is not new, and we just summarize the known results and explain how to put them together to get the arm exponents of the critical Ising model.

5.2. *Quasi-Multiplicativity.* In this section, we first introduce a stronger version of RSW inequality—Proposition 5.1—for the critical Ising model and then define quasi-multiplicativity for the model. The quasi-multiplicativity is a consequence of the RSW inequality and, roughly speaking, it guarantees that we can use the crossing events of SLE₃ to approximate the crossing events of the Ising model.

A discrete topological rectangle $(\Omega; a, b, c, d)$ is a bounded simply-connected subdomain of \mathbb{Z}^2 with four marked boundary points. The four points are in counterclockwise order and (ab) denotes the arc of $\partial\Omega$ from a to b . We denote by $d_\Omega((ab), (cd))$ the discrete external distance between (ab) and (cd) in Ω , see [Che16, Section 6]. The discrete extremal distance is uniformly comparable to and converges to its continuous counterpart—the classical extremal distance. The rectangle $(\Omega; a, b, c, d)$ is crossed by \oplus in an Ising configuration σ if there exists a path of \oplus going from (ab) to (cd) in Ω . We denote this event by $(ab) \xleftrightarrow{\oplus} (cd)$. We have the following RSW-type estimate on the crossing probability at critical.

PROPOSITION 5.1. [CDCH16, Corollary 1.7] *For each $L > 0$ there exists $c(L) > 0$ such that the following holds: for any topological rectangle $(\Omega; a, b, c, d)$ with $d_\Omega((ab), (cd)) \leq L$,*

$$\mu_{\beta_c, \Omega}^{\text{mixed}} \left[(ab) \xleftrightarrow{\oplus} (cd) \right] \geq c(L),$$

where the boundary conditions are free on $(ab) \cup (cd)$ and \ominus on $(bc) \cup (da)$.

As a consequence of Proposition 5.1, we have the following spatial mixing property at criticality.

COROLLARY 5.2. *There exists $\alpha > 0$ such that for any $2k \leq n$, for any event \mathcal{A} depending only on edges in Λ_k , and for any boundary conditions τ, ξ , we have*

$$|\mu_{\beta_c, \Lambda_n}^\tau[\mathcal{A}] - \mu_{\beta_c, \Lambda_n}^\xi[\mathcal{A}]| \leq \left(\frac{k}{n}\right)^\alpha \mu_{\beta_c, \Lambda_n}^\tau[\mathcal{A}].$$

In particular, this implies that, for any boundary conditions τ , for any $2k \leq n \leq m$, for any event \mathcal{A} depending only on $\{\sigma_x, x \in \Lambda_k\}$, and for any event \mathcal{B} depending only on $\{\sigma_x, x \in \Lambda_m \setminus \Lambda_n\}$, we have

$$|\mu_{\beta_c, \Lambda_m}^\tau[\mathcal{A} \cap \mathcal{B}] - \mu_{\beta_c, \Lambda_m}^\tau[\mathcal{A}] \mu_{\beta_c, \Lambda_m}^\tau[\mathcal{B}]| \leq \left(\frac{k}{n}\right)^\alpha \mu_{\beta_c, \Lambda_m}^\tau[\mathcal{A}] \mu_{\beta_c, \Lambda_m}^\tau[\mathcal{B}].$$

Fix $n < N$ and consider the annulus $\Lambda_N \setminus \Lambda_n$. A simple path of \oplus or of \ominus connecting $\partial\Lambda_n$ to $\partial\Lambda_N$ is called an *arm*. Fix an integer $j \geq 1$ and $\omega = (\omega_1, \dots, \omega_j) \in \{\ominus, \oplus\}^j$. For $n < N$, define $\mathcal{A}_\omega(n, N)$ to be the event that there are j disjoint arms $(\gamma_k)_{1 \leq k \leq j}$ connecting $\partial\Lambda_n$ to $\partial\Lambda_N$ in the annulus $\Lambda_N \setminus \Lambda_n$ which are of types $(\omega_k)_{1 \leq k \leq j}$, where we identify two sequences ω and ω' if they are the same up to cyclic permutation and the arms are indexed in clockwise order. For each $j \geq 1$, there exists a smallest integer $n_0(j)$ such that, for all $N \geq n_0(j)$, we have $\mathcal{A}_\omega(n_0(j), N) \neq \emptyset$.

PROPOSITION 5.3. *Assume that ω is alternating with even length. For all $n_0(j) \leq n_1 < n_2 < n_3 \leq m/2$, and for all boundary conditions τ , we have*

$$\mu_{\beta_c, \Lambda_m}^\tau[\mathcal{A}_\omega(n_1, n_3)] \asymp \mu_{\beta_c, \Lambda_m}^\tau[\mathcal{A}_\omega(n_1, n_2)] \mu_{\beta_c, \Lambda_m}^\tau[\mathcal{A}_\omega(n_2, n_3)],$$

where the constants in \asymp are uniform over n_1, n_2, n_3, m and τ .

Proposition 5.3 is called the quasi-multiplicativity. We will introduce several auxiliary subevents of $\mathcal{A}_\omega(n, N)$ which are both important for the proof of Proposition 5.3 and also important for us to derive the arm exponents of the Ising model. Fix $\omega = (\omega_1, \dots, \omega_j) \in \{\ominus, \oplus\}^j$. Fix some $\delta > 0$ small. Suppose $Q = [-1, 1]^2$ is the unit square. A landing sequence $(I_k)_{1 \leq k \leq j}$ is a sequence of disjoint sub-intervals on ∂Q in clockwise order. We denote by $z(I_k)$ the center of I_k . We say $(I_k)_{1 \leq k \leq j}$ is δ -separated if

- the intervals are at distance at least 2δ from each other, and they are at distance at least 2δ from the four corners of ∂Q

- for each I_k , the length of I_k is at least 2δ .

We say that two sets are ω_k -connected if there is a path of type ω_k connecting them. Fix two δ -separated landing sequences $(I_k)_{1 \leq k \leq j}$ and $(I'_k)_{1 \leq k \leq j}$. We say that the arms $(\gamma_k)_{1 \leq k \leq j}$ are δ -well-separated with landing sequence $(I_k)_{1 \leq k \leq j}$ on $\partial\Lambda_n$ and landing sequence $(I'_k)_{1 \leq k \leq j}$ on $\partial\Lambda_N$ if

- for each k , the arm γ_k connects nI_k to NI'_k ;
- for each k , the arm γ_k can be ω_k -connected to distance δn of $\partial\Lambda_n$ inside $\Lambda_{\delta n}(z(I_k))$;
- for each k , the arm γ_k can be ω_k -connected to distance δN of $\partial\Lambda_N$ inside $\Lambda_{\delta N}(z(I'_k))$.

We denote this event by $\mathcal{A}_\omega^{I/I'}(n, N)$.

LEMMA 5.4. *Fix $j \geq 1$ and $\delta > 0$ and two δ -separated landing sequences $(I_k)_{1 \leq k \leq j}$ and $(I'_k)_{1 \leq k \leq j}$. Assume that ω is alternating with length $2j$. For all $n < N \leq m/2$ such that $\mathcal{A}_\omega^{I/I'}(n, N)$ is not empty, and for all boundary conditions τ , we have*

$$\mu_{\beta_c, \Lambda_m}^\tau \left[\mathcal{A}_\omega^{I/I'}(n, N) \right] \asymp \mu_{\beta_c, \Lambda_m}^\tau \left[\mathcal{A}_\omega(n, N) \right],$$

where the constants in \asymp depend only on δ .

We have similar results for the boundary arm events. Denote by

$$\Lambda_n^+(x) = [-n, n] \times [0, n] + x, \quad \Lambda_n^+ = \Lambda_n^+(0).$$

We consider the arm events in the semi-annulus $\Lambda_N^+ \setminus \Lambda_n^+$ and extend the definition of arm events and arm events with landing sequences in the obvious way, and denote them by $\mathcal{A}_\omega^+(n, N)$ and $\mathcal{A}_\omega^{+, I/I}(n, N)$.

We need to restrict to the cases that the arms together with the boundary conditions are alternating. Precisely, in the statements of Proposition 5.5 and Lemma 5.6, we restrict to the cases where the arm patterns and the boundary conditions are listed in Theorem 1.1.

PROPOSITION 5.5. *For all $n_0^+(j) \leq n_1 < n_2 < n_3 \leq m/2$, we have*

$$\mu_{\beta_c, \Lambda_m^+}^\tau \left[\mathcal{A}_\omega^+(n_1, n_3) \right] \asymp \mu_{\beta_c, \Lambda_m^+}^\tau \left[\mathcal{A}_\omega^+(n_1, n_2) \right] \mu_{\beta_c, \Lambda_m^+}^\tau \left[\mathcal{A}_\omega^+(n_2, n_3) \right],$$

where the constants in \asymp are uniform over n_1, n_2, n_3 and m .

LEMMA 5.6. Fix $j \geq 1$, $\delta > 0$ and two δ -separated landing sequences $(I_k)_{1 \leq k \leq j}$ and $(I'_k)_{1 \leq k \leq j}$. For all $n < N \leq m/2$ such that $\mathcal{A}_\omega^{+,I/I'}(n, N)$ is not empty, we have


$$\mu_{\beta_c, \Lambda_m^+}^\tau \left[\mathcal{A}_\omega^{+,I/I'}(n, N) \right] \asymp \mu_{\beta_c, \Lambda_m^+}^\tau \left[\mathcal{A}_\omega^+(n, N) \right],$$

where the constants in \asymp depend only on δ .

We do not give the proof of the quasi-multiplicativity in this paper, because the proof is exactly the same as the proof of the quasi-multiplicativity for FK-Ising model proved in [CDCH16].

5.3. *Proof of Theorems 1.1 and 1.2.* In this section, we first define the interfaces of the Ising model; then explain the convergence results on the interfaces; and finally, complete the proof of Theorems 1.1 and 1.2.

The *dual square lattice* $(\mathbb{Z}^2)^*$ is the dual graph of \mathbb{Z}^2 . The vertex set is $(1/2, 1/2) + \mathbb{Z}^2$ and the edges are given by nearest neighbors. The vertices and edges of $(\mathbb{Z}^2)^*$ are called dual-vertices and dual-edges. In particular, for each edge e of \mathbb{Z}^2 , it is associated to a dual edge, denoted by e^* . The dual edge e^* crosses e in the middle. For a finite subgraph G , we define G^* to be the subgraph of $(\mathbb{Z}^2)^*$ with edge-set $E(G^*) = \{e^* : e \in E(G)\}$ and vertex set given by the end-points of these dual-edges. The *medial lattice* $(\mathbb{Z}^2)^\diamond$ is the graph with the centers of edges of \mathbb{Z}^2 as vertex set, and edges connecting nearest vertices. This lattice is a rotated and rescaled version of \mathbb{Z}^2 , see Fig 5.1. The vertices and edges of $(\mathbb{Z}^2)^\diamond$ are called medial-vertices and medial-edges. We identify the faces of $(\mathbb{Z}^2)^\diamond$ with the vertices of \mathbb{Z}^2 and $(\mathbb{Z}^2)^*$. A face of $(\mathbb{Z}^2)^\diamond$ is said to be black if it corresponds to a vertex of \mathbb{Z}^2 and white if it corresponds to a vertex of $(\mathbb{Z}^2)^*$.



(a) The square lattice.

(b) The dual lattice.

(c) The medial lattice.

Fig 5.1: The lattices.




Fig 5.2: The Ising interface.

For $u > 0$, we consider the rescaled square lattice $u\mathbb{Z}^2$. The definitions of dual lattice, medial lattice and Dobrushin domains extend to this context, and they will be denoted by $(\Omega_u; a_u, b_u)$, $(\Omega_u^*; a_u^*, b_u^*)$, $(\Omega_u^\diamond; a_u^\diamond, b_u^\diamond)$ respectively. Consider the critical Ising model on $(\Omega_u^*; a_u^*, b_u^*)$. The boundary $\partial\Omega_u^*$ is divided into two parts $(a_u^*b_u^*)$ and $(b_u^*a_u^*)$. We fix the boundary conditions to be \ominus on $(b_u^*a_u^*)$ and \oplus on $(a_u^*b_u^*)$, or \ominus on $(b_u^*a_u^*)$ and free on $(a_u^*b_u^*)$. Define the *interface* as follows. It starts from a_u^\diamond , lies on the primal lattice and turns at every vertex of Ω_u is such a way that it has always dual vertices with spin \ominus on its left and \oplus on its right. If there is an indetermination when arriving at a vertex (this may happen on the square lattice), turn left. See Fig 5.2.

Let $(\Omega; a, b)$ be a simply connected domain with two marked points on its boundary. Consider a sequence of Dobrushin domains $(\Omega_u; a_u, b_u)$. We say that $(\Omega_u; a_u, b_u)$ converges to $(\Omega; a, b)$ in the *Carathéodory sense* if $f_u \rightarrow f$ on any compact subset $K \subset \mathbb{H}$, where f_u (resp. f) is the unique conformal map from \mathbb{H} to Ω_u (resp. Ω) satisfying $f_u(0) = a_u, f_u(\infty) = b_u$ and $f'_u(\infty) = 1$ (resp. $f(0) = a, f(\infty) = b, f'(\infty) = 1$).

Let X be the set of continuous parameterized curves and d be the distance on X defined for $\eta_1 : I \rightarrow \mathbb{C}$ and $\eta_2 : J \rightarrow \mathbb{C}$ by


$$d(\eta_1, \eta_2) = \inf_{\varphi_1: [0,1] \rightarrow I, \varphi_2: [0,1] \rightarrow J} \sup_{t \in [0,1]} |\eta_1(\varphi_1(t)) - \eta_2(\varphi_2(t))|,$$

where the infimum is over increasing bijective functions φ_1, φ_2 . Note that I and J can be equal to $\mathbb{R}_+ \cup \{\infty\}$. The topology on (X, d) gives rise to a notion of weak convergence for random curves on X .

THEOREM 5.7. [CDCH⁺14]. Let $(\Omega_u^\diamond; a_u^\diamond, b_u^\diamond)$ be a family of Dobrushin domains converging to a Dobrushin domain $(\Omega; a, b)$ in the Carathéodory sense. The interface of the critical Ising model in $(\Omega_u^*; a_u^*, b_u^*)$ with $(\ominus\oplus)$ boundary conditions converges weakly to SLE_3 as $u \rightarrow 0$.

THEOREM 5.8. Let $(\Omega_u^\diamond; a_u^\diamond, b_u^\diamond)$ be a family of Dobrushin domains converging to a Dobrushin domain $(\Omega; a, b)$ in the Carathéodory sense. The interface of the critical Ising model in $(\Omega_u^*; a_u^*, b_u^*)$ with $(\ominus \text{free})$ boundary conditions converges weakly to $\text{SLE}_3(-3/2)$ as $u \rightarrow 0$.

PROOF. It is proved in [HK13, BDCH16] that the interface with (free free) boundary conditions converges weakly to $\text{SLE}_3(-3/2; -3/2)$ as $u \rightarrow 0$. The same proof works here. \square



(a) $\mathcal{A}^{I/I}(n, N)$ is the well-separated arm event. (b) The four gray parts are R_1 to R_4 respectively.

Fig 5.3: The explanation of the proof of Theorem 1.2.

PROOF OF THEOREM 1.2. We only give the proof for α_4 and the other cases can be proved similarly. Consider Λ_m with two boundary points $a_m = (-m, 0)$ and $b_m = (m, 0)$. Fix $(\ominus\oplus)$ boundary conditions: the vertices along $\partial\Lambda_m$ from b_m to a_m (counterclockwise) are \oplus and the vertices from a_m to b_m are \ominus . Since we fix $\beta = \beta_c$ and the boundary conditions, and $\omega = (\oplus\ominus\oplus\ominus)$, we eliminate them from the notations. We will prove that, for $n < N \leq m/2$,

$$(5.2) \quad \mu_{\Lambda_m}[\mathcal{A}(n, N)] = N^{-\alpha_4 + o(1)}, \quad \text{as } N \rightarrow \infty.$$

Fix the landing sequence $I = (I_1, I_2, I_3, I_4)$ where $I_1 = [-1/2, 1/2] \times \{-1\}$, $I_2 = \{-1\} \times [-1/2, 1/2]$, $I_3 = [-1/2, 1/2] \times \{1\}$, and $I_4 = \{1\} \times [-1/2, 1/2]$. Recall that $\mathcal{A}^{I/I}(n, N)$ is the 1/8-well-separated arm event with the landing sequence nI on $\partial\Lambda_n$ and NI on $\partial\Lambda_N$. The four arms in $\mathcal{A}(n, N)$ are denoted by $(\gamma_1, \gamma_2, \gamma_3, \gamma_4)$ where γ_1 and γ_3 are \oplus and γ_2 and γ_4 are \ominus . Consider the critical Ising model in Λ_{2N} . Let R_1 to be the rectangle $[-3N/4, 3N/4] \times [-2N, -N]$, and define \mathcal{C}_1^\oplus to be the event that γ_1 is connected by path of \oplus in R_1 to the bottom of R_1 . Let R_2 to be the rectangle $[-9N/8, -N] \times [-N/2, 2N]$, and define \mathcal{C}_2^\ominus to be the event that γ_2 is connected by path of \ominus in R_2 to the top of R_2 . Let R_3 be the rectangle $[-3n/4, 3n/4] \times [-n, n]$, and define \mathcal{C}_3^\oplus to be the event that γ_3 is connected to γ_1 by path of \oplus in R_3 . Let R_4 be the rectangle $[N, 9N/8] \times [-N/2, 2N]$, and define \mathcal{C}_4^\ominus to be the event that γ_4 is connected by path of \ominus in R_4 to the top of R_4 . See Fig 5.3.

By (5.1), Proposition 5.1 and Corollary 5.2, we deduce

$$(5.3) \quad \mu_{\Lambda_{2N}} \left[\mathcal{A}^{I/I}(n, N) \right] \asymp \mu_{\Lambda_{2N}} \left[\mathcal{A}^{I/I}(n, N) \cap \mathcal{C}_1^\oplus \cap \mathcal{C}_2^\ominus \cap \mathcal{C}_3^\oplus \cap \mathcal{C}_4^\ominus \right],$$

where the constants in \asymp are uniform over n, N .

Let \mathbb{P}_N be the probability measure $\mu_{\Lambda_{2N}}$ where the square lattice is scaled by $1/N$ and let \mathbb{P}_∞ be the law of SLE₃ in $[-2, 2] \times [-2, 2]$ from $(-2, 0)$ to $(2, 0)$. On the event $\mathcal{A}^{I/I}(n, N) \cap \mathcal{C}_1^\oplus \cap \mathcal{C}_2^\ominus \cap \mathcal{C}_3^\oplus \cap \mathcal{C}_4^\ominus$, consider the interface η from a_{2N} to b_{2N} . Let τ_1 be the first time that η hits $\partial\Lambda_n$. The event $\mathcal{C}_1^\oplus \cap \mathcal{C}_2^\ominus$ guarantees that $\eta[0, \tau_1]$ is bounded away from the target b_{2N} . The event \mathcal{C}_3^\oplus guarantees that, after τ_1 , the path η hits the neighborhood of $(0, 2N)$ at some time σ_1 . The event \mathcal{C}_4^\ominus guarantees that, after σ_1 , the path η hits $\partial\Lambda_n$ again. Therefore, by (4.1), we have, for $\epsilon > 0$,

$$\begin{aligned} & \limsup_{N \rightarrow \infty} \mathbb{P}_N \left[\mathcal{A}^{I/I}(\epsilon N, N) \cap \mathcal{C}_1^\oplus \cap \mathcal{C}_2^\ominus \cap \mathcal{C}_3^\oplus \cap \mathcal{C}_4^\ominus \right] \\ & \leq \epsilon^{\alpha_4 + o(1)} \leq \liminf_{N \rightarrow \infty} \mathbb{P}_N[\mathcal{A}(\epsilon N, N)]. \end{aligned}$$

Combining with Lemma 5.4 and (5.3), we have

$$\liminf_{N \rightarrow \infty} \mathbb{P}_N[\mathcal{A}(\epsilon N, N)] \asymp \limsup_{N \rightarrow \infty} \mathbb{P}_N[\mathcal{A}(\epsilon N, N)] \asymp \epsilon^{\alpha_4 + o(1)}.$$

By Corollary 5.2, we know that

$$(5.4) \quad \liminf_{N \rightarrow \infty} \mu_{\Lambda_m}[\mathcal{A}(\epsilon N, N)] \asymp \limsup_{N \rightarrow \infty} \mu_{\Lambda_m}[\mathcal{A}(\epsilon N, N)] \asymp \epsilon^{\alpha_4 + o(1)},$$

where the constants in \asymp are uniform over ϵ and $m \geq 2N$.

Suppose $N = n\epsilon^{-K}$ for some integer K . By Proposition 5.3, for $m \geq 2N$, we have

$$\mu_{\Lambda_m}[\mathcal{A}(n, N)] \leq C^K \prod_{j=1}^K \mu_{\Lambda_m}[\mathcal{A}(n\epsilon^{-j+1}, n\epsilon^{-j})],$$

where C is some universal constant. Thus

$$\frac{\log \mu_{\Lambda_m}[\mathcal{A}(n, N)]}{\log N} \leq \frac{K \log C}{\log N} + \frac{1}{\log N} \sum_{j=1}^K \log \mu_{\Lambda_m}[\mathcal{A}(n\epsilon^{-j+1}, n\epsilon^{-j})].$$

By (5.4), we have

$$\limsup_{j \rightarrow \infty} \mu_{\Lambda_m}[\mathcal{A}(n\epsilon^{-j+1}, n\epsilon^{-j})] \asymp \epsilon^{\alpha_4 + o(1)}.$$

Therefore,

$$\limsup_{K \rightarrow \infty} \frac{\log \mu_{\Lambda_m}[\mathcal{A}(n, N)]}{\log N} \leq \frac{\tilde{C}}{\log(1/\epsilon)} - \alpha_4,$$

where \tilde{C} is some universal constant. Let $\epsilon \rightarrow 0$, we have

$$\limsup_{N \rightarrow \infty} \frac{\log \mu_{\Lambda_m}[\mathcal{A}(n, N)]}{\log N} \leq -\alpha_4.$$

We deduce the lower bound similarly:


$$\liminf_{N \rightarrow \infty} \frac{\log \mu_{\Lambda_m}[\mathcal{A}(n, N)]}{\log N} \geq -\alpha_4.$$

These imply (5.2) and complete the proof. \square

To prove Theorem 1.1, we will show the proof for γ_{2j-1}^+ , and the results for $\alpha_{2j-1}^+, \gamma_{2j}^+$ can be proved similarly; and we will show the proof for β_{2j}^+ , and the results for $\alpha_{2j}^+, \beta_{2j-1}^+$ can be proved similarly.

PROOF OF THEOREM 1.1–(1.5). We will prove the conclusion for γ_3^+ and the other cases can be proved similarly. Consider Λ_m^+ with two boundary points $a_m = (-m, m/2)$ and $b_m = (m, m/2)$. Fix $(\ominus \text{ free})$ boundary conditions: the vertices along $\partial\Lambda_m$ from b_m to a_m (counterclockwise) are free and the vertices from a_m to b_m are \ominus . Since we fix $\beta = \beta_c$ and the boundary conditions, and $\omega = (\ominus \oplus \ominus)$, we eliminate them from the notations. We will prove that, for $n < N \leq m/2$,

$$(5.5) \quad \mu_{\Lambda_m^+}[\mathcal{A}^+(n, N)] = N^{-\gamma_3^+ + o(1)}, \quad \text{as } N \rightarrow \infty.$$



(a) $\mathcal{A}^{+,I/I}(n, N)$ is the well-separated arm event. (b) The four gray parts are R_1 to R_4 respectively.

Fig 5.4: The explanation of the proof of (1.5).

Fix the landing sequence $I = (I_1, I_2, I_3)$ where $I_1 = \{-1\} \times [1/2, 3/4]$, $I_2 = [-1/2, 1/2] \times \{1\}$, and $I_3 = \{1\} \times [1/2, 3/4]$. Recall that $\mathcal{A}^{+,I/I}(n, N)$ is the $1/8$ -well-separated arm event with the landing sequence nI on $\partial\Lambda_n^+$ and NI on $\partial\Lambda_N^+$. The three arms in $\mathcal{A}^{+,I/I}(n, N)$ are denoted by $(\gamma_1, \gamma_2, \gamma_3)$ where γ_1 and γ_3 are \ominus and γ_2 is \oplus . Consider the critical Ising model in Λ_{2N}^+ . Let R_1 be the rectangle $[-9N/8, -N] \times [N/2, N]$ and define \mathcal{C}_1^\ominus to be the event that γ_1 is connected by path of \ominus in R_1 to the top of R_1 . Let R_2 be the rectangle $[-3n/4, 3n/4] \times [0, n]$ and define \mathcal{C}_2^\oplus to be the event that γ_2 is connected by path of \oplus in R_2 to the bottom of R_2 . Let R_3 be the rectangle $[N, 9N/8] \times [N/2, N]$ and define \mathcal{C}_3^\ominus to be the event that γ_3 is connected by path of \ominus in R_3 to the top of R_3 . For $\delta > 0$, let R_4 be the semi-annulus $[3n/4, 4n] \times [0, n/4] \setminus [n, 3n] \times [0, \delta n]$ and define $\mathcal{C}_4^\oplus(\delta)$ to be the event that there is a path of \oplus in R_4 connecting the left bottom to the right bottom.

By (5.1), Proposition 5.1 and Corollary 5.2, we can prove, for $\delta > 0$ small enough,

$$(5.6) \quad \mu_{\Lambda_{2N}} \left[\mathcal{A}^{+,I/I}(n, N) \right] \asymp \mu_{\Lambda_{2N}} \left[\mathcal{A}^{+,I/I}(n, N) \cap \mathcal{C}_1^\ominus \cap \mathcal{C}_2^\oplus \cap \mathcal{C}_3^\ominus \cap \mathcal{C}_4^\oplus(\delta) \right],$$


where the constants in \asymp are uniform over n, N .

Let \mathbb{P}_N be the probability measure $\mu_{\Lambda_{2N}}$ where the square lattice is scaled by $1/N$ and let \mathbb{P}_∞ be the law of $\text{SLE}_3(-3/2)$ in $[-2, 2] \times [0, 2]$ from $(-2, 1)$ to $(2, 1)$. On the event $\mathcal{A}^{+,I/I}(n, N) \cap \mathcal{C}_1^\ominus \cap \mathcal{C}_2^\oplus \cap \mathcal{C}_3^\ominus \cap \mathcal{C}_4^\oplus(\delta)$, consider the interface η from a_{2N} to b_{2N} . Let τ_1 be the first time that η hits $\partial\Lambda_n$. The event \mathcal{C}_1^\ominus guarantees that $\eta[0, \tau_1]$ is bounded away from the target b_{2N} . The event $\mathcal{C}_4^\oplus(\delta)$ guarantees that $\eta[0, \tau_1]$ is bounded away from the segment $[n, 3n]$. The event \mathcal{C}_2^\oplus guarantees that, after τ_1 , the interface η hits the neighborhood of the point $(0, N)$ at some time σ_1 . The event \mathcal{C}_3^\ominus guarantees that, after σ_1 ,

the interface η hits $\partial\Lambda_n$ again. See Fig 5.4. Therefore, by (3.11), we have, for $\epsilon > 0$,

$$\begin{aligned} \limsup_{N \rightarrow \infty} \mathbb{P}_N \left[\mathcal{A}^{+,I/I}(\epsilon N, N) \cap \mathcal{C}_1^\ominus \cap \mathcal{C}_2^\oplus \cap \mathcal{C}_3^\ominus \cap \mathcal{C}_4^\oplus(\delta) \right] \\ \lesssim \epsilon \gamma_3^+ \leq \liminf_{N \rightarrow \infty} \mathbb{P}_N \left[\mathcal{A}^+(\epsilon N, N) \right]. \end{aligned}$$

Now we can repeat the same proof of Theorem 1.2 to obtain (5.5). \square



(a) $\mathcal{A}^{+,I/I}(n, N)$ is the well-separated arm event. (b) The two gray parts are R_1 and R_2 respectively.

Fig 5.5: The explanation of the proof of (1.4).

PROOF OF THEOREM 1.1–(1.4). We will prove the conclusion for β_2^+ and the other cases can be proved similarly. Consider Λ_m^+ with two boundary points $a_m = (0, 0)$ and $b_m = (0, m)$. Fix $(\ominus \text{ free})$ boundary conditions: the vertices along $\partial\Lambda_m$ from b_m to a_m (counterclockwise) are free and the vertices from a_m to b_m are \ominus . Since we fix $\beta = \beta_c$ and the boundary conditions, and $\omega = (\oplus \ominus)$, we eliminate them from the notations. We will prove that, for $n < N \leq m/2$,

$$(5.7) \quad \mu_{\Lambda_m^+}[\mathcal{A}^+(n, N)] = N^{-\beta_2^+ + o(1)}, \quad \text{as } N \rightarrow \infty.$$

Fix the landing sequence $I = (I_1, I_2)$ where $I_1 = \{-1\} \times [1/2, 3/4]$ and $I_2 = [-1/2, 1/2] \times \{1\}$. Recall that $\mathcal{A}^{+,I/I}(n, N)$ is the $1/8$ -well-separated arm event with the landing sequence nI on $\partial\Lambda_n^+$ and NI on $\partial\Lambda_N^+$. The two arms in $\mathcal{A}^{+,I/I}(n, N)$ are denoted by (γ_1, γ_2) where γ_1 is \oplus and γ_2 is \ominus . Consider the critical Ising model in Λ_{2N}^+ . Let R_1 be the tube $[-n, 3n/4] \times [0, 3n/4] \setminus [-n, n/2] \times [0, n/4]$ and define \mathcal{C}_1^\oplus to be the event that γ_1 is connected by path of \oplus in R_1 to the bottom of R_1 . Let R_2 be the rectangle

$[-N, N/2] \times [N/2, 5N/8]$ and define \mathcal{C}_2^\ominus to be the event that γ_2 is connected by path of \ominus in R_2 to the left side of R_2 .

By (5.1), Proposition 5.1 and Corollary 5.2, we can prove, for $\delta > 0$ small enough,

$$(5.8) \quad \mu_{\Lambda_{2N}} \left[\mathcal{A}^{+,I/I}(n, N) \right] \asymp \mu_{\Lambda_{2N}} \left[\mathcal{A}^{+,I/I}(n, N) \cap \mathcal{C}_1^\oplus \cap \mathcal{C}_2^\ominus \right],$$

where the constants in \asymp are uniform over n, N .

Let \mathbb{P}_N be the probability measure $\mu_{\Lambda_{2N}}$ where the square lattice is scaled by $1/N$ and let \mathbb{P}_∞ be the law of $\text{SLE}_3(-3/2)$ in $[-2, 2] \times [0, 2]$ from $(0, 0)$ to $(0, 1)$. On the event $\mathcal{A}^{+,I/I}(n, N) \cap \mathcal{C}_1^\oplus \cap \mathcal{C}_2^\ominus$, consider the interface η from a_{2N} to b_{2N} , the event guarantees that the interface hits the neighborhood of the point $(-N, N/2)$, and then comes back to Λ_n^+ . See Fig 5.5. Therefore, by (3.4) and (3.6) (taking $\rho = -3/2$), we have, for $\epsilon > 0$,

$$\limsup_{N \rightarrow \infty} \mathbb{P}_N \left[\mathcal{A}^{+,I/I}(n, N) \cap \mathcal{C}_1^\oplus \cap \mathcal{C}_2^\ominus \right] \lesssim \epsilon^{\beta_2^+} \leq \liminf_{N \rightarrow \infty} \mathbb{P}_N \left[\mathcal{A}^+(\epsilon N, N) \right].$$

Now we can repeat the same proof of Theorem 1.2 to obtain (5.7). \square

6. Appendix: One-Point Estimate of the Intersection of SLE with the Boundary.

THEOREM 6.1. *Fix $\kappa > 0, \rho^L, \rho^{1,R}, \rho^{2,R} \in \mathbb{R}$ such that*

$$(6.1) \quad \rho^L > -2, \quad \rho^{1,R} > (-2) \vee (\kappa/2 - 4), \quad \rho^{1,R} + \rho^{2,R} > \kappa/2 - 4.$$

Suppose η is an $\text{SLE}_\kappa(\rho^L; \rho^{1,R}, \rho^{2,R})$ process with force points $(x^L; x^R, 1)$ where $x^L \leq 0$ and $x^R \in [0, 1)$. Define

$$\alpha = (\rho^{1,R} + 2)(\rho^{1,R} + \rho^{2,R} + 4 - \kappa/2)/\kappa, \quad \beta = 2(\rho^{1,R} + \rho^{2,R} + 4 - \kappa/2)/\kappa.$$

For $\epsilon \in (0, 1/2)$ and $r \geq 4$, define $\tau_\epsilon = \inf\{t : \eta(t) \in \partial B(1, \epsilon(1 - x^R))\}$, and $S_r = \inf\{t : \eta(t) \in \partial B(0, r)\}$. Then we have

$$\epsilon^\alpha (1 - x^R)^\beta \lesssim \mathbb{P}[\tau_\epsilon \leq S_r] = \epsilon^{\alpha + o(1)},$$

where the constant in \lesssim is uniform and the $o(1)$ term goes to zero as $\epsilon \rightarrow 0$ at a rate which depends only on x^R and r .

COROLLARY 6.2. *Fix $\kappa > 0, \rho^L, \rho^{1,R}, \rho^{2,R} \in \mathbb{R}$ such that*

$$(6.2) \quad \rho^L \in (-2, 0], \quad \rho^{1,R} > (-2) \vee (\kappa/2 - 4), \quad \rho^{1,R} + \rho^{2,R} > \kappa/2 - 4.$$

Assume the same notations as in Theorem 6.1. Then we have

$$\epsilon^\alpha (1 - x^R)^\beta \lesssim \mathbb{P}[\tau_\epsilon < \infty] = \epsilon^{\alpha+o(1)},$$

where the constant in \lesssim is uniform and the $o(1)$ term goes to zero as $\epsilon \rightarrow 0$ at a rate which depends only on x^R .

We also expect that Corollary 6.2 holds for all $\rho^L > -2$, but we do not have a proof yet for $\rho^L \geq 0$. Before proving the theorem, we first summarize the existing related results. For standard SLE_κ with $\kappa \in (0, 8)$, a stronger conclusion is known [AK08]:

$$(6.3) \quad \mathbb{P}[\eta \text{ hits } B(1, \epsilon)] \asymp \epsilon^\alpha, \quad \text{where } \alpha = (8 - \kappa)/\kappa.$$

For $\text{SLE}_\kappa(\rho)$ with one force point at $x^R \in [0, 1)$, a stronger conclusion is known [Law15, Proposition 5.4]:

$$\mathbb{P}[\tau_\epsilon < \infty] \asymp \epsilon^\alpha (1 - x^R)^\beta,$$

where $\alpha = (\rho+2)(\rho+4-\kappa/2)/\kappa$, and $\beta = 2(\rho+4-\kappa/2)/\kappa$. For $\text{SLE}_\kappa(\rho^{1,R}, \rho^{2,R})$ processes, the conclusion in Theorem 6.1 is proved in [MW17, Theorem 3.1]. The proof here is simpler than the one in [MW17].

LEMMA 6.3. *Assume the same notation as in Theorem 6.1. For $\delta \in (0, 1/4)$ and $r \geq 4$, we have*

$$\epsilon^\alpha (1 - x^R)^\beta \lesssim \mathbb{P}[\tau_\epsilon \leq S_r, \text{Im } \eta(\tau_\epsilon) \geq \delta\epsilon(1 - x^R)] \lesssim \epsilon^\alpha (1 - x^R)^\beta \delta^{-\beta} r^B,$$

where $B = 0 \vee (\beta\rho^L/2)$ and the constants in \lesssim are uniform over $\epsilon, \delta, x^L, x^R, r$.

PROOF. Let V_t^L be the evolution of x^L and V_t^R be the evolution of x^R . Set $\nu = -\beta\kappa \leq 0$ and

$$\begin{aligned} M_t &= g'_t(1)^{\nu(\nu+2\rho^{2,R}+4-\kappa)/(4\kappa)} (g_t(1) - W_t)^{\nu/\kappa} \\ &\quad \times \left(\frac{g_t(1) - V_t^R}{1 - x^R} \right)^{\nu\rho^{1,R}/(2\kappa)} \left(\frac{g_t(1) - V_t^L}{1 - x^L} \right)^{\nu\rho^L/(2\kappa)}. \end{aligned}$$

By Lemma 2.4, the process M is a local martingale and the law of η weighted by M becomes the law of $\text{SLE}_\kappa(\rho^L; \rho^{1,R}, \rho^{2,R} + \nu)$ with force points $(x^L; x^R, 1)$.

On $E_\epsilon(\delta, r) := \{\tau_\epsilon \leq S_r, \text{Im } \eta(\tau_\epsilon) \geq \delta\epsilon(1 - x^R)\}$, let us estimate the terms in M_t one by one for $t = \tau_\epsilon$. Let O_t be the image of the rightmost point of $\eta[0, t] \cap \mathbb{R}$ under g_t . By the Koebe 1/4 theorem, we have $g_t(1) - O_t \asymp g'_t(1)\epsilon(1 - x^R)$.

- Consider the term $g_t(1) - W_t$. Since $\text{Im } \eta(t) \geq \delta\epsilon(1 - x^R)$, combining with [MW17, Lemma 3.4], we have

$$g'_t(1)\epsilon(1-x^R) \asymp g_t(1) - O_t \leq g_t(1) - W_t \lesssim (g_t(1) - O_t)/\delta \asymp g'_t(1)\epsilon(1-x^R)/\delta.$$

- Consider the term $g_t(1) - V_t^R$. If x^R is swallowed by $\eta[0, t]$, then we have $g_t(1) - V_t^R = g_t(1) - O_t \asymp g'_t(1)\epsilon(1 - x^R)$. If not, by Keobe 1/4 theorem, we have $g_t(1) - V_t^R \asymp g'_t(1)\epsilon(1 - x^R)$. In any case, we have

$$g_t(1) - V_t^R \asymp g'_t(1)\epsilon(1 - x^R).$$

- Consider the term $g_t(1) - V_t^L$. Since $g_t(1) - V_t^L$ is increasing in t , we have $g_t(1) - V_t^L \geq 1 - x^L$ for all t . Suppose \mathcal{B}^{yi} is a Brownian motion starting from yi , from [Law05, Remark 3.50], we know that

$$g_t(1) - V_t^L = \lim_{y \rightarrow \infty} \pi y \mathbb{P} [\mathcal{B}^{yi} \text{ exits } \mathbb{H} \setminus \eta[0, t] \text{ through the union } [x^L, 0] \cup \eta[0, t] \cup [0, 1]].$$

Since $t = \tau_\epsilon \leq S_r$, we have

$$g_t(1) - V_t^L \leq \lim_{y \rightarrow \infty} \pi y \mathbb{P} [\mathcal{B}^{yi} \text{ exits } \mathbb{H} \setminus B(0, r) \text{ through the union } [x^L, 0] \cup B(0, r)].$$

If $|x^L| \leq r$, then $g_t(1) - V_t^L \leq 4r$; if $|x^L| \geq r$, we have $g_t(1) - V_t^L \leq |x^L| + 3r$. Thus we have

$$1 \leq \frac{g_t(1) - V_t^L}{1 - x^L} \leq 4r.$$

Combining the above three parts, on $E_\epsilon(\delta, r)$, we have

$$\epsilon^{-\alpha}(1 - x^R)^{-\beta} \delta^\beta r^{-B} \lesssim M_{\tau_\epsilon} \lesssim \epsilon^{-\alpha}(1 - x^R)^{-\beta} r^{0 \vee (-\beta\rho^L/2)}.$$

Therefore, we have the lower bound: $\mathbb{P}[E_\epsilon(\delta, r)] \geq \mathbb{P}[E_\epsilon(4, 1/4)]$ and

$$\mathbb{P}[E_\epsilon(4, 1/4)] \gtrsim \epsilon^\alpha(1 - x^R)^\beta \mathbb{E}[M_{\tau_\epsilon} \mathbb{1}_{E_\epsilon(4, 1/4)}] = \epsilon^\alpha(1 - x^R)^\beta \mathbb{P}^*[E_\epsilon^*(4, 1/4)],$$

where η^* is an $\text{SLE}_\kappa(\rho^L; \rho^{1,R}, \rho^{2,R} + \nu)$ with force points $(x^L; x^R, 1)$ and \mathbb{P}^* is its law, and $E_\epsilon^*(r, \delta)$ is defined for η^* . Note that

$$\rho^{1,R} + \rho^{2,R} + \nu = \kappa - 8 - \rho^{1,R} - \rho^{2,R} < \kappa/2 - 4.$$

Thus η^* converges to the point 1 at finite time. Let $\phi(z) = z/(1 - z)$ be the Mobius transform of \mathbb{H} that sends $(0, 1, \infty)$ to $(0, \infty, -1)$ and let $\hat{\eta}$ be

the image of η^* under ϕ . Then $\hat{\eta}$ is an $\text{SLE}_\kappa(\rho^{1,R} + \rho^{2,R} + 2 - \rho^L, \rho^L; \rho^{1,R})$ with force points $(-1, \phi(x^L); \phi(x^R))$. Define \hat{E} to be the event that $\hat{\eta}$ never hits $B(-1, 1/3)$ and $\hat{\eta}$ exits the ball of radius $1/(\epsilon(1 - x^R))$ through the angle interval $[\pi/4, 3\pi/4]$. It is clear that $\mathbb{P}^*[E_\epsilon^*(4, 1/4)] \geq \hat{\mathbb{P}}[\hat{E}] \asymp 1$ (see for instance [MW17, Lemma 2.3]), since $\rho^{1,R} + \rho^{2,R} + 2 > \kappa/2 - 2$. This gives the lower bound. For the upper bound, we have

$$\mathbb{P}[E_\epsilon(\delta, r)] \lesssim \epsilon^\alpha (1 - x^R)^\beta \delta^{-\beta} r^B \mathbb{E}[M_{\tau_\epsilon} \mathbb{1}_{E_\epsilon(\delta, r)}] \leq \epsilon^\alpha (1 - x^R)^\beta \delta^{-\beta} r^B,$$

as desired. \square

LEMMA 6.4. *Fix $\kappa > 0$ and $\rho^L, \rho^R \in \mathbb{R}$. Suppose that x_n^L (resp. x_n^R) is a sequence of negative (resp. positive) real numbers converging to $x^L \leq 0$ (resp. $x^R \geq 0$) as $n \rightarrow \infty$. For each n , suppose that (W_n, V_n^L, V_n^R) is the driving triple for an $\text{SLE}_\kappa(\rho^L; \rho^R)$ process in \mathbb{H} with force points $(x_n^L; x_n^R)$. Then (W_n, V_n^L, V_n^R) converges weakly in law with respect to the local uniform topology to the driving triple (W, V^L, V^R) of an $\text{SLE}_\kappa(\rho^L; \rho^R)$ with force points $(x^L; x^R)$ as $n \rightarrow \infty$. The same likewise holds in the setting of multiple force points $\text{SLE}_\kappa(\underline{\rho})$ processes.*

PROOF. Proof of [MS16, Theorem 2.4]. \square

LEMMA 6.5. *Assume (6.1) holds. Suppose that η is an $\text{SLE}_\kappa(\rho^L; \rho^{1,R}, \rho^{2,R})$ with force points $(x^L; x^{1,R}, x^{2,R})$ where $x^L \leq 0$ and $0 \leq x^{1,R} \leq 1$ and $x^{1,R} \leq x^{2,R}$. Then there exists a function $p(\delta) \rightarrow 0$ as $\delta \rightarrow 0$ which depends only on $\kappa, \rho^L, \rho^{1,R}, \rho^{2,R}$ such that*

$$\mathbb{P}[\eta \text{ hits } B(1, \delta)] \leq p(\delta).$$

We emphasize that $p(\delta)$ is uniform over $x^L \leq 0 \leq x^{1,R} \leq 1$ and $x^{2,R} \geq x^{1,R}$.

PROOF. Define $f(x^L, x^{1,R}, x^{2,R}, \delta) = \mathbb{P}[\eta \text{ hits } B(1, \delta)]$. We argue that f is continuous. Suppose η_n is an $\text{SLE}_\kappa(\rho^L; \rho^{1,R}, \rho^{2,R})$ with force points $(x_n^L; x_n^{1,R}, x_n^{2,R})$ and assume $(x_n^L, x_n^{1,R}, x_n^{2,R}, \delta_n) \rightarrow (x^L, x^{1,R}, x^{2,R}, \delta)$. Denote by $g_t^n, W_n(t)$ the conformal map and the driving function for η_n . By Lemma 6.4, we see that W_n converges to W in local uniform topology. Combining with [Law05, Section 4.7], we obtain that g^n converges to g in Carathéodory topology. This gives the continuity of f .

Define $p(\delta) := \sup f(x^L, x^{1,R}, x^{2,R}, \delta)$ where the supremum is taken over $x^L \leq 0, x^{1,R} \in [0, 1], x^{2,R} \geq x^{1,R}$. Since (6.1) holds, for each $x^L, x^{1,R}, x^{2,R}$ we have $f(x^L, x^{1,R}, x^{2,R}, \delta) \rightarrow 0$ as $\delta \rightarrow 0$. When $|x^L|, x^{2,R} \rightarrow \infty$, the law of η becomes the law of $\text{SLE}_\kappa(\rho^{1,R})$. This implies $p(\delta) \rightarrow 0$ as $\delta \rightarrow 0$. \square

PROOF OF THEOREM 6.1. Lemma 6.3 gives the lower bound, and we only need to show the upper bound. Pick an integer n such that $2^n \epsilon \leq 1/4$. For $1 \leq k \leq n$, let T_k be the first time that η hits the ball centered at 1 with radius $2^{n+1-k} \epsilon (1 - x^R)$. Define

$$\mathcal{F}_k = \{\text{Im } \eta(T_k) \geq \delta 2^{n+1-k} \epsilon (1 - x^R)\}.$$

By Lemma 6.3, we know that

$$\begin{aligned} \mathbb{P}[\tau_\epsilon \leq S_r] &\leq \sum_1^n \mathbb{P}[\{T_k < \infty\} \cap \mathcal{F}_k] + \mathbb{P}[\{\tau_\epsilon \leq S_r\} \cap_1^n \mathcal{F}_k^c] \\ &\lesssim 2^{n\alpha} \epsilon^\alpha (1 - x^R)^\beta \delta^{-\beta} r^B + \mathbb{P}[\{\tau_\epsilon \leq S_r\} \cap_1^n \mathcal{F}_k^c]. \end{aligned}$$

By a similar argument as in Lemma 4.4, there is a function $q(\delta) \rightarrow 0$ as $\delta \rightarrow 0$ which depends only on $\kappa, \rho^L, \rho^{1,R}$ and $\rho^{2,R}$ such that $\mathbb{P}[\cap_1^n \mathcal{F}_k^c] \leq q(\delta)^n$. Thus we have

$$\mathbb{P}[\tau_\epsilon \leq S_r] \lesssim 2^{n\alpha} \epsilon^\alpha (1 - x^R)^\beta \delta^{-\beta} r^B + q(\delta)^n.$$

This implies the conclusion. \square

PROOF OF COROLLARY 6.2. Assume the same notations as in the proof of Theorem 6.1. When $\rho^L \leq 0$, we have $B = 0$, thus

$$\mathbb{P}[\tau_\epsilon < \infty] \lesssim 2^{n\alpha} \epsilon^\alpha (1 - x^R)^\beta \delta^{-\beta} + q(\delta)^n.$$

This implies the conclusion. \square

REFERENCES

- [Ahl06] Lars V Ahlfors. Lectures on quasiconformal mappings. *American Mathematical Society, Providence, RI*, 1(4):12, 2006.
- [Ahl10] Lars Valerian Ahlfors. *Conformal invariants: topics in geometric function theory*, volume 371. American Mathematical Soc., 2010.
- [AK08] Tom Alberts and Michael J. Kozdron. Intersection probabilities for a chordal SLE path and a semicircle. *Electron. Commun. Probab.*, 13:448–460, 2008.
- [BDCH16] Stéphane Benoist, Hugo Duminil-Copin, and Clément Hongler. Conformal invariance of crossing probabilities for the Ising model with free boundary conditions. *Ann. Inst. Henri Poincaré Probab. Stat.*, 52(4):1784–1798, 2016.
- [Bef08] Vincent Beffara. The dimension of the SLE curves. *Ann. Probab.*, 36(4):1421–1452, 2008.
- [CDCH⁺14] Dmitry Chelkak, Hugo Duminil-Copin, Clément Hongler, Antti Kemppainen, and Stanislav Smirnov. Convergence of Ising interfaces to Schramms SLE curves. *Comptes Rendus Mathématique*, 352(2):157–161, 2014.
- [CDCH16] Dmitry Chelkak, Hugo Duminil-Copin, and Clément Hongler. Crossing probabilities in topological rectangles for the critical planar FK-Ising model. *Electron. J. Probab.*, 21:Paper No. 5, 28, 2016.

- [Che16] Dmitry Chelkak. Robust discrete complex analysis: A toolbox. *Ann. Probab.*, 44(1):628–683, 2016.
- [CS12] Dmitry Chelkak and Stanislav Smirnov. Universality in the 2D Ising model and conformal invariance of fermionic observables. *Invent. Math.*, 189(3):515–580, 2012.
- [Dub09] Julien Dubédat. Duality of Schramm-Loewner evolutions. *Ann. Sci. Éc. Norm. Supér. (4)*, 42(5):697–724, 2009.
- [Dup03] Bertrand Duplantier. Conformal fractal geometry and boundary quantum gravity. 2003.
- [HK13] Clément Hongler and Kalle Kytölä. Ising interfaces and free boundary conditions. *Journal of the American Mathematical Society*, 26(4):1107–1189, 2013.
- [Law05] Gregory F. Lawler. *Conformally invariant processes in the plane*, volume 114 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2005.
- [Law15] Gregory F. Lawler. Minkowski content of the intersection of a Schramm-Loewner evolution (SLE) curve with the real line. *J. Math. Soc. Japan*, 67(4):1631–1669, 2015.
- [LSW01] Gregory F. Lawler, Oded Schramm, and Wendelin Werner. Values of Brownian intersection exponents. I. Half-plane exponents. *Acta Math.*, 187(2):237–273, 2001.
- [LSW02] Gregory F. Lawler, Oded Schramm, and Wendelin Werner. One-arm exponent for critical 2D percolation. *Electron. J. Probab.*, 7:no. 2, 13 pp. (electronic), 2002.
- [MS16] Jason Miller and Scott Sheffield. Imaginary geometry I: Interacting SLEs. *Probab. Theory Related Fields*, 164(3-4):553–705, 2016.
- [MW17] Jason Miller and Hao Wu. Intersections of SLE Paths: the double and cut point dimension of SLE. *Probab. Theory Related Fields*, 167(1-2):45–105, 2017.
- [Pom92] Ch. Pommerenke. *Boundary behaviour of conformal maps*, volume 299. Springer-Verlag, Berlin, 1992.
- [RS05] Steffen Rohde and Oded Schramm. Basic properties of SLE. *Ann. of Math. (2)*, 161(2):883–924, 2005.
- [SW01] Stanislav Smirnov and Wendelin Werner. Critical exponents for two-dimensional percolation. *Math. Res. Lett.*, 8(5-6):729–744, 2001.
- [SW05] Oded Schramm and David B. Wilson. SLE coordinate changes. *New York J. Math.*, 11:659–669 (electronic), 2005.
- [VL12] Fredrik Johansson Viklund and Gregory F. Lawler. Almost sure multifractal spectrum for the tip of an SLE curve. *Acta Math.*, 209(2):265–322, 2012.
- [Wu17+] Hao Wu. Polychromatic arm exponents for the critical planar FK-Ising model. *To appear in Journal of Statistical Physics*. 2017+
- [WW13] Wendelin Werner and Hao Wu. From CLE(κ) to SLE(κ, ρ). *Electron. J. Probab.*, 18:no. 36, 1–20, 2013.
- [WZ16] Hao Wu and Dapeng Zhan. Boundary arm exponents for SLE. *Electron. J. Probab.*, 22:no. 89, 1–26, 2017.