

# Gross-Zagier formula for $GL(2)$ II

Shou-Wu Zhang  
Department of Mathematics  
Columbia University  
New York, NY 10027

March 4, 2003

## Contents

1	Introduction and notations	2
2	Automorphic forms	6
3	Weights and levels	9
4	Automorphic L-series	11
5	Rankin-Selberg L-series	13
6	Odd case	14
7	Even case	17
8	Idea of Gross and Zagier	19
9	Calculus on arithmetic surfaces	22
10	Decomposition of Heights	25
11	Construction of the kernels	28
12	Geometric pairing	33

<b>13 Local Gross-Zagier formula</b>	<b>37</b>
<b>14 Gross-Zagier formula in level <math>ND</math></b>	<b>40</b>
<b>15 Green's functions of Heegner points</b>	<b>41</b>
<b>16 Spectral decomposition</b>	<b>46</b>
<b>17 Lowering levels</b>	<b>48</b>
<b>18 Continuous spectrum</b>	<b>55</b>
<b>19 Periods of Eisenstein series</b>	<b>58</b>

## 1 Introduction and notations

Let  $A$  be an abelian variety defined over a number field  $F$  and let

$$\rho : \text{Gal}(\bar{F}/F) \longrightarrow \text{GL}_n(\mathbb{C})$$

be a finite dimensional representation of the Galois group of  $F$ . Then the Birch and Swinnerton-Dyer conjecture predicts the following identity

$$\text{ord}_{s=1} L(s, \rho, A) = \dim(A(\bar{F}) \otimes \rho)^{\text{Gal}(\bar{F}/F)}.$$

Here  $L(s, \rho, A)$  denotes an Euler product over all places of  $F$ :

$$L(s, \rho, A) := \prod_v L_v(s, \rho, A), \quad (\text{Re}(s) \gg 0)$$

with good local factors given by

$$L_v(s, \rho, A) = \det(1 - q_v^{-s} \text{Frob}_v |_{T_\ell(A) \otimes \rho})^{-1},$$

where  $\ell$  is a prime different than the residue characteristic of  $v$ , and  $\mathbb{Z}_\ell$  has been embedded into  $\mathbb{C}$ . Moreover precisely, Birch and Swinnerton-Dyer conjecture predicts that the leading term of  $L(s, \rho, A)$  in the Taylor expansion in  $(s - 1)$  is given in terms of periods, Tate-Sharfarevich groups, and Mordell-Weil group. We refer to Tate's Bourbaki talk [9] for the details of the formulation.

In this paper, we will restrict ourself to the following very special situation:

- $A/F$  is an abelian variety associated to a Hilbert newform  $\phi$  over a totally real field  $F$  with trivial central character;
- $\rho$  is a representation induced from a ring class character  $\chi$  of  $\text{Gal}(\bar{K}/K)$  where  $K/F$  is a totally imaginary quadratic extension;
- the conductor  $N$  of  $\phi$ , the conductor  $c$  of  $\chi$ , and the discriminant  $d_{K/F}$  of  $K/F$  are coprime to each other.

In this case,  $L(s + 1/2, \rho, A)$  is a product of the Rankin L-series  $L(s, \chi, \phi^\sigma)$ , where  $\phi^\sigma$  are the Galois conjugates of  $\phi$ . Moreover,  $L(s, \chi, \phi)$  has a *symmetric functional equation*:

$$L(s, \chi, \phi) = \epsilon(\chi, \phi) \cdot N_{F/\mathbb{Q}}(ND)^{1-2s} \cdot L(1-s, \chi, \phi)$$

where

$$\epsilon(\chi, \phi) = \pm 1, \quad D = c^2 d_{K/F}.$$

The main result in our papers in Asia Journal and Annals [16, 17] is to express  $L'(1, \chi, \phi)$  (resp.  $L(1, \chi, \phi)$ ) when  $\epsilon(\chi, \phi) = -1$  (resp.  $\epsilon(\chi, \phi) = +1$ ) in terms of Heegner cycles in certain Shimura varieties of dimension 1 (resp. 0) of level  $ND$ . This result is a generalization of the landmark work of Gross and Zagier in their Inventiones paper [6] on Heegner points on  $X_0(N)/\mathbb{Q}$  with square free discriminant  $D$ .

The aim of this paper is to review the proofs in our previous papers [16, 17]. We also take this opportunity to deduce a new formula for Shimura varieties of level  $N$ . In odd case, the formula reads as

$$L'(1/2, \chi, \phi) = \frac{2^{g+1}}{\sqrt{N(D)}} \|\phi\|^2 \|x_\phi\|^2$$

where  $x_\phi$  is certain Heegner point in the Jacobian of Shimura curve. See Theorem 6.1 for details. In even case, the formula reads as

$$L(1/2, \chi, \phi) = \frac{2^g}{\sqrt{N(D)}} \|\phi\|^2 |(\tilde{\phi}, P_\chi)|^2$$

where  $(\tilde{\phi}, P_\chi)$  is the evaluation of certain test form on a CM-cycle  $P_\chi$  on a Shimura variety of dimension 0. See Theorem 7.1 for details. These results have more direct applications to the Birch and Swinnerton-Dyer conjecture

and  $p$ -adic L-series and Iwasawa theory. See papers ([1, 11]) of Bertolini-Darmon and Vatsal for details.

To do so, we need to compute various constants arising in the comparisons of normalizations of newforms or test vectors. This will follow from a comparison of two different ways to compute the periods of Eisenstein series. One is an extension of the method for cusp form in our Asia Journal paper [16], and another one is a direct evaluation by unfolding the integrals. Notice that the residue and constant term of Dedekind zeta function can be computed by the periods formula for Eisenstein series. Thus, the Gross-Zagier formula can be considered as an extension of class number formula and Kronecker limit formula not only in its statement but also in its method of proof.

Notice that Waldspurger has obtained a formula (when  $\chi$  is trivial [12]) and a criterion (when  $\chi$  is non trivial [13]) in the general situation where

- $K/F$  is any quadratic extension of number fields, and
- $\phi$  is any cusp form for  $\mathrm{GL}_2(\mathbb{A}_F)$ , and
- $\chi$  is any automorphic character of  $\mathrm{GL}_1(\mathbb{A}_K)$  such that the central character is reciprocal to  $\chi|_{\mathbb{A}_F^\times}$ .

We refer to papers of Gross and Vatsal ([5, 10]) in this volume for the explanation of connections between our formula and his work. There seems to be a lot of rooms left to generalize our formula to the case considered by Waldspurger. In this direction, Hui Xue in his thesis ([15]) has obtained a formula for the central values for L-series attached to a holomorphic Hilbert modular form of parallel weight  $2k$ .

This paper is organized as follows. In the first part (§2-7), we will give the basic definitions of forms, L-series, Shimura varieties, CM-points, and state our main formula (Theorem 6.1 and Theorem 7.1) in level  $N$ . The definitions here are more or less standard and can be found from our previous work as well as the work of Jacquet, Langlands, Waldspurger, Deligne, Carayol, Gross, and Prasad. Forms has been normalized as *newforms or test vectors* according to the action of unimportant or torus subgroup.

In the second part (§8-10), we will review the original ideas of Gross-Zagier in their Inventiones paper ([6]) on  $X_0(N)$  with square free  $D$  and its generalization to Shimura curves of  $(N, K)$ -type in our Annals paper ([17]). The central idea is to compare the Fourier coefficients of certain *natural* kernel functions of level  $N$  with certain *natural* CM-points on Shimura curves

$X(N, K)$  of  $(N, K)$ -type. This idea only works perfectly when  $D$  is square free and when  $X(N, D)$  has regular integral model but has essential difficulty for the general case.

In the third part (§11-16), we review the basic construction and the proof in our Asia Journal paper ([16]) for formulas in level  $ND$ . The kernel function and CM-points we pick are good for computation but have level  $ND$ . Their correspondence is given by local Gross-Zagier formula which is of course the key of the whole proof. The final formulas involve the notion of *quasi-newforms* or *toric newforms* as variations of newforms or test vectors.

In the last part (§17-19) which is our new contribution in addition to our previous papers, we will deduce the formula in level  $N$  from level  $ND$ . The plan of proof is stated in the beginning of §17 as three steps. The central ideal is to use Eisenstein series to compute certain local constants. This is one more example in number theory that local questions can be solved by global method, as in the early development of local class field theory and in the current work of Harris-Taylor on local Langlands conjecture.

The first three parts (§2-18) are simply review of ideas used in our previous papers. For details one may need to go to the original papers. For an elementary expository of the Gross-Zagier formula (or its variants as Gross formula or Kohen-Gross-Zagier formula) and its applications to Birch and Swinnerton-Dyer conjecture, we refer to our paper for Harvard-MIT conference on current developments of mathematics ([18]).

I would like to thank N. Vatsal and H. Xue for pointing out many inaccuracies in our previous paper [16] ( especially the missing of the first Fourier coefficient of the quasi-newform in the main formulas); to B. Gross for his belief of the existence of a formula in level  $N$  and for his many very useful suggestions in preparation of this note; to D. Goldfeld and H. Jacquet for their constant supports and encouragements.

## Notations

The notations of this note are mainly adopted from our Asia Journal paper [16] with some simplifications.

1. Let  $F$  denote a totally real field of degree  $g$  with ring of integers  $\mathcal{O}_F$ , and adèles  $\mathbb{A}$ . For each place  $v$  of  $F$ , let  $F_v$  denote the completion of  $F$  at  $v$ . When  $v$  is finite, let  $\mathcal{O}_v$  denote the ring of integers and let  $\pi_v$  denote a uniformizer of  $\mathcal{O}_v$ . We write  $\widehat{\mathcal{O}}_F$  for the product of  $\mathcal{O}_v$  in  $\mathbb{A}$ .
2. Let  $\psi$  denote a fixed nontrivial additive character of  $F \backslash \mathbb{A}$ . For each

place  $v$ , let  $\psi_v$  denote the component of  $\psi$  and let  $\delta_v \in F_v^\times$  denote the conductor of  $\psi_v$ . When  $v$  is finite,  $\delta_v^{-1}\mathcal{O}_v$  is the maximal fractional ideal of  $F_v$  over which  $\psi_v$  is trivial. When  $v$  is infinite,  $\psi_v(x) = e^{2\pi\delta_v x}$ . Let  $\delta$  denote  $\prod \delta_v \in \mathbb{A}^\times$ . Then the norm  $|\delta|^{-1} = d_F$  is the discriminant of  $F$ .

3. Let  $dx$  denote a Haar measure on  $\mathbb{A}$  such that the volume of  $F \backslash \mathbb{A}$  is one. This measure has a decomposition  $dx = \otimes dx_v$  into local measures  $dx_v$  on  $F_v$  which are self-dual with respect to characters  $\psi_v$ . Let  $d^\times x$  denote a Haar measure on  $\mathbb{A}^\times$  which has a decomposition  $d^\times x = \otimes d^\times x_v$  such that  $d^\times x_v = dx_v/x_v$  on  $F_v^\times = \mathbb{R}^\times$  when  $v$  is infinite, and such that the volume of  $\mathcal{O}_v^\times$  is one when  $v$  is finite. Notice that our setting of multiplicative measures is different than those in Tate's thesis, where the volume of  $\mathcal{O}_v^\times$  is  $|\delta_v|^{1/2}$ .

4. Let  $K$  denote a totally imaginary quadratic extension of  $F$  and  $T$  denote the algebraic group  $K^\times/F^\times$  over  $F$ . We will fix a Haar measure  $dt$  and its decomposition  $dt = \otimes dt_v$  such that  $T(F_v)$  has volume 1 when  $v$  is infinite.

5. Let  $B$  denote a quaternion algebra over  $F$  and let  $G$  denote the algebraic group  $B^\times/F^\times$  over  $F$ . We will fix a Haar measure  $dg$  on  $G(\mathbb{A})$  and a decomposition  $dg = \otimes dg_v$  such that at an infinite place  $G(F_v)$  has volume one if it is compact, and that when  $G(F_v) \simeq \mathrm{PGL}_2(\mathbb{R})$ ,

$$dg_v = \frac{|dxdy|}{2\pi y^2} d\theta,$$

with respect to the decomposition

$$g_v = z \begin{pmatrix} y & x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}.$$

In this way, the volume  $|U|$  of the compact open subgroup  $U$  of  $G(\mathbb{A}_f)$  (or  $G(F_v)$  for some  $v \nmid \infty$ ) is well defined. We write  $(f_1, f_2)_U$  for the hermitian product

$$(f_1, f_2)_U = |U|^{-1} \int_{G(\mathbb{A})} f_1 \bar{f}_2 dg$$

for functions  $f_1, f_2$  on  $G(\mathbb{A})$  (or  $G(\mathbb{A}_f)$ , or  $G(F_v)$ ). This product depends only on the choice of  $U$  but not on  $dg$ .

## 2 Automorphic forms

Let  $F$  be a totally real field of degree  $g$ , with ring of adeles  $\mathbb{A}$ , and discriminant  $d_F$ . Let  $\omega$  be a (unitary) character of  $F^\times \backslash \mathbb{A}^\times$ . By an *automorphic*

form on  $\mathrm{GL}_2(\mathbb{A})$  with central character  $\omega$  we mean a continuous function  $\phi$  on  $\mathrm{GL}_2(\mathbb{A})$  such that the following properties hold:

- $\phi(z\gamma g) = \omega(z)\phi(g)$  for  $z \in Z(\mathbb{A}), \gamma \in \mathrm{GL}_2(F)$ ;
- $\phi$  is invariant under right action of some open subgroup of  $\mathrm{GL}_2(\mathbb{A}_f)$ ;
- for a place  $v \mid \infty$ ,  $\phi$  is smooth in  $g_v \in \mathrm{GL}_2(F_v)$ , and the vector space generated by

$$\phi(gr_v), \quad r_v \in \mathrm{SO}_2(F_v) \subset \mathrm{GL}_2(\mathbb{A})$$

is finite dimensional;

- for any compact subset  $\Omega$  there are positive numbers  $C, t$  such that

$$\left| \phi \left( \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} g \right) \right| \leq C(|a| + |a^{-1}|)^t$$

for all  $g \in \Omega$ .

Let  $\mathcal{A}(\omega)$  denote the space of automorphic forms with central character  $\omega$ . Then  $\mathcal{A}(\omega)$  admits an *admissible representation*  $\rho$  by  $\mathrm{GL}_2(\mathbb{A})$ . This is a combination of a representation  $\rho_f$  of  $\mathrm{GL}_2(\mathbb{A}_f)$  via right action:

$$\rho_f(h)\phi(g) = \phi(gh), \quad h \in \mathrm{GL}_2(\mathbb{A}_f), \phi \in \mathcal{A}(\omega), g \in \mathrm{GL}_2(\mathbb{A}),$$

and an action  $\rho_\infty$  by pairs

$$(M_2(F_v), O_2(F_v)), \quad v \mid \infty.$$

Here the action of  $O_2(F_v)$  is the same as above while the action of  $M_2(F_v)$  is given by

$$\rho_\infty(x)\phi(g) = \frac{d\phi}{dt}(ge^{tx})|_{t=0}, \quad x \in M_2(F_v), \phi \in \mathcal{A}(\omega), g \in \mathrm{GL}_2(\mathbb{A}), v \mid \infty.$$

An admissible and irreducible representation  $\Pi$  of  $\mathrm{GL}_2(\mathbb{A})$  is called *automorphic* if it is isomorphic to a sub-representation of  $\mathcal{A}(\omega)$ . It is well-known that the multiplicity of any irreducible representation in  $\mathcal{A}(\omega)$  is at most 1. Moreover, if we decompose such a representation into local representations  $\Pi = \otimes \Pi_v$  then *the strong multiplicity one* says that  $\Pi$  is determined by all but finitely many  $\Pi_v$ .

Fix an additive character  $\psi$  on  $F \backslash \mathbb{A}$ . Then any automorphic form will have a Fourier expansion:

$$(2.1) \quad \phi(g) = C_\phi(g) + \sum_{\alpha \in F^\times} W_\phi \left( \begin{pmatrix} \alpha & 0 \\ 0 & 1 \end{pmatrix} g \right),$$

where  $C_\phi$  is the constant term:

$$(2.2) \quad C_\phi(g) := \int_{F \backslash \mathbb{A}} \phi \left( \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} g \right) dx,$$

and  $W_\phi(g)$  is the Whittaker function:

$$(2.3) \quad W_\phi(g) = \int_{F \backslash \mathbb{A}} \phi \left( \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} g \right) dx.$$

It is not difficult to show that a form with vanishing Whittaker function will have the form  $\alpha(\det g)$  where  $\alpha$  is a function on  $F^\times \backslash \mathbb{A}^\times$ . Every automorphic representation of dimension 1 appears in this space and corresponds to a characters  $\mu$  of  $F^\times \backslash \mathbb{A}^\times$  such that  $\mu^2 = \omega$ .

We say that an automorphic form  $\phi$  is *cuspidal* if the constant term  $C_\phi(g) = 0$ . The space of cuspidal forms is denoted by  $\mathcal{A}_0(\omega)$ . We call an automorphic representation *cuspidal* if it appears in  $\mathcal{A}_0(\omega)$ .

An irreducible automorphic representation which is neither one dimensional nor cuspidal must be isomorphic to the space  $\Pi(\mu_1, \mu_2)$  of Eisenstein series associated to two quasi characters  $\mu_1, \mu_2$  of  $F^\times \backslash \mathbb{A}^\times$  such that  $\mu_1 \mu_2 = \omega$ . To construct an Eisenstein series, let  $\Phi$  be a Schwartz-Bruhat function on  $\mathbb{A}^2$ . For  $s$  a complex number, define

$$(2.4) \quad f_\Phi(s, g) := \mu_1(\det g) |\det g|^{s+1/2} \int_{\mathbb{A}^\times} \Phi[(0, t)g] \mu_1 \mu_2^{-1}(t) |t|^{1+2s} d^\times t.$$

Then  $f_\Phi(s, g)$  belongs to the space  $\mathcal{B}(\mu_1 \cdot |\cdot|^s, \mu_2 \cdot |\cdot|^{-s})$  of functions on  $\mathrm{GL}_2(\mathbb{A})$  satisfying

$$(2.5) \quad f_\Phi \left( s, \begin{pmatrix} a & x \\ 0 & b \end{pmatrix} g \right) = \mu_1(a) \mu_2(b) \left| \frac{a}{b} \right|^{1/2+s} f_\Phi(s, g).$$

The Eisenstein series  $E(s, g, \Phi)$  is defined as follows:

$$(2.6) \quad E(s, g, \Phi) = \sum_{\gamma \in P(F) \backslash \mathrm{GL}_2(F)} f_\Phi(s, \gamma g).$$

One can show that  $E(s, g, \Phi)$  is absolutely convergent when  $\operatorname{Re}(s)$  is sufficiently large, and has a meromorphic continuation to the whole complex plane. The so-defined meromorphic function  $E(s, g, \Phi)$  has at most simple poles with constant residue. The space  $\Pi(\mu_1, \mu_2)$  consists of the following Eisenstein series:

$$(2.7) \quad E(g, \Phi) := \lim_{s \rightarrow 0} (E(s, g, \Phi) - (\text{residue})s^{-1}).$$

### 3 Weights and levels

Let  $N$  be an ideal of  $\mathcal{O}_F$  and let  $U_0(N)$  and  $U_1(N)$  be the following subgroups of  $\operatorname{GL}_2(\widehat{\mathbb{A}}_f)$ :

$$(3.1) \quad U_0(N) := \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \operatorname{GL}_2(\widehat{\mathcal{O}}_F) : c \equiv 0 \pmod{N} \right\},$$

$$(3.2) \quad U_1(N) := \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in U_0(N) : d \equiv 1 \pmod{N} \right\}.$$

For each infinite place  $v$  of  $F$ , let  $k_v$  be an integer such that  $\omega_v(-1) = (-1)^{k_v}$ .

An automorphic form  $\phi \in \mathcal{A}(\omega)$  is said to have *level*  $N$ , *weight*  $k = (k_v : v \mid \infty)$ , if the following conditions are satisfied:

- $\phi(gu) = \phi(g)$  for  $u \in U_1(N)$ ;
- for a place  $v \mid \infty$ ,

$$\phi(gr_v(\theta)) = \phi(g)e^{2\pi i k_v \theta}$$

where  $r_v(\theta)$  is an element in  $\operatorname{SO}_2(F_v) \subset \operatorname{GL}_2(\mathbb{A})$  of the form

$$r_v(\theta) = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}.$$

Let  $\mathcal{A}_k(N, \omega)$  denote the space of forms of weight  $k$ , level  $N$ , and central character  $\omega$ . For any level  $N' \mid N$  of  $N$  and weight  $k' \leq k$  by which we mean that  $k - k'$  has non-negative components, we may define embeddings

$$\mathcal{A}_{k'}(N', \omega) \longrightarrow \mathcal{A}_k(N, \omega)$$

by applying some of the following operators:

$$\phi \mapsto \rho_v \begin{pmatrix} \pi_v^{-1} & 0 \\ 0 & 1 \end{pmatrix} \phi, \quad (v \nmid \infty)$$

$$\phi \mapsto \rho_v \begin{pmatrix} 1 & i \\ i & -1 \end{pmatrix} \phi. \quad (v \mid \infty)$$

The first operator increases level by order 1 at a finite place  $v$ ; while the second operator increases weight by 2 at an infinite place  $v$ . Let  $\mathcal{A}_k^{\text{old}}(N, \omega)$  denote the subspace of forms obtained from lower level  $N'$  or lower weight  $k'$  by applying *at least one of the above operators*.

For any ideal  $a$  prime to  $N$ , the *Hecke operator*  $T_a$  on  $\mathcal{A}_k(N, \omega)$  is defined as follows:

$$(3.3) \quad T_a \phi(g) = \sum_{\substack{\alpha\beta=a \\ x \pmod{\alpha}}} \phi \left( g \begin{pmatrix} \alpha & x \\ 0 & \beta \end{pmatrix} \right)$$

where  $\alpha$  and  $\beta$  runs through representatives of integral ideles modulo  $\widehat{\mathcal{O}}_F^\times$  with trivial component at the place dividing  $N$  such that  $\alpha\beta$  generates  $a$ . One has the following formula for the Whittaker function:

$$(3.4) \quad W_\phi \left( g \begin{pmatrix} a\delta^{-1} & 0 \\ 0 & 1 \end{pmatrix} \right) = |a| W_{T_a \phi}(g)$$

where  $g \in \text{GL}_2(\mathbb{A})$  with component 1 at places  $v \nmid N \cdot \infty$ .

We say that  $\phi$  is an *eigenform* if for any ideal  $a$  prime to  $N$ ,  $\phi$  is an eigenform under the Hecke operator  $T_a$ . We say an eigenform  $\phi$  is *new* if all  $k_v \geq 0$ , and if there is no old eigenform with the same eigenvalues as  $\phi$ . One can show that two new eigenforms are proportional if and only if they share the same eigenvalues for all but finitely many  $T_v$ .

For  $\phi \in \mathcal{A}_k(N, \omega)$ , let's write  $\Pi(\phi)$  for the space of forms in

$$\mathcal{A}(\omega) = \cup_{k,N} \mathcal{A}_k(N, \omega)$$

generated by  $\phi$  by right action of  $\text{GL}_2(\mathbb{A})$ . Then one can show that  $\Pi(\phi)$  is irreducible if and only if  $\phi$  is an eigenform. Conversely, any irreducible representation  $\Pi$  of  $\text{GL}_2(\mathbb{A})$  in  $\mathcal{A}(\omega)$  contains a unique line of new eigenform. An eigenform  $\phi$  with  $\dim \Pi(\phi) < \infty$  will have vanishing Whittaker function and is a multiple of a character.

It can be shown that an eigen new form  $\phi$  with  $\dim \Pi(\phi) = \infty$  will have Whittaker function non-vanishing and decomposable:

$$(3.5) \quad W_\phi(g) = \otimes W_v(g_v)$$

where  $W_v(g_v)$  at finite places can be normalized such that

$$(3.6) \quad W_v \begin{pmatrix} \delta_v^{-1} & 0 \\ 0 & 1 \end{pmatrix} = 1.$$

Each local components  $\Pi_v$  is realized in the subspace

$$\mathcal{W}(\Pi_v, \psi_v) = \Pi(W_v)$$

generated by  $W_v$  under action right action of  $\mathrm{GL}_2(F_v)$  (or  $(M_2(F_v), O_2(F_v))$  when  $v$  is infinite.)

## 4 Automorphic L-series

For an automorphic form  $\phi$ , let us define its L-series by

$$(4.1) \quad \begin{aligned} L(s, \phi) &:= d_F^{1/2-s} \int_{F^\times \backslash \mathbb{A}^\times} (\phi - C_\phi) \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} |a|^{s-1/2} d^\times a \\ &= d_F^{1/2-s} \int_{\mathbb{A}^\times} W_\phi \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} |a|^{s-1/2} d^\times a \end{aligned}$$

which is absolutely convergent for  $\mathrm{Re}(s) \gg 0$ , and has a meromorphic continuation to the entire complex plane, and satisfies a functional equation.

Assume that  $\phi$  is an eigen new form. Then its Whittaker function is decomposable. The L-series  $L(s, \phi)$  is then an Euler product

$$(4.2) \quad L(s, \phi) = \prod_v L_v(s, \phi)$$

where

$$(4.3) \quad L_v(s, \phi) = |\delta_v|^{s-1/2} \int_{F_v^\times} W_v \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} |a|^{s-1/2} d^\times a.$$

For a finite place  $v$ , the L-factor has the usual expression:

$$(4.4) \quad L_v(s, \phi) = \begin{cases} (1 - \lambda_v |\pi_v|^s + \omega(\pi_v) |\pi_v|^{2s})^{-1}, & \text{if } v \nmid N, \\ (1 - \lambda_v |\pi_v|^s)^{-1}, & \text{if } v \mid N, \end{cases}$$

where  $\lambda_v \in \mathbb{C}$  is such that  $\lambda_v |\pi_v|^{-1/2}$  is the eigenvalue of  $T_v$  if  $v \nmid N$ .

For an archimedean place  $v$ , the local factor  $L_v(s, \phi)$  is certain product of Gamma functions and is determined by analytic properties of  $\phi$  at  $v$ . For the purpose of this paper, we will only consider new forms such that at an infinite place which is either *holomorphic* or *even of weight 0*, i.e. invariant under  $O_2(F_v)$  rather than  $SO_2(F_v)$ . More precisely, at an infinite place  $v$  lets consider the function on  $\mathcal{H} \times \mathrm{GL}_2(\mathbb{A}^v)$  defined by

$$(4.5) \quad f(z, g^v) := |y|^{-(k_v+w_v)/2} \phi \left( \begin{pmatrix} y & x \\ 0 & 1 \end{pmatrix}, g^v \right), \quad z = x + yi$$

where  $w_v = 0$  or  $1$  is such that  $\omega_v(-1) = (-1)^{w_v}$ . Then we require that  $f(z, g^v)$  is holomorphic in  $z$  if  $k_v \geq 1$ , and that  $f(z, g^v) = f(-\bar{z}, g^v)$  if  $k_v = 0$ . If  $\phi$  is of weight 0, then  $\phi$  is an eigen form for the Laplacien

$$(4.6) \quad \Delta = -y^2 \left( \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right).$$

We write eigenvalues as  $1/4 + t_v^2$  and call  $t_v$  the *parameter* of  $\phi$  at  $v$ . Let's define *the standard Whittaker function* at archiemdean places  $v$  of weight  $k_v$  in the following way: if  $k_v > 0$ ,

$$(4.7) \quad W_v \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} = \begin{cases} 2a^{(w_v+k_v)/2} e^{-2\pi a} & \text{if } a > 0, \\ 0 & \text{if } a < 0, \end{cases}$$

and if  $k_v = 0$ , then

$$(4.8) \quad W_v \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} = |a|^{1/2} \int_0^\infty e^{-\pi|a|(y+y^{-1})} y^{it_v} d^\times y.$$

In this manner, (up to a constant  $c \neq 0$ ),  $\phi$  will have a Whittaker function decomposable as in (3.5) with local function  $W_v$  normalized as in (3.6), (4.7), (4.8). We say that  $\phi$  is a *newform* if  $c = 1$ . Equivalently,  $\phi$  is a newform if and only if  $L(s, \phi_v)$  has decomposition (4.2) with local factors given by (4.4) when  $v \nmid \infty$ , and the following when  $v \mid \infty$ :

$$(4.9) \quad L_v(s, \phi) = \begin{cases} G_2(s + k_v + w_v), & \text{if } k_v > 0, \\ G_1(s + it_v)G_1(s - it_v), & \text{if } k_v = 0, \end{cases}$$

where

$$(4.10) \quad G_1(s) = \pi^{-s/2} \Gamma(s/2), \quad G_2(s) = 2(2\pi)^{-s} \Gamma(s) = G_1(s)G_1(s+1).$$

If  $\Pi$  is an automorphic representation generated by a newform  $\phi$ , we write  $L(s, \Pi)$  and  $L(s, \Pi_v)$  for  $L(s, \phi)$  and  $L_v(s, \phi)$ , respectively.

## 5 Rankin-Selberg L-series

Let  $K$  be a totally imaginary quadratic extension of  $F$ , and let  $\omega$  be the non-trivial quadratic character of  $\mathbb{A}^\times/F^\times N\mathbb{A}_K^\times$ . The conductor  $c(\omega)$  is the relative discriminant of  $K/F$ . Let  $\chi$  be a character of finite order of  $\mathbb{A}_K^\times/K^\times\mathbb{A}^\times$ . The conductor  $c(\chi)$  is an ideal of  $\mathcal{O}_F$  which is maximal such that  $\chi$  is factorized through

$$\mathbb{A}_K^\times/K^\times\mathbb{A}^\times\widehat{\mathcal{O}}_{c(\chi)}^\times K_\infty^\times = \text{Gal}(H_{c(\chi)}/K),$$

where  $\mathcal{O}_c = \mathcal{O}_F + c(\chi)\mathcal{O}_K$  and  $H_c$  is the ring class field of conductor  $c(\chi)$ . We define the ideal  $D = c(\chi)^2c(\omega)$ , and call  $\chi$  a *ring class character of conductor*  $c(\chi)$ .

Let  $\phi$  be a newform with trivial central character and of level  $N$ . The Rankin-Selberg convolution L-function  $L(s, \chi, \phi)$  is defined by an Euler product over primes  $v$  of  $F$ :

$$(5.1) \quad L(s, \chi, \phi) := \prod_v L_v(s, \chi, \phi)$$

where the factors have degree  $\leq 4$  in  $|\pi_v|^s$ . This function has an analytic continuation to the entire complex plane, and satisfies a functional equation. We will assume that the ideals  $c(\omega)$ ,  $c(\chi)$ ,  $N$  are coprime each other. Then the local factors can be defined explicitly as follows.

For  $v$  a finite place, let's write

$$L_v(s, \phi) = (1 - \alpha_1|\pi_v|^s)^{-1}(1 - \alpha_2|\pi_v|^s)^{-1},$$

$$\prod_{w|v} L(s, \chi_w) = (1 - \beta_1|\pi_v|^s)^{-1}(1 - \beta_2|\pi_v|^s)^{-1},$$

then

$$(5.2) \quad L_v(s, \chi, \phi) = \prod_{i,j} (1 - \alpha_i\beta_j|\pi_v|^s)^{-1}.$$

Here for a place  $w$  of  $K$ , the local factor  $L(s, \chi_w)$  is defined as follows:

$$(5.3) \quad L(s, \chi_w) = \begin{cases} (1 - \chi(\pi_w)|\pi_w|^s)^{-1}, & \text{if } w \nmid c \cdot \infty, \\ G_2(s), & \text{if } v \mid \infty, \\ 1, & \text{if } v \mid c. \end{cases}$$

At an infinite place  $v$ , using formula  $G_2(s) = G_1(s)G_1(s + 1)$  we may write

$$\begin{aligned} L_v(s, \phi) &= G_1(s + \sigma_1)G_1(s + \sigma_2), \\ L_v(s, \chi) &= G_1(s + \tau_1)G_1(s + \tau_2). \end{aligned}$$

Then the L-factor  $L_v(s, \chi, \phi)$  is defined as follows:

$$(5.4) \quad \begin{aligned} L_v(s, \chi, \phi) &= \prod_{i,j} G_1(s + \sigma_i + \tau_j) \\ &= \begin{cases} G_2(s + (k_v - 1)/2)^2, & \text{if } k_v \geq 2, \\ G_2(s + it_v)G_2(s - it_v), & \text{if } k_v = 0. \end{cases} \end{aligned}$$

where  $t_v$  is the parameter associated to  $\phi$  at a place  $v$  where the weight is 0.

The functional equation is then

$$(5.5) \quad L(1 - s, \chi, \phi) = (-1)^{\#\Sigma} N_{F/\mathbb{Q}}(ND)^{1-2s} L(s, \chi, \phi),$$

where  $\Sigma = \Sigma(N, K)$  is the following set of places of  $F$ :

$$(5.6) \quad \Sigma(N, K) = \left\{ v \left| \begin{array}{l} v \text{ is infinite, and } \phi \text{ has weight } k_v > 0 \text{ at } v, \text{ or} \\ v \text{ is finite, and } \omega_v(N) = -1. \end{array} \right. \right\}$$

## 6 Odd case

Now we assume that all  $k_v = 2$  and that the sign of the functional equation (5.5) is  $-1$ , so  $\#\Sigma$  is odd. Our main formula expresses the central derivative  $L'(1/2, \chi, \phi)$  in terms of the heights of CM-points on a Shimura curve. Let  $\tau$  be any real place of  $F$ , and let  $B$  be the quaternion algebra over  $F$  which ramified exactly at the places in  $\Sigma - \{\tau\}$ . Let  $G$  be the algebraic group over  $F$ , which is an inner form of  $\mathrm{PGL}_2$ , and has  $G(F) = B^\times / F^\times$ .

The group  $G(F_v) \simeq \mathrm{PGL}_2(\mathbb{R})$  acts on  $\mathcal{H}^\pm = \mathbb{C} - \mathbb{R}$ . If  $U \subset G(\mathbb{A}_f)$  is open and compact, we get an analytic space

$$(6.1) \quad M_U(\mathbb{C}) = G(F)_+ \backslash \mathcal{H} \times G(\mathbb{A}_f) / U$$

where  $G(F)_+$  denote the subgroup of elements of  $G(F)$  with totally positive determinants. Shimura proved these were the complex points of an algebraic curve  $M_U$ , which descends canonically to  $F$  (embedded in  $\mathbb{C}$ , by the place  $\tau$ ). The curve  $M_U$  over  $F$  is independent of the choice of  $\tau$  in  $\Sigma$ .

To specify  $M_U$ , we must define  $U \subset G(\mathbb{A}_f)$ . To do this, we fix an embedding  $K \rightarrow B$ , which exists, as all places in  $\Sigma$  are either inert or ramified in  $K$ . One can show that there is an order  $R$  of  $B$  containing  $\mathcal{O}_K$  with reduced discriminant  $N$ . For an explicit description of such an order, we fix a maximal ideal  $\mathcal{O}_B$  of  $B$  containing  $\mathcal{O}_K$  and an ideal  $\mathcal{N}$  of  $\mathcal{O}_K$  such that

$$(6.2) \quad N_{K/F}\mathcal{N} \cdot \text{disc}_{B/F} = N,$$

where  $\text{disc}_{B/F}$  is the reduced discriminant of  $\mathcal{O}_B$  over  $\mathcal{O}_F$ . Then we take

$$(6.3) \quad R = \mathcal{O}_K + \mathcal{N} \cdot \mathcal{O}_B.$$

We call  $R$  an order of  $(N, K)$ -type. Define an open compact subgroup  $U_v$  of  $G(F_v)$  by

$$(6.4) \quad U_v = R_v^\times / \mathcal{O}_v^\times.$$

Let  $U = \prod_v U_v$ . This defines the curve  $M_U$  up to  $F$ -isomorphism. Let  $X$  be its compactification over  $F$ , so  $X = M_U$  unless  $F = \mathbb{Q}$  and  $\Sigma = \{\infty\}$ , where  $X$  is obtained by adding many cusps. We call  $X$  a Shimura curve of  $(N, K)$ -type. We write  $R(N, K)$ ,  $U(N, K)$ ,  $X(N, K)$  when types need to be specified.

We will now construct points in  $\text{Jac}(X)$ , the connected component of  $\text{Pic}(X)$ , from CM-points on the curve  $X$ . The CM-points corresponding to  $K$  on  $M_U(\mathbb{C})$  form a set

$$(6.5) \quad G(F)_+ \backslash G(F)_+ \cdot h_0 \times G(\mathbb{A}_f) / U = T(F) \backslash G(\mathbb{A}_f) / U,$$

where  $h_0 \in \mathcal{H}$  is the unique fixed point of the torus points  $T(F) = K^\times / F^\times$ . Let  $P_c$  denote a point in  $X$  represented by  $(h_0, i_c)$  where  $i_c \in G(\mathbb{A}_f)$  such that

$$(6.6) \quad U_T := i_c U i_c^{-1} \cap T(\mathbb{A}_f) \simeq \widehat{\mathcal{O}}_c^\times / \widehat{\mathcal{O}}_F^\times.$$

By Shimura's theory,  $P_c$  is defined over the ring class field  $H_c$  of conductor  $c$  corresponding to the Artin map

$$\text{Gal}(H_c/K) \simeq T(F) \backslash T(\mathbb{A}_f) / T(F_\infty) U_T.$$

Let  $P_\chi$  be a divisor on  $X$  with complex coefficients defined by

$$(6.7) \quad P_\chi = \sum_{\sigma \in \text{Gal}(H_c/K)} \chi^{-1}(\sigma) [P_c^\sigma].$$

If  $\chi$  is not of form  $\chi = \nu \cdot N_{K/F}$  with  $\nu$  a quadratic character of  $F^\times \mathbb{A}^\times$ , then  $P_\chi$  has degree 0 on each connected component of  $X$ . Thus  $P_\chi$  defines a class  $x$  in  $\text{Jac}(X) \otimes \mathbb{C}$ . Otherwise we need a reference divisor to send  $P_\chi$  to  $\text{Jac}(X)$ . In the modular curve case, one uses cusps. In the general case, we use the *Hodge class*  $\xi \in \text{Pic}(X) \otimes \mathbb{Q}$ : the unique class whose degree is 1 on each connected component and such that

$$T_m \xi = \deg(T_m) \xi$$

for all integral nonzero ideal  $m$  of  $\mathcal{O}_F$  prime to  $ND$ . The Heegner class we want now is the class difference

$$(6.8) \quad x := [P_\chi - \deg(P_\chi)\xi] \in \text{Jac}(X)(H_c) \otimes \mathbb{C},$$

where  $\deg(P_\chi)$  is the multi-degree of  $P_\chi$  on geometric components.

Notice that the curve  $X$  and its Jacobian have an action by the ring of good Hecke operators. Thus  $x$  is a sum of eigen vectors of the Hecke operators.

**Theorem 6.1.** *Let  $x_\phi$  denote the  $\phi$ -typical component of  $x$ . Then*

$$L'(1/2, \chi, \phi) = \frac{2^{g+1}}{\sqrt{N(D)}} \cdot \|\phi\|^2 \cdot \|x_\phi\|^2.$$

Here,

- $\|\phi\|^2$  is computed using the invariant measure on

$$\text{PGL}_2(F) \backslash \mathcal{H}^g \times \text{PGL}_2(\mathbb{A}_f) / U_0(N)$$

induced by  $dxdy/y^2$  on  $\mathcal{H}$ ;

- $\|x_\phi\|^2$  is the Neron-Tate pairing of  $x_\phi$  with itself.

To see the application to the Birch and Swinnerton-Dyer conjecture, we just notice that  $x_\phi$  actually lives in a unique abelian subvariety  $A_\phi$  of the Jacobian  $\text{Jac}(X)$  such that

$$(6.9) \quad L(s, A_\phi) = \prod_{\sigma: \mathbb{Z}[\phi] \rightarrow \mathbb{C}} L(s, \phi^\sigma).$$

Y. Tian [14] has recently generalized the work of Kolyvagin and Bertolini-Darmon to our setting and showed that the rank conjecture of Birch and Swinnerton-Dyer for  $A$  in the case  $\text{ord}_{s=1/2} L(s, \chi, \phi) \leq 1$ .

Notice that  $\|\phi\|^2$  is not exactly the periods of  $A_\phi$  appearing in the Birch and Swinnerton-Dyer conjecture, but it has expression in L-series:

$$(6.10) \quad \|\phi\|^2 = 2N(N) \cdot d_F \cdot L(1, \text{Sym}^2 \phi)$$

where  $L(s, \text{Sym}^2 \phi)$  is the L-series defined by an Euler product with local factors  $L_v(s, \text{Sym}^2 \phi)$  given by

$$(6.11) \quad L_v(s, \text{Sym}^2 \phi) = G_2(s + 1/2)^2 G_1(s)^{-1},$$

if  $v \mid \infty$ , and by

$$(6.12) \quad L_v(s, \text{Sym}^2 \phi) = (1 - \alpha^2 |\pi_v|^s)^{-1} (1 - \beta^2 |\pi_v|^s)^{-1} (1 - \alpha\beta |\pi_v|^s)^{-1},$$

if  $v \nmid \infty$ , and  $\alpha$  and  $\beta$  are given as follows:

$$L_v(s, \phi) = (1 - \alpha |\pi_v|^s)^{-1} (1 - \beta |\pi_v|^s)^{-1}.$$

It will be an interesting question to see how this relates the periods in  $A_\phi$ .

## 7 Even case

We now return to the case where  $\phi$  has possible nonholomorphic components, but we assume that all weights be either 0 or 2 and that the sign of the functional equation of  $L(s, \chi, \phi)$  is  $+1$ , or equivalently,  $\Sigma$  is even. In this case, we have an explicit formula for  $L(1/2, \chi, \phi)$  in terms of CM-points on locally symmetric varieties covered by  $\mathcal{H}^n$  where  $n$  is the number of real places of  $F$  where  $\phi$  has weight 0.

More precisely, let  $B$  be the quaternion algebra over  $F$  ramified at  $\Sigma$ , and  $G$  the algebraic group associated to  $B^\times / F^\times$ . Then

$$(7.1) \quad G(F \otimes \mathbb{R}) \simeq \text{PGL}_2(\mathbb{R})^n \times \text{SO}_3^{g-n}$$

acts on  $(\mathcal{H}^\pm)^n$ . The locally symmetric variety we will consider is

$$(7.2) \quad M_U = G(F)_+ \backslash \mathcal{H}^n \times G(\mathbb{A}_f) / U,$$

where  $U = \prod U_v$  was defined in the previous §. Again we call  $M_U$  or its compactification  $X$  a *quaternion Shimura variety of  $(N, K)$ -type*. We will also have a CM-point  $P_c$  and a CM-cycle  $P_\chi$  defined as in (6.6) and (6.7) but with  $\text{Gal}(H_c/K)$  replaced by  $T(F)\backslash T(\mathbb{A}_f)/T(F_\infty)U_T$ .

By some results of Waldspurger, Tunnel, and Gross-Prasad ([17], Theorem 3.2.2), there is a unique line of cuspidal functions  $\tilde{\phi}$  on  $M_U$  such that for each finite place  $v$  not dividing  $N \cdot D$ ,  $\tilde{\phi}$  is the eigenform for Hecke operators  $T_v$  with the same eigenvalues as  $\phi$ . We call any such a form a *test form* of  $(N, K)$ -type.

**Theorem 7.1.** *Let  $\tilde{\phi}$  be a test form of norm 1 with respect to the measure on  $X$  induced by  $dx dy/y^2$  on  $\mathcal{H}$ . Then*

$$L(1/2, \chi, \phi) = \frac{2^{g+n}}{\sqrt{N(D)}} \cdot \|\phi\|^2 \cdot |(\tilde{\phi}, P_\chi)|^2.$$

Here

$$(\tilde{\phi}, P_\chi) = \sum_{t \in T(F)\backslash T(\mathbb{A}_f)/U_T} \chi^{-1}(t) \tilde{\phi}(tP_c).$$

### Remark

There is a naive analogue between even and odd cases via Hodge theory which is actually a starting point to believe that there will be a simultaneous proof for both cases. To see this, let's consider the space  $Z(\Omega_X^1)$  of closed smooth 1-forms on a Shimura curve  $X$  of  $(N, K)$ -type with hermitian product defined by

$$(\alpha, \beta) = \frac{i}{2} \int \alpha \bar{\beta}.$$

The Hodge theory gives a decomposition of this space into a direct sum

$$Z(\Omega_X^1) = \bigoplus_{\alpha} \mathbb{C}\alpha \oplus (\text{continuous spectrum})$$

where  $\alpha$  runs through eigenforms under the Hecke operators and the Laplacian. Each  $\alpha$  is either holomorphic, anti-holomorphic or exact. In either case,  $\alpha$  corresponds to a test form  $\tilde{\phi}$  of weight 2,  $-2$ , or 0 on  $X$  in the following sense

$$\alpha = \begin{cases} \tilde{\phi} dz, & \text{if } \alpha \text{ is holomorphic,} \\ \tilde{\phi} d\bar{z}, & \text{if } \alpha \text{ is anti-holomorphic,} \\ d\tilde{\phi}, & \text{if } \alpha \text{ is exact.} \end{cases}$$

We may take integration  $c \mapsto \int_{\tilde{c}} \alpha$  to define a map

$$\pi_\alpha : \text{Div}^0(X) \otimes \mathbb{C} \longrightarrow \begin{cases} A_\phi \otimes \mathbb{C}, & \text{if } \alpha \text{ is holomorphic,} \\ \mathbb{C}, & \text{if } \alpha \text{ is exact.} \end{cases}$$

Here  $\tilde{c}$  is an 1-cycle on  $X$  with boundary  $c$ . In this manner, we have

$$\pi_\alpha(P_{\tilde{\chi}}) = \begin{cases} z_\phi, & \text{if } \alpha \text{ is holomorphic,} \\ (\phi, P_\chi), & \text{if } \alpha \text{ is exact.} \end{cases}$$

Thus we can think of  $\mathbb{C}$  as an abelian variety corresponding to  $\phi$  in the even case with Neron-Tate heights given by absolute value. This gives a complete analogue of the right hand side of the Gross-Zagier formulas in the even and odd case.

On the other hand, in the even case, one can define L-series  $L(s, \chi, \partial\phi/\partial z_v)$  by (4.1) where  $v$  is the only archimedean place where  $k_v = 0$ . It is not difficult to see that this L-series is essentially  $(s - 1/2)L(s, \chi, \phi)$ . Its derivative is given by  $L(1/2, \chi, \phi)$ . Thus we have an analogue of the left hand side as well!

## 8 Idea of Gross and Zagier

Let us describe the original idea of Gross and Zagier in the proof of a central derivative formula (for Heegner points on  $X_0(N)/\mathbb{Q}$  with square free discriminant  $D$ ) in their famous Inventiones paper. For simplicity, we fix  $N$ ,  $\chi$  and assume that  $\Sigma(N, K)$  is odd. For a holomorphic form  $\phi$  of weight 2, we define its *Fourier coefficient*  $\widehat{\phi}(a)$  at an integral idele  $a$  by the equation

$$(8.1) \quad W_\phi \begin{pmatrix} ay_\infty \delta^{-1} & 0 \\ 0 & 1 \end{pmatrix} = \widehat{\phi}(a) W_\infty \begin{pmatrix} y_\infty & 0 \\ 0 & 1 \end{pmatrix},$$

where  $W_\infty = \prod_{v \nmid \infty} W_v$  is the standard Whittaker function for weight 2 defined in (4.7).

With the notation of §6, there is a cusp form  $\Psi$  of level  $N$  whose Fourier coefficient is given by

$$(8.2) \quad \widehat{\Psi}(a) = |a| \langle x, T_a x \rangle.$$

This follows from the following two facts:

- the subalgebra  $\mathbb{T}'$  of  $\text{Jac}(X) \otimes \mathbb{C}$  generated by Hecke operators  $T_a$  is a quotient of the subalgebra  $\mathbb{T}$  in  $\text{End}(S_2(N))$  generated by Hecke operators  $T_a$ . Here  $S_2(N)$  is the space of holomorphic cusp forms of weight  $(2, \dots, 2)$ , level  $N$ , with trivial central character;
- any linear functional  $\ell$  of  $\mathbb{T}$  is represented by a cusp form  $f \in S_2(N)$  in the sense that  $|a|\ell(T_a) = \widehat{f}(a)$ .

(This form  $\Psi$  is not unique in general. But it is if we can normalize it to be a sum of new forms.)

It is then easy to see that

$$(8.3) \quad (\phi, \Psi) = \langle x_\phi, x_\phi \rangle(\phi, \phi).$$

Here  $(\phi, \Psi)$  is the inner product as in Theorem 6.1 which is the same as  $(\phi, \Psi)_{U_0(N)}$  in our notations in Introduction.

Thus, the question is reduced to showing that

$$(8.4) \quad L'(1/2, \chi, \phi) = \frac{2^{g+1}}{\sqrt{N(D)}}(\Psi, \phi).$$

On the other hand, one can express  $L(s, \chi, \phi)$  using a method of Rankin and Selberg:

$$(8.5) \quad L(s, \chi, \phi) = \frac{d_F^{1/2-s}}{|U_0(ND)|} \int_{\text{PGL}_2(F) \backslash \text{PGL}_2(\mathbb{A})} \phi(g)\theta(g)E(s, g)dg.$$

We need to explain various term in this integration.

First of all,  $\theta$  is a theta series associated to  $\chi$ . More precisely,  $\theta$  is an eigen form of weight  $(-1, \dots, -1)$ , level  $D$ , and central character  $\omega$  such that its local Whittaker functions  $W_v(g)$  produces the local L-functions for  $\chi$ :

$$(8.6) \quad |\delta_v|^{s-1/2} \int_{F_v^\times} W_v \begin{pmatrix} -a & 0 \\ 0 & 1 \end{pmatrix} |a|^{s-1/2} d^\times a = \prod_{w|v} L(s, \chi_w).$$

Here  $L(s, \chi_w)$  is defined in (5.3). It follows that the automorphic representation  $\Pi(\chi) := \Pi(\theta)$  generated by  $\theta$  is irreducible with newform  $\theta_\chi(g) = \theta(g\epsilon)$  where  $\epsilon = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$ .

Secondly,  $E(s, g) = E(s, g, \mathcal{F})$  is an Eisenstein series (2.6) for the quasi-characters  $|\cdot|^{s-1/2}$  and  $|\cdot|^{1/2-s}\omega$  with following specific  $\mathcal{F} = \otimes \mathcal{F}_v \in \mathcal{S}(\mathbb{A}^2)$ . If  $v$  is a finite place, then

$$(8.7) \quad \mathcal{F}_v(x, y) = \begin{cases} 1, & \text{if } v \nmid c(\omega), |x| \leq |ND|_v, |y| \leq 1, \\ \omega_v^{-1}(y), & \text{if } v \mid c(\omega), |x| \leq |ND|_v, |y| = 1, \\ 0, & \text{otherwise.} \end{cases}$$

If  $v$  is an infinite place, then

$$(8.8) \quad \mathcal{F}_v(x, y) = (\pm ix + y)e^{-\pi(x^2+y^2)}$$

where we take + sign (resp. -) if  $k_v = 2$  (resp.  $k_v = 0$ ).

Taking a trace, we obtain a form of level  $N$ :

$$(8.9) \quad \Phi_s(g) = \text{tr}_D \Phi_s^\sharp(g) = \sum_{\gamma \in U_0(D)/U_0(ND)} \overline{d_F^{1/2-s} \theta(g\gamma) E(s, g\gamma)}.$$

This form has the property:

$$(8.10) \quad L(s, \chi, \phi) = (\phi, \Phi_s)_{U_0(N)}.$$

The idea of Gross and Zagier (in the odd case) is to compute the derivative  $\Phi'_{1/2}$  of  $\Phi_s$  with respect to  $s$  at  $s = 1/2$  and take a holomorphic projection to obtain a holomorphic form  $\Phi$  so that

$$(8.11) \quad L'(1/2, \chi, \phi) = (\phi, \Phi)_{U_0(N)}.$$

(See [4] for a direct construction of the kernel using Poincaré series instead of Rankin-Selberg method and holomorphic projection.) Now the problem is reduced to proving that

$$\Phi - \frac{2^{g+1}}{\sqrt{N(D)}} \Psi$$

is an old form. In other words, we need to show that the Fourier coefficients of  $\Phi$  are given by height pairings of Heegner points on  $\text{Jac}(X)$ :

$$(8.12) \quad \widehat{\Phi}(a) = |a| \langle x, T_a x \rangle$$

for any finite integral ideles  $a$  prime to  $ND$ .

One expects to prove the above equality by explicit computations for both sides respectively. These computations have been successfully carried out by Gross and Zagier [6] when  $F = \mathbb{Q}$ ,  $D$  is square free, and  $X(N, K) = X_0(N)$ . The computation of Fourier coefficients of  $\Phi$  is essentially straightforward and has been carried out for totally real fields [17]. For the computation of  $\langle x, T_a x \rangle$ , Gross and Zagier represented the Hodge class  $\xi$  by cusps 0 and  $\infty$  on  $X_0(N)$ :

$$(8.13) \quad \langle x, T_a x \rangle = \langle P_\chi - h_\chi 0, T_a(P_\chi) - h_{\chi,a} \infty \rangle$$

where  $h_\chi$  and  $h_{\chi,a}$  are integers to make both divisors to have degree 0. The right hand side can be further decomposed into local height pairing by deforming self-intersections using Dedekind  $\eta$ -functions. These local height pairings can be finally computed by a modular interpretation in terms of deformation of formal groups.

When  $F$  is arbitrary (even when  $D$  is square free), the computation of heights has a lot of problems as there is no canonical representatives for the Hodge class, and no canonical modular form for self-intersections. In §9-10, we will see how Arakelov theory been used to compute the heights.

When  $D$  is arbitrary (even when  $F = \mathbb{Q}$ ), the computations of both kernels and heights for Theorem 6.1 seem impossible to carry out directly because of *singularities* in both analysis and geometry. Alternatively, we will actually prove a Gross-Zagier formula for level  $ND$  (§11-16) and try to reduce the level by using continuous spectrum (§17-19).

## 9 Calculus on arithmetic surfaces

The new idea in our Annals paper ([17]) is to use Arakelov theory to decompose the heights of Heegner points as locally as possible, and to show that the contribution of these terms which we don't know how to compute are *negligible*.

Let  $F$  be a number field. By an arithmetic surface over  $\text{Spec} \mathcal{O}_F$ , we mean a projective and flat morphism  $\mathcal{X} \rightarrow \text{Spec} \mathcal{O}_F$  such that that  $\mathcal{X}$  is a regular scheme of dimension 2. Let  $\widehat{\text{Div}}(\mathcal{X})$  denote the group of *arithmetic divisors* on  $\mathcal{X}$ . Recall that an arithmetic divisor on  $\mathcal{X}$  is a pair  $\widehat{D} := (D, g)$  where  $D$  is a divisor on  $\mathcal{X}$  and  $g$  is a function on

$$X(\mathbb{C}) = \coprod X_\tau(\mathbb{C})$$

with some logarithmic singularities on  $|D|$ . The form  $-\frac{\partial\bar{\partial}}{\pi i}g$  on  $X(\mathbb{C})-|D|$  can be extended to a smooth form  $c_1(\widehat{D})$  on  $X(\mathbb{C})$  which is called the *curvature* of the divisor  $\widehat{D}$ . If  $f$  is a nonzero rational function on  $\mathcal{X}$  then we can define the corresponding *principal arithmetic divisor* by

$$(9.1) \quad \widehat{\operatorname{div}} f = (\operatorname{div} f, -\log |f|).$$

An arithmetic divisor  $(D, g)$  is called *vertical* (resp. *horizontal*) if  $D$  is supported in the special fibers (resp.  $D$  does not have component supported in the special fiber).

The group of arithmetic divisors is denoted by  $\widehat{\operatorname{Div}}(\mathcal{X})$  while the subgroup of principal divisor is denoted by  $\widehat{\operatorname{Pr}}(\mathcal{X})$ . The quotient  $\widehat{\operatorname{Cl}}(\mathcal{X})$  of these two groups is called the *arithmetic divisor class group* which is actually isomorphic to the group  $\widehat{\operatorname{Pic}}(\mathcal{X})$  of hermitian line bundles on  $\mathcal{X}$ . Recall that a hermitian line bundle on  $\mathcal{X}$  is a pair  $\overline{\mathcal{L}} = (\mathcal{L}, \|\cdot\|)$ , where  $\mathcal{L}$  is a line bundle on  $\mathcal{X}$  and  $\|\cdot\|$  is hermitian metric on  $\mathcal{L}(\mathbb{C})$  over  $X(\mathbb{C})$ . For a rational section  $\ell$  of  $\mathcal{L}$ , we can define the corresponding divisor by

$$(9.2) \quad \widehat{\operatorname{div}}(\ell) = (\operatorname{div}\ell, -\log \|\ell\|).$$

It is easy to see that the divisor class of  $\widehat{\operatorname{div}}(\ell)$  does not depend on the choice of  $\ell$ . Thus one has a well defined map from  $\widehat{\operatorname{Pic}}(\mathcal{X})$  to  $\widehat{\operatorname{Cl}}(\mathcal{X})$ . This map is actually an isomorphism.

Let  $\widehat{D}_i = (D_i, g_i)$  ( $i = 1, 2$ ) be two arithmetic divisors on  $\mathcal{X}$  with disjoint support in the generic fiber:

$$|D_{1F}| \cap |D_{2F}| = \emptyset.$$

Then one can define an *arithmetic intersection pairing*

$$(9.3) \quad \widehat{D}_1 \cdot \widehat{D}_2 = \sum_v (\widehat{D}_1 \cdot \widehat{D}_2)_v,$$

where  $v$  runs through the set of places of  $F$ . The intersection pairing only depends on the divisor class. It follows that we have a well defined pairing on  $\widehat{\operatorname{Pic}}(\mathcal{X})$ :

$$(9.4) \quad (\overline{\mathcal{L}}, \overline{\mathcal{M}}) \longrightarrow \widehat{c}_1(\overline{\mathcal{L}}) \cdot \widehat{c}_1(\overline{\mathcal{M}}) \in \mathbb{R}.$$

Let  $V(\mathcal{X})$  be the group of *vertical metrized line bundles*: namely  $\overline{\mathcal{L}} \in \widehat{\text{Pic}}(\mathcal{X})$  with  $\mathcal{L} \simeq \mathcal{O}_X$ . Then we have an exact sequence

$$0 \longrightarrow V(\mathcal{X}) \longrightarrow \widehat{\text{Pic}}(\mathcal{X}) \longrightarrow \text{Pic}(\mathcal{X}_F) \longrightarrow 0.$$

Define the group of *flat bundles*  $\widehat{\text{Pic}}^0(\mathcal{X})$  as the orthogonal complement of  $V(\mathcal{X})$ . Then we have an exact sequence

$$0 \longrightarrow \widehat{\text{Pic}}(\mathcal{O}_F) \longrightarrow \widehat{\text{Pic}}^0(\mathcal{X}) \longrightarrow \text{Pic}^0(X_F) \longrightarrow 0.$$

The following formula, which is referred as *Hodge index theorem*, gives a relation between intersection pairing and height pairing: for  $\overline{\mathcal{L}}, \overline{\mathcal{M}} \in \widehat{\text{Pic}}^0(\mathcal{X})$ ,

$$(9.5) \quad \langle \mathcal{L}_F, \mathcal{M}_F \rangle = -\widehat{c}_1(\overline{\mathcal{L}}) \cdot \widehat{c}_1(\overline{\mathcal{M}}),$$

where the left hand side denotes the Neron-Tate height pairing on  $\text{Pic}^0(X) = \text{Jac}(X)(F)$ .

For  $X$  a curve over  $F$ , let  $\widehat{\text{Pic}}(X)$  denote the direct limit of  $\widehat{\text{Pic}}(\mathcal{X})$  over all models over  $X$ . Then the intersection pairing can be extended to  $\widehat{\text{Pic}}(X)$ . Let  $\overline{F}$  be an algebraic closure of  $F$  and let  $\widehat{\text{Pic}}(X_{\overline{F}})$  be the direct limit of  $\text{Pic}(X_L)$  for all finite extensions  $L$  of  $F$ , then the intersection pairing on  $\widehat{\text{Pic}}(X_L)$  times  $[L : F]^{-1}$  can be extended to an intersection pairing on  $\widehat{\text{Pic}}(X_{\overline{F}})$ .

Let  $\overline{\mathcal{L}} \in \widehat{\text{Pic}}(\mathcal{X})_{\mathbb{Q}}$  be a fixed class with degree 1 at the generic fiber. Let  $x \in X(F)$  be a rational point and let  $\bar{x}$  be the corresponding section  $\mathcal{X}(\mathcal{O}_F)$ . Then  $\bar{x}$  can be extended to a unique element  $\widehat{x} = (x + D, g)$  in  $\widehat{\text{Div}}(\mathcal{X})_{\mathbb{Q}}$  satisfying the following conditions:

- the bundle  $\mathcal{O}(\widehat{x}) \otimes \overline{\mathcal{L}}^{-1}$  is flat;
- for any finite place  $v$  of  $F$ , the component  $D_v$  of  $D$  on the special fiber of  $\mathcal{X}$  over  $v$  satisfies

$$D_v \cdot c_1(\mathcal{L}) = 0;$$

- for any infinite place  $v$ ,

$$\int_{X_v(\mathbb{C})} gc_1(\overline{\mathcal{L}}) = 0.$$

We define now *Green's function*  $g_v(x, y)$  on

$$X(F) \times X(F) - \text{diagonal}$$

by

$$(9.6) \quad g_v(x, y) = (\widehat{x} \cdot \widehat{y})_v / \log q_v,$$

where  $\log q_v = 1$  or  $2$  if  $v$  is real or complex. It is easy to see that  $g_v(x, y)$  is symmetric, and does not depend on the model  $\mathcal{X}$  of  $X$ , and is stable under base change. Thus we have a well-defined Green's function on  $X(\bar{F})$  for each place  $v$  of  $F$ .

## 10 Decomposition of Heights

We now want to apply the general theory of the previous section to intersections of CM-points to Shimura curves  $X = X(N, K)$  over a totally real field  $F$  as defined in §6. Recall that  $X$  has the form

$$(10.1) \quad X = G(F)_+ \backslash \mathcal{H} \times G(\mathbb{A}_f) / U(N, K) \cup \{\text{cusps}\}$$

which is a smooth and projective curve over  $F$  but may not be connected.

To define Green's function we need to extend the Hodge class  $\xi$  in  $\text{Pic}(X)_{\mathbb{Q}}$  to a class in  $\widehat{\text{Pic}}(X) \otimes \mathbb{Q}$ . Notice that  $\xi \in \text{Pic}(X)_{\mathbb{Q}}$  is *Eisenstein* under the action of Hecke operators:

$$(10.2) \quad T_a \xi = \sigma_1(a) \cdot \xi, \quad \sigma_1(a) := \deg T_a = \sum_{b|a} N(b),$$

for any integral idele  $a$  prime to the level of  $X$ .

It is an interesting question to construct a class  $\widehat{\xi}$  to extend  $\xi$  such that the above equation holds for  $\widehat{\xi}$ . But in [17], Corollary 4.3.3, we have constructed an extension  $\widehat{\xi}$  of  $\xi$  such that

$$(10.3) \quad T_a \widehat{\xi} = \sigma_1(a) \widehat{\xi} + \phi(a)$$

where  $\phi(a) \in \widehat{\text{Pic}}(F)$  is a  $\sigma_1$ -derivation, i.e., for any coprime  $a', a''$

$$\phi(a'a'') = \sigma(a')\phi(a'') + \sigma(a'')\phi(a').$$

We have the following a general definition.

**Definition 10.1.** Let  $\mathbb{N}_F$  denote the semigroup of nonzero ideals of  $\mathcal{O}_F$ . For each  $a \in \mathbb{N}_F$ , let  $|a|$  denote the inverse norm of  $a$ :

$$|a|^{-1} = \#\mathcal{O}_F/a.$$

For a fixed ideal  $M$ , let  $\mathbb{N}_F(M)$  denote the sub-semigroup of ideals prime to  $M$ .

A function  $f$  on  $\mathbb{N}_F(M)$  is called quasi-multiplicative if

$$f(a_1a_2) = f(a_1) \cdot f(a_2)$$

for all coprime  $a_1, a_2 \in \mathbb{N}_F(M)$ . For a quasi-multiplicative function  $f$ , let  $\mathcal{D}(f)$  denote the set of all  $f$ -derivations, that is the set of all a linear combinations

$$g = cf + h$$

where  $c$  is a constant, and where  $h$  satisfies

$$h(a_1a_2) = h(a_1)f(a_2) + h(a_2)f(a_1)$$

for all  $a_1, a_2 \in \mathbb{N}_F(M)$  with  $(a_1, a_2) = 1$ .

For a representation  $\Pi$ , the Fourier coefficients  $\widehat{\Pi}(a)$  is defined to be

$$\widehat{\Pi}(a) := W_{\Pi, f} \begin{pmatrix} a\delta^{-1} & 0 \\ 0 & 1 \end{pmatrix},$$

where  $W_{\Pi, f}$  is the product of Whittaker newvectors at finite places. In other words,  $\widehat{\Pi}(a)$  is defined such that the finite part of  $L$ -series has expansion

$$L_f(s, \Pi) = \sum \widehat{\Pi}(a)|a|^{s-1/2}.$$

Then  $\widehat{\Pi}(a)$  is quasi-multiplicative.

We can now define Green's functions  $g_v$  on divisors on  $X(\bar{F})$  which are disjoint at the generic fiber for each place  $v$  of  $F$ . Let's try to decompose the heights of our Heegner points. The linear functional

$$a \longrightarrow |a|\langle x, \mathbb{T}_a x \rangle$$

is now the Fourier coefficient of a cuspform  $\Psi$  of weight 2:

$$(10.4) \quad \widehat{\Psi}(a) = |a|\langle x, \mathbb{T}_a x \rangle.$$

In the following we want to express this height in terms of intersections modulo some Eisenstein series and theta series.

Let  $\widehat{P}_\chi$  be the arithmetic closure of  $P_\chi$  with respect to  $\widehat{\xi}$ . Then the Hodge index formula (9.5) gives

$$(10.5) \quad \begin{aligned} |a|\langle x, T_a x \rangle &= -|a| \left( \widehat{P}_\chi - \deg(P_\chi)\widehat{\xi}, T_a \widehat{P}_\chi - \deg(T_a \widehat{P}_\chi)\widehat{\xi} \right) \\ &= -|a|(\widehat{P}_\chi, T_a \widehat{P}_\chi) + \widehat{E}(a), \end{aligned}$$

where  $\widehat{E}$  is a certain derivations of Eisenstein series.

The divisor  $P_\chi$  and  $T_a P_\chi$  have some common components. We want to compute its contribution in the intersections. Let  $r_\chi(a)$  denote the Fourier coefficients of the theta series associated to  $\chi$ :

$$(10.6) \quad r_\chi(a) = \sum_{b|a} \chi(b).$$

Then the divisor

$$(10.7) \quad T_a^0 P_\chi := T_a P_\chi - r_\chi(a) P_\chi$$

is disjoint with  $P_\chi$ .

It follows that  $\widehat{\Psi}(a)$  is essentially given by a sum of local intersections

$$-\frac{1}{[L:F]} \sum_v \sum_{\iota \in \text{Gal}(H_c/F)} g_v(T_a^0 P_\chi^\iota, P_\chi^\iota) |a| \log q_v$$

modulo some derivations of Eisenstein series, and theta series of weight 1. We can further simplify this sum by using the fact that the Galois action of  $\text{Gal}(K^{\text{ab}}/F)$  is given by the composition of the class field theory map

$$\nu : \text{Gal}(K^{\text{ab}}/F) \longrightarrow N_T(F) \backslash N_T(\mathbb{A}_f),$$

and the left multiplication of the group  $N_T(\mathbb{A}_f)$ . Finally we obtain the following:

$$(10.8) \quad \widehat{\Psi}(a) = -|a| \sum_v g_v(P_\chi, T_a^0 P_\chi) \log q_v \pmod{\mathcal{D}(\sigma_1) + \mathcal{D}(r_\chi)}.$$

Assume that  $D$  is square free, and that  $X(N, K)$  has regular canonical integral model over  $\mathcal{O}_K$ , which is the case when  $\text{ord}_v(N) = 1$  if  $v$  is not split

in  $K$  or  $v \mid 2$ . We can use the theory of Gross on canonical or quasi-canonical lifting to compute  $g_v(P_\chi, T_a^0 P_\chi)$  and to prove that the functional

$$\widehat{\Phi} - \frac{2^{g+1}}{\sqrt{N(D)}} \widehat{\Psi}$$

vanishes modulo derivations of Eisenstein series or theta series. It then follows that this functional is actually zero by the following lemma:

**Lemma 10.2.** *Let  $f_1, \dots, f_r$  be distinct quas-multiplicative functions on  $\mathbb{N}_F(ND)$  then the sum*

$$\mathcal{D}(f_1) + \mathcal{D}(f_2) + \dots + \mathcal{D}(f_r)$$

*is a direct sum.*

Thus we can prove the Gross-Zagier formula in the case  $D$  is square free and  $X(N, K)$  has regular canonical model over  $\mathcal{O}_K$ . This is the main result in our Annals paper [17].

## 11 Construction of the kernels

From this section to the end, we want to explain the idea to prove the Gross-Zagier formula in §6-7 for the general case. We will start with a construction of kernels. As explained early, there is no good construction of kernels in level  $N$ . The best we can do is to construct some *nice kernel* in level  $ND$  in the sense that the Fourier coefficients are *symmetric* and easy to compute, and that the projection of this form in  $\Pi(\phi)$  is *recognizable*.

Recall from (8.5) that we have an integral expression of Rankin-Selberg convolution:

$$(11.1) \quad L(s, \chi, \phi) = \frac{|\delta|^{s-1/2}}{|U_0(ND)|} \int \phi(g)\theta(g)E(s, g)dg$$

To obtain a more symmetric kernel we have to apply Atkin-Lehner operators to  $\theta(g)E(s, g)$ . Let  $S$  be the set of finite places ramified in  $K$ . For each such set  $T$  of  $S$ , let  $h_T$  be an element in  $\mathrm{GL}_2(\mathbb{A})$  which has component 1 outside  $T$  and has component  $\begin{pmatrix} 0 & 1 \\ -t_v & 0 \end{pmatrix}$  where  $t_v$  has the same order as  $c(\omega_v)$  and such that  $\omega_v(t_v) = 1$ . Now one can show that

$$(11.2) \quad L(s, \chi, \phi) = \frac{\gamma_T(s)}{|U_0(ND)|} \int \phi(g)\theta(gh_T^{-1}\epsilon)E(s, gh_T^{-1})dg$$

where  $\gamma_T$  is certain exponential function of  $s$ . Finally we define the kernel function

$$\Theta(s, g) = 2^{-|S|} \sum_{T \subset S} \gamma_T(s) \theta(gh_T^{-1}) E(s, gh_T^{-1}).$$

By our construction,

$$L(s, \chi, \phi) = (\phi, \bar{\Theta}(s, -))_{U_0(ND)}.$$

Now the functional equation of  $L(s, \chi, \phi)$  follows from the following equation of the kernel function which can be proved by a careful analysis of Atkin-Lehner operators:

$$(11.3) \quad \Theta(s, g) = \epsilon(s, \chi, \phi) \Theta(1-s, g).$$

Assume that  $\phi$  is cuspidal, then we may define the projection of  $\bar{\Theta}$  in  $\Pi(\phi)$  as a form  $\varphi \in \Pi(\phi)$  such that

$$\int f \Theta dg = \int f \bar{\varphi} dg, \quad \forall f \in \Pi.$$

Since the kernel  $\Theta(s, g)$  constructed above has level  $ND$ , its projection onto  $\Pi$  will have level  $DN$  and thus is a linear combination of the forms

$$(11.4) \quad \phi_a := \rho \begin{pmatrix} a^{-1} & 0 \\ 0 & 1 \end{pmatrix} \phi, \quad (a \mid D).$$

**Proposition 11.1.** *The projection of  $\bar{\Theta}(s, g)$  on  $\Pi$  is given by*

$$\frac{L(s, \chi, \phi)}{(\phi_s^\sharp, \phi_s^\sharp)_{U_0(ND)}} \cdot \phi_s^\sharp,$$

where  $\phi_s^\sharp$  is the unique nonzero form in the space of  $\Pi(\phi)$  of level  $ND$  satisfying the following identities:

$$(\phi_s^\sharp, \phi_a) = \nu^*(a) (\phi_s^\sharp, \phi_s^\sharp), \quad (a \mid D),$$

where

$$\nu^*(a)_s = \prod_{v \mid S} \frac{|a|_v^{s-1/2} + |a|_v^{1/2-s}}{2} \begin{cases} \nu(a), & \text{if } a \mid c(\omega), \\ 0, & \text{otherwise.} \end{cases}$$

Write  $\phi^\sharp = \phi_{1/2}^\sharp$  and call it the *quasi-newform* with respect to  $\chi$ .  
The function  $\Theta(s, g)$  has a Fourier expansion

$$(11.5) \quad \Theta(s, g) = C(s, g) + \sum_{\alpha \in F^\times} W \left( s, \begin{pmatrix} \alpha & 0 \\ 0 & 1 \end{pmatrix} g \right).$$

Since  $\Theta(s, g)$  is a linear combination of the form

$$\Theta(s, g) = \sum_i \theta_i(g) E_i(g)$$

with  $\theta_i \in \Pi(\chi)$  and  $E_i(g) \in \Pi(|\cdot|^{s-1/2}, |\cdot|^{1/2-s}\omega)$ , the constant and Whittaker function of  $\Theta(s, g)$  can be expressed precisely in terms of Fourier expansion of  $\theta_i$  and  $E_i(g)$ .

More precisely, let

$$(11.6) \quad \theta_i(g) = \sum_{\xi \in F} W_{\theta_i}(\xi, g), \quad E_i(g) = \sum_{\xi \in F} W_{E_i}(\xi, g),$$

be Fourier expansions of  $\theta_i$  and  $E_i$  respectively into characters  $\begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \mapsto \psi(\xi x)$  on  $N(\mathbb{A})$ . Then

$$(11.7) \quad C(s, g) = \sum_{\xi \in F} C(s, \xi, g),$$

$$(11.8) \quad W(s, g) = \sum_{\xi \in F} W(s, \xi, g),$$

where

$$(11.9) \quad C(s, \xi, g) = \sum_i W_{\theta_i}(-\xi, g) W_{E_i}(\xi, g),$$

and

$$(11.10) \quad W(s, \xi, g) = \sum_i W_{\theta_i}(1 - \xi, g) W_{E_i}(\xi, g).$$

The behavior of the degenerate term  $C(s, \xi, g)$  can be understood very well. The computation shows that the complex conjugation of  $\Theta(s, g)$  is finite

at each cusp unless  $\chi$  is a form  $\nu \circ N_{K/F}$  in which case, we need to remove two Eisenstein series in the space

$$E_1 \in \Pi(\|\cdot\|^s, \|\cdot\|^{-s}) \otimes \nu, \quad E_2 \in \Pi(\|\cdot\|^{1-s}, \|\cdot\|^{s-1}) \otimes \nu\omega.$$

We let  $\Phi(s, g)$  denote  $\bar{\Theta}(1/2, g)$  if  $\chi$  is not of form  $\nu \circ N_{K/F}$ , or

$$\bar{\Theta}(s, g) - E_1 - E_2$$

if it is. Then  $\Phi(s, g)$  is a form with following growth:

$$\Phi\left(s, \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix}\right) = c_1(g)|a|^{s-1/2} + c_2(g)|a|^{1/2-s} + O_g(e^{-\epsilon|a|})$$

where  $c_1(g)$ ,  $c_2(g)$ , and  $O_g$  term are all smooth functions of  $g$  and  $s$ . It follows that the value or all derivatives of  $\Phi(s, g)$  at  $s = 1/2$  are  $L^2$ -forms.

With our very definition of  $\Theta(s, g)$  in the last section, we are able to decompose the non-degenerate term:

$$(11.11) \quad W(s, \xi, g) = \otimes W_v(s, \xi_v, g_v).$$

An explicit computation has given the following local functional equation:

$$(11.12) \quad W_v(s, \xi_v, g_v) = \omega_v(1 - \xi_v^{-1})(-1)^{\#\Sigma \cap \{v\}} W_v(1 - s, \xi_v, g_v).$$

If  $\Sigma$  is even, then we can compute the Fourier coefficients of  $\Theta(1/2, g)$  for  $g = \begin{pmatrix} a\delta^{-1} & 0 \\ 0 & 1 \end{pmatrix}$  very explicitly. The computation of non-degenerate term  $W(s, \xi, g)$  is reduced to local terms  $W_v(s, \xi, g)$ . By the functional equation, we need only consider those  $\xi$  such that Equivalently,

$$(11.13) \quad 1 - \xi^{-1} \in N(K_v^\times) \iff v \notin \Sigma.$$

The form  $\Phi$  is holomorphic of weight 2 at infinite places where  $\Pi$  is of weight 2.

If  $\Sigma$  is odd, then by functional equation,  $\Theta(1/2, g) = 0$ . We want to compute its derivative  $\Theta'(1/2, g)$  at  $s = 1/2$ . Let's now describe the central derivative for  $W(s, \xi, g)$  for  $g$  of the form  $\begin{pmatrix} a\delta^{-1} & 0 \\ 0 & 1 \end{pmatrix}$ . Recall that  $W(s, \xi, g)$  is

a product of  $W_v(s, \xi, g)$ , and that  $W_v(s, \xi, g)$  satisfies the functional equation (11.12). It follows that

$$(11.14) \quad W'(1/2, g) = \sum_v W'(1/2, g)_v,$$

where  $v$  runs through the places which are not split in  $K$  with

$$(11.15) \quad W'(1/2, g)_v = \sum_{\xi} W^v(1/2, \xi, g^v) \cdot W'_v(1/2, \xi, g).$$

Here

- $W^v$  is the product of  $W_{\ell}$  over places  $\ell \neq v$ , and
- $W'_v$  is the derivative for the variable  $s$ , and
- $\xi \in F - \{0, 1\}$  satisfies

$$(11.16) \quad 1 - \xi^{-1} \in N(K_w^{\times}) \iff w \notin_v \Sigma,$$

with  $_v\Sigma$  given by

$${}_v\Sigma = \begin{cases} \Sigma \cup \{v\}, & \text{if } v \notin \Sigma, \\ \Sigma - \{v\}, & \text{if } v \in \Sigma. \end{cases}$$

All these terms can be computed explicitly. We need to find the holomorphic projection of  $\bar{\Theta}'(1/2, g)$ . That is a holomorphic cusp form  $\Phi$  of weight 2 such that  $\bar{\Theta}'(1/2, g) - \Phi$  is perpendicular to any holomorphic form.

**Proposition 11.2.** *With respect to the standard Whittaker function for holomorphic weight 2 forms, the  $a$ -th Fourier coefficients  $\widehat{\Phi}(a)$  of the holomorphic projection  $\Phi$  of  $\bar{\Theta}'(1/2, g)$  is a sum*

$$\widehat{\Phi}(a) = A(a) + B(a) + \sum_v \widehat{\Phi}_v(a)$$

where

$$A \in \mathcal{D}(\widehat{\Pi}(\chi) \otimes \alpha^{1/2}),$$

$$B \in \mathcal{D}(\widehat{\Pi}(\alpha^{1/2}\nu, \alpha^{-1/2}\nu)) + \mathcal{D}(\widehat{\Pi}(\alpha^{1/2}\nu\omega, \alpha^{-1/2}\nu\omega)),$$

and the sum is over places of  $F$  which are not split in  $K$ , with  $\widehat{\Phi}_v(a)$  given by the following formulas:

1. if  $v$  is a finite place then  $\widehat{\Phi}_v(a)$  is a sum over  $\xi \in F$  with  $0 < \xi < 1$  of the following terms:

$$(2i)^g |(1 - \xi)\xi|_\infty^{1/2} \cdot \bar{W}_f^v \left( 1/2, \xi, \begin{pmatrix} a\delta_f^{-1} & 0 \\ 0 & 1 \end{pmatrix} \right) \cdot \bar{W}'_v \left( 1/2, \xi, \begin{pmatrix} a\delta_f^{-1} & 0 \\ 0 & 1 \end{pmatrix} \right).$$

2. if  $v$  is an infinite place, then  $\widehat{\Phi}_v(a)$  is the constant term at  $s = 0$  of a sum over  $\xi \in F$  such that  $0 < \xi_w < 1$  for all infinite place  $w \neq v$  and  $\xi_v < 0$  of the following terms:

$$(2i)^g |\xi(1 - \xi)|_\infty^{1/2} \cdot \bar{W}_f \left( 1/2, \xi, \begin{pmatrix} a\delta_f^{-1} & 0 \\ 0 & 1 \end{pmatrix} \right) \cdot \int_1^\infty \frac{-dx}{x(1 + |\xi|_v x)^{1+s}}.$$

## 12 Geometric pairing

The key to prove the Gross-Zagier formula is to compare the Fourier coefficients of the kernel functions and the local heights of CM-points. These local heights are naturally grouped by definite quaternion algebras which are the endomorphism rings of the supersingular points in the reductions of modular or Shimura curves. Furthermore, the intersection of two CM-points at supersingular points is given by a *multiplicity function* which depends only on the *relative position* of these two CM-points. In this section, we would like to abstractly define this kind of pairing with respect to an arbitrary multiplicity function. We will describe the relative position of two CM-points by a certain parameter  $\xi$  which will relate the same parameter in the last section by a *local Gross-Zagier formula*.

Let  $G$  be an inner form of  $\mathrm{PGL}_2$  over  $F$ . This means that  $G = B^\times / F^\times$  with  $B$  a quaternion algebra over  $F$ . Let  $K$  be a totally imaginary quadratic extension of  $F$  which is embedded into  $B$ . Let  $T$  denote the subgroup of  $G$  given by  $K^\times / F^\times$ . Then the set

$$(12.1) \quad C := T(F) \backslash G(\mathbb{A}_f)$$

is called the *set of CM-points*. This set admits a natural action by  $T(\mathbb{A}_f)$  (resp.  $G(\mathbb{A}_f)$ ) by left (resp. right) multiplications.

There is a map from  $C$  to the Shimura variety defined by  $G$

$$(12.2) \quad \iota : C \longrightarrow M := G(F)_+ \backslash \mathcal{H}^n \times G(\mathbb{A}_f)$$

as in §6-7 which sending the class of  $g \in G(\mathbb{A}_f)$  to the class of  $(h_0, g)$ , where  $h_0 \in \mathcal{H}^n$  is fixed by  $T$ . This map is an embedding if  $G$  is not totally definite.

The set of CM-points has a topology induced from  $G(\mathbb{A}_f)$  and has a unique  $G(\mathbb{A}_f)$ -invariant measure  $dx$  induced from the one on  $G(\mathbb{A}_f)$ . The space

$$\mathcal{S}(C) = \mathcal{S}(T(F)\backslash G(\mathbb{A}_f))$$

of locally constant functions with compact support is called the space of *CM-cycles* which admits a natural action by  $T(\mathbb{A}_f) \times G(\mathbb{A}_f)$ . There is a natural pairing between functions  $f$  on Shimura variety  $M$  and CM-cycles  $\alpha$  by

$$(12.3) \quad (f, \alpha) = \int_C \bar{\alpha}(x) f(\iota x) dx.$$

Thus CM-cycles may serve as distributions or functionals on the space of functions on  $M$ . Of course this pairing is invariant under the action by  $G(\mathbb{A}_f)$ .

Since  $T(F)\backslash T(\mathbb{A}_f)$  is compact, one has a natural decomposition

$$\mathcal{S}(C) = \oplus_{\chi} \mathcal{S}(\chi, C)$$

where the sum is over the characters of  $T(F)\backslash T(\mathbb{A}_f)$ . There is also a local decomposition for each character  $\chi$ :

$$(12.4) \quad \mathcal{S}(\chi, C) = \otimes_v \mathcal{S}(\chi_v, G(F_v)).$$

In the following we will define a class of pairings on CM-cycles which are *geometric* since it appears naturally in the local intersection pairing of CM-points on Shimura curves. To do this, let's write CM-points in a slightly different way,

$$(12.5) \quad C = G(F)\backslash(G(F)/T(F)) \times G(\mathbb{A}_f),$$

then the topology and measure of  $C$  is still induced by those of  $G(\mathbb{A}_f)$  and the *discrete* ones of  $G(F)/T(F)$ .

Let  $m$  be a *real valued* function on  $G(F)$  which is  $T(F)$ -invariant and such that  $m(\gamma) = m(\gamma^{-1})$ . Then  $m$  can be extended to  $G(F)/T(F) \times G(\mathbb{A}_f)$  such that

$$(12.6) \quad m(\gamma, g_f) = \begin{cases} m(\gamma), & \text{if } g_f = 1, \\ 0, & \text{otherwise.} \end{cases}$$

We now have a kernel function

$$(12.7) \quad k(x, y) = \sum_{\gamma \in G(F)} m(x^{-1}\gamma y)$$

on  $C \times C$ . Then we can define a pairing on  $\mathcal{S}(C)$  by

$$(12.8) \quad \begin{aligned} \langle \alpha, \beta \rangle &= \int_{C^2} \alpha(x)k(x, y)\bar{\beta}(y)dx dy \\ &:= \lim_{U \rightarrow 1} \int_{C^2} \alpha(x)k_U(x, y)\bar{\beta}(y)dx dy \end{aligned}$$

where  $U$  runs through the open subgroup of  $G(\mathbb{A}_f)$  and

$$k_U(x, y) = \text{vol}(U)^{-2} \int_{U^2} k(xu, yv)dudv.$$

This pairing is called a *geometric pairing with multiplicity function  $m$* . For two function  $\alpha$  and  $\beta$  in  $\mathcal{S}(T(F)\backslash G(\mathbb{A}_f))$ , one has

$$(12.9) \quad \langle \alpha, \beta \rangle = \sum_{\gamma \in T(F)\backslash G(F)/T(F)} m(\gamma)\langle \alpha, \beta \rangle_\gamma$$

where

$$(12.10) \quad \langle \alpha, \beta \rangle_\gamma = \int_{T_\gamma(F)\backslash G(\mathbb{A}_f)} \alpha(\gamma y)\bar{\beta}(y)dy,$$

and where

$$(12.11) \quad T_\gamma := \gamma^{-1}T\gamma \cap T = \begin{cases} T & \text{if } \gamma \in N_T, \\ 1 & \text{otherwise,} \end{cases}$$

and where  $N_T$  is the normalizer of  $T$  in  $G$ . The integral  $\langle \alpha, \beta \rangle_\gamma$  is called the *linking number* of  $\alpha$  and  $\beta$  at  $\gamma$ .

Since both  $\alpha$  and  $\beta$  are invariant under left-translation by  $T(F)$ , the linking number at  $\gamma$  depends only on the class of  $\gamma$  in  $T(F)\backslash G(F)/T(F)$ . Lets define a parameterization of this sets by the following function by writing  $B = K + K\epsilon$  where  $\epsilon \in B$  is an element such that  $\epsilon^2 \in F^\times$  and  $\epsilon x = \bar{x}\epsilon$ . Then the function

$$\xi(a + b\epsilon) = \frac{N(b\epsilon)}{N(a + b\epsilon)}$$

defines an embedding

$$(12.12) \quad \xi : T(F) \backslash G(F) / T(F) \longrightarrow F.$$

Write

$$(12.13) \quad \langle \alpha, \beta \rangle_\gamma = \langle \alpha, \beta \rangle_\xi.$$

Notice that  $\xi(\gamma) = 0$  (resp. 1) iff  $\xi \in T$  (resp.  $\xi \in N_T - T$ ). The image of  $G(F) - N_T$  is the set of  $\xi \in F$  such that  $\xi \neq 0, 1$  and where for any place  $v$  of  $F$ ,

$$(12.14) \quad 1 - \xi^{-1} \in \begin{cases} \mathbf{N}(K_v^\times), & \text{if } B_v \text{ is split,} \\ F_v^\times - \mathbf{N}(K_v^\times) & \text{if } B_v \text{ is not split.} \end{cases}$$

We may write  $m(\xi)$  for  $m(\gamma)$  when  $\xi(\gamma) = \xi$ , and extend  $m(\xi)$  to all  $F$  by setting  $m(\xi) = 0$  if  $\xi$  is not in the image of the map in (12.12). Then we have the following:

$$(12.15) \quad \langle \alpha, \beta \rangle = \sum_{\xi \in F} m(\xi) \langle \alpha, \beta \rangle_\xi.$$

Let  $\chi$  be a character of  $T(F) \backslash T(\mathbb{A}_f)$ . The linking number is easy to compute if  $\xi = 0$  or 1. The difficult problem is to compute  $\langle \alpha, \beta \rangle_\xi$  when  $\xi \neq 0, 1$ .

If both  $\alpha$  and  $\beta$  are decomposable,

$$\alpha = \otimes \alpha_v, \quad \beta = \otimes \beta_v,$$

then we have a decomposition of linking numbers into local *linking numbers* when  $\xi \neq 0, 1$ :

$$(12.16) \quad \langle \alpha, \beta \rangle_\xi = \prod \langle \alpha_v, \beta_v \rangle_\xi,$$

where

$$(12.17) \quad \langle \alpha_v, \beta_v \rangle_\xi = \int_{G(F_v)} \alpha_v(\gamma y) \bar{\beta}_v(y) dy.$$

Notice that when  $\gamma \notin N_T$ , these local linking numbers depend on the choice of  $\gamma$  in its class in  $T(F) \backslash G(F) / T(F)$  while their product does not. This problem can be solved by taking  $\gamma$  to be a *trace free element* in its class which is unique up to conjugation by  $T(F)$ .

## Notations

For a compact open subgroup  $U$  of  $G(\mathbb{A}_f)$  (or  $G(F_v)$ ) and two CM-cycles  $\alpha$  and  $\beta$ , we write  $\langle \alpha, \beta \rangle_U$  and  $\langle \alpha, \beta \rangle_{\xi, U}$  for

$$\langle \alpha, \beta \rangle_U = |U|^{-1} \langle \alpha, \beta \rangle, \quad \langle \alpha, \beta \rangle_{\xi, U} = |U|^{-1} \langle \alpha, \beta \rangle_{\xi}.$$

Similarly, for a CM-cycle  $\alpha$  and a function  $f$  on  $M$ , we write

$$(f, \alpha)_U = |U|^{-1} (f, \alpha).$$

## 13 Local Gross-Zagier formula

In this section, we would like to compute the linking numbers for some special CM-cycles and then compare with Fourier coefficients of the kernel functions. The construction of CM-cycles is actually quite simple and is given as follows.

We will fix one order  $A$  of  $B$  such that for each finite place  $v$ ,

$$(13.1) \quad A_v = \mathcal{O}_{K,v} + \mathcal{O}_{K,v} \lambda_v c(\chi_v),$$

where  $\lambda_v \in B_v^{\times}$  such that

- $\lambda_v x = \bar{x} \lambda_v$  for all  $x \in K$ , and
- $\text{ord}(\det \lambda_v) = \text{ord}_v(N)$ .

Let  $\Delta$  be a subgroup of  $G(\mathbb{A}_f)$  generated by images of  $\widehat{A}^{\times}$  and  $K_v^{\times}$  for  $v$  ramified in  $K$ :

$$(13.2) \quad \Delta = \prod_{v|c(\omega_v)} A_v^{\times} F_v^{\times} / F_v^{\times} \cdot \prod_{v|c(\omega_v)} A_v^{\times} K_v^{\times} / F_v^{\times}.$$

The character can be naturally extended to a character of  $\Delta$ . The CM-cycle we need is defined by the following function:

$$(13.3) \quad \eta = \prod \eta_v$$

with  $\eta_v$  supported on  $T(F_v) \cdot \Delta_v$  and such that

$$(13.4) \quad \eta_v(tu) = \chi_v(t) \chi_v(u), \quad t \in T(F_v), u \in \Delta_v.$$

Take an  $a \in \mathbb{A}_f^\times$  integral and prime to  $ND$ . We would like to compute the pairing  $\langle T_a \eta, \eta \rangle$ . The Hecke operator here is defined as

$$(13.5) \quad T_a \eta = \prod_{v|a} T_{a_v} \eta_v, \quad T_{a_v} \eta_v(x) = \int_{H(a_v)} \eta_v(xg) dg,$$

where

$$(13.6) \quad H(a_v) := \{g \in M_2(\mathcal{O}_v) : |\det g| = |a_v|\},$$

and  $dg$  is a measure such that  $\mathrm{GL}_2(\mathcal{O}_v)$  has volume 1. Then we have the following decomposition:

$$(13.7) \quad \begin{aligned} \langle T_a \eta, \eta \rangle_\Delta &= \mathrm{vol}(T(F) \backslash T(\mathbb{A}_f) \Delta) (m(0) T_a \eta(e) + m(1) T_a \eta(\epsilon) \delta_{\chi^2=1}) \\ &+ \sum_{\xi \neq 0,1} m(\xi) \prod_v \langle T_a \eta_v, \eta_v \rangle_{\xi, \Delta_v}, \end{aligned}$$

where  $\epsilon \in N_T(F) - T(F)$ .

Let  $v$  be a fixed finite place of  $F$ . We want to compute all terms in the right hand side involving  $\eta_v$ . Notice that we have extended the definition to all  $\xi \in F - \{0, 1\}$  by insisting that  $\langle T_a \eta_v, \eta_v \rangle_{\xi, \Delta_v} = 0$  when  $\xi$  is not in the image of (12.12).

The computation of degenerate terms is easy. The non-degenerate term is given by the following local Gross-Zagier formula:

**Proposition 13.1.** *Let  $g = \begin{pmatrix} a\delta_v^{-1} & 0 \\ 0 & 1 \end{pmatrix}$ . Then*

$$\bar{W}_v \left( \frac{1}{2}, \xi, g \right) = |c(\omega_v)|^{1/2} \cdot \epsilon(\omega_v, \psi_v) \chi_v(u) \cdot |(1 - \xi)\xi|_v^{1/2} |a| \cdot \langle T_a \eta_v, \eta_v \rangle_{\xi, \Delta_v},$$

where  $u$  is any trace free element in  $K^\times$ .

**Corollary 13.2.** *Let  $\langle \cdot, \cdot \rangle$  be the geometric pairing on the CM-cycle with multiplicity function  $m$  on  $F$  such that  $m(\xi) = 0$  if  $\xi$  is not in the image of (12.12). Assume that  $\delta_v = 1$  for  $v | \infty$ . Then there are constants  $c_1, c_2$  such that for an integral idele  $a$  prime to  $ND$ ,*

$$\begin{aligned} |c(\omega)|^{1/2} |a| \langle T_a \eta, \eta \rangle_\Delta &= (c_1 m(0) + c_1 m(1)) |a|^{1/2} W_f(g) \\ &+ i^{[F:\mathbb{Q}]} \sum_{\xi \in F - \{0,1\}} |\xi(1 - \xi)|_\infty^{1/2} \bar{W}_f(1/2, \xi, g) m(\xi), \end{aligned}$$

where  $g = \begin{pmatrix} a\delta_f^{-1} & 0 \\ 0 & 1 \end{pmatrix}$ .

## Remarks

If the kernel  $\Theta$  had level  $N$  as in the original approach §8, the CM-cycle we should consider is  $P_\chi$  as in §6 corresponding to the function  $\zeta$  supported in  $T(\mathbb{A}_f) i_c U(N, K)$  such that  $\zeta(ticu) = \chi(t)$ . The computation of linking numbers for this divisor seems very difficult!

Our local formula is the key to the proof of the Gross-Zagier formula. But the formula is only proved under the condition that  $c(\chi)$ ,  $c(\omega)$ , and  $N$  are coprime to each other. One may still expect that this local formula is still true in the general case considered by Waldspurger but with more than one term on the right hand side. The main problem is to construct elements in

$$\mathcal{S}(\chi_v, G(F_v)) \quad \text{and} \quad \mathcal{W}(\Pi(\chi_v), \psi_v) \otimes \mathcal{W}(\Pi(|\cdot|^{s-1/2}, |\cdot|^{1/2-s}\omega)).$$

More precisely, we need to find an element

$$W = \sum W_{i1} \otimes W_{2i} \in \mathcal{W}(\Pi(\chi_v), \psi_v) \otimes \mathcal{W}(\Pi(|\cdot|^{s-1/2}, |\cdot|^{1/2-s}\omega))$$

which satisfies the following properties:

- Let  $\Phi_i$  be the element in  $\mathcal{S}(F_v^2)$  such that  $W_{2i} = W_{\Phi_i}$ . Then for any representation  $\Pi_v$  of  $\text{GL}_2(F_v)$  with a newform  $W_v \in \mathcal{W}(\Pi, \psi_v)$ ,

$$L(s, \chi_v, \phi_v) = \sum \Psi(s, W_v, W_{1i}, \Phi_i)$$

with notation in [16] §2.5.

- Let's define

$$W(s, \xi, g) = \sum_i W_{1i} \left( \begin{pmatrix} 1 - \xi & 0 \\ 0 & 1 \end{pmatrix} g \right) W_{2i} \left( \begin{pmatrix} \xi & 0 \\ 0 & 1 \end{pmatrix} g \right).$$

Then  $W(s, \xi, g)$  satisfies the following functional equation

$$W(s, \xi, g) = \omega_v(1 - \xi^{-1}) \epsilon_v(\Pi_v \otimes \chi_v, s) W(1 - s, \xi, g).$$

The next step is to find elements  $q_j \in \mathcal{S}(\chi_v, G(F_v))$  such that the above local Gross-Zagier formula is true with  $\langle T_a \eta_v, \eta_v \rangle_\xi$  replaced by

$$\sum_j \langle T_a q_j, q_j \rangle_\xi.$$

We may even assume that  $a = 1$  in time. Thus, what really varies is the parameter  $\xi$ .

## 14 Gross-Zagier formula in level $ND$

The study of kernel functions, geometric pairing, and local Gross-Zagier formula in the last three sections suggests that it may be easier to prove a Gross-Zagier formula in level  $ND$  instead of level  $N$  directly. This is the main result in our Asia Journal paper [16].

Let's start with the case where  $\Sigma$  is odd. Thus, we are in the situation of §6. Let  $U$  be any open compact subgroup of  $\Delta$  over which  $\chi$  is trivial. Let  $X_U$  be corresponding Shimura curve or its compactification over  $F$ .

Recall that the CM-points corresponding to  $K$  on  $X_U(\mathbb{C})$  form a set

$$C_U := G(F)_+ \backslash G(F)_+ \cdot h_0 \times G(\mathbb{A}_f)/U = T(F) \backslash G(\mathbb{A}_f)/U,$$

where  $h_0 \in \mathcal{H}^+$  is the unique fixed point of the torus points  $K^\times/F^\times$ . Let  $\eta_U$  be a divisor on  $X_U$  with complex coefficient defined by

$$\eta_U = \sum_{x \in C_U} \eta(x)[x].$$

The Heegner class we want now is the class difference

$$y := [\eta_U - \deg(\eta_U)\xi] \in \text{Jac}(X_U)(H_c) \otimes \mathbb{C}.$$

Notice that this class has character  $\chi_\Delta$  under the action by  $\Delta$  on  $\text{Jac}(H_c)$ . Let  $y_\phi$  denote the  $\phi$ -typical component of  $y$ . The main theorem in our Asia Journal paper [16] is now the following

**Theorem 14.1.** *Let  $\phi^\sharp$  be the quasi-newform as in §11. Then*

$$\widehat{\phi^\sharp}(1)L'(1, \chi, \phi) = 2^{g+1}d_{K/F}^{-1/2} \cdot \|\phi^\sharp\|_{U_0(ND)}^2 \cdot \|y_\phi\|_\Delta^2$$

where

- $d_{K/F}$  is the relative discriminant of  $K$  over  $F$ ;
- $\|\phi^\sharp\|_{U_0(ND)}^2$  is the  $L^2$ -norm with respect to the Haar measure  $dg$  as in Introduction normalized such that  $\text{vol}(U_0(ND)) = 1$
- $\|y_\phi\|_\Delta$  is the Neron-Tate height of  $y_\phi$  on  $X_U$  times  $[\Delta : U]^{-1}$  which is independent of choice of  $U$ ;
- $\widehat{\phi^\sharp}(1)$  is the first Fourier coefficient of  $\phi^\sharp$  as defined (8.1).

We now move to the situation in §7 where  $\phi$  has possible nonholomorphic components, but we assume that the sign of the functional equation of  $L(s, \chi, \phi)$  is  $+1$ , or equivalently,  $\Sigma$  is even. We have the variety  $X_U$  which is defined in the same way as in odd case. Then we have a unique line of cuspidal functions  $\phi_\chi$  on  $X_U$  with the following properties:

- $\phi_\chi$  has character  $\chi_\Delta$  under the action of  $\Delta$ ;
- for each finite place  $v$  not dividing  $N \cdot D$ ,  $\phi_\chi$  is the eigenform for Hecke operators  $T_v$  with the same eigenvalues as  $\phi$ .

We call  $\phi_\chi$  a *toric newform* associated to  $\phi$ . See [16], §2.3 for more details.

The CM-points on  $X_U$ , associated to the embedding  $K \longrightarrow B$ , form the infinite set

$$C_U := G(F)_+ \backslash G(F)_+ h_0 \times G(\mathbb{A}_f) / U \simeq H \backslash G(\mathbb{A}_f) / U,$$

where  $h_0$  is a point in  $\mathcal{H}^n$  fixed by  $T$  and  $H \subset G$  is the stabilizer of  $z$  in  $G$ . Notice that  $H$  is either isomorphic to  $T$  if  $n \neq 0$  or  $H = G$  if  $n = 0$ . In any case there is a finite map

$$\iota : C_U = T(F) \backslash G(\mathbb{A}_f) / U \longrightarrow M_U.$$

The Gross-Zagier formula for central value in level  $ND$  is the following:

**Theorem 14.2.** *Let  $\phi_\chi$  be a toric newform such that  $\|\phi_\chi\|_\Delta = 1$ . Then*

$$\widehat{\phi}^\sharp(1)L(1, \chi, \phi) = 2^{g+n} d_{K/F}^{-1/2} \cdot \|\phi^\sharp\|_{U_0(ND)}^2 \cdot |(\phi, \eta)_\Delta|^2,$$

where  $\widehat{\phi}^\sharp(1)$  is the first Fourier coefficient by the same formula as (8.1) with respect the standard Whittaker function defined in (4.7) and (4.8), and where

$$(\phi_\chi, \eta)_\Delta := [\Delta : U]^{-1} \sum_{x \in C_U} \bar{\eta}(x) \phi_\chi(\iota(x)).$$

## 15 Green's functions of Heegner points

In this section we want to explain the proof of the central derivative formula for level  $ND$  stated in the last section. Just as explained in §8, the question is reduced to a comparison of the Fourier coefficients of the kernel and heights

of CM-points. We need to show that, up to a constant and modulo some *negligible forms*, the new form  $\Psi$  with Fourier coefficient

$$(15.1) \quad \widehat{\Psi}(a) := |a| \langle \eta, T_a \eta \rangle$$

is equal to the holomorphic cusp form  $\Phi$  defined in §11 which represents the derivative of Rankin L-function  $L'(1/2, \chi, \phi)$ . Thus we need to show that the functional  $\widehat{\Psi}$  on  $\mathbb{N}_F(ND)$  is equal to the Fourier coefficient  $\widehat{\Phi}(a)$ .

As in §9, we would like to decompose the height pairing to Green's functions. It is more convenient work on the tower of Shimura curves than a single one. Let's first try to extend the theory of heights to the projective limit  $X_\infty$  of  $X_U$ . Let  $\widehat{\text{Pic}}(X_\infty)$  denote the direct limit of  $\widehat{\text{Pic}}(X_U)$  with respect to the pull-back maps. Then the intersection pairing can be extended to  $\widehat{\text{Pic}}(X_\infty)$  if we multiply the pairings on  $\text{Pic}(X_U)$  by the scale  $\text{vol}(U)$ . Of course, this pairing depends on the choice of measure  $dg$  on  $G(\mathbb{A})$  as in Introduction. For some fixed open compact subgroup  $U$  of  $G(\mathbb{A}_f)$ , we write  $\langle z_1, z_2 \rangle_U$  for the measure

$$\langle z_1, z_2 \rangle_U = |U|^{-1} \langle z_1, z_2 \rangle = [U : U']^{-1} \langle z_1, z_2 \rangle_{X_{U'}}.$$

where  $z_1, z_2$  are certain elements in  $\widehat{\text{Pic}}(X_\infty)$  realized on  $X_{U'}$  for some  $U' \subset U$ . So defined pairing will depends only on the choice of  $U$  and gives the exact pairing on  $X_U$ .

Similarly, we can modify the local intersection pairing and extend the height pairing to  $\text{Jac}(X_\infty) = \text{Pic}^0(X_\infty)$ , which is the direct limit of  $\text{Pic}^0(X_U)$  where  $\text{Pic}^0(X_U)$  is the subgroup of  $\text{Pic}(X_U)$  of classes whose degrees are 0 on each connected component.

We can now define Green's functions  $g_v$  on divisors on  $X_\infty(\bar{F})$  which are disjoint at the generic fiber for each place  $v$  of  $F$  by multiplying the Green's functions on  $X_U$  by  $\text{vol}(U)$ . Notice that for two CM-divisors  $A$  and  $B$  on  $X_U$  with disjoint support represented by two functions  $\alpha$  and  $\beta$  on  $T(F) \backslash G(\mathbb{A}_f)$ , the Green's function at a place  $v$  depends only on  $\alpha$  and  $\beta$ . Thus, we may simply denote it as

$$g_v(A, B)_{U_v} = g_v(\alpha, \beta)_{U_v}.$$

Recall that  $\eta$  is a divisor on  $X_\infty$  defined by (13.3). As in §10, with  $P_\chi$  replaced by  $\eta_U$ , we have

$$(15.2) \quad \widehat{\Psi} = \sum_v \widehat{\Psi}_v \pmod{\mathcal{D}(\sigma_1) + \mathcal{D}(r_\chi)},$$

where  $v$  runs through the set of places of  $F$ , and

$$(15.3) \quad \widehat{\Psi}_v(a) := -|a|g_v(\eta, \Gamma_a^0\eta)_{\Delta_v} \log q_v.$$

Thus, it suffices to compare these local terms for each place  $v$  of  $F$ . We need only consider  $v$  which is not split in  $K$ , since  $\widehat{\Phi}_v = 0$  and  $\widehat{\Psi}_v$  is a finite sum of derivations of Eisenstein series when  $v$  is split in  $K$ .

Our main tool is the local Gross-Zagier formula in §13 for quaternion algebra  ${}_vB$  with the ramification set

$$(15.4) \quad {}_v\Sigma = \begin{cases} \Sigma \cup \{v\}, & \text{if } v \notin \Sigma, \\ \Sigma - \{v\}, & \text{if } v \in \Sigma. \end{cases}$$

Let  ${}_vG$  denote the algebraic group  ${}_vB^\times/F^\times$ .

**Lemma 15.1.** *For  $v$  an infinite place,*

$$\widehat{\Phi}_v(a) = 2^{g+1}|c(\omega)|^{1/2}\widehat{\Psi}_v(a).$$

The idea of proof is to use the local Gross-Zagier formula to write both sides as the constant terms at  $s = 0$  of two geometric pairings of divisors  $T_a\eta$  and  $\eta$  with two multiplicity functions:

$$m_s^v(\xi) = \int_1^\infty \frac{dx}{x(1 + |\xi|_v x)^{1+s}}, \quad 2Q_s(\xi) = \int_1^\infty \frac{(1-x)^s dx}{x^{1+s}(1 + |\xi|_v x)^{1+s}}.$$

It follows that the difference of two sides will be the constant term of a geometric pairing on  $T(F)\backslash {}_vG(\mathbb{A}_f)$  with multiplicity function

$$m_s^v - 2Q_s$$

which has no singularity and converges to 0 as  $s \rightarrow 0$ . Notice that the Legendre function  $Q_s$  appears here because an explicit construction of Green's function at archimedean place.

We now consider unramified cases.

**Lemma 15.2.** *Let  $v$  be a finite place prime to  $ND$ . Then there is a constant  $c$  such that*

$$\widehat{\Phi}_v(a) - 2^{g+1}|c(\omega)|^{1/2}\widehat{\Psi}_v(a) = c \log |a|_v \cdot |a|^{1/2}\widehat{\Pi}(\chi)(a).$$

The proof is similar to the archimedean case. Write  $a = \pi_v^n a'$  ( $\pi_v \nmid a'$ ). Since the Shimura curve and CM-points all have good reduction, using Gross' theory of canonical lifting, we can show that  $\widehat{\Psi}_v(a)$  is the geometric pairing of  $T_{a'}\eta$  and  $\eta$  on  $T(F)\backslash_v G(\mathbb{A}_f)$  with multiplicity function

$$m_n(\xi) = \begin{cases} \frac{1}{2}\text{ord}_v(\xi\pi_v^{1+n}), & \text{if } \xi \neq 0 \text{ and } \text{ord}_v(\xi\pi_v^n) \text{ is odd,} \\ n/2, & \text{if } \xi \neq 0 \text{ and } n \text{ is even,} \\ 0, & \text{otherwise.} \end{cases}$$

On other hand, by using the local Gross-Zagier formula, we may also write  $\Phi$ , up to a multiple of  $|a|^{1/2}\widehat{\Pi}(\chi) \log |a|_v$ , as a geometric pairing  $\langle T_{a'}\eta, \eta \rangle$  with multiplicity

$$-2m_n(\xi) \log q_v.$$

It remains to treat the case where  $v$  is place dividing  $ND$ . In this case we will not be able to prove the identity as in the archimedean case, or in the unramified case, since there is no explicit regular model of Shimura curves we can use. But we can classify these contributions:

**Lemma 15.3.** *For  $v$  a finite place dividing  $ND$ , we have*

$$\widehat{\Phi}_v(a) - 2^{g+1}|c(\omega)|^{1/2}\widehat{\Psi}_v(a) = c|a|^{1/2}\widehat{\Pi}(\chi)(a) +_v \widehat{f}$$

where  $c$  is a constant, and  $_v \widehat{f}$  is a form on  $_v G(F)\backslash_v G(\mathbb{A}_f)$ . Moreover, the function  $_v f$  has character  $\chi$  under the right translation by  $K_v^\times$ .

Using the local Gross-Zagier formula, we still can show that  $\widehat{\Phi}_v$  is equal the geometric local pairing

$$2^g |c(\omega)|^{1/2} |a| \langle \eta, T_a \eta \rangle$$

for a multiplicity function  $m(g)$  on  $_v G(F)$  with singularity

$$\log |\xi|_v.$$

On other hand, it is not difficult to show that Green's function

$$\widehat{\Psi}_v(a) = -g_v(\eta, T_a^0 \eta) \log q_v$$

is also a geometric pairing for a multiplicity function with singularity

$$\frac{1}{2} \log |\xi|_v.$$

(This is equivalent to saying that  $\xi^{1/2}$  is a local parameter in the  $v$ -adic space of CM-points). Thus the difference

$$\widehat{\Phi}_v(a) - 2^{g+1}|c(\omega)|^{1/2}\widehat{\Psi}_v(a),$$

is a geometric pairing without *singularity*. In other words, it is given by

$$\int_{[T(F)\backslash_v G(\mathbb{A}_f)]^2} \eta(x)k(x, y)\mathrm{T}_a\eta(y)dx dy,$$

for  $k(x, y)$  a locally constant function of  $({}_v G(F)\backslash_v G(\mathbb{A}_f))^2$  which has a decomposition

$$k(x, y) = \sum_i c_i(x)f_i(y)$$

into eigenfunctions  $f_j$  for Hecke operators on  ${}_v G(F)\backslash_v G(\mathbb{A}_f)$ . It follows that the difference of two sides in the lemma is given by

$$\sum_i \lambda_i(a) \int_{T(F)\backslash_v G(\mathbb{A}_f)} \eta(x)c_i(x)dx \cdot \int_{T(F)\backslash_v G(\mathbb{A}_f)} f_i(y)\bar{\eta}(y)dy,$$

where  $\lambda_i(a)$  is the eigenvalue of  $\mathrm{T}_a$  for  $f_i$ . Thus, we may take

$${}_v f = \sum_i \int_{T(F)\backslash_v G(\mathbb{A}_f)} \eta(x)c_i(x)dx \cdot \int_{T(F)\backslash_v G(\mathbb{A}_f)} f_i(y)\bar{\eta}(y)dy.$$

In summary, at this stage we have shown that the quasi-newform

$$\Phi - 2^{g+1}|c(\omega)|^{1/2}\Psi$$

has Fourier coefficients which are a sum of the following terms:

- derivations  $A$  of Eisenstein series,
- derivations  $B$  of theta series  $\Pi(\chi) \otimes \alpha^{1/2}$ ,
- functions  ${}_v f$  appearing in  ${}_v G(F)\backslash_v G(\mathbb{A}_f)$  with character  $\chi$  under the right translation of  $K_v^\times$ , where  $v$  are places dividing  $DN$ .

By linear independence of Fourier coefficients of derivations of forms in Lemma 10.3, we may conclude that  $A = B = 0$ .

Let  $\Pi$  now be the representation generated by the form  $\phi$ , then all the projections of  ${}_v f$ 's in  $\Pi$  must vanish by some local results of Waldspurger, Gross-Prasad. See §2.3 in [16].

In summary we have shown that  $\Phi - 2^{g+1}|c(\omega)|^{1/2}\Psi$  is an old form. By Proposition 11.1, the projection of this difference on  $\Pi(\phi)$  is

$$\frac{L'(1/2, \chi, \phi)}{(\phi^\#, \phi^\#)_{U_0(ND)}} \cdot \phi^\# - 2^{g+1}|c(\omega)|^{1/2} \cdot \langle y_\phi, y_\phi \rangle_\Delta \cdot \phi.$$

This is again an old form and has vanishing first Fourier coefficient. Theorem 14.1 follows by taking the first Fourier coefficient.

## 16 Spectral decomposition

In this section we want to explain the proof for the central value formula, Theorem 14.2. The idea is copied from the odd case. Thus, we need to define a *height pairing of CM-cycles*. Since there is no natural arithmetic and geometric setting for heights corresponding to non-holomorphic forms, we would like to use the local Gross-Zagier formula to *suggest* a definition of height. Indeed, by corollary 13.2, modulo some Einstein series of type  $\Pi(\|\cdot\|^{1/2}, \|\cdot\|^{-1/2}) \otimes \eta$  with  $\eta$  quadratic, the kernel  $\Phi(g) := \Phi(1/2, g)$  has a Whittaker function satisfying

$$(16.1) \quad W_\Phi \left( g_\infty \cdot \begin{pmatrix} a\delta^{-1} & 0 \\ 0 & 1 \end{pmatrix} \right) = |c(\omega)|^{1/2} |a| \langle T_a \eta, \eta \rangle_\Delta (g_\infty),$$

where  $g_\infty \in \mathrm{GL}_2(F_\infty)$  is viewed as a parameter, and  $a$  is a finite integral idele which prime to  $ND$ , and the pairing  $\langle \cdot, \cdot \rangle_\Delta$  is defined by the multiplicity function

$$(16.2) \quad m(\xi, g_\infty) := \prod_{v|\infty} m_v(\xi, g_v),$$

with each  $m_v(\xi, g)$  the Whittaker function of weight  $k_v$  whose value at  $\begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix}$  given as follows:

$$(16.3) \quad m_v \left( \xi, \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \right) = \begin{cases} 4|a|e^{-2\pi a}, & \text{if } 1 \geq \xi \geq 0, a > 0, k_v = 2, \\ 4|a|e^{2\pi a(2\xi-1)}, & \text{if } a\xi \leq \min(0, a), k_v = 0, \\ 0, & \text{otherwise.} \end{cases}$$

This suggests a definition of the height pairing for CM-cycles by the above multiplicity function. This height pairing is no longer valued in numbers but in Whittaker functions. In the case where all  $k_v = 2$ , then this Whittaker function is twice the standard one. Thus, we can get a pairing with values in  $\mathbb{C}$ . We would like to have a good understanding of decomposition of this height pairing according to eigenforms on  $X_U$ . By (12. 8), we need only decompose the kernel  $k_U$  defined in (12. 7). This decomposition is actually very simple: As Whittaker functions on  $\mathrm{GL}_2(F_\infty)$ ,

$$(16.4) \quad k_U(x, y)(g_\infty) = 2^{[F:\mathbb{Q}]+n} \sum_{\phi_i} W_i(g_\infty) \cdot \phi_i(x) \bar{\phi}_i(y) \\ + 2^{[F:\mathbb{Q}]+n} \int_{\mathfrak{M}} W_m(g_\infty) E_m(x) \bar{E}_m(y) dm$$

where  $n$  is the number of places where  $k_v = 0$ , and the sum is over all cuspidal eigenforms  $\phi_i$  of Laplacian and Hecke operators on  $G(F) \backslash G(\mathbb{A})/U$  such that  $\|\phi_i\|_\Delta = 1$ , and  $W_i$  are standard Whittaker function for  $\phi_i$ . Here the integration is nontrivial only when  $n = g$  then  $\mathfrak{M}$  is a measured space parameterizing an orthogonal basis of Eisenstein series of norm 1. (See §18 for more details). Thus for a cuspidal eigenform  $\phi$ ,

$$\frac{1}{|\Delta|} \int_{G(F) \backslash G(\mathbb{A})} k(x, y) \phi(y) dy = 2^{[F:\mathbb{Q}]+n} W_\phi(g_\infty) \phi(x).$$

It follows that for any two CM-cycles  $\alpha$  and  $\beta$  on  $X_U$ , the height pairing has a decomposition

$$(16.5) \quad \langle \alpha, \beta \rangle = 2^{[F:\mathbb{Q}]+n} \sum_{\phi_i} W_i(g_\infty) \cdot (\phi_i, \bar{\alpha})_\Delta (\bar{\phi}_i, \beta)_\Delta \\ + 2^{[F:\mathbb{Q}]+n} \int_{\mathfrak{M}} W_m(g_\infty) (E_m, \bar{\alpha})_\Delta (\bar{E}_m, \beta)_\Delta dm.$$

This leads to define the following form of  $\mathrm{PGL}_2(\mathbb{A})$  of weight  $k_v$  at  $v$ :

$$(16.6) \quad H(\alpha, \beta) = 2^{[F:\mathbb{Q}]+n} \sum_{\phi_i} \phi_i^{\mathrm{new}}(g_\infty) \cdot (\phi_i, \bar{\alpha})_\Delta (\bar{\phi}_i, \beta)_\Delta \\ + 2^{[F:\mathbb{Q}]+n} \int_{\mathfrak{M}} E_m^{\mathrm{new}}(g_\infty) (E_m, \bar{\alpha})_\Delta (\bar{E}_m, \beta)_\Delta dm,$$

where  $\phi_i^{\text{new}}$  (resp.  $E_\lambda^{\text{new}}$ ) is the *newform* of weight  $(2, \dots, 2, 0, \dots, 0)$  in the representation  $\Pi_i$  of  $\text{PGL}_2(\mathbb{A})$  corresponding to the representation  $\Pi'_i$  of  $G(\mathbb{A})$  generated by  $\phi_i$  (resp.  $E_m$ ) via Jacquet-Langlands theory. With  $\alpha$  replaced by  $T_a\alpha$  in (16.5), one obtains the usual relation between height pairing and Fourier coefficient:

$$(16.7) \quad |a| \langle T_a\alpha, \beta \rangle_\Delta(g_\infty) = 2^{[F:\mathbb{Q}]+n} W_{H(\alpha,\beta)} \left( g_\infty \cdot \begin{pmatrix} a\delta^{-1} & 0 \\ 0 & 1 \end{pmatrix} \right).$$

Let  $\Psi$  denote the form  $2^{[F:\mathbb{Q}]+n} |c(\omega)|^{1/2} H(\eta, \eta)$  which has a decomposition:

$$(16.8) \quad \Psi = 2^{[F:\mathbb{Q}]+n} |c(\omega)|^{1/2} \sum_i \phi_i^{\text{new}} |(\phi_i, \eta)_\Delta|^2 \\ + 2^{[F:\mathbb{Q}]+n} |c(\omega)|^{1/2} \int_{\mathfrak{M}} E_m^{\text{new}} |(E_m, \eta)|^2 dm.$$

Since  $\eta$  has a character  $\chi$  under the action by  $\Delta$ , we may require that  $\phi_i$  (resp.  $E_{m,\chi}$ ) has character  $\chi$  under the action by  $\Delta$ . For a given  $\phi^{\text{new}}$  of level  $N$ , then  $\phi_i$  with  $\phi_i^{\text{new}} = \phi^{\text{new}}$  must be the toric newform as in §14.

The equations (16.1) and (16.7) shows that, modulo certain Eisenstein series in the space  $\Pi(\|\cdot\|^{1/2}, \|\cdot\|^{-1/2}) \otimes \eta$  with  $\eta^2 = 1$ , the forms  $\Phi - \Psi$  has vanishes Fourier coefficient at  $g$  such that  $g_f = \begin{pmatrix} \delta^{-1}a & 0 \\ 0 & 1 \end{pmatrix}$  with integral  $a$  prime to  $ND$ . Thus  $\Phi - \Psi$  is an old form.

Let  $\phi$  be the newform as in Theorem 14.2. By Proposition 11.1 and formula (16.6), the projection of  $\Phi - \Psi$  in  $\Pi(\phi)$  is given by

$$\frac{L(1/2, \chi, \phi)}{\|\phi^\sharp\|_{U_0(ND)}^2} \phi^\sharp - 2^{[F:\mathbb{Q}]+n} |c(\omega)|^{1/2} \cdot |(\eta, \phi_\chi)_\Delta|^2 \phi.$$

Thus, we have proven the Gross-Zagier formula, Theorem 14.2, by computing the first Fourier coefficient of the above form.

## 17 Lowering levels

Now it remains to deduce GZF(N) (the Gross-Zagier formulas for level  $N$  in §6-7) from GZF(ND) (formulas in §14). Our plan is as follows:

1. show GZF (N) up to a certain universal function of local parameters.

2. prove GZF(ND) for Eisenstein series for level  $ND$  thus get GZF(N) for Eisenstein series with the same universal function.
3. prove GZF(N) for Eisenstein series directly by evaluating the periods thus get the triviality of the universal function.

In this section we are doing the first step.

**Proposition 17.1.** *For each  $v \mid D$ , there is a rational function  $Q_v(t) \in \mathbb{C}(t)$  depending only on  $\chi_v$  which takes 1 at  $t = 0$  and is regular for*

$$|t| < |\pi_v|^{1/2} + |\pi_v|^{-1/2},$$

such that both Gross-Zagier formulas in §6-7 are true after multiplying the left hand side by

$$C(\chi) \prod_{\text{ord}_v(D) > 0} Q_v(\lambda_v),$$

where  $C(\chi)$  is a constant depends only  $\chi$ , and  $\lambda_v$  the parameter appeared in the  $L$ -function:

$$L_v(s, \phi) = \frac{1}{1 - \lambda_v |\pi_v|^s + |\pi_v|^{2s}}.$$

The idea of proof is to show that, in the comparison of GZF (N) and GZF (ND), all 4 quantities

$$\widehat{\phi}^\sharp(1), \quad \frac{\|\phi^\sharp\|_{U_0(ND)}^2}{\|\phi\|_{U_0(N)}^2}, \quad \frac{|(\eta, \phi_\chi)_\Delta|^2}{|i_\chi(\widetilde{\phi})|^2}, \quad \frac{\|x_\phi\|_\Delta^2}{\|y_\phi\|_{U_0(N)}^2},$$

are universal functions described in the Proposition, and that the last two quantities have the same functions. Here the last two fractions are considered as ratios since the denominators may be 0.

Lets try to localize the definition of quasi-newform in §11. For each finite place  $v$ , let  $\Pi_v$  be the local component of  $\Pi(\phi)$  at  $v$ . Then  $\Pi_v$  is a unitary representation as  $\Pi = \otimes \Pi_v$  is. Lets fix an Hermitian form for the Whittaker model  $\mathcal{W}(\Pi_v, \psi_v)$  such that the norm of the new vector is 1 for almost all  $v$ . The product of this norm induces a norm on  $\Pi$  which is proportional to the  $L^2$ -norm on  $\Pi$ . Now we can define the quasi-newform

$$(17.1) \quad W_v^\sharp \in \mathcal{W}(\Pi_v, \psi_v)$$

to be a certain form of level  $D_v$ . Recall that the space of forms of level  $D_v$  has a basis consisting of forms

$$(17.2) \quad W_{vi}(g) = W_v \left( g \begin{pmatrix} \pi_v^{-i} & 0 \\ 0 & 1 \end{pmatrix} \right), \quad 0 \leq i \leq \text{ord}_v(D_v),$$

where  $W_{v0} = W_v$  is the newform. Then  $W_v^\sharp$  is the unique nonzero form of level  $D_v$  satisfying the following equations:

$$(17.3) \quad (W_v^\sharp, W_{vi} - \nu_v^i W_v^\sharp) = 0, \quad (0 \leq i \leq \text{ord}_v(D_v)),$$

where  $\nu_v = 0$  if  $v$  is not ramified in  $K$ ; otherwise  $\nu_v = \chi_v(\pi_{K,v})$ . It is not difficult to show that if we write

$$W_v^\sharp = \sum c_{vi} W_{v,i},$$

then  $c_{vi}$  is rational function of quantities

$$\alpha_{vi} := (W_{vi}, W_v) / (W_v, W_v).$$

Notice that quantities  $\alpha_{vi}$  does not depend on the choice of pairing  $(\cdot, \cdot)$  on Whittaker models. On other hand, it is easy to show that  $\phi^\sharp$  has Whittaker function as product of  $W_v^\sharp$ .

It follows that both quantities

$$\widehat{\phi}^\sharp(1), \quad \text{and} \quad \frac{\|\phi^\sharp\|_{U_0(ND)}^2}{\|\phi\|_{U_0(N)}^2} = \frac{\|\phi^\sharp\|_{U_0(N)}^2}{\|\phi\|_{U_0(N)}^2} \cdot [U_0(1) : U_0(D)],$$

are the products of some rational functions at  $v$  of quantities  $\alpha_{vi}$ . It remains to show that  $\alpha_{vi}$  are rational functions of  $\lambda_v$ . Let  $U_v = \text{GL}_2(\mathcal{O}_v)$ . Then we have

$$\begin{aligned} \alpha_{vi} &= (W_v, W_v)^{-1} \text{vol}(U_v)^{-1} \int_{U_v} (\rho(u) W_{vi}, \rho(u) W_v) du \\ &= (\rho(t_{vi}) W_v, W_v) / (W_v, W_v), \end{aligned}$$

where  $t_{vi}$  is the Hecke operator corresponding the constant function  $\text{vol}(H_{vi})^{-1}$  on

$$H_{vi} = U_v \begin{pmatrix} \pi_v^{-i} & 0 \\ 0 & 1 \end{pmatrix} U_v.$$

It is well known that  $W_v$  is an eigenform under  $t_{vi}$  with eigenvalue a rational function of  $\lambda_v$ . This shows that  $\alpha_{vi}$  are rational function of  $\lambda_v$ . Since we have used only the unitary property of local representation  $\Pi_v$  for  $v \mid D$ , the so obtained rational functions as in theorem for these quantities are regular for any  $\lambda_v$  as long as  $\Pi_v$  is unitary. In other words, these functions are regular at  $\lambda_v$  satisfying

$$|\lambda_v| < |\pi_v|^{1/2} + |\pi_v|^{-1/2}.$$

It remains to compare the last two quantities in both odd and even case respectively. Obviously the ratio of normalizations of measures is given by

$$|U(N, K)|/|\Delta|$$

which equals a product of constants at places dividing  $D$ . Thus we may take the same measure in the comparison. Let's define a function  $\zeta$  on CM-points  $T(F)\backslash G(\mathbb{A}_f)$  supported on  $T(\mathbb{A})i_c U(N, K)$  such that

$$\zeta(ti_c u) = \chi(t), \quad t \in T(\mathbb{A}_f), u \in U(N, K).$$

Then the CM-Points in GZF(N) (resp. GZF (ND)) is defined by  $\zeta$  (resp.  $\eta$ ). The key to proving our result is to compare these CM-cycles.

Recall that for a finite place  $v$ , and a compactly supported, and locally constant function  $h$  on  $G(F_v)$ , one defines the Hecke operator  $\rho(h)$  on CM-cycles by

$$\rho(h) = \int_{G(F_v)} h(g)\rho(g)dg.$$

Let  $U_1$  and  $U_2$  be the compact subgroup of  $G(\mathbb{A}_f)$  defines as products  $U_i = \prod U_{iv}$  and

$$\begin{aligned} U_{1v} &= (\mathcal{O}_{c_v} + c_v \mathcal{O}_{K,v} \lambda_v)^\times \\ U_{2v} &= U(N, K)_v^\times. \end{aligned}$$

**Lemma 17.2.** *For each finite place  $v$  let  $h_v$  denote the constant function  $\text{vol}(U_{1v})^{-1}$  on  $G(F_v)$  supported on  $U_{2,v}i_{c,v}^{-1}$ . Then*

$$\zeta_v = \rho(h_v)\eta_v.$$

Before we prove this lemma, let us see how to use this lemma to finish the proof of our Proposition. First assume we are in the even case. Then we have

$$(\tilde{\phi}, P_\chi) = (\tilde{\phi}, \zeta)_{U(N,K)}.$$

Here the product is taken as pairings between CM-cycles and functions. Let  $\Psi_\zeta$  be the form  $H(\zeta, \zeta)$  defined in (16. 6). Then  $\Psi_\zeta$  has a decomposition

$$(17.4) \quad \Psi_\zeta = \sum_i \phi_i^{\text{new}} |(\zeta, \phi_i)|^2 + \text{Eisenstein series},$$

where  $\phi_i$  is an orthonormal basis of eigenforms on  $X(N, K)$ . For  $\phi_i^{\text{new}} = \phi$  with level  $N$ ,  $\phi_i$  must be the test form  $\tilde{\phi}$  as in §7.

Now the equation (16. 7) implies that the Hecke operator the adjoint of  $\rho(h)$  is  $\rho(h^\vee)$  with  $h^\vee(g) = \bar{h}(g^{-1})$ . It follows that,

$$H(\zeta, \zeta) = H(\rho(h)\eta, \rho(h)\eta) = H(\rho(h^\vee * h)\eta, \eta).$$

Since  $\eta$  has character  $\chi$  under the action by  $\Delta$ , we may replace  $h^\vee * h$  by a function  $h_0$  which has character  $(\chi, \chi^{-1})$  by actions of  $\Delta$  from both sides and invariant under conjugation. Now

$$(17.5) \quad \begin{aligned} \Psi_\zeta &= \sum \phi_i^{\text{new}}(\rho(h_0)\eta, \phi_i) \overline{(\eta, \phi_i)} + \dots \\ &= \sum \phi_i(\eta, \rho(h_0)\phi_i) \overline{(\eta, \phi_i)} + \dots \end{aligned}$$

Since  $\eta$  and  $\rho(h_0)\eta$  both have character  $\tilde{\chi}$  under the action by  $\Delta$ , we may replacing  $\phi_i$  by functions  $\phi_{i,\chi}$  which has character  $\chi$  under  $\chi$ . For  $\phi_i^{\text{new}} = \phi^{\text{new}}$  with level  $N$ ,  $\phi_{i,\chi}$  must be the toric new form  $\phi_\chi$  as in §14. Of course,

$$(17.6) \quad \rho(h_0)\phi_\chi = \prod_{v|D} P_v(\lambda_v)\phi_\chi$$

as  $\phi_\chi$  with  $P_v$  some polynomial functions. From (17.4 -6), we obtain

$$|(\zeta, \tilde{\phi})|^2 = \prod_{v|D} P_v(\lambda_v) \cdot |(\eta, \phi_\chi)|^2.$$

In the odd case, the proof is same but simpler with  $H(\zeta, \zeta)$  defined as a holomorphic cusp form of weight 2 with Fourier coefficients given by the height pairings of  $\langle T_a x, x \rangle$  for two CM-divisors after minus some multiple of Hodge class. Then we end up with expression:

$$\Psi_\zeta = \sum_i \phi_i \|x_{\phi_i}\|^2,$$

where  $\phi_i$  are newforms of level dividing  $N$ . The same reasoning as above shows that

$$\Psi_\zeta = H(\rho(h_0)y, y) = \sum \phi_i \cdot \langle y_{\phi_i}, \rho(h_0)y_{\phi_i} \rangle.$$

Thus we have

$$\|x_\phi\|^2 = \langle y_\phi, \rho(h_0)y_\phi \rangle = \prod_{v|D} P_v(\lambda_v) \|y_\phi\|^2.$$

It remains to prove the lemma. By definition,

$$\rho(h_v)\eta_v(g) = \text{vol}(U_{1v})^{-1} \int_{U_{2v}} \eta_v(gui_{c,v}^{-1})du.$$

If  $\rho(h_v)\eta_v(g) \neq 0$ , then

$$gU_{1,v}i_{c,v}^{-1} \in T(F_v)U_{1,v}$$

or equivalently,

$$g \in T(F_v)U_{1,v}i_{c,v}U_{2,v}.$$

By (iii) in the following lemma, we have  $g \in T(F_v)i_{c,v}U_{2,v}$ . Lets write  $g = ti_{c,v}u_g$ . It follows that

$$\rho(h_v)\eta_v(g) = \chi(t)\text{vol}(U_{1v})^{-1} \int_{U_{2v}} \eta_v(i_{c,v}ui_{c,v}^{-1})du.$$

By (iii) in the following lemma again, the integral is the same as

$$\int_{U_{1v}} \eta_v(u)du.$$

Thus we have

$$\rho(h_v)\eta_v = \zeta_v.$$

**Lemma 17.3.** *For  $v$  split in  $B$ , there is an isomorphism*

$$\mu : M_2(F_v) \longrightarrow \text{End}_{F_v}(K_v)$$

*with compatible embedding of  $K_v$  and such that*

$$\mu(M_2(\mathcal{O}_v)) = \text{End}_{\mathcal{O}_v}(\mathcal{O}_{K,v}).$$

Moreover, let  $i_{\pi^n} \in \mathrm{GL}_2(F_v)$  such that

$$\mu(i_{\pi^n})(\mathcal{O}_{K,v}) = \mathcal{O}_{\pi^n} = \mathcal{O}_v + \pi_v^n \mathcal{O}_{K,v}.$$

Then

$$(i) \quad \mathrm{GL}_2(F_v) = \coprod_{n \geq 0} K_v^\times i_{\pi^n} \mathrm{GL}_2(\mathcal{O}_v),$$

$$(ii) \quad i_{\pi^n} \mathrm{GL}_2(\mathcal{O}_v) i_{\pi^n}^{-1} \cap K_v^\times = \mathcal{O}_{\pi^n}^\times,$$

$$(iii) \quad i_{\pi^n} U_{2,v} i_{\pi^n}^{-1} \cap K_v^\times U_{1,v} = U_{1,v}.$$

*Proof.* Indeed for any given embedding  $K_v \longrightarrow M_2(F_v)$  such that  $\mathcal{O}_{K,v}$  maps to  $M_2(\mathcal{O}_v)$ , then  $F_v^2$  becomes a  $K_v$ -module of rank 1 such that  $\mathcal{O}_v^2$  is stable under  $\mathcal{O}_{K,v}$ . Then we find an isomorphism  $\mathcal{O}_v^2 \simeq \mathcal{O}_{K,v}$  as  $\mathcal{O}_{K,v}$  module. This induces the required isomorphism  $\mu : B_v \longrightarrow \mathrm{End}_{F_v}(K_v)$ .

Now, for any  $g \in \mathrm{GL}_2(F_v)$ , let  $t \in g(\mathcal{O}_{K,v})$  be the elements with minimal order. Then  $\mu(t^{-1}g)(\mathcal{O}_{K,v})$  will be an order of  $\mathcal{O}_{K,v}$ , say  $\mathcal{O}_{\pi^n}$ . Thus

$$\mu(t^{-1}g)(\mathcal{O}_{K,v}) = \mathcal{O}_{\pi^n} = \mu(i_{\pi^n})(\mathcal{O}_{K,v}).$$

It follows that

$$g \in t i_{\pi^n} \mathrm{GL}_2(\mathcal{O}_v).$$

The first equality follows.

For the second equality, let  $t \in K_v$ . Then

$$i_{\pi^n}^{-1} t i_{\pi^n} \in \mathrm{GL}_2(\mathcal{O}_v)$$

if and only if

$$\mu(i_{\pi^n}^{-1} t i_{\pi^n}) \mathcal{O}_K = \mathcal{O}_K,$$

or equivalently,

$$\mu(t i_{\pi^n}) \mathcal{O}_K = \mu(i_{\pi^n}) \mathcal{O}_K,$$

$$t \mathcal{O}_{\pi^n} = \mathcal{O}_{\pi^n}.$$

This is equivalent to the fact that  $t \in \mathcal{O}_{\pi^n}^\times$ .

It remains to show the last equality. First we want to show

$$i_{c,v}^{-1} U_{1,v} i_{c,v} \subset U_{2,v}.$$

To see this, we need to show that

$$\mu(i_{c,v}^{-1}ui_{c,v})\mathcal{O}_{K,v} = \mathcal{O}_{K,v}$$

for each  $u \in U_{1,v}$ . This is equivalent to

$$\mu(ui_{c,v})\mathcal{O}_{K,v} = i_{c,v}\mathcal{O}_{K,v},$$

or equivalently,

$$\mu(u)\mathcal{O}_{c,v} = \mathcal{O}_{c,v}.$$

Thus is clear from the fact that  $u = t(1 + cM_2(\mathcal{O}_v))$  for some  $t \in \mathcal{O}_c^\times$ .

Now the last equality follows easily: since

$$i_{\pi_v^n}U_{2,v}i_{\pi_v^n}^{-1} \cap K_v^\times = \mathcal{O}_{\pi_v^n}$$

and

$$i_{\pi_v^n}U_{2,v}i_{\pi_v^n}^{-1} \supset U_{1,v},$$

it follows that

$$i_{\pi_v^n}U_{2,v}i_{\pi_v^n}^{-1} \cap K_v^\times U_{1,v} = U_{1,v}.$$

□

## 18 Continuous spectrum

In this section, we would like to extend GZF(ND) (the Gross-Zagier formula in level  $ND$ , Theorem 14.2) to Eisenstein series in the continuous spectrum. Recall that the space of  $L^2$ -forms on  $\mathrm{PGL}_2(F)\backslash\mathrm{PGL}_2(\mathbb{A})$  is a direct sum of cusp forms, characters, and Eisenstein series corresponding to characters  $(\mu, \mu^{-1})$ . We say two characters  $\mu_1, \mu_2$  are connected if  $\mu_1 \cdot \mu_2^\pm$  is trivial on the subgroup  $\mathbb{A}^1$  of norm 1. Thus each connected component is a homogeneous space of  $\mathbb{R}$  or  $\mathbb{R}/\pm 1$ . See [3] for more details.

We now fix a component containing a character  $(\mu, \mu^{-1})$ . Without loss of generality, we assume that  $\mu^2$  is not of form  $|\cdot|^t$  for some  $t \neq 0$ . Then the space  $\mathrm{Eis}(\mu)$  of  $L^2$ -form corresponding to this component consists of the forms

$$(18.1) \quad E(g) = \int_{-\infty}^{\infty} E_t(g)dt$$

where  $E_t(g)$  is the Eisenstein series corresponding to characters  $(\mu|\cdot|^{it}, \mu^{-1}|\cdot|^{-it})$ . For the uniqueness of this integration we assume that  $E_t(g) = 0$  if  $t < 0$  and  $\mu^2 = 1$ . Now the two elements  $E_1(g)$  and  $E_2(g)$  has inner product given by

$$(18.2) \quad (E_1, E_2) = \int_{-\infty}^{\infty} (E_{1t}, E_{2t})_t dt,$$

where  $(\cdot, \cdot)_t$  is some Hermitian form on the space

$$\Pi_t := \Pi(\mu|\cdot|^{it}, \mu^{-1}|\cdot|^{-it}).$$

This Hermitian norm is unique up to constant multiple as the representation is irreducible. The precise definition of this norm is not important to us.

Now we want to compute the Rankin-Selberg convolution of  $E \in \text{Eis}(\mu)$  with  $\theta$  as in §8. Assume that  $\chi$  is not of form  $\nu \cdot \mathbf{N}_{K/F}$ . Then  $\theta$  is a cusp form and the kernel function  $\Theta$  is of  $L^2$ -form as its constant term has exponential decay near cusp. Thus it makes sense to compute  $(E, \Theta)$ .

For  $\phi$  a function on  $\mathbb{R}$  (or  $\mathbb{R}_+$  when  $\mu^2 = 1$ ), lets write  $E_\phi$  for element in  $\text{Eis}(\mu)$  with form  $E_t(g) = \phi(t)E_t^{\text{new}}(g)$  with  $E_t^{\text{new}}(g)$  a newform in  $\Pi(\mu|\cdot|^s, \mu^{-1}|\cdot|^{-s})$  and  $\phi(s) \in \mathbb{C}$ , then we still have

$$\begin{aligned} (E_\phi, \bar{\Theta}_s) &= \int_0^\infty L(s, \Pi_t \otimes \chi) \phi(t) dt \\ &= \int_0^\infty L(s + it, \mu \otimes \chi) L(s + it, \mu^{-1} \otimes \chi) \phi(t) dt. \end{aligned}$$

If  $s = 1/2$ , we obtain

$$(18.3) \quad (E_\phi, \bar{\Theta}_{1/2}) = \int |L(1/2 + it, \mu \otimes \chi)|^2 \phi(t) dt.$$

The form  $\Theta$  has level  $D$ . For any  $a$  dividing  $D$ , lets define

$$E_{\phi,a} = \rho \begin{pmatrix} a^{-1} & 0 \\ 0 & 1 \end{pmatrix} E_\phi.$$

Then the space of Eisenstein series is generated by  $E_{\phi,a}$ . We can define so called *quasi-newforms* by the formula

$$(18.4) \quad E_\phi^\sharp = \int_0^\infty E_t^\sharp \phi(t) dt.$$

One can show that the projection of  $\Phi := \bar{\Theta}_{1/2}$  on the continuous spectrum corresponding to  $\mu$  is  $E_\phi^\sharp$  with  $\phi$  given by

$$(18.5) \quad \phi(t) = \frac{|L(1/2 + it, \mu \otimes \chi)|^2}{\|E_t^\sharp\|^2}.$$

Now let us study the geometric pairing in §16. The formula (16.8) shows that the continuous contribution for representation  $\Pi_t$  in the form  $\Psi$  is given by

$$2^{2g}|c(\omega)|^2 E_\psi$$

where

$$(18.6) \quad \psi(t) = (E_{t,\chi}, \eta).$$

Here  $E_{t,\chi}$  is a form toric form of norm 1 with respect to  $\Delta$ .

Again  $E_\phi^\sharp - 2^{2g}|c(\omega)|^{1/2} E_\psi$  will be an old form. Its first Fourier coefficient vanishes. Thus the Gross-Zagier formula can be extended to Eisenstein series:

**Proposition 18.1.** *Assume that  $\chi$  is not of form  $\nu \circ N_{K/F}$  with  $\nu$  a character of  $F^\times \backslash \mathbb{A}^\times$ . Then*

$$\widehat{E}_t^\sharp(1) |L(1/2 + it, \chi)|^2 = 2^{2g}|c(\omega)|^{1/2} \cdot \|E_t^\sharp\|^2 |(E_{\chi,t}, \eta)|^2.$$

Also the proof of Proposition 17.1 is purely local, thus can be extended to Eisenstein series:

**Proposition 18.2.** *Assume that  $\chi$  is not of form  $\nu \circ N_{K/F}$  with  $\nu$  a character of  $F^\times \backslash \mathbb{A}^\times$ . Let*

$$\lambda_v(t) = \mu_v(\pi_v) |\pi_v|^{it} + \mu_v(\pi_v)^{-1} |\pi_v|^{-it}, \quad \text{and} \quad E_t^* := \|E_t\| \cdot \widetilde{E}_t,$$

then

$$c(\chi) \prod_{\text{ord}_v(D) > 0} Q_v(\lambda_v(t)) = \frac{2^{2g}}{\sqrt{N(D)}} \left| \frac{(E_t^*, P_\chi)}{L(1/2 + it, \chi)} \right|^2.$$

Notice that when  $\mu$  is unramified then  $\widehat{R}$  is conjugate to  $M_2(\widehat{\mathcal{O}}_F)$ . The form  $E_t^*$  is obtained from  $E_t$  by  $\rho(j)$  for a certain  $j \in G(\mathbb{A})$  satisfying (19.1) and (19.2) below. Thus the formula does not involve the definition of hermitian forms on  $\Pi_t$ .

## 19 Periods of Eisenstein series

In this section we want to compute the periods of Eisenstein series appearing in the GZF(N) (up to a universal function), Proposition 18.2. Our result shows that all the universal functions are trivial, and thus end up the proof of GZF(N).

First let's describe the main result. Let  $\mu$  be a unramified quasi-character of  $F^\times \backslash \mathbb{A}^\times$ . Let  $R$  be a maximal order of  $M_2(F)$  containing  $\mathcal{O}_K$ . Let  $E$  be the newform in  $\Pi(\mu, \mu^{-1})$ . Let  $j \in G(\mathbb{A})$  such that

$$(19.1) \quad j_\infty \mathrm{SO}_2(\mathbb{R}) j_\infty^{-1} = T(\mathbb{R})$$

and

$$(19.2) \quad j_f \mathrm{GL}_2(\widehat{\mathcal{O}}) j_f^{-1} = R.$$

Then the form  $E^*(g) := E(gj)$  is invariant under  $T(\mathbb{R}) \cdot \widehat{R}^\times$ . Let  $\lambda \in K$  be a non-zero trace free element. Then one can show that  $\mathrm{ord}_v(\lambda/D)$  for all finite place  $v$  is always even. We thus assume that  $4\lambda/D$  has square root at finite place and that  $D_v = -1$  when  $v \mid \infty$ .

**Proposition 19.1.** *Assume that  $\chi \neq \mu_K := \mu \circ N_{K/F}$ . Then*

$$(E^*, P_\chi) = 2^{-g} \mu \left( \delta^{-1} \sqrt{4\lambda/D} \right) |4\lambda/D|^{1/4} L(1/2, \bar{\chi} \cdot \mu_K).$$

Before we go to the proof of this result, let's see how to use Proposition 19.1 to complete the proof of GZF(N). Combined Propositions 19.1 and 18.2 with  $\mu(x) = |x|^{it}$  ( $t \in \mathbb{R}$ ), we obtain the following

$$C(\chi) \prod_{\mathrm{ord}_v(D) > 0} Q_v(\lambda_v(t)) = 1, \quad \forall t \in \mathbb{R}.$$

Notice that each  $\lambda_v(t)$  is a rational function of  $p^{ti}$  where  $p$  is prime number divisible by  $v$ . Since functions  $p^{ti}$  for different primes  $p$  are rationally independent, we obtain that for each prime  $p$

$$\prod_{\mathrm{ord}_v(D_p) > 0} Q_v(\lambda_v(t)) = \mathrm{const}$$

where  $D_p = \prod_{v|p} D_v$ .

It is not difficult to show that for each  $\chi_v$  we can find a finite character  $\chi'$  of  $\mathbb{A}_K^\times / K^\times \mathbb{A}^\times$  such that the following conditions are verified:

- $c(\chi')$  is prime to  $N$ ,  $c(\omega)$ ;
- $\chi'$  is unramified at all  $w \mid p, w \neq v$ ;
- $\chi'$  is not of form  $\nu \circ N_{K/F}$ .

If we apply the above result to  $\chi'$  then we found that  $Q_v$  is constant thus is 1. Thus we have shown the following:

**Proposition 19.2.** *All polynomials  $Q_v(t)$  and  $C(\chi)$  are constant 1.*

Now GZF (N) follows from Proposition 17.1.

We now start the proof of Proposition 19.1 from the following integral:

$$(E^*, P_\chi) = \int_{T(F) \backslash T(\mathbb{A}_f)} \chi^{-1}(x) E^*(x_\infty x i_c) dx$$

where  $x_\infty \in \mathcal{H}^g$  is fixed by  $T(\mathbb{R})$  and  $i_c \in G(\mathbb{A}_f)$  is an element such that

$$i_c R i_c^{-1} \cap K = \mathcal{O}_c.$$

Here we pick up a measure  $dt$  on  $T(\mathbb{A})$  with local decomposition  $dt = \otimes_v dt_v$  such that  $T(\mathbb{R})$  and  $T(\mathcal{O}_{c,v})$  all have volume 1. Write  $h = i_c j$ . It follows that

$$(E^*, P_\chi) = \int_{T(F) \backslash T(\mathbb{A})} \bar{\chi}(x) E(xh) dx.$$

Since  $E$  is obtained by analytic continuation from the newform in the Eisenstein series in  $\Pi(\mu | \cdot |^s, \mu^{-1} | \cdot |^s)$  with  $\text{Re}(s) \gg 0$ , we thus need only compute the periods for quasi-character  $\mu$  with big exponent. In this case,

$$E(g) = \sum_{\gamma \in P(F) \backslash G(F)} f(\gamma g)$$

with

$$f(g) = \mu^{-1}(\delta) f_\Phi,$$

where  $\Phi = \otimes \Phi_v \in \mathcal{S}(\mathbb{A}^2)$  is the standard element:  $\Phi_v$  is the characteristic function of  $\mathcal{O}_v^2$  if  $v \nmid \infty$ , and  $\Phi_v(x, y) = e^{-\pi(x^2+y^2)}$  if  $v \mid \infty$ . It is not difficult to show that the embedding  $T \rightarrow G$  defines an bijective map

$$T(F) \simeq P(F) \backslash G(F).$$

Thus

$$(E^*, P_\chi) = \int_{T(\mathbb{A})} \chi^{-1}(x) f(xh) dx.$$

This is of course the product of local integrals

$$i_{\chi_v}(f_v) = \int_{T(F_v)} \chi_v(x^{-1}) f_v(xh_v) dx.$$

Recall that  $f_v$  is defined as follows:

$$f_v(g) = \mu(\delta_v^{-1} \cdot \det g) |\det g|^{1/2} \int_{F_v^\times} \Phi[(0, t)g] \mu^2(t) |t| d^\times t.$$

It follows that

$$\begin{aligned} i_{\chi_v}(f_v) &= \int_{T(F_v)} \chi_v(x^{-1}) \mu(\delta_v^{-1} \det xh_v) |\det xh_v|^{1/2} \int_{F_v^\times} \Phi[(0, t)xh_v] \mu^2(t) |t| d^\times t dx \\ &= \mu(\delta_v^{-1} \det h_v) |\det h_v|^{1/2} \int_{K_v^\times} \bar{\chi}_v \mu_K(x) |x|_K^{1/2} \Phi_{K_v}(x) dx \\ &= \mu(\delta_v^{-1} \det h_v) |\det h_v|^{1/2} Z(1/2, \bar{\chi} \cdot \mu_K, \Phi_{K_v}) \end{aligned}$$

where for  $x \in K_v^\times$ ,

$$\Phi_{K_v}(x) = \Phi_v[(0, 1)xh_v].$$

Thus the computation of period is reduced to compute the local Zeta functions.

Let  $v$  be a finite place. The map  $x \rightarrow (0, 1)x$  defines an isomorphism between  $K$  and  $F^2$  with compatible actions by  $K$ . Thus we have two lattices  $\mathcal{O}_F^2$  and  $\mathcal{O}_c$  in  $K$ . The element  $h_f$  as a class in

$$\widehat{K}^\times \backslash \mathrm{GL}_2(\mathbb{A}_f) / \mathrm{GL}_2(\widehat{\mathcal{O}}_F)$$

is determined by the property that

$$h_f M_2(\widehat{\mathcal{O}}_F) h_f^{-1} \cap K = \mathcal{O}_c.$$

We may take  $h_f$  such that

$$(0, 1)\mathcal{O}_c h_f = \mathcal{O}_F^2.$$

It follows that  $\Phi_{K,v}$  is the characteristic function of  $\widehat{\mathcal{O}}_c$ . Now the Zeta function is easy to compute:

$$Z(1/2, \bar{\chi} \cdot \mu_K, \Phi_{K,c}) = \int_{\mathcal{O}_{c_v}} \chi \cdot \mu_K(x) |x|_K^{1/2} d^\times x.$$

We get the standard L-function if  $c = 0$ .

We assume now that  $c > 0$  thus  $K_v/F_v$  is an unramified extension. First we assume that  $K_v$  is a field. We decompose the set  $\mathcal{O}_c$  into the disjoint union of  $\mathcal{O}_{c,n}$  of subset of elements of order  $n$ . Then

$$Z(1/2, \bar{\chi} \cdot \mu_K, \Phi_{K,c}) = \sum_{n \geq 0} \mu(\pi_v)^{2n} |\pi_v|^n \int_{\mathcal{O}_{c,n}} \chi(x) d^\times x.$$

Write  $\mathcal{O}_{K,v} = \mathcal{O}_v + \mathcal{O}_v \lambda$  then

$$\mathcal{O}_c = \mathcal{O}_v + \pi_v^c \mathcal{O}_v \lambda.$$

If  $n \geq c$ , then  $\mathcal{O}_{c,n} = \pi^n \mathcal{O}_K^\times$ . The integral vanishes as  $\chi$  has conductor  $\pi^c$ . If  $n < c$  then

$$\mathcal{O}_{c,n} = |\pi_v|^n \mathcal{O}_v^\times (1 + \pi_v^{c-n} \mathcal{O}_K).$$

The integration on  $\mathcal{O}_{n,c}$  vanishes unless  $n = 0$  as  $\chi$  has conductor  $\pi^c$ . Thus the total contribution is

$$\text{vol}(\mathcal{O}_{c_v}^\times) = 1.$$

We assume now that  $K_v/F_v$  is split. Then  $K_v = F_v^2$  and  $\mathcal{O}_c$  consists of integral elements  $(a, b)$  such that  $a \equiv b \pmod{\pi_v^c}$ . Write  $\chi = (\nu, \nu^{-1})$  then  $\nu$  has conductor  $\pi_v^c$ . It follows that

$$Z(1/2, \bar{\chi} \cdot \mu_K, \Phi_{K,c}) = \int_{(a,b) \in \mathcal{O}_c} \nu(a/b) \mu(ab) |ab|^{1/2} d^\times a d^\times b.$$

For a fixed  $b \in \mathcal{O}_v$ , the condition in  $a$  is as follows:

$$\begin{cases} a \in \pi^c \mathcal{O}_F, & \text{if } b \in \pi^c \mathcal{O}_v, \\ a \in b(1 + \pi^{c-n} \mathcal{O}_v), & \text{if } b \in \pi^n \mathcal{O}_v^\times \text{ with } n < c. \end{cases}$$

Since  $\nu$  has conductor  $c$ , the only case gives nontrivial contribution is when  $b \in \mathcal{O}_v^\times$  and  $a \in b(1 + \pi^c \mathcal{O}_v)$ . The contribution is given by

$$\text{vol}(\mathcal{O}_{c_v}^\times) = 1.$$

To compute  $\det h_v$ , we write  $K = F + F\sqrt{\lambda}$  and make the following embedding  $K \rightarrow M_2(F)$ :

$$(19.3) \quad a + b\lambda \mapsto \begin{pmatrix} a & b\lambda \\ b & a \end{pmatrix}.$$

Then  $\mathcal{O}_F^2$  corresponding to the lattice

$$\mathcal{O}_F + \mathcal{O}_F\sqrt{\lambda}.$$

Thus  $h_f$  satisfies

$$(0, 1)(\mathcal{O}_v + \mathcal{O}_v\sqrt{\lambda}) = (0, 1)\mathcal{O}_{c_v}h_f.$$

It follows that

$$\text{disc}(\mathcal{O}_v + \mathcal{O}_v\sqrt{\lambda}) = \text{disc}(\mathcal{O}_{c_v}) \det h_v^2.$$

Thus

$$\det h_v = \sqrt{4\lambda/D_v}$$

for a suitable  $D_v$  in its class modulo  $\mathcal{O}_v^\times$  such that  $4\lambda/D_v$  does have a square root in  $F_v^\times$ . In summary we have shown that

$$(19.4) \quad i_v(f_v) = L(1/2, \bar{\chi} \otimes \mu) \mu(\delta_v \sqrt{4\lambda/D_v}) |4\lambda/D_v|^{1/4}.$$

It remains to compute the periods at archimedean places  $v$ . For equation (19.1), we may take

$$h_v = \begin{pmatrix} |\lambda_v|^{1/2} & 0 \\ 0 & 1 \end{pmatrix}.$$

Then it is easy to see that

$$\Phi_{K,v}(x) = e^{-\pi|x|^2}.$$

Assume that  $\mu(x) = |x|^t$ ,  $\chi = 1$ , and notice that the measure on  $K_v^\times = \mathbb{C}^\times$  is induced from the standard  $d^\times x$  from  $\mathbb{R}^\times$  and one from  $\mathbb{C}^\times/\mathbb{R}^\times$  with volume one. Thus the measure has the form  $\frac{drd\theta}{\pi r}$  for polar coordinates  $re^{i\theta}$ . It follows that

$$\begin{aligned} Z(1/2, \bar{\chi} \cdot \mu_K, \Phi_{K,v}) &= \int_{\mathbb{C}^\times} e^{-\pi r^2} r^{2t+1} \frac{drd\theta}{\pi r} \\ &= \pi^{-1/2-t} \Gamma(t+1/2) \\ &= \mu(2) 2^{-1/2} L(1/2, \bar{\chi} \cdot \mu_K). \end{aligned}$$

The period at  $v$  is then given by

$$(19.5) \quad i_v(f_v) = 2^{-1} \mu_v(|4\lambda|^{1/2}) |4\lambda_v|^{1/4} L(1/2, \bar{\chi} \mu_K).$$

Let us set  $D_v = -1$  for archimedean places, then we obtain the same formula as (19.4). The proof of the Proposition is completed.

## References

- [1] M. Bertolini and H. Darmon, *Heegner points on Mumford-Tate curves*, Invent. Math. **126** (1996) pp. 413-456.
- [2] H. Carayol, *Sur la mauvaise réduction des courbes de Shimura*, Compositio Math. **59** (1986), pp 151-230.
- [3] S. Gelbart and H. Jacquet, *Forms of  $GL(2)$  from the analytic point of view*, Proc. Sympos. Pure Math. (Corvallis), **33**, Part I, Amer. Math. Soc., Providence, RI, (1979) pp. 213-251.
- [4] D. Goldfeld and S. Zhang, *The holomorphic kernel of the Rankin-Selberg convolution*, Asian J. Math. **3** (1999) 729–747.
- [5] B. H. Gross, *Heegner points and representation theory*, in this volume.
- [6] B. H. Gross and D. Zagier, *Heegner points and derivatives of  $L$ -series*, Invent. Math., **84** (1986), pp. 225-320.
- [7] H. Jacquet, *Automorphic forms on  $GL_2$  II*, Lect. Notes. Math. **289**, Springer-Verlag (1972).
- [8] H. Jacquet and R. Langlands, *Automorphic forms on  $GL(2)$* , Lect. Notes. Math. **114**, Springer-Verlag (1971).
- [9] J. Tate, *On the conjectures of Birch and Swinnerton-Dyer and a geometric analog*, Séminaire Bourbaki, Vol **9**, Exp. 306, 415-440, S.M.F, 1995.
- [10] V. Vatsal, *Special values for Rankin  $L$ -functions*, in this volume.
- [11] V. Vatsal, *Uniform distribution of Heegner points*, Invent. Math. (2002).

- [12] J. L. Waldspurger, *Sur les valeurs de certaines fonctions  $L$  automorphes en leur centre de symmétre*, Compositio Math. **54** (1985), 173-242.
- [13] J. L. Waldspurger, *Correspondence de Shimura et quaternions*, Forum Math., **3** (1991), 219-307.
- [14] Y. Tian, *Euler systems on Shimura curves*, Thesis at Columbia University (2003).
- [15] H. Xue, *Central values for twisted Rankin  $L$ -functions*, Thesis at Columbia University (2002).
- [16] S. Zhang, *Gross-Zagier formula for  $GL_2$* , Asian J. Math. **5**, (2001), pp. 183-290
- [17] S. Zhang, *Heights of Heegner points on Shimura curves*, Annals of mathematics (2), 153:1 (2000), pp 27-147.
- [18] S. Zhang, *Elliptic curves,  $L$ -functions, and  $CM$ -points*, in proceedings for Harvard-MIT joint conference on “current development in mathmetics” (2001), pp. 179-219.