

ON TWO INEQUALITY CONJECTURES FOR THE k -TH YAU NUMBERS OF ISOLATED HYPERSURFACE SINGULARITIES

NAVEED HUSSAIN, STEPHEN S.-T. YAU, AND HUAIQING ZUO

ABSTRACT. Let $(V, 0)$ be an isolated hypersurface singularity defined by the holomorphic function $f : (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}, 0)$. In our previous work, we introduced a series of novel Lie algebras associated to $(V, 0)$, i.e., k -th Yau algebra $L^k(V), k \geq 0$. It was defined to be the Lie algebra of derivations of the k -th moduli algebras $A^k(V) = \mathcal{O}_n/(f, m^k J(f)), k \geq 0$, where m is the maximal ideal of \mathcal{O}_n . I.e., $L^k(V) := \text{Der}(A^k(V), A^k(V))$. The dimension of $L^k(V)$ was denoted by $\lambda^k(V)$. The number $\lambda^k(V)$, which was called k -th Yau number, is a subtle numerical analytic invariant of $(V, 0)$. Furthermore, we formulated two conjectures for these k -th Yau number invariants: a sharp upper estimate conjecture of $\lambda^k(V)$ for weighted homogeneous isolated hypersurface singularities (see Conjecture 1.2) and an inequality conjecture $\lambda^{(k+1)}(V) > \lambda^k(V), k \geq 0$ (see Conjecture 1.1). In this article, we verify these two conjectures when k is small for large class of singularities.

Keywords. Derivation, Lie algebra, isolated singularity, Yau algebra.

MSC(2010). 14B05, 32S05.

1. INTRODUCTION

For any isolated hypersurface singularity $(V, 0) \subset (\mathbb{C}^n, 0)$ defined by the holomorphic function $f : (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}, 0)$, one has the moduli algebra $A(V) := \mathcal{O}_n/(f, \frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n})$ which is finite dimensional. Its dimension $\tau(V)$ is called Tyurina number. The order of the lowest nonvanishing term in the power series expansion of f at 0 is called the multiplicity (denoted by $\text{mult}(f)$) of the singularity $(V, 0)$. A polynomial $f \in \mathbb{C}[x_1, \dots, x_n]$ is said to be weighted homogeneous if there exist positive rational numbers w_1, \dots, w_n (weights of x_1, \dots, x_n) and d such that, $\sum a_i w_i = d$ for each monomial $\prod x_i^{a_i}$ appearing in f with nonzero coefficient. The number d is called weighted homogeneous degree (w -degree) of f with respect to weights w_j . The weight type of f is denoted as $(w_1, \dots, w_n; d)$. By a beautiful result of Saito [19], we shall always assume without loss of generality that $2w_i \leq d$ for all $1 \leq i \leq n$. Without loss of generality, we can assume that $w\text{-deg} f = 1$. and $w_i \leq \frac{1}{2}$.

The well-known Mather-Yau theorem [17] stated that: Let V_1 and V_2 be two isolated hypersurface singularities and, $A(V_1)$ and $A(V_2)$ be the moduli algebras, then

Both Yau and Zuo are supported by NSFC Grants 11961141005 and 11531007. Zuo is supported by NSFC Grant 11771231. Yau is supported by Tsinghua university start-up fund and Tsinghua university education foundation fund (042202008).

$(V_1, 0) \cong (V_2, 0) \iff A(V_1) \cong A(V_2)$. Motivated from the Mather-Yau theorem, the second author introduced the Lie algebra of derivations of moduli algebra $A(V)$, i.e., $L(V) = \text{Der}(A(V), A(V))$. The finite dimensional Lie algebra $L(V)$ was called Yau algebra and its dimension $\lambda(V)$ was called Yau number in [26]. The Yau algebra plays an important role in singularity theory (cf. [21], [11]). Yau and his collaborators have been systematically studying the Yau algebras of isolated hypersurface singularities begin from eighties ([23, 24, 25], [1, 2], [21], [27, 28], [7], [22], [5], [6], [4], [13]). In particular, Yau algebras of simple singularities and simple elliptic singularities were computed and a number of elaborate applications to deformation theory were presented in [1] and [21]. However, the Yau algebra can not characterize the simple singularities completely. In [8], it was shown that if X and Y are two simple singularities except the pair A_6 and D_5 , then $L(X) \cong L(Y)$ as Lie algebras if and only if X and Y are analytically isomorphic. Therefore, a natural question is to find new Lie algebras which can be used to distinguish singularities (at least for the simple singularities) completely. In our previous work [14], we introduced the series of new k -th Yau algebra associated to isolated hypersurface singularities. We defined this new k -th Yau algebra as follows.

Recall that we have the following theorem.

Theorem 1.1. ([10], Theorem 2.26) *Let $f, g \in m \subset \mathcal{O}_n$. The following are equivalent:*

- 1) $(V(f), 0) \cong (V(g), 0)$.
- 2) For all $k \geq 0$, $\mathcal{O}_n/(f, m^k J(f)) \cong \mathcal{O}_n/(g, m^k J(g))$ as \mathbb{C} -algebra.
- 3) There is some $k \geq 0$ such that $\mathcal{O}_n/(f, m^k J(f)) \cong \mathcal{O}_n/(g, m^k J(g))$ as \mathbb{C} -algebra, where $J(f) = (\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n})$.

In particular, if $k = 0$ and $k = 1$ above, and f, g define isolated singularities, then the claim of the equivalence of 1) and 3) is exactly the same as the Mather-Yau theorem.

Based on Theorem 1.1, it is natural for us to introduce the new series of k -th Yau algebras $L^k(V)$ which are defined to be the Lie algebra of derivations of the k -th moduli algebra $A^k(V) = \mathcal{O}_n/(f, m^k J(f))$, $k \geq 0$, i.e., $L^k(V) = \text{Der}(A^k(V), A^k(V))$. Its dimension is denoted as $\lambda^k(V)$. This number $\lambda^k(V)$ is a new numerical analytic invariant of a singularity. We call it k -th Yau number. In particular, $L^0(V)$ is exactly the Yau algebra, thus $L^0(V) = L(V)$, $\lambda^0(V) = \lambda(V)$.

On the one hand, since $L(V)$ can not characterize the simple singularities completely, so there is a natural question: whether these simple singularities (or which classes of more general singularities) can be characterized completely by the Lie algebra $L^k(V)$, $k \geq 1$?

In [16], we have proven that the simple singularities V can be characterized completely using the $L^1(V)$.

Theorem 1.2. [16] *If X and Y are two simple hypersurface singularities, then $L^1(X) \cong L^1(Y)$ as Lie algebras, if and only if X and Y are analytically isomorphic.*

We believe Theorem 1.2 is also true for $L^k(V)$, $k > 1$. Therefore the k -th Yau algebra $L^k(V)$, $k \geq 1$, is more subtle comparing to the Yau algebra $L(V)$ in some sense.

Furthermore, since derivations of moduli algebras are analogs of vector fields on smooth manifolds, such direction of research is in the spirit of the classical theorem of Pursell and Shanks stating that the Lie algebra of smooth vectors fields on a smooth manifold determines the diffeomorphism type of the manifold [18]. It is interesting to investigate the structure of k -th Yau algebras and find out, for which classes of singularities those Lie algebras determine the analytic or topological structures of singularities by analogy with the mentioned result of Pursell and Shanks. In fact, Theorem 1.2 yields a similar for the simple singularities.

On the other hand, it is well known that finite dimensional Lie algebras are semi-direct product of the semi-simple Lie algebras and solvable Lie algebras. Brieskorn gave the connection between simple Lie algebras and simple singularities. Simple Lie algebras have been well understood, but not the solvable (nilpotent) Lie algebras. It is extremely important to establish connections between singularities and solvable (nilpotent) Lie algebras. In fact $L^k(V)$ are finite dimensional solvable (nilpotent) Lie algebras naturally from isolated hypersurface singularities. These objects $L^k(V)$ help us to understand the solvable (nilpotent) Lie algebras from the geometric point of view. Moreover, it is known that the classification of nilpotent Lie algebras in higher dimensions (> 7) remains to be a vast open area. There are one-parameter families of non-isomorphic nilpotent Lie algebras (but no two-parameter families) in dimension seven. Dimension seven is the watershed of the existence of such families. It is well-known that no such family exists in dimension less than seven, while it is hard to construct one-parameter family in dimension greater than seven. However, such examples are hard to construct [20].

Recall that Griffiths has studied the Torelli problem when a family of complex projective hypersurfaces in $\mathbb{C}P^n$ is given and his school asks whether the period map is injective on that family, i.e., whether the family of complex hypersurfaces can be distinguished by means of their Hodge structures. A complex projective hypersurface in $\mathbb{C}P^n$ can be viewed as a complex hypersurface with isolated singularity in \mathbb{C}^{n+1} . Let $V = \{z \in \mathbb{C}^{n+1} : f(z) = 0\}$ be a complex hypersurface with isolated singularity at the origin. Seeley and Yau [21]

investigated the family of isolated complex hypersurface singularities using Yau algebras $L(V)$ and obtained two strong Torelli-type theorems for simple elliptic singularities \tilde{E}_7 and \tilde{E}_8 . We obtained the following similar result for $L^k(V)$.

Theorem 1.3. [16] $L^2(\tilde{E}_6)$, $L^1(\tilde{E}_7)$, $L^2(\tilde{E}_7)$, $L^1(\tilde{E}_8)$ and $L^2(\tilde{E}_8)$ are non-trivial one-parameter families. Thus the weak Torelli-type theorems hold for simple elliptic singularities \tilde{E}_6 , \tilde{E}_7 and \tilde{E}_8 .

The second author [23] showed that the Yau algebra $L(V)$ of the family \tilde{E}_6 is constant, i.e., it does not depend on the parameter. In a recent paper [12], we have shown that the first Yau algebra $L^1(V)$ is also constant,. However, the strong Torelli-type theorem holds for \tilde{E}_6 using k -th Yau algebra $L^k(V)$, $k \geq 2$.

Theorem 1.4. [12] Let $\{V_t\}$ represent a family of simple elliptic singularities \tilde{E}_6 . If $k \geq 2$, then L_t^k and L_s^k are isomorphic as Lie algebras if and only if V_t is biholomorphic to V_s .

As a corollary of Theorem 1.3 and Theorem 1.4, we can obtain many non-trivial one-parameter families of solvable (nilpotent) Lie algebras in dimension greater than seven.

Since the k -th Yau algebras $L^k(V)$ are more subtle invariants than the Yau algebras, we believe that these new Lie algebras $L^k(V)$ and numerical invariants $\lambda^k(V)$ will also play an important role in the study of singularities.

In this paper, we investigate the new analytic invariants $\lambda^k(V)$. A natural question arises: are there any numerical relations between the invariants $\lambda^k(V)$, $k \geq 0$? We proposed the following conjecture:

Conjecture 1.1. ([15]) With the above notations, let $(V, 0)$ be an isolated hypersurface singularity defined by $f \in \mathcal{O}_n$, $n \geq 2$, and $\text{mult}(f) \geq 3$. Then

$$\lambda^{(k+1)}(V) > \lambda^k(V), k \geq 0.$$

The Conjecture 1.1 has already been verified for binomial singularities (see Definition 2.4) when $k = 0, 1$, and trinomial singularities (see Definition 2.4) when $k = 0$ by the authors in [15]. In this paper we shall prove this conjecture for trinomial singularities when $k = 1$ (see Theorem A).

It is also interesting to bound the k -th Yau number of weighted homogeneous singularities with a number that depends on weight type. In [28], Yau and Zuo firstly proposed the sharp upper estimate conjecture that bound the Yau number. They also proved that this conjecture holds in the case of binomial isolated hypersurface singularities. Furthermore,

in [13], this conjecture was verified for trinomial singularities. We proposed the following sharp upper estimate conjecture which is a generalization of the conjecture in [28].

Conjecture 1.2. ([14]) Assume that $\lambda^k(\{x_1^{a_1} + \cdots + x_n^{a_n} = 0\}) = h_k(a_1, \cdots, a_n)$, ($k \geq 0$). Let $(V, 0) = \{(x_1, x_2, \cdots, x_n) \in \mathbb{C}^n : f(x_1, x_2, \cdots, x_n) = 0\}$, ($n \geq 2$) be an isolated singularity defined by the weighted homogeneous polynomial $f(x_1, x_2, \cdots, x_n)$ of weight type $(w_1, w_2, \cdots, w_n; 1)$. Then $\lambda^k(V) \leq h_k(1/w_1, \cdots, 1/w_n)$.

The Conjecture 1.2 tells, when fixing a weight type, the Brieskorn singularity $x_1^{a_1} + \cdots + x_n^{a_n}$ has maximal k -th Yau number. It has already been verified for binomial singularities and trinomial singularities when $k = 0$ in [28, 13] respectively and when $k = 1$ [14]. In this paper we shall prove this conjecture for binomial singularities and trinomial singularities when $k = 2$ (see Theorem B, Theorem C, and Theorem D). We obtain the following main results.

Theorem A. *Let $(V, 0)$ be a fewnomial singularity defined by the weighted homogeneous polynomial $f(x_1, x_2, x_3)$ (see proposition 2.2) with $\text{mult}(f) \geq 3$, then*

$$\lambda^2(V) > \lambda^1(V).$$

Theorem B. *Let $(V, 0)$ be a binomial singularity defined by the weighted homogeneous polynomial $f(x_1, x_2)$ (see Corollary 2.1) with weight type $(w_1, w_2; 1)$ and $\text{mult}(f) \geq 3$, then*

$$\lambda^2(V) \leq h_2\left(\frac{1}{w_1}, \frac{1}{w_2}\right) = \begin{cases} \frac{2}{w_1 w_2} - 3\left(\frac{1}{w_1} + \frac{1}{w_2}\right) + 17; & w_1 \leq \frac{1}{5}, w_2 \leq \frac{1}{5} \\ \frac{3}{w_2} + 5; & w_1 = \frac{1}{3}, w_2 \leq \frac{1}{4} \\ 13; & w_1 = \frac{1}{3}, w_2 = \frac{1}{3} \\ \frac{5}{w_2} + 4; & w_1 = \frac{1}{4}, w_2 \leq \frac{1}{5} \\ 23; & w_1 = \frac{1}{4}, w_2 = \frac{1}{4}. \end{cases}$$

Remark 1.1. *If $\text{mult}f(x_1, x_2) = 2$, then $f(x_1, x_2)$ is contact equivalent [10] to $x_1^2 + x_2^b$ where $b \geq 2$. Thus our Conjecture 1.2 is obviously true for this case. Therefore in Theorem B, we only need to consider $\text{mult}(f) \geq 3$.*

Theorem C. *Let $(V, 0)$ be a fewnomial singularity defined by the weighted homogeneous polynomial $f(x_1, x_2, x_3)$ (see Proposition 2.2) with weight type $(w_1, w_2, w_3; 1)$ and $\text{mult}(f) \geq 3$, then*

$$\lambda^2(V) \leq h_2\left(\frac{1}{w_1}, \frac{1}{w_2}, \frac{1}{w_3}\right) = \frac{3}{w_1 w_2 w_3} + 5\left(\frac{1}{w_1} + \frac{1}{w_2} + \frac{1}{w_3}\right) - 4\left(\frac{1}{w_1 w_2} + \frac{1}{w_1 w_3} + \frac{1}{w_2 w_3}\right) + 34, \quad w_1 \leq \frac{1}{3}, w_2 \leq \frac{1}{3}, w_3 \leq \frac{1}{3}.$$

Theorem D. Let $(V, 0)$ be a fewnomial singularity defined by the weighted homogeneous polynomial $f(x_1, x_2, x_3)$ (see Proposition 2.2) with weight type $(w_1, w_2, w_3; 1)$ and $\text{mult}(f) = 2$ (see Remark 1.2), then

$$\lambda^2(V) \leq h_2\left(\frac{1}{w_1}, \frac{1}{w_2}, 2\right) = \begin{cases} \frac{2}{w_1 w_2} - 3\left(\frac{1}{w_1} + \frac{1}{w_2}\right) + 36; & w_1 \leq \frac{1}{5}, w_2 \leq \frac{1}{5} \\ \frac{3}{w_2} + 22; & w_1 = \frac{1}{3}, w_2 \leq \frac{1}{4} \\ 30; & w_1 = \frac{1}{3}, w_2 = \frac{1}{3} \\ \frac{5}{w_2} + 23; & w_1 = \frac{1}{4}, w_2 \leq \frac{1}{5} \\ 42; & w_1 = \frac{1}{4}, w_2 = \frac{1}{4}. \end{cases}$$

Remark 1.2. If $f(x_1, x_2, x_3)$ is fewnomial and $\text{mult}(f(x_1, x_2, x_3)) = 2$, then $f(x_1, x_2, x_3)$ is contact equivalent to one of the following cases:

Case 1: A) $x_1^{a_1} + x_2^{a_2} + x_3^2$, $a_1, a_2 \geq 3$; B) $x_1^{a_1} x_2 + x_2^{a_2} + x_3^2$, $a_1 \geq 2, a_2 \geq 3$; C) $x_1^{a_1} x_2 + x_2^{a_2} x_1 + x_3^2$, $a_1, a_2 \geq 2$;

Case 2. A) $x_1^{a_1} + x_2^2 + x_3^2$, $a_1 \geq 2$.

The Conjecture 1.2 is true for case 2. Thus in Theorem D, we only need to verify the Conjecture 1.2 for case 1.

2. GENERALITIES ON DERIVATION LIE ALGEBRAS OF ISOLATED SINGULARITIES

In this section, we shall briefly define the basic definitions and important results that are helpful to solve the problem. The following basic concepts and results will be used to compute the derivation Lie algebras of isolated hypersurface singularities.

Let A, B be associative algebras over \mathbb{C} . The subalgebra of endomorphisms of A generated by the identity element and left and right multiplications by elements of A is called multiplication algebra $M(A)$ of A . The centroid $C(A)$ is defined as the set of endomorphisms of A which commute with all elements of $M(A)$. Obviously, $C(A)$ is a unital subalgebra of $\text{End}(A)$. The following statement is a particular case of a general result from Proposition 1.2 of [3]. Let $S = A \otimes B$ be a tensor product of finite dimensional associative algebras with units. Then

$$\text{Der}S \cong (\text{Der}A) \otimes C(B) + C(A) \otimes (\text{Der}B).$$

We will only use this result for commutative associative algebras with unit, in which case the centroid coincides with the algebra itself and one has following result for commutative associative algebras A, B :

Theorem 2.1. ([3]) For commutative associative algebras A, B ,

$$\text{Der}S \cong (\text{Der}A) \otimes B + A \otimes (\text{Der}B). \quad (1)$$

We shall use this formula in the sequel.

Definition 2.1. Let J be an ideal in an analytic algebra S . Then $\text{Der}_J S \subseteq \text{Der}_{\mathbb{C}} S$ is Lie subalgebra of all $\sigma \in \text{Der}_{\mathbb{C}} S$ for which $\sigma(J) \subset J$.

We shall use the following well-known result to compute the derivations.

Theorem 2.2. ([28]) *Let J be an ideal in $R = \mathbb{C}\{x_1, \dots, x_n\}$. Then there is a natural isomorphism of Lie algebras*

$$(\text{Der}_J R)/(J \cdot \text{Der}_{\mathbb{C}} R) \cong \text{Der}_{\mathbb{C}}(R/J).$$

Recall that a derivation of commutative associative algebra A is defined as a linear endomorphism D of A satisfying the Leibniz rule: $D(ab) = D(a)b + aD(b)$. Thus for such an algebra A one can consider the Lie algebra of its derivations $\text{Der}(A, A)$ with the bracket defined by the commutator of linear endomorphisms.

Definition 2.2. Let $f(x_1, \dots, x_n)$ be a complex polynomial and $V = \{f = 0\}$ be a germ of an isolated hypersurface singularity at the origin in \mathbb{C}^n . Let $A^k(V) = \mathcal{O}_n/(f, m^k J(f))$, $1 \leq k \leq n$ be a moduli algebra. Then $\text{Der}(A^k(V), A^k(V))$ defined the derivation Lie algebras $L^k(V)$. The $\lambda^k(V)$ is the dimension of derivation Lie algebra $L^k(V)$.

It is noted that when $k = 0$, then derivation Lie algebra is called Yau algebra.

Definition 2.3. A polynomial $f \in \mathbb{C}[x_1, x_2, \dots, x_n]$ is called quasi-homogeneous (or weighted homogeneous) if there exist positive rational numbers w_1, \dots, w_n (called weights of indeterminates x_j) and d such that, for each monomial $\prod x_j^{k_j}$ appearing in f with non-zero coefficient, one has $\sum w_j k_j = d$. The number d is called the quasi-homogeneous degree (w -degree) of f with respect to weights w_j and is denoted $\text{deg } f$. The collection $(w; d) = (w_1, \dots, w_n; d)$ is called the quasi-homogeneity type (qh-type) of f .

Definition 2.4. An isolated hypersurface singularity in \mathbb{C}^n is fewnomial if it can be defined by a n -nomial in n variables and it is a weighted homogeneous fewnomial isolated singularity if it can be defined by a weighted homogeneous fewnomial. 2-nomial (resp. 3-nomial) isolated hypersurface singularity is also called binomial (resp. trinomial) singularity.

Proposition 2.1. *Let f be a weighted homogeneous fewnomial isolated singularity with $\text{mult}(f) \geq 3$. Then f analytically equivalent to a linear combination of the following three series:*

Type A. $x_1^{a_1} + x_2^{a_2} + \dots + x_{n-1}^{a_{n-1}} + x_n^{a_n}$, $n \geq 1$,

Type B. $x_1^{a_1} x_2 + x_2^{a_2} x_3 + \dots + x_{n-1}^{a_{n-1}} x_n + x_n^{a_n}$, $n \geq 2$,

Type C. $x_1^{a_1} x_2 + x_2^{a_2} x_3 + \dots + x_{n-1}^{a_{n-1}} x_n + x_n^{a_n} x_1$, $n \geq 2$.

Proposition 2.1 has an immediate corollary.

Corollary 2.1. *Each binomial isolated singularity is analytically equivalent to one from the three series: A) $x_1^{a_1} + x_2^{a_2}$, B) $x_1^{a_1}x_2 + x_2^{a_2}$, C) $x_1^{a_1}x_2 + x_2^{a_2}x_1$.*

Wolfgang and Atsushi [9] give the following classification of weighted homogeneous fewnomial singularities.

Proposition 2.2. ([9]) *Let $f(x_1, x_2, x_3)$ be a weighted homogeneous fewnomial isolated singularity with $\text{mult}(f) \geq 3$. Then f is analytically equivalent to following five types:*

Type 1. $x_1^{a_1} + x_2^{a_2} + x_3^{a_3}$,

Type 2. $x_1^{a_1}x_2 + x_2^{a_2}x_3 + x_3^{a_3}$,

Type 3. $x_1^{a_1}x_2 + x_2^{a_2}x_3 + x_3^{a_3}x_1$,

Type 4. $x_1^{a_1} + x_2^{a_2} + x_3^{a_3}x_2$,

Type 5. $x_1^{a_1}x_2 + x_2^{a_2}x_1 + x_3^{a_3}$.

In order to prove the Theorem A, we need to use the following propositions.

Proposition 2.3. ([14]) *Let $(V, 0)$ be a fewnomial surface isolated singularity of type 1 which is defined by $f = x_1^{a_1} + x_2^{a_2} + x_3^{a_3}$ ($a_1 \geq 3, a_2 \geq 3, a_3 \geq 3$) with weight type $(\frac{1}{a_1}, \frac{1}{a_2}, \frac{1}{a_3}; 1)$. Then*

$$\lambda^1(V) = 3a_1a_2a_3 + 5(a_1 + a_2 + a_3) - 4(a_1a_2 + a_1a_3 + a_2a_3) + 6.$$

Proposition 2.4. ([14]) *Let $(V, 0)$ be a fewnomial surface isolated singularity of type 2 which is defined by $f = x_1^{a_1}x_2 + x_2^{a_2}x_3 + x_3^{a_3}$ ($a_1 \geq 2, a_2 \geq 2, a_3 \geq 3$) with weight type $(\frac{1-a_3+a_2a_3}{a_1a_2a_3}, \frac{a_3-1}{a_2a_3}, \frac{1}{a_3}; 1)$. Then*

$$\lambda^1(V) = \begin{cases} 4a_1a_3 - 2a_1 - 3a_3 + 11; & a_1 \geq 3, a_2 = 2, a_3 \geq 3 \\ 5a_3 + 7; & a_1 = 2, a_2 = 2, a_3 \geq 3 \\ 3a_1a_2a_3 - 2a_1a_2 - 2a_1a_3 - 4a_2a_3 + 2a_1 + 2a_2 + 6a_3 + 5; & \text{Otherwise.} \end{cases}$$

Proposition 2.5. ([14]) *Let $(V, 0)$ be a fewnomial surface isolated singularity of type 3 which is defined by $f = x_1^{a_1}x_2 + x_2^{a_2}x_3 + x_3^{a_3}x_1$ ($a_1 \geq 2, a_2 \geq 2, a_3 \geq 2$) with weight type*

$$\left(\frac{1 - a_3 + a_2a_3}{1 + a_1a_2a_3}, \frac{1 - a_1 + a_1a_3}{1 + a_1a_2a_3}, \frac{1 - a_2 + a_1a_2}{1 + a_1a_2a_3}; 1 \right).$$

Then

$$\lambda^1(V) = \begin{cases} 24; & a_1 = 2, a_2 = 2, a_3 = 2 \\ 3a_1a_2a_3 + 2(a_1 + a_2 + a_3) - 2(a_1a_2 + a_1a_3 + a_2a_3) + 11; & \text{Otherwise.} \end{cases}$$

Proposition 2.6. ([14]) *Let $(V, 0)$ be a fewnomial surface isolated singularity of type 4 which is defined by $f = x_1^{a_1} + x_2^{a_2} + x_3^{a_3}x_2$ ($a_1 \geq 3, a_2 \geq 3, a_3 \geq 2$) with weight type $(\frac{1}{a_1}, \frac{1}{a_2}, \frac{a_2-1}{a_2a_3}; 1)$. Then*

$$\lambda^1(V) = \begin{cases} 5a_1a_2 - a_1 - 7a_2 + 15; & a_1 \geq 3, a_2 \geq 3, a_3 = 3 \\ 3a_1a_2a_3 - 4a_1a_2 - 3a_1a_3 - 4a_2a_3 + 8a_1 + 5a_2 + 5a_3 - 1; & \text{Otherwise.} \end{cases}$$

Proposition 2.7. ([14]) *Let $(V, 0)$ be a fewnomial surface isolated singularity of type 5 which is defined by $f = x_1^{a_1}x_2 + x_2^{a_2}x_1 + x_3^{a_3}$ ($a_1 \geq 2, a_2 \geq 2, a_3 \geq 3$) with weight type $(\frac{a_2-1}{a_1a_2-1}, \frac{a_1-1}{a_1a_2-1}, \frac{1}{a_3}; 1)$. Then*

$$\lambda^1(V) = \begin{cases} 4a_2a_3 - 6a_2 + 12; & a_1 = 2, a_2 \geq 2, a_3 \geq 3 \\ 3a_1a_2a_3 - 4a_1a_2 - 2a_2a_3 - 2a_1a_3 + 2a_1 + 2a_2 + 6a_3 + 6; & \text{Otherwise.} \end{cases}$$

3. PROOF OF MAIN THEOREMS

Proposition 3.1. *Let $(V, 0)$ be a weighted homogeneous fewnomial isolated singularity of type A which is defined by $f = x_1^{a_1} + x_2^{a_2}$ ($a_1 \geq 2, a_2 \geq 2$) with weight type $(\frac{1}{a_1}, \frac{1}{a_2}; 1)$. Then*

$$\lambda^2(V) = \begin{cases} 2a_1a_2 - 3(a_1 + a_2) + 17; & a_1 \geq 5, a_2 \geq 5 \\ 3a_2 + 5; & a_1 = 3, a_2 \geq 4 \\ 13; & a_1 = 3, a_2 = 3 \\ 5a_2 + 4; & a_1 = 4, a_2 \geq 5 \\ 23; & a_1 = 4, a_2 = 4 \\ a_2 + 5; & a_1 = 2, a_2 \geq 3 \\ 6; & a_1 = 2, a_2 = 2. \end{cases}$$

Proof. It follows that the generalized moduli algebra

$$A^2(V) = \mathbb{C}\{x_1, x_2\}/(f, m^2J(f)),$$

has dimension $a_1a_2 - (a_1 + a_2) + 6$ and has a monomial basis of the form:

$$(1) \text{ if } a_1 \geq 3, \{x_1^{i_1}x_2^{i_2}, 1 \leq i_1 \leq a_1 - 2; 0 \leq i_2 \leq a_2 - 2; x_1^{a_1-1}; x_1^{a_1-1}x_2; x_1x_2^{a_2-1}; x_2^{i_2}, 0 \leq i_2 \leq a_2\}, \quad (2)$$

$$(2) \text{ if } a_1 = 2, a_2 \geq 3, \{x_2^{i_2}, 0 \leq i_2 \leq a_2; x_1x_2; x_1\}, \quad (3)$$

$$(3) \text{ if } a_1 = 2, a_2 = 2, \{1; x_1; x_1x_2; x_2; x_2^2\}, \quad (4)$$

$$(4) \text{ if } a_1 = 1, a_2 \geq 1, \{1; x_2\}, \quad (5)$$

with the following relations:

$$x_1^{a_1+1} = 0, \quad (6)$$

$$x_1^{a_1-1}x_2^2 = 0, \quad (7)$$

$$x_1^{a_1} x_2 = 0, \quad (8)$$

$$x_1^2 x_2^{a_2-1} = 0, \quad (9)$$

$$x_2^{a_2+1} = 0, \quad (10)$$

$$x_1 x_2^{a_2} = 0. \quad (11)$$

In order to compute a derivation D of $A^2(V)$ it suffices to indicate its values on the generators x_1, x_2 which can be written in terms of the basis (2), (3), (4) or (5). Without loss of generality, we write

$$Dx_j = \sum_{i_1=1}^{a_1-2} \sum_{i_2=0}^{a_2-2} c_{i_1, i_2}^j x_1^{i_1} x_2^{i_2} + c_{a_1-1, 0}^j x_1^{a_1-1} + c_{a_1-1, 1}^j x_1^{a_1-1} x_2 + c_{1, a_2-1}^j x_1 x_2^{a_2-1} + \sum_{i_2=0}^{a_2} c_{0, i_2}^j x_2^{i_2}, \quad j = 1, 2.$$

Using the relations (6)-(11) one easily finds the necessary and sufficient conditions defining a derivation of $A^2(V)$ as follows:

$$c_{0,0}^1 = c_{0,1}^1 = \cdots = c_{0, a_2-4}^1 = 0; \quad (12)$$

$$c_{0,0}^2 = c_{1,0}^2 = \cdots = c_{a_1-4,0}^2 = 0; \quad (13)$$

$$a_1 c_{1,0}^1 = a_2 c_{0,1}^2. \quad (14)$$

Using (12)-(14) we obtain the following description of the Lie algebras in question. The following derivations form a basis of $\text{Der}A^2(V)$:

$$\begin{aligned} & x_1^{i_1} x_2^{i_2} \partial_1, 1 \leq i_1 \leq a_1 - 2, 1 \leq i_2 \leq a_2 - 2; x_1^{a_1-1} x_2 \partial_1; x_1 x_2^{a_2-1} \partial_1; x_2^{i_2} \partial_1, a_2 - 3 \leq i_2 \leq a_2; \\ & x_1^{i_1} \partial_1, 2 \leq i_1 \leq a_1 - 1; x_2^{i_2} \partial_2, 2 \leq i_2 \leq a_2; x_1^{i_1} \partial_2, a_1 - 3 \leq i_1 \leq a_1 - 1; \\ & x_1^{i_1} x_2^{i_2} \partial_2, 1 \leq i_1 \leq a_1 - 2, 1 \leq i_2 \leq a_2 - 2; x_1^{a_1-1} x_2 \partial_2; \\ & x_1 x_2^{a_2-1} \partial_2; x_1 \partial_1 + \frac{a_1}{a_2} x_2 \partial_2. \end{aligned}$$

Therefore we have the following formula

$$\lambda^2(V) = 2a_1 a_2 - 3(a_1 + a_2) + 17.$$

In case of $a_1 = 3, a_2 \geq 4$, we have following derivations form a basis of $\text{Der}A^2(V)$:

$$\begin{aligned} & x_1 x_2^{i_2} \partial_1, 1 \leq i_2 \leq a_2 - 1; x_1^2 x_2 \partial_1; x_1^2 \partial_1; x_2^{i_2} \partial_1, a_2 - 2 \leq i_2 \leq a_2; \\ & x_1 x_2^{i_2} \partial_2, 1 \leq i_2 \leq a_2 - 1; x_1^2 x_2 \partial_2; x_1^2 \partial_2; x_2^{i_2} \partial_2, 2 \leq i_2 \leq a_2; x_1 \partial_1 + \frac{3}{a_2} x_2 \partial_2. \end{aligned}$$

Therefore we have the following formula

$$\lambda^2(V) = 3a_2 + 5.$$

In case of $a_1 = 3, a_2 = 3$, we have following derivations form a basis of $\text{Der}A^2(V)$:

$$x_2^2 \partial_1; x_2^3 \partial_1; x_1 x_2 \partial_1; x_1 x_2^2 \partial_1; x_1^2 \partial_1; x_1^2 x_2 \partial_1; x_2^2 \partial_2; x_2^3 \partial_2; x_1 x_2 \partial_2; x_1 x_2^2 \partial_2; x_1^2 \partial_2; x_1^2 x_2 \partial_2; x_1 \partial_1 + x_2 \partial_2.$$

In case of $a_1 = 4, a_2 \geq 5$, we have following derivations form a basis of $\text{Der}A^2(V)$:

$$x_2^{i_2}\partial_1, a_2-3 \leq i_2 \leq a_2; x_1\partial_1 + \frac{4}{a_2}x_2\partial_2; x_1^{i_1}x_2\partial_1, 1 \leq i_1 \leq 2, 1 \leq i_2 \leq a_2-2; x_1x_2^{a_2-1}\partial_1; x_1^2\partial_1; x_1^3\partial_1; x_1^3x_2\partial_1; x_2^{i_2}\partial_2, 2 \leq i_2 \leq a_2; x_1^{i_1}x_2^{i_2}\partial_2, 1 \leq i_1 \leq 2, 1 \leq i_2 \leq a_2-2; x_1x_2^{a_2-1}\partial_2; x_1^2\partial_2; x_1^3\partial_2; x_1^3x_2\partial_2.$$

Therefore we have the following formula

$$\lambda^2(V) = 5a_2 + 4.$$

In case of $a_1 = 4, a_2 = 4$, we have following derivations form a basis of $\text{Der}A^2(V)$:

$$x_2^2\partial_1; x_2^3\partial_1; x_2^4\partial_1; x_1\partial_1 + x_2\partial_2; x_1x_2\partial_1; x_1x_2^2\partial_1; x_1x_2^3\partial_1; x_1^2\partial_1; x_1^2x_2\partial_1; x_1^2x_2^2\partial_1; x_1^3\partial_1; x_1^3x_2\partial_1; x_2^2\partial_2; x_2^3\partial_2; x_2^4\partial_2; x_1x_2\partial_2; x_1x_2^2\partial_2; x_1x_2^3\partial_2; x_1^2\partial_2; x_1^2x_2\partial_2; x_1^2x_2^2\partial_2; x_1^3\partial_2; x_1^3x_2\partial_2.$$

Therefore we have the following formula

$$\lambda^2(V) = 23.$$

In case of $a_1 = 2, a_2 \geq 3$, we have following derivations which form a basis of $L^2(V)$:

$$x_2^{i_2}\partial_1, a_2 - 1 \leq i_2 \leq a_2; x_1\partial_1 + \frac{2}{a_2}x_2\partial_2; x_1x_2\partial_1; x_2^{i_2}\partial_2, 2 \leq i_2 \leq a_2; x_1\partial_2; x_1x_2\partial_2.$$

Therefore we get following formula

$$\lambda^2(V) = a_2 + 5.$$

In case of $a_1 = 2, a_2 = 2$, we have following derivations which form a basis of $L^2(V)$:

$$x_2\partial_1 + x_1\partial_2; x_2^2\partial_1; x_1\partial_1 + x_2\partial_2; x_1x_2\partial_1; x_2^2\partial_2; x_1x_2\partial_2.$$

Therefore we get following formula

$$\lambda^2(V) = 6.$$

□

Proposition 3.2. *Let $(V, 0)$ be a binomial isolated singularity of type B which is defined by $f = x_1^{a_1}x_2 + x_2^{a_2}$ ($a_1 \geq 1, a_2 \geq 2$) with weight type $(\frac{a_2-1}{a_1a_2}, \frac{1}{a_2}; 1)$. Then*

$$\lambda^2(V) = \begin{cases} 2a_1a_2 - 2a_1 - 3a_2 + 20; & a_1 \geq 5, a_2 \geq 5 \\ 5a_2 + 12; & a_1 = 4, a_2 \geq 5 \\ 31; & a_1 = 4, a_2 = 4 \\ 4a_1 + 7; & a_1 \geq 3, a_2 = 3 \\ 2a_1 + 5; & a_1 \geq 2, a_2 = 2 \\ a_2 + 11; & a_1 = 2, a_2 \geq 4 \\ 13; & a_1 = 2, a_2 = 3 \\ 6; & a_1 = 1, a_2 \geq 2. \end{cases}$$

Furthermore, we need to show that when $a_1 \geq 5, a_2 \geq 5$, then $2a_1a_2 - 2a_1 - 3a_2 + 20 \leq \frac{2a_1a_2^2}{a_2-1} - 3(\frac{a_1a_2}{a_2-1} + a_2) + 17$.

Proof. It follows that the generalized moduli algebra

$$A^2(V) = \mathbb{C}\{x_1, x_2\}/(f, m^2J(f)),$$

has dimension $a_1a_2 - a_2 + 6$ and has a monomial basis of the form:

$$(1) \text{if } a_1 \geq 2, a_2 \geq 3, \{x_1^{i_1}x_2^{i_2}, 1 \leq i_1 \leq a_1 - 2; 1 \leq i_2 \leq a_2 - 2; x_1^{i_1}x_2^{a_2-1}, 1 \leq i_1 \leq a_1 - 2; x_1^{a_1-1}x_2^{i_2}, 1 \leq i_2 \leq 2; x_2^{i_2}, 0 \leq i_2 \leq a_2; x_1^{i_1}, 1 \leq i_1 \leq a_1 + 1\}, \quad (15)$$

$$(2) \text{if } a_1 \geq 2, a_2 = 2, \{x_1^{i_1}x_2^{i_2}, 1 \leq i_1 \leq a_1 - 1; 0 \leq i_2 \leq 1; x_2^{i_2}, 0 \leq i_2 \leq 2; x_1^{i_1}, a_1 \leq i_1 \leq a_1 + 1\}, \quad (16)$$

$$(3) \text{if } a_1 = 1, a_2 \geq 2, \{1; x_1; x_1^2; x_2; x_2^2\}, \quad (17)$$

$$(4) \text{if } a_1 \geq 1, a_2 = 1, \{x_1^{i_1}, 0 \leq i_1 \leq a_1 + 1\}, \quad (18)$$

with the following relations:

$$x_1^{a_1+1}x_2 = 0, \quad (19)$$

$$x_1^{a_1-1}x_2^3 = 0, \quad (20)$$

$$x_1^{a_1}x_2^2 = 0, \quad (21)$$

$$x_1^{a_1+2} + a_2x_1^2x_2^{a_2-1} = 0, \quad (22)$$

$$x_1^{a_1}x_2^2 + a_2x_2^{a_2+1} = 0, \quad (23)$$

$$x_1^{a_1+1}x_2 + a_2x_1x_2^{a_2} = 0. \quad (24)$$

In order to compute a derivation D of $A^2(V)$ it suffices to indicate its values on the generators x_1, x_2 which can be written in terms of the basis (15), (16), (17) or (18). Without loss of generality, we write

$$Dx_j = \sum_{i_1=1}^{a_1-2} \sum_{i_2=1}^{a_2-2} c_{i_1, i_2}^j x_1^{i_1} x_2^{i_2} + \sum_{i_1=1}^{a_1-2} c_{i_1, a_2-1}^j x_1^{i_1} x_2^{a_2-1} + \sum_{i_2=1}^2 c_{a_1-1, i_2}^j x_1^{a_1-1} x_2^{i_2} + \sum_{i_2=0}^{a_2} c_{0, i_2}^j x_2^{i_2} + \sum_{i_1=1}^{a_1+1} c_{i_1, 0}^j x_1^{i_1}, \quad j = 1, 2.$$

Using the relations (19)-(24) one easily finds the necessary and sufficient conditions defining a derivation of $A^2(V)$ as follows:

$$c_{0,0}^1 = c_{0,1}^1 = \cdots = c_{0,a_2-4}^1 = 0; \quad (25)$$

$$c_{0,0}^2 = c_{1,0}^2 = \cdots = c_{a_1-3,0}^2 = 0; \quad (26)$$

$$c_{1,0}^1 = c_{0,1}^2, c_{2,0}^1 = c_{1,1}^2, c_{3,0}^1 = c_{2,1}^2, \cdots, c_{a_1-3,0}^1 = c_{a_1-4,1}^2. \quad (27)$$

Using (25)-(27) we obtain the following description of the Lie algebras in question. The following derivations form a basis of $\text{Der}A^2(V)$:

$$\begin{aligned} & x_1^{i_1} x_2^{i_2} \partial_1, 1 \leq i_1 \leq a_1 - 2, 1 \leq i_2 \leq a_2 - 1; x_1^{i_1} \partial_1 + \frac{a_1}{a_2 - 1} x_1^{i_1-1} x_2 \partial_2, 1 \leq i_1 \leq a_1 - 3; \\ & x_1^{i_1} \partial_1, a_1 - 2 \leq i_1 \leq a_1 + 1; x_1^{a_1-1} x_2^{i_2} \partial_1, 1 \leq i_2 \leq 2; x_1^{i_1} x_2^{i_2} \partial_2, 1 \leq i_1 \leq a_1 - 2, 2 \leq i_2 \leq a_2 - 1; \\ & x_2^{i_2} \partial_1, a_2 - 3 \leq i_2 \leq a_2; x_2^{i_2} \partial_2, 2 \leq i_2 \leq a_2; x_1^{i_1} x_2 \partial_2, a_1 - 3 \leq i_1 \leq a_1 - 1; x_1^{i_1} \partial_2, a_1 - 2 \leq i_1 \leq a_1 + 1. \end{aligned}$$

Therefore we have the following formula

$$\lambda^2(V) = 2a_1 a_2 - 2a_1 - 3a_2 + 20.$$

In case of $a_1 = 4, a_2 \geq 5$, we have following derivations which form a basis of $L^2(V)$:

$$\begin{aligned} & x_2^{i_2} \partial_1, a_2 - 3 \leq i_2 \leq a_2; x_1 \partial_1 + \frac{4}{a_2 - 1} x_2 \partial_1; x_1^{i_1} x_2^{i_2} \partial_1, 1 \leq i_1 \leq 2, 1 \leq i_2 \leq a_2 - 1; x_1^{i_1} \partial_1, 2 \leq i_1 \leq 5; \\ & x_1^3 x_2^{i_2} \partial_1, 1 \leq i_2 \leq 2; x_2^{i_2} \partial_2, 2 \leq i_2 \leq a_2; x_1^{i_1} x_2^{i_2} \partial_2, 1 \leq i_1 \leq 2, 1 \leq i_2 \leq a_2 - 1; x_1^{i_1} \partial_2, 2 \leq i_1 \leq 5; \\ & x_1^3 x_2^{i_2} \partial_2, 1 \leq i_2 \leq 2. \end{aligned}$$

Therefore we get following formula

$$\lambda^2(V) = 5a_2 + 12.$$

In case of $a_1 = 4, a_2 = 4$, we have following derivations which form a basis of $L^2(V)$:

$$\begin{aligned} & x_2^2 \partial_1; x_2^3 \partial_1; x_2^4 \partial_1; x_1 \partial_1 + \frac{4}{3} x_2 \partial_1; x_1 x_2 \partial_1; x_1 x_2^2 \partial_1; x_1 x_2^3 \partial_1; x_1^2 \partial_1; x_1^2 x_2 \partial_1; x_1^2 x_2^2 \partial_1; x_1^2 x_2^3 \partial_1; x_1^3 \partial_1; x_1^3 x_2 \partial_1; \\ & x_1^3 x_2^2 \partial_1; x_1^4 \partial_1; x_1^5 \partial_1; x_2^2 \partial_2; x_2^3 \partial_2; x_2^4 \partial_2; x_1 \partial_1 + \frac{4}{3} x_2 \partial_2; x_1 x_2 \partial_2; x_1 x_2^2 \partial_2; x_1 x_2^3 \partial_2; x_1^2 \partial_2; x_1^2 x_2 \partial_2; x_1^2 x_2^2 \partial_2; \\ & x_1^2 x_2^3 \partial_2; x_1^3 \partial_2; x_1^3 x_2 \partial_2; x_1^3 x_2^2 \partial_2; x_1^4 \partial_2; x_1^5 \partial_2 \end{aligned}$$

Therefore we get following formula

$$\lambda^2(V) = 31.$$

In case of $a_1 \geq 3, a_2 = 3$, we have following derivations which form a basis of $L^2(V)$:

$$\begin{aligned} & x_2^2 \partial_1; x_2^3 \partial_1; x_1^{i_1} \partial_1 + \frac{a_1}{2} x_1^{i_1-1} x_2 \partial_2, 1 \leq i_1 \leq a_1 - 2; x_1^{i_1} x_2^{i_2} \partial_1, 1 \leq i_1 \leq a_1 - 1, 1 \leq i_2 \leq 2; \\ & x_1^{i_1} \partial_1, a_1 - 1 \leq i_1 \leq a_1 + 1; x_2^2 \partial_2; x_2^3 \partial_2; x_1^{i_1} \partial_2, a_1 - 1 \leq i_1 \leq a_1 + 1; x_1^{i_1} x_2^2 \partial_2, 1 \leq i_1 \leq a_1 - 1; \\ & x_1^{i_1} x_2 \partial_2, a_1 - 2 \leq i_1 \leq a_1 - 1. \end{aligned}$$

Therefore we get following formula

$$\lambda^2(V) = 4a_1 + 7.$$

In case of $a_1 \geq 2, a_2 = 2$, we have following derivations which form a basis of $L^2(V)$:

$$x_2^{i_2} \partial_1, 1 \leq i_2 \leq 2; x_1^{i_1} \partial_1 + a_1 x_1^{i_1-1} x_2 \partial_2, 1 \leq i_1 \leq a_1 - 1; x_1^{i_1} x_2 \partial_1, 1 \leq i_1 \leq a_1 - 1;$$

$$x_1^{i_1} \partial_1, a_1 \leq i_1 \leq a_1 + 1; x_2^2 \partial_2; \left(\frac{x_1^{a_1+1}}{2} + x_1 x_2\right) \partial_2; x_1^{a_1-1} x_2 \partial_2.$$

Therefore we get following formula

$$\lambda^2(V) = 2a_1 + 5.$$

In case of $a_1 = 2, a_2 \geq 3$, it is easy to check that we get following basis:

$$x_2^{i_2} \partial_1, 2 \leq i_2 \leq a_2; x_1 \partial_1 + \frac{2}{a_2 - 1} x_2 \partial_2; x_2^{i_2} \partial_1, a_2 - 2 \leq i_2 \leq a_2;$$

$$x_1 x_2 \partial_1; x_1 x_2^2 \partial_1; x_1^2 \partial_1; x_1^3 \partial_1; x_1 x_2 \partial_2; x_1 x_2^2 \partial_2; x_1^2 \partial_2; x_1^3 \partial_2.$$

Therefore

$$\lambda^2(V) = a_2 + 11.$$

In case of $a_1 = 2, a_2 = 3$, we have following derivations which form a basis of $L^2(V)$:

$$x_2^2 \partial_1; x_2^3 \partial_1; x_1 \partial_1 + x_2 \partial_2; x_1 x_2 \partial_1; x_1 x_2^2 \partial_1; x_1^2 \partial_1; x_1^3 \partial_1; x_2^2 \partial_2; x_2^3 \partial_2; x_1 x_2 \partial_2; x_1 x_2^2 \partial_2; x_1^2 \partial_2; x_1^3 \partial_2.$$

Therefore we get following formula

$$\lambda^2(V) = 13.$$

In case of $a_1 = 1, a_2 \geq 2$, we have following derivations which form a basis of $L^2(V)$:

$$x_2^2 \partial_1; x_1 \partial_1; x_1^2 \partial_1; x_2 \partial_2; x_2^2 \partial_2; x_1^2 \partial_2.$$

Therefore we get following formula

$$\lambda^2(V) = 6.$$

We have

$$h_2(a_1, a_2) = \begin{cases} 2a_1 a_2 - 3(a_1 + a_2) + 17; & a_1 \geq 5, a_2 \geq 5 \\ 3a_2 + 5; & a_1 = 3, a_2 \geq 4 \\ 13; & a_1 = 3, a_2 = 3 \\ 5a_2 + 4; & a_1 = 4, a_2 \geq 5 \\ 23; & a_1 = 4, a_2 = 4 \\ a_2 + 5; & a_1 = 2, a_2 \geq 3 \\ 6; & a_1 = 2, a_2 = 2. \end{cases}$$

After putting the weight type $(\frac{a_2-1}{a_1a_2}, \frac{1}{a_2}; 1)$ of binomial isolated singularity of type B we have

$$h_2\left(\frac{1}{w_1}, \frac{1}{w_2}\right) = \begin{cases} \frac{2a_1a_2^2}{a_2-1} - 3\left(\frac{a_1a_2}{a_2-1} + a_2\right) + 17; & \frac{a_1a_2}{a_2-1} \geq 5, a_2 \geq 5 \\ 13; & a_1 = 2, a_2 = 3 \\ 23; & a_1 = 3, a_2 = 4 \\ 6; & a_1 = 1, a_2 = 2. \end{cases}$$

It is easy to see that when $a_1 \geq 5, a_2 \geq 5$, then

$$2a_1a_2 - 2a_1 - 3a_2 + 20 \leq \frac{2a_1a_2^2}{a_2-1} - 3\left(\frac{a_1a_2}{a_2-1} + a_2\right) + 17.$$

After solving above inequality we get $(a_2-2)(a_1-1) - 3 \geq 0$. It is noted that $(a_2-2) \geq 3$ and $(a_1-1) \geq 4$. \square

Proposition 3.3. *Let $(V, 0)$ be a binomial isolated singularity of type C which is defined by $f = x_1^{a_1}x_2 + x_2^{a_2}x_1$ ($a_1 \geq 1, a_2 \geq 1$) with weight type $(\frac{a_2-1}{a_1a_2-1}, \frac{a_1-1}{a_1a_2-1}; 1)$. Then*

$$\lambda^2(V) = \begin{cases} 2a_1a_2 - 2(a_1 + a_2) + 21; & a_1 \geq 4, a_2 \geq 4 \\ 4a_2 + 13; & a_1 = 3, a_2 \geq 4 \\ 2a_2 + 10; & a_1 = 2, a_2 \geq 3 \\ 23; & a_1 = 3, a_2 = 3 \\ 13; & a_1 = 2, a_2 = 2 \\ 6; & a_1 = 1, a_2 \geq 1. \end{cases}$$

Furthermore, we need to show that when $a_1 \geq 5, a_2 \geq 5$, then

$$2a_1a_2 - 2(a_1 + a_2) + 21 \leq \frac{2(a_1a_2-1)^2}{(a_1-1)(a_2-1)} - 3\left(\frac{a_1a_2-1}{a_2-1} + \frac{a_1a_2-1}{a_1-1}\right) + 17.$$

Proof. It follows that the generalized moduli algebra

$$A^2(V) = \mathbb{C}\{x_1, x_2\}/(f, m^2J(f)),$$

has dimension $a_1a_2 + 5$ and has a monomial basis of the form:

(1) if $a_1 \geq 2, a_2 \geq 2$, $\{x_1^{i_1}x_2^{i_2}, 0 \leq i_1 \leq a_1 - 2; 0 \leq i_2 \leq a_2 - 1; x_2^{i_2}, a_2 \leq i_2 \leq 2a_2 - 2;$

$$x_1^{i_1}, a_1 - 1 \leq i_1 \leq a_1 + 1; x_1x_2^{a_2}; x_1^{a_1-1}x_2; x_1^{a_1-1}x_2^2\}, \quad (28)$$

(2) if $a_1 = 1, a_2 \geq 1$, $\{x_2^{i_2}, 0 \leq i_2 \leq a_2 + 1; x_1; x_1^2\}$, (29)

with the following relations:

$$a_1x_1^{a_1+1} + x_1^2x_2^{a_2} = 0, \quad (30)$$

$$a_1x_1^{a_1-1}x_2^3 + x_2^{a_2+2} = 0, \quad (31)$$

$$a_1x_1^{a_1}x_2^2 + x_1x_2^{a_2+1} = 0, \quad (32)$$

$$x_1^{a_1+2} + a_2x_1^3x_2^{a_2-1} = 0, \quad (33)$$

$$x_1^{a_1}x_2^2 + a_2x_1x_2^{a_2+1} = 0, \quad (34)$$

$$x_1^{a_1+1}x_2 + a_2x_1^2x_2^{a_2} = 0. \quad (35)$$

In order to compute a derivation D of $A^2(V)$ it suffices to indicate its values on the generators x_1, x_2 which can be written in terms of the basis (28) or (29). Without loss of generality, we write

$$\begin{aligned} Dx_j = & \sum_{i_1=0}^{a_1-2} \sum_{i_2=0}^{a_2-1} c_{i_1, i_2}^j x_1^{i_1} x_2^{i_2} + \sum_{i_2=a_2}^{2a_2-2} c_{0, i_2}^j x_2^{i_2} + \sum_{i_1=a_1-1}^{a_1+1} c_{i_1, 0}^j x_1^{i_1} + c_{1, a_2}^j x_1 x_2^{a_2} \\ & + c_{a_1-1, 1}^j x_1^{a_1-1} x_2 + c_{a_1-1, 2}^j x_1^{a_1-1} x_2^2, \quad j = 1, 2. \end{aligned}$$

Using the relations (30)-(35) one easily finds the necessary and sufficient conditions defining a derivation of $A^2(V)$ as follows:

$$c_{0,0}^1 = c_{0,1}^1 = \cdots = c_{0, a_2-3}^1 = 0; \quad (36)$$

$$c_{0,0}^2 = c_{1,0}^2 = \cdots = c_{a_1-3,0}^2 = 0; \quad (37)$$

$$(a_1 - 1)c_{1,0}^1 = (a_2 - 1)c_{0,1}^2, (a_1 - 1)c_{1,1}^1 = (a_2 - 1)c_{0,2}^2, \cdots, (a_1 - 1)c_{1, a_2-4}^1 = (a_2 - 1)c_{0, a_2-3}^2; \quad (38)$$

$$(a_1 - 1)c_{a_1-3,0}^1 = (a_2 - 1)c_{a_1-4,1}^2. \quad (39)$$

Using (36)-(39) we obtain the following description of the Lie algebras in question. The following derivations form a basis of $\text{Der} A^2(V)$:

$$\begin{aligned} & x_1^{i_1} x_2^{i_2} \partial_1, 2 \leq i_1 \leq a_1 - 2, 1 \leq i_2 \leq a_2 - 1; x_2^{i_2} \partial_1, a_2 - 2 \leq i_2 \leq 2a_2 - 2; \\ & x_1^{i_1} \partial_1 + \frac{a_2 - 1}{a_1} x_1^{i_1-1} x_2 \partial_2, 1 \leq i_1 \leq a_1 - 3; x_1^{a_1-1} x_2 \partial_1; x_1^{a_1-1} x_2^2 \partial_1; \\ & x_1 x_2^{i_2} \partial_1 + \frac{a_2 - 1}{a_1} x_2^{i_2+1} \partial_2, 1 \leq i_2 \leq a_2 - 4; x_1 x_2^{i_2} \partial_1, a_2 - 3 \leq i_2 \leq a_2; x_1^{i_1} \partial_1, a_1 - 2 \leq i_1 \leq a_1 + 1; \\ & x_2^{i_2} \partial_2, a_2 - 2 \leq i_2 \leq 2a_2 - 2; x_1^{i_1} x_2^{i_2} \partial_2, 1 \leq i_1 \leq a_1 - 2, 2 \leq i_2 \leq a_2 - 1; \\ & x_1^{i_1} \partial_2, a_1 - 2 \leq i_1 \leq a_1 + 1; x_1^{i_1} x_2 \partial_2, a_1 - 3 \leq i_1 \leq a_1 - 1; x_1 x_2^{a_2} \partial_2; x_1^{a_1-1} x_2^2 \partial_2. \end{aligned}$$

Therefore we have the following formula

$$\lambda^2(V) = 2a_1 a_2 - 2(a_1 + a_2) + 21.$$

In case of $a_1 = 3, a_2 \geq 4$, we have following derivations which form a basis of $L^2(V)$:

$$\begin{aligned} & x_2^{i_2} \partial_1, a_2 - 2 \leq i_2 \leq 2a_2 - 2; x_1 \partial_1 + \frac{2}{a_2 - 1} x_2 \partial_2; x_1 x_2^{i_2} \partial_1 + \frac{2}{a_2 - 1} x_2^{i_2+1} \partial_2, 1 \leq i_2 \leq a_2 - 4; \\ & x_1 x_2^{i_2} \partial_1, a_2 - 3 \leq i_2 \leq a_2; x_1^2 x_2 \partial_1; x_1^2 x_2^2 \partial_1; x_2^{i_2} \partial_2, a_2 - 2 \leq i_2 \leq 2a_2 - 2; x_1^{i_1} \partial_2, 2 \leq i_1 \leq 4; \\ & x_1^{i_1} \partial_1, 2 \leq i_1 \leq 4; x_1 x_2^{i_2} \partial_2, 1 \leq i_2 \leq a_2; x_1^2 x_2 \partial_2; x_1^2 x_2^2 \partial_2. \end{aligned}$$

Therefore we get following formula

$$\lambda^2(V) = 4a_2 + 13.$$

In case of $a_1 = 2, a_2 \geq 3$, we have following derivations which form a basis of $L^2(V)$:

$$\begin{aligned} & x_2^{(a_2-2)-i_2} \partial_1 - \frac{2}{a_2-1} x_2^{i_2} \partial_2, 2 \leq i_2 \leq a_2 - 2; x_2^{i_2} \partial_1, 2a_2 - 3 \leq i_2 \leq 2a_2 - 1; \\ & x_1^2 \partial_1; x_1^3 \partial_1; x_2^{i_2} \partial_2, a_2 - 1 \leq i_2 \leq 2a_2 - 1; x_1^2 \partial_2; x_1^3 \partial_2; x_1 x_2 \partial_2; x_1 x_2^2 \partial_2; \\ & x_1 x_2^{i_2} \partial_1 + \frac{x_2^{i_2+1}}{a_2-1} \partial_2, 0 \leq i_2 \leq 2. \end{aligned}$$

Therefore we get following formula

$$\lambda^2(V) = 2a_2 + 10.$$

In case of $a_1 = 3, a_2 = 3$, we have following derivations which form a basis of $L^2(V)$:

$$\begin{aligned} & x_2^2 \partial_1; x_2^3 \partial_1; x_2^4 \partial_1; x_1 \partial_1 + x_2 \partial_2; x_1 x_2 \partial_1; x_1 x_2^2 \partial_1; x_1 x_2^3 \partial_1; x_1^2 \partial_1; x_1^2 x_2 \partial_1; x_1^2 x_2^2 \partial_1; x_1^3 \partial_1; x_1^4 \partial_1; x_2^2 \partial_2; x_2^3 \partial_2; \\ & x_2^4 \partial_2; x_1 x_2 \partial_2; x_1 x_2^2 \partial_2; x_1 x_2^3 \partial_2; x_1^2 \partial_2; x_1^2 x_2 \partial_2; x_1^2 x_2^2 \partial_2; x_1^3 \partial_2; x_1^4 \partial_2. \end{aligned}$$

Therefore we get following formula

$$\lambda^2(V) = 23.$$

In case of $a_1 = 2, a_2 = 2$, we have following derivations which form a basis of $L^2(V)$:

$$x_2^2 \partial_1; x_2^3 \partial_1; x_1 \partial_1 + x_2 \partial_2; x_1 x_2 \partial_1; x_1 x_2^2 \partial_1; x_1^2 \partial_1; x_1^3 \partial_1; x_2^2 \partial_2; x_2^3 \partial_2; x_1 x_2 \partial_2; x_1 x_2^2 \partial_2; x_1^2 \partial_2; x_1^3 \partial_2.$$

Therefore we get following formula

$$\lambda^2(V) = 13.$$

In case of $a_1 = 1, a_2 \geq 1$, we have following derivations which form a basis of $L^2(V)$:

$$(x_2^{a_2+1} + x_2^2) \partial_1; x_1 \partial_1; x_1^2 \partial_1; (-x_2^{a_2+1} + x_2) \partial_2; (x_2^{a_2+1} + x_2^2) \partial_2; x_1^2 \partial_2.$$

Therefore we get following formula

$$\lambda^2(V) = 6.$$

We have

$$h_2\left(\frac{1}{w_1}, \frac{1}{w_2}\right) = \begin{cases} \frac{2(a_1 a_2 - 1)^2}{(a_1 - 1)(a_2 - 1)} - 3\left(\frac{a_1 a_2 - 1}{a_2 - 1} + \frac{a_1 a_2 - 1}{a_1 - 1}\right) + 17; & \frac{a_1 a_2 - 1}{a_2 - 1} \geq 5, \frac{a_1 a_2 - 1}{a_1 - 1} \geq 5 \\ 13; & a_1 = 2, a_2 = 2 \\ 23; & a_1 = 3, a_2 = 3. \end{cases}$$

It is easy to see that when $a_1 \geq 5, a_2 \geq 5$, then

$$2a_1 a_2 - 2(a_1 + a_2) + 21 \leq \frac{2(a_1 a_2 - 1)^2}{(a_1 - 1)(a_2 - 1)} - 3\left(\frac{a_1 a_2 - 1}{a_2 - 1} + \frac{a_1 a_2 - 1}{a_1 - 1}\right) + 17.$$

After solving above inequality it is easy to see that

$$a_1(a_2 - 1)(a_1 - 3) + a_2(a_1 - 1)(a_2 - 3) + 2(a_1 - 3) + 2a_1 + 4a_2 - 1 \geq 0.$$

□

Proposition 3.4. *Let $(V, 0)$ be a fewnomial surface isolated singularity of type 1 which is defined by $f = x_1^{a_1} + x_2^{a_2} + x_3^{a_3}$ ($a_1 \geq 3, a_2 \geq 3, a_3 \geq 3$) with weight type $(\frac{1}{a_1}, \frac{1}{a_2}, \frac{1}{a_3}; 1)$. Then*

$$\lambda^2(V) = 3a_1a_2a_3 + 5(a_1 + a_2 + a_3) - 4(a_1a_2 + a_1a_3 + a_2a_3) + 34.$$

Proof. It is easy to see that the moduli algebra $A^2(V) = \mathbb{C}\{x_1, x_2, x_3\}/(f, m^2J(f))$ has dimension $(a_1a_2a_3 - (a_1a_2 + a_1a_3 + a_2a_3) + (a_1 + a_2 + a_3) + 10)$ and has a monomial basis of the form:

$$\{x_1^{i_1}x_2^{i_2}x_3^{i_3}, 0 \leq i_1 \leq a_1 - 2; 0 \leq i_2 \leq a_2 - 2; 0 \leq i_3 \leq a_3 - 2; x_2^{i_2}, a_2 - 1 \leq i_2 \leq a_2; x_3^{i_3}, a_3 - 1 \leq i_3 \leq a_3; x_1^{a_1-1}; x_1^{a_1-1}x_2; x_1^{a_1-1}x_3; x_2^{a_2-1}x_3; x_2x_3^{a_3-1}; x_1x_3^{a_3-1}; x_1x_2^{a_2-1}\}.$$

In order to compute a derivation D of $A^2(V)$ it suffices to indicate its values on the generators x_1, x_2, x_3 which can be written in terms of the basis. Thus we can write

$$\begin{aligned} Dx_j = & \sum_{i_1=0}^{a_1-2} \sum_{i_2=0}^{a_2-2} \sum_{i_3=0}^{a_3-2} c_{i_1, i_2, i_3}^j x_1^{i_1} x_2^{i_2} x_3^{i_3} + \sum_{i_2=a_2-1}^{a_2} c_{0, i_2, 0}^j x_2^{i_2} + \sum_{i_3=a_3-1}^{a_3} c_{0, 0, i_3}^j x_3^{i_3} + c_{a_1-1, 0, 0}^j x_1^{a_1-1} \\ & + c_{a_1-1, 1, 0}^j x_1^{a_1-1} x_2 + c_{0, a_2-1, 1}^j x_2^{a_2-1} x_3 + c_{0, 1, a_3-1}^j x_2 x_3^{a_3-1} + c_{1, 0, a_3-1}^j x_1 x_3^{a_3-1} \\ & + c_{1, a_2-1, 0}^j x_1 x_2^{a_2-1}, \quad j = 1, 2, 3. \end{aligned}$$

Using the above derivations we obtain the following description of Lie algebras in question. The derivations represented by the following vector fields form a basis in $\text{Der}A^2(V)$:

$$\begin{aligned} & x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_1, \quad 1 \leq i_1 \leq a_1 - 2, 0 \leq i_2 \leq a_2 - 2, 0 \leq i_3 \leq a_3 - 2; \\ & x_1 \partial_1 + \frac{a_1}{a_2} x_2 \partial_2 + \frac{a_1}{a_3} x_3 \partial_3; x_1^{a_1-1} x_3 \partial_1; x_1^{a_1-1} x_2 \partial_1; x_2^{i_2} \partial_1, \quad a_2 - 1 \leq i_2 \leq a_2; \\ & x_3^{i_3} \partial_1, \quad a_3 - 1 \leq i_3 \leq a_3; x_1 x_3^{a_3-1} \partial_1; x_1 x_2^{a_2-1} \partial_1; x_2^{a_2-1} x_3 \partial_1; x_2 x_3^{a_3-1} \partial_1; x_2^{a_2-3} x_3^{a_3-2} \partial_1; \\ & x_2^{a_2-2} x_3^{i_3} \partial_1, \quad a_3 - 3 \leq i_3 \leq a_3 - 2; x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_2, \quad 1 \leq i_1 \leq a_1 - 2, 1 \leq i_2 \leq a_2 - 2, 0 \leq i_3 \leq a_3 - 2; \\ & x_3^{i_3} \partial_2, \quad a_3 - 1 \leq i_3 \leq a_3; x_2^{i_2} \partial_2, \quad 2 \leq i_2 \leq a_2; x_2^{i_2} x_3^{i_3} \partial_2, \quad 1 \leq i_2 \leq a_2 - 2, 1 \leq i_3 \leq a_3 - 2; \\ & x_2 x_3^{a_3-1} \partial_2; x_2^{a_2-1} x_3 \partial_2; x_1 x_3^{i_3} \partial_2, \quad a_3 - 2 \leq i_3 \leq a_3 - 1; x_1^{a_1-1} \partial_2; x_1^{a_1-1} x_2 \partial_2; x_1^{a_1-1} x_3 \partial_2; \\ & x_1^{a_1-2} x_3^{i_3} \partial_2, \quad a_3 - 3 \leq i_3 \leq a_3 - 2; x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_3, \quad 1 \leq i_1 \leq a_1 - 2, 1 \leq i_2 \leq a_2 - 2, 1 \leq i_3 \leq a_3 - 2; \\ & x_1 x_2^{a_2-1} \partial_2; x_3^{i_3} \partial_3, \quad 2 \leq i_3 \leq a_3; x_1^{a_1-1} \partial_3; x_1^{a_1-1} x_2 \partial_3; x_1^{a_1-1} x_3 \partial_3; x_1 x_3^{a_3-1} \partial_3; \\ & x_1^{i_1} x_3^{i_3} \partial_3, \quad 1 \leq i_1 \leq a_1 - 2, 1 \leq i_3 \leq a_3 - 2; x_2^{i_2} x_3^{i_3} \partial_3, \quad 1 \leq i_2 \leq a_2 - 2, 1 \leq i_3 \leq a_3 - 2; x_2 x_3^{a_3-1} \partial_3; \\ & x_2^{a_2-1} x_3 \partial_3; x_2^{i_2} \partial_3, \quad a_2 - 1 \leq i_2 \leq a_2; x_1 x_2^{i_2} \partial_3, \quad a_2 - 2 \leq i_2 \leq a_2 - 1; x_1^{a_1-2} x_2^{i_2} \partial_3, \quad a_2 - 3 \leq i_2 \leq a_2 - 2. \end{aligned}$$

Therefore we have

$$\lambda^2(V) = 3a_1a_2a_3 + 5(a_1 + a_2 + a_3) - 4(a_1a_2 + a_1a_3 + a_2a_3) + 34.$$

□

Proposition 3.5. *Let $(V, 0)$ be a fewnomial surface isolated singularity of type 2 which is defined by $f = x_1^{a_1}x_2 + x_2^{a_2}x_3 + x_3^{a_3}$ ($a_1 \geq 2, a_2 \geq 2, a_3 \geq 3$) with weight type $(\frac{1-a_3+a_2a_3}{a_1a_2a_3}, \frac{a_3-1}{a_2a_3}, \frac{1}{a_3}; 1)$. Then*

$$\lambda^2(V) = \begin{cases} 3a_1a_2a_3 + 2a_1 + 2a_2 + 6a_3 - 4a_2a_3 - 2a_1a_2 - 2a_1a_3 + 37; & a_1 \geq 3, a_2 \geq 3, a_3 \geq 3 \\ 4a_2 + 42; & a_1 = 2, a_2 \geq 3, a_3 = 3 \\ 2a_2a_3 - 2a_2 + 2a_3 + 38; & a_1 = 2, a_2 \geq 3, a_3 \geq 4 \\ 4a_1a_3 - 3a_3 - 2a_1 + 42; & a_1 \geq 3, a_2 = 2, a_3 \geq 4 \\ 5a_3 + 35; & a_1 = 2, a_2 = 2, a_3 \geq 4 \\ 46; & a_1 = 2, a_2 = 2, a_3 = 3 \\ 10a_1 + 32; & a_1 \geq 3, a_2 = 2, a_3 = 3. \end{cases}$$

Furthermore, we need to show that when $a_1 \geq 3, a_2 \geq 3, a_3 \geq 3$, then

$$3a_1a_2a_3 - 2a_1a_2 - 2a_1a_3 - 4a_2a_3 + 2a_1 + 2a_2 + 6a_3 + 37 \leq 3\frac{a_1a_2^2a_3^3}{(1-a_3+a_2a_3)(a_3-1)} - 4\left(\frac{a_1a_2^2a_3^3}{(1-a_3+a_2a_3)(a_3-1)} + \frac{a_1a_2a_3^2}{1-a_3+a_2a_3} + \frac{a_2a_3^2}{a_3-1}\right) + 5\left(\frac{a_1a_2a_3}{1-a_3+a_2a_3} + \frac{a_2a_3}{a_3-1} + a_3\right) + 34.$$

Proof. It is easy to see that the moduli algebra $A^2(V) = \mathbb{C}\{x_1, x_2, x_3\}/(f, m^2J(f))$ has dimension $(a_1a_2a_3 - a_2a_3 + a_3 + 10)$ and has a monomial basis of the form:

$$\{x_1^{i_1}x_2^{i_2}x_3^{i_3}, 0 \leq i_1 \leq a_1 - 2; 0 \leq i_2 \leq a_2 - 1; 0 \leq i_3 \leq a_3 - 1; x_2^{i_2}, a_2 \leq i_2 \leq a_2 + 1; x_1^{i_1}, a_1 \leq i_1 \leq a_1 + 1; x_2^{a_2}x_3; x_1x_2^{a_2}; x_3^{a_3}; x_1^{a_1-1}x_3^{i_3}, 0 \leq i_3 \leq a_3 - 2; x_1^{a_1-1}x_2x_3; x_1^{a_1}x_3; x_1^{a_1-1}x_2^{i_2}; 1 \leq i_2 \leq 2\}.$$

In order to compute a derivation D of $A^2(V)$ it suffices to indicate its values on the generators x_1, x_2, x_3 which can be written in terms of the basis. Thus we can write

$$\begin{aligned} Dx_j &= \sum_{i_1=0}^{a_1-2} \sum_{i_2=0}^{a_2-1} \sum_{i_3=0}^{a_3-1} c_{i_1, i_2, i_3}^j x_1^{i_1} x_2^{i_2} x_3^{i_3} + \sum_{i_2=a_2}^{a_2+1} c_{0, i_2, 0}^j x_2^{i_2} + \sum_{i_1=a_1}^{a_1+1} c_{i_1, 0, 0}^j x_1^{i_1} + c_{0, a_2, 1}^j x_2^{a_2} x_3 + c_{1, a_2, 0}^j x_1 x_2^{a_2} \\ &+ c_{0, 0, a_3}^j x_3^{a_3} + \sum_{i_3=0}^{a_3-2} c_{a_1-1, 0, i_3}^j x_1^{a_1-1} x_3^{i_3} + c_{a_1-1, 1, 1}^j x_1^{a_1-1} x_2 x_3 + c_{a_1, 0, 1}^j x_1^{a_1} x_3 \\ &+ \sum_{i_2=1}^2 c_{a_1-1, i_2, 0}^j x_1^{a_1-1} x_2^{i_2}, \quad j = 1, 2, 3. \end{aligned}$$

Using the above derivations we obtain the following description of Lie algebras in question.

The derivations represented by the following vector fields form a basis in $\text{Der}A^2(V)$:

$$\begin{aligned} &x_3^{a_3} \partial_1; x_1^{i_1} \partial_1 + \frac{a_1(a_3-1)}{a_2(a_3-1)+1} x_1^{i_1-1} x_2 \partial_2 + \frac{a_1 a_2}{a_2(a_3-1)+1} x_1^{i_1-1} x_3 \partial_3, \quad 1 \leq i_1 \leq a_1 - 1; \\ &\left(\frac{1}{a_3} x_2^{i_2} + x_2 x_3^{a_3-1}\right) \partial_1, \quad a_2 \leq i_2 \leq a_2 + 1; x_1^{i_1} x_3^{i_3} \partial_1 + a_1 x_1^{i_1-1} x_3^{i_3+1} \partial_3, \quad 1 \leq i_1 \leq a_1 - 2; 1 \leq i_3 \leq a_3 - 1; \\ &x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_1, \quad 1 \leq i_1 \leq a_1 - 2; 1 \leq i_2 \leq a_2 - 1; 0 \leq i_3 \leq a_3 - 1; x_1^{a_1-1} x_2 x_3 \partial_1; \end{aligned}$$

$$\begin{aligned}
& x_2^{a_2-2} x_3 \partial_1 + \frac{a_1-1}{a_2} x_1^{a_1-1} \partial_2; x_1^{i_1} \partial_1, \quad a_1 \leq i_1 \leq a_1+1; x_1^{a_1-1} x_2^{i_2} \partial_1, \quad 1 \leq i_2 \leq 2; \\
& x_1^{a_1-1} x_3^{i_3} \partial_1, \quad 1 \leq i_3 \leq a_3-2; x_1^{a_1} x_3 \partial_1; x_1 x_2^{a_2} \partial_1; x_2^{i_2} x_3^{i_3} \partial_1, \quad a_2-2 \leq i_2 \leq a_2-1; 2 \leq i_3 \leq a_3-1; \\
& x_2^{a_2-3} x_3^{a_3-1} \partial_1; x_2^{a_3} x_3 \partial_1; x_2^{a_2-1} x_3 \partial_1; x_3^{a_3-1} \partial_2 + \frac{1}{a_2(a_2-1)} x_1^{a_1} \partial_3; x_3^{a_3} \partial_2; \\
& x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_2, \quad 1 \leq i_1 \leq a_1-2; 1 \leq i_2 \leq a_2-1; 1 \leq i_3 \leq a_3-1; x_1^{i_1} \partial_1, \quad a_1 \leq i_1 \leq a_1+1; \\
& x_1^{a_1-1} x_3^{i_3} \partial_2, \quad a_3-3 \leq i_3 \leq a_3-2; x_1^{a_1-1} x_2^{i_2} \partial_2, \quad 1 \leq i_2 \leq 2; x_1^{a_1-1} x_2 x_3 \partial_2; x_1^{a_1} x_3 \partial_2; \\
& x_2^{i_2} x_3^{i_3} \partial_2, \quad 1 \leq i_2 \leq a_2-1; 1 \leq i_3 \leq a_3-1; x_2^{i_2} \partial_2, \quad 2 \leq i_2 \leq a_2+1; x_2^{a_2} x_3 \partial_2; \\
& x_1^{i_1} x_2^{i_2} \partial_2, \quad 1 \leq i_1 \leq a_1-2; 2 \leq i_2 \leq a_2-1; x_1 x_2^{a_2} \partial_2; x_1^{a_1-2} x_3^{i_3} \partial_2, \quad a_3-2 \leq i_3 \leq a_3-1; \\
& x_1^{i_1} x_3^{a_3-1} \partial_2, \quad 1 \leq i_1 \leq a_1-3; \left(\frac{1}{a_3} x_2^{a_2} + x_3^{a_3-1}\right) \partial_3; x_3^{a_3} \partial_3; \left(\frac{1}{a_2} x_1^{a_1} + x_2^{a_2-1} x_3\right) \partial_3; \\
& x_2^{i_2} x_3^{i_3} \partial_3, \quad 1 \leq i_2 \leq a_2-1; 2 \leq i_3 \leq a_3-1; x_2^{a_2+1} \partial_3; x_2^{a_2} x_3 \partial_3; x_1 x_3^{a_3-1} \partial_3; \\
& x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_3, \quad 1 \leq i_1 \leq a_1-2; 1 \leq i_2 \leq a_2-1; 2 \leq i_3 \leq a_3-1; x_1^{a_1+1} \partial_3; x_1^{a_1-1} x_3^{i_3} \partial_3, \quad 1 \leq i_3 \leq a_3-2; \\
& x_1 x_2^{a_2} \partial_3; x_1^{a_1-2} x_3^{i_3} \partial_3, \quad a_3-2 \leq i_3 \leq a_3-1; x_1^{a_1-1} x_2 \partial_3; x_1^{a_1-1} x_2 x_3 \partial_3; x_1^{a_1} x_3 \partial_3; \\
& x_1^{a_1-1} x_2^2 \partial_3; x_1^{a_1-2} x_2^{a_2-1} \partial_3; x_1^{a_1-2} x_2^{a_2-2} x_3 \partial_3; x_1^{i_1} x_2^{a_2-1} x_3 \partial_3, \quad 1 \leq i_1 \leq 2.
\end{aligned}$$

Therefore we have

$$\lambda^2(V) = 3a_1 a_2 a_3 + 2a_1 + 2a_2 + 6a_3 - 4a_2 a_3 - 2a_1 a_2 - 2a_1 a_3 + 37.$$

In case of $a_1 = 2, a_2 \geq 3, a_3 = 3$, we obtain the basis of derivation represented by following derivations form a basis of $\text{Der} A^2(V)$:

$$\begin{aligned}
& \left(\frac{1}{3} x_2^{i_2} + x_2^{i_2-a_2} x_3^2\right) \partial_1, \quad a_2 \leq i_2 \leq a_2+1; x_3^3 \partial_1; x_2^{i_2} x_3^2 \partial_1; \quad a_2-3 \leq i_2 \leq a_2-1; x_2^{i_2} x_3 \partial_1, \quad a_2-2 \leq i_2 \leq a_2; \\
& \left(\frac{1}{3} x_2^{i_2} + x_2^{i_2-a_2} x_3^2\right) \partial_1, \quad a_2 \leq i_2 \leq a_2+1; x_3^3 \partial_1; x_2^{i_2} x_3^2 \partial_1; \quad a_2-3 \leq i_2 \leq a_2-1; x_2^{i_2} x_3 \partial_1, \quad a_2-2 \leq i_2 \leq a_2; \\
& x_1 \partial_1 + \frac{4}{3(a_2-1)+1} x_2 \partial_2 + \frac{2a_2}{3(a_2-1)+1} x_3 \partial_3; x_1 x_3 \partial_1; x_1 x_3^2 \partial_1; x_1^2 \partial_1; x_1^2 x_3 \partial_1; x_1^3 \partial_1; x_1 x_2 x_3 \partial_1; \\
& x_1 x_2 \partial_1; x_1 x_2^2 \partial_1; x_3^3 \partial_2; x_3^3 \partial_2; x_2^{i_2} x_3^{i_3} \partial_2, \quad 1 \leq i_2 \leq a_2-1; 1 \leq i_3 \leq 2; x_2^{i_2} \partial_2, \quad 2 \leq i_2 \leq a_2+1; \\
& x_2^{a_2} x_3 \partial_2; x_1 x_3 \partial_2; x_1 x_3^2 \partial_2; x_1 x_2 x_3 \partial_2; x_1 x_2 \partial_2; x_1 x_2^2 \partial_2; x_1^2 \partial_2; x_1^2 x_3 \partial_2; x_1^3 \partial_2; x_3^2 \partial_3; x_3^3 \partial_3; \\
& x_2^{i_2} x_3^2 \partial_3, \quad 1 \leq i_2 \leq a_2-3; x_2^{i_2} x_3^{i_3} \partial_3, \quad a_2-2 \leq i_2 \leq a_2-1; 1 \leq i_3 \leq 2; x_2^{i_2} \partial_3, \quad a_2-1 \leq i_2 \leq a_2+1; \\
& x_1^{a_1} x_3 \partial_3; x_1 x_3 \partial_3; x_1 x_3^2 \partial_3; x_1 x_2 \partial_3; x_1 x_2 x_3 \partial_3; x_1 x_2^2 \partial_3; x_1^2 \partial_3; x_1^2 x_3 \partial_3; x_1^3 \partial_3.
\end{aligned}$$

Therefore we have

$$\lambda^2(V) = 4a_2 + 42.$$

In case of $a_1 = 2, a_2 \geq 3, a_3 \geq 4$, we obtain the basis of derivation represented by following derivations form a basis of $\text{Der} A^2(V)$:

$$\left(\frac{1}{a_3} x_2^{a_2} + x_3^{a_3-1}\right) \partial_1; x_3^{a_3} \partial_1; x_2^{a_2-3} x_3^{a_3-1} \partial_1; x_2^{i_2} x_3^{i_3} \partial_1, \quad a_2-2 \leq i_2 \leq a_2-1; 2 \leq i_3 \leq a_3-1;$$

$$\begin{aligned}
& x_2^{i_2} x_3 \partial_1, \quad a_2 - 1 \leq i_1 \leq a_2; x_2^{a_2+1} \partial_1; x_1 \partial_1 + \frac{2(a_3 - 1)}{(a_2 - 1)(a_3 - 1) + 1} x_2 \partial_2 + \frac{2(a_3 - 1)}{(a_2 - 1)(a_3 - 1)} x_3 \partial_3; \\
& x_1 x_3^{i_3} \partial_1, \quad 1 \leq i_3 \leq a_3 - 1; x_1 x_2 \partial_1; x_1 x_2 x_3 \partial_1; x_1 x_2^2 \partial_1; x_1^2 \partial_1; x_1^2 x_3 \partial_1; x_1^3 \partial_1; x_3^{i_3} \partial_2, \quad a_3 - 2 \leq i_3 \leq a_3; \\
& x_2^{i_2} x_3^{i_3} \partial_2, \quad 1 \leq i_2 \leq a_2 - 1; 1 \leq i_3 \leq a_3 - 1; x_2^{i_2} \partial_2, \quad 2 \leq i_2 \leq a_2 + 1; x_2^{a_2} x_3 \partial_2; x_1 x_3^{i_3} \partial_2, \quad 2 \leq i_3 \leq a_3 - 1; \\
& \quad x_1 x_2 x_3 \partial_2; x_1^2 \partial_2; x_1^2 x_3 \partial_2; x_1^3 \partial_2; x_1 x_2 \partial_2; x_1 x_2^2 \partial_2; x_3^{i_3} \partial_3, \quad a_3 - 2 \leq i_3 \leq a_3; \\
& x_2^{i_2} x_3^{i_3} \partial_3, \quad 1 \leq i_2 \leq a_2 - 1; 2 \leq i_3 \leq a_3 - 1; x_2^{i_2} x_3 \partial_3, \quad a_2 - 2 \leq i_2 \leq a_2; x_2^{i_2} \partial_3, \quad a_2 - 1 \leq i_2 \leq a_2 + 1; \\
& \quad x_1 x_3^{i_3} \partial_3, \quad 1 \leq i_3 \leq a_3 - 1; x_1 x_2 \partial_3; x_1 x_2^2 \partial_3; x_1 x_2 x_3 \partial_3; x_1^2 x_3 \partial_3; x_1^2 \partial_3; x_1^3 \partial_3.
\end{aligned}$$

Therefore we have

$$\lambda^2(V) = 2a_2 a_3 - 2a_2 + 2a_3 + 38.$$

In case of $a_1 \geq 3, a_2 = 2, a_3 \geq 4$, we obtain the basis of derivation represented by following derivations form a basis of $\text{Der}A^2(V)$:

$$\begin{aligned}
& x_3^{a_3-2} \partial_1 - \frac{a_1}{2} x_1^{a_1-1} x_3^{a_3-3} \partial_2; x_3^{i_3} \partial_1, \quad a_3 - 1 \leq i_3 \leq a_3; x_2 x_3^{i_3} \partial_1, \quad 1 \leq i_3 \leq a_3 - 1; x_2^2 \partial_1; \\
& \quad x_2^2 x_3 \partial_1; x_2^3 \partial_1; x_1^{i_1} \partial_1 + \frac{a_1(a_3 - 1)}{a_3 + 1} x_1^{i_1-1} x_2 \partial_2 + \frac{a_1 a_2}{a_3 + 1} x_1^{i_1-1} x_3 \partial_3, \quad 1 \leq i_1 \leq a_1 - 2; \\
& x_1^{i_1} x_3^{i_3} \partial_1, \quad 1 \leq i_1 \leq a_1 - 1; 1 \leq i_3 \leq a_3 - 1; x_1^{i_1} x_2 x_3^{i_3} \partial_1, \quad 1 \leq i_1 \leq a_1 - 2; 0 \leq i_3 \leq a_3 - 1; \\
& \quad x_1 x_2^2 \partial_1; x_1^{i_1} \partial_1, \quad a_1 - 1 \leq i_1 \leq a_1 + 1; x_1^{a_1-1} x_2 \partial_1; x_1^{a_1-1} x_2 x_3 \partial_1; x_1^{a_1} x_3 \partial_1; \\
& \quad x_3^{a_3-1} \partial_2 + \frac{1}{a_3(a_3 - 1)} x_1^{a_1} \partial_3; x_3^{a_3} \partial_2; x_2 x_3^{i_3} \partial_2, \quad 1 \leq i_3 \leq a_3 - 1; x_2^2 \partial_2; x_2^3 \partial_2; x_2^2 x_3 \partial_2; \\
& x_1^{i_1} x_3^{a_3-1} \partial_2, \quad 1 \leq i_1 \leq a_1 - 3; x_1^{i_1} x_2 x_3^{i_3} \partial_2, \quad 1 \leq i_1 \leq a_1 - 2; 1 \leq i_3 \leq a_3 - 1; x_1^{i_1} \partial_2, \quad a_1 \leq i_1 \leq a_1 + 1; \\
& x_1^{i_1} x_3^{i_3} \partial_2, \quad a_1 - 2 \leq i_1 \leq a_1 - 1; a_3 - 2 \leq i_3 \leq a_3 - 1; x_1^{a_1-1} x_2 x_3 \partial_2; x_1^{a_1} x_3 \partial_2; \left(\frac{1}{a_3} x_2^2 + x_3^{a_3-1}\right) \partial_2; \\
& \quad x_1 x_2^2 \partial_2; x_1^{i_1} x_2 \partial_2, \quad a_1 - 2 \leq i_1 \leq a_1 - 1; x_3^{a_3} \partial_3; \left(\frac{1}{2} x_1^{a_1} + x_2 x_3\right) \partial_3; x_2 x_3^{i_3} \partial_3, \quad 2 \leq i_3 \leq a_3 - 1; \\
& x_2^2 x_3 \partial_3; x_2^3 \partial_3; x_1^{i_1} x_3^{a_3-1} \partial_3, \quad a_1 - 3 \leq i_1 \leq a_1 - 2; x_1^{i_1} x_2 x_3^{i_3} \partial_3, \quad 1 \leq i_1 \leq a_1 - 2; 2 \leq i_3 \leq a_3 - 1; \\
& x_1 x_2^2 \partial_3; x_1^{a_1-1} x_3^{i_3} \partial_3, \quad 1 \leq i_3 \leq a_3 - 1; x_1^{a_1+1} \partial_3; x_1^{a_1} x_3 \partial_3; x_1^{a_1-1} x_2 x_3 \partial_3; x_1^{i_1} x_2 \partial_3, \quad a_1 - 2 \leq i_1 \leq a_1 - 1; \\
& \quad x_1^{a_1-2} x_3^{i_3} \partial_3, \quad a_3 - 2 \leq i_3 \leq a_3 - 1; x_1^{i_1} x_2 x_3 \partial_3, \quad a_1 - 3 \leq i_1 \leq a_1 - 2.
\end{aligned}$$

Therefore we have

$$\lambda^2(V) = 4a_1 a_3 - 3a_3 - 2a_1 + 42.$$

In case of $a_1 = 2, a_2 = 2, a_3 \geq 4$, we obtain the basis of derivation represented by following derivations form a basis of $\text{Der}A^2(V)$:

$$\begin{aligned}
& x_3^{i_3} \partial_1, \quad a_3 - 2 \leq i_3 \leq a_3; x_2 x_3^{i_3} \partial_1, \quad 1 \leq i_3 \leq a_3 - 1; x_2^2 \partial_1; x_1 \partial_1 + \frac{2(a_3 - 1)}{a_3 + 1} x_2 \partial_2 + \frac{4}{a_3 + 1} x_3 \partial_3; \\
& \quad x_2^2 x_3 \partial_1; x_2^3 \partial_1; x_1 x_3^{i_3} \partial_1, \quad 1 \leq i_3 \leq a_3 - 1; x_1 x_2 \partial_1; x_1 x_2 x_3 \partial_1; x_1 x_2^2 \partial_1; x_1^2 \partial_1; x_1^2 x_3 \partial_1; x_1^3 \partial_1; \\
& x_3^{i_3} \partial_2, \quad a_3 - 2 \leq i_3 \leq a_3; x_2 x_3^{i_3} \partial_2, \quad 1 \leq i_3 \leq a_3 - 1; x_2^2 \partial_2; x_2^2 x_3 \partial_2; x_2^3 \partial_2; x_1 x_3^{i_3} \partial_2, \quad a_3 - 3 \leq i_3 \leq a_3 - 1;
\end{aligned}$$

$x_1x_2\partial_2; x_1x_2^2\partial_2; x_1x_2x_3\partial_2; x_1^2\partial_2; x_1^2x_3\partial_2; x_1^3\partial_2; x_3^{i_3}\partial_3, a_3-2 \leq i_3 \leq a_3; x_2x_3^{i_3}\partial_3, 1 \leq i_3 \leq a_3-1;$
 $x_2^2\partial_3; x_2^2x_3\partial_3; x_1x_3^{i_3}\partial_3, 1 \leq i_3 \leq a_3-1; x_1x_2\partial_3; x_1x_2x_3\partial_3; x_1x_2^2\partial_3; x_1^2\partial_3; x_1^2x_3\partial_3; x_1^3\partial_3.$

Therefore we have

$$\lambda^2(V) = 5a_3 + 35.$$

In case of $a_1 = 2, a_2 = 2, a_3 = 3$, we obtain the basis of derivation represented by following derivations form a basis of $\text{Der}A^2(V)$:

$x_3^2\partial_1; x_3^3\partial_1; x_2x_3\partial_1; x_2x_3^2\partial_1; x_2^2\partial_1; x_2^2x_3\partial_1; x_2^3\partial_1; x_1\partial_1 + x_2\partial_2 + x_3\partial_3; x_1x_3\partial_1; x_1x_3^2\partial_1; x_1x_2\partial_1;$
 $x_1x_2x_3\partial_1; x_1x_2^2\partial_1; x_1^2\partial_1; x_1^2x_3\partial_1; x_1^3\partial_1; x_3^2\partial_2; x_3^3\partial_2; x_2x_3\partial_2; x_2x_3^2\partial_2; x_2^2\partial_2; x_2^2x_3\partial_2; x_2^3\partial_2; x_1x_3\partial_2;$
 $x_1x_3^2\partial_2; x_1x_2\partial_2; x_1x_2x_3\partial_2; x_1x_2^2\partial_2; x_1^2\partial_2; x_1^2x_3\partial_2; x_1^3\partial_2; x_3^2\partial_3; x_3^3\partial_3; x_2x_3\partial_3; x_2x_3^2\partial_3; x_2^2\partial_3;$
 $x_2^2x_3\partial_3; x_2^3\partial_3; x_1x_3\partial_3; x_1x_3^2\partial_3; x_1x_2\partial_3; x_1x_2x_3\partial_3; x_1x_2^2\partial_3; x_1^2\partial_3; x_1^2x_3\partial_3; x_1^3\partial_3.$

Therefore we have

$$\lambda^2(V) = 46.$$

In case of $a_1 \geq 3, a_2 = 2, a_3 = 3$, we obtain the basis of derivation represented by following derivations form a basis of $\text{Der}A^2(V)$:

$x_3^2\partial_1; x_3^3\partial_1; x_2x_3\partial_1; x_2x_3^2\partial_1; x_2^2\partial_1; x_2^2x_3\partial_1; x_2^3\partial_1; x_1^{i_1}\partial_1 + \frac{a_1}{2}x_1^{i_1-1}x_2\partial_2 + \frac{a_1}{2}x_1^{i_1-1}x_3\partial_3, 1 \leq i_1 \leq a_1-2;$
 $x_1^{i_1}x_2x_3^{i_3}\partial_1, 1 \leq i_1 \leq a_1-2; 0 \leq i_3 \leq 2; x_1^{i_1}x_3^{i_3}\partial_1, 1 \leq i_1 \leq a_1-1; 1 \leq i_3 \leq 2; x_1x_2^2\partial_1;$
 $x_1^{i_1}\partial_1, a_1-1 \leq i_1 \leq a_1+1; x_1^{a_1-1}x_2\partial_1; x_1^{a_1-1}x_2x_3\partial_1; x_1^{a_1}x_3\partial_1; x_3^2\partial_2 + \frac{1}{6}x_1^{a_1}\partial_3; x_3^3\partial_2;$
 $x_2x_3\partial_2 + \frac{1}{3}x_2^2\partial_3; x_2x_3^2\partial_3; x_2^2\partial_2 - \frac{1}{2}x_1^{a_1}\partial_3; x_2^2x_3\partial_2; x_2^3\partial_2; x_1x_2^2\partial_2;$
 $x_1^{i_1}x_2x_3^{i_3}\partial_2, 1 \leq i_1 \leq a_1-2; 1 \leq i_3 \leq 2; x_1^{a_1-3}x_3^2\partial_2; x_1^{i_1}x_3^{i_3}\partial_2, 1 \leq i_1 \leq a_1-1; 1 \leq i_3 \leq 2;$
 $x_1^{i_1}x_2\partial_2, a_1-2 \leq i_1 \leq a_1-1; x_1^{a_1-1}x_2x_3\partial_2; x_1^{a_1}\partial_2 - \frac{2}{3}x_1^2\partial_3; x_1^{a_1-1}x_3\partial_2; x_1^{a_1+1}\partial_2 - \frac{2}{3}x_1x_2^2\partial_3;$
 $(\frac{1}{3}x_2^2+x_3^2)\partial_3; x_3^3\partial_3; (\frac{1}{2}x_1^{a_1}+x_2x_3)\partial_3; x_2x_3^2\partial_3; x_2^2x_3\partial_3; x_2^3\partial_3; (\frac{1}{3}x_1x_2^2+x_1x_3^2)\partial_3; (\frac{1}{2}x_1^{a_1+1}+x_1x_2x_3)\partial_3;$
 $x_1^{i_1}x_2x_3^{i_3}\partial_3, a_1-3 \leq i_1 \leq a_1-2; 1 \leq i_3 \leq 2; x_1^{i_1}x_3^{i_3}\partial_3, a_1-2 \leq i_1 \leq a_1-1; 1 \leq i_3 \leq 2;$
 $x_1^{i_1}x_2\partial_3, a_1-2 \leq i_1 \leq a_1-1; x_1x_2x_3^2\partial_3; x_1^{a_1-3}x_3^2\partial_3; x_1^{a_1-1}x_2x_3\partial_3; x_1^{a_1}x_3\partial_3.$

Therefore we have

$$\lambda^2(V) = 10a_1 + 32.$$

It is follows from proposition 3.4 and weight types of proposition 3.5 we have

$$h_2\left(\frac{1}{w_1}, \frac{1}{w_2}, \frac{1}{w_3}\right) = \begin{cases} 3\frac{a_1a_2a_3}{1-a_3+a_2a_3}\frac{a_2a_3}{a_3-1}a_3 - 4\left(\frac{a_1a_2a_3}{1-a_3+a_2a_3}\frac{a_2a_3}{a_3-1} + \frac{a_1a_2a_3}{1-a_3+a_2a_3}a_3 + \frac{a_2a_3}{a_3-1}a_3\right) \\ + 5\left(\frac{a_1a_2a_3}{1-a_3+a_2a_3} + \frac{a_2a_3}{a_3-1} + a_3\right) + 34; \frac{a_1a_2a_3}{1-a_3+a_2a_3} \geq 3, \frac{a_2a_3}{a_3-1} \geq 3, a_3 \geq 3. \end{cases}$$

We need to prove following cases:

Case(i): When $a_1 \geq 3, a_2 \geq 3, a_3 \geq 3$, then

$$3a_1a_2a_3 - 2a_1a_2 - 2a_1a_3 - 4a_2a_3 + 2a_1 + 2a_2 + 6a_3 + 37 \leq 3\frac{a_1a_2^2a_3^3}{(1-a_3+a_2a_3)(a_3-1)} - 4\left(\frac{a_1a_2^2a_3^3}{(1-a_3+a_2a_3)(a_3-1)} + \frac{a_1a_2a_3^2}{1-a_3+a_2a_3} + \frac{a_2a_3^2}{a_3-1}\right) + 5\left(\frac{a_1a_2a_3}{1-a_3+a_2a_3} + \frac{a_2a_3}{a_3-1} + a_3\right) + 34.$$

After simplification we get

$$(a_3-1)^3(a_2-3) + (a_1-1)a_2a_3((a_3-1)(a_2-3) + (a_3-2)) + a_2(3a_3-2)(a_1-1) + a_2(a_1-1) - 4 \geq 0.$$

Case(ii): When $a_1 = 2, a_2 \geq 3, a_3 \geq 4$, then

$$2a_2a_3 - 2a_2 + 2a_3 + 38 \leq 6\frac{a_2^2a_3^3}{(1-a_3+a_2a_3)(a_3-1)} - 4\left(\frac{2a_2^2a_3^3}{(1-a_3+a_2a_3)(a_3-1)} + \frac{2a_2a_3^2}{1-a_3+a_2a_3} + \frac{a_2a_3^2}{a_3-1}\right) + 5\left(\frac{2a_2a_3}{1-a_3+a_2a_3} + \frac{a_2a_3}{a_3-1} + a_3\right) + 34.$$

After simplification we get

$$(a_3-1)^3(a_2-3) + a_2a_3((a_3-1)(a_2-3) + (a_3-2)) + a_2(3a_3-2) + a_2 - 7 \geq 0.$$

□

Proposition 3.6. *Let $(V, 0)$ be a fewnomial surface isolated singularity of type 3 which is defined by $f = x_1^{a_1}x_2 + x_2^{a_2}x_3 + x_3^{a_3}x_1$ ($a_1 \geq 2, a_2 \geq 2, a_3 \geq 2$) with weight type*

$$\left(\frac{1-a_3+a_2a_3}{1+a_1a_2a_3}, \frac{1-a_1+a_1a_3}{1+a_1a_2a_3}, \frac{1-a_2+a_1a_2}{1+a_1a_2a_3}; 1\right).$$

Then

$$\lambda^2(V) = \begin{cases} 3a_1a_2a_3 + 2(a_1 + a_2 + a_3) - 2(a_1a_2 + a_1a_3 + a_2a_3) + 43; & a_1 \geq 3, a_2 \geq 3, a_3 \geq 3 \\ 4a_2a_3 - 2(a_2 + a_3) + 46; & a_1 = 2, a_2 \geq 3, a_3 \geq 3. \end{cases}$$

Furthermore, we need to show that when $a_1 \geq 3, a_2 \geq 3, a_3 \geq 3$, then

$$3a_1a_2a_3 + 2(a_1 + a_2 + a_3) - 2(a_1a_2 + a_1a_3 + a_2a_3) + 43 \leq \frac{3(1+a_1a_2a_3)^3}{(1-a_3+a_2a_3)(1-a_1+a_1a_3)(1-a_2+a_1a_2)} + 5\left(\frac{1+a_1a_2a_3}{1-a_3+a_2a_3} + \frac{1+a_1a_2a_3}{1-a_1+a_1a_3} + \frac{1+a_1a_2a_3}{1-a_2+a_1a_2}\right) - 4\left(\frac{(1+a_1a_2a_3)^2}{(1-a_3+a_2a_3)(1-a_1+a_1a_3)} + \frac{(1+a_1a_2a_3)^2}{(1-a_1+a_1a_3)(1-a_2+a_1a_2)} + \frac{(1+a_1a_2a_3)^2}{(1-a_3+a_2a_3)(1-a_2+a_1a_2)}\right) + 34.$$

Proof. It is easy to see that the moduli algebra $A^2(V) = \mathbb{C}\{x_1, x_2, x_3\}/(f, m^2J(f))$ has dimension $(a_1a_2a_3 + 11)$ and has a monomial basis of the form:

$$\begin{aligned} & \{x_1^{i_1}x_2^{i_2}x_3^{i_3}, 1 \leq i_1 \leq a_1 - 2; 1 \leq i_2 \leq a_2 - 1; 0 \leq i_3 \leq a_3 - 2; x_2^{i_2}, a_2 \leq i_2 \leq 2a_2 - 1; \\ & x_2^{i_2}x_3^{i_3}, 0 \leq i_2 \leq a_2 - 2; 0 \leq i_3 \leq 2a_3 - 1; x_2^{a_2-1}x_3^{i_3}, 0 \leq i_3 \leq a_3 - 1; x_2^{a_2}x_3; \\ & x_1^{i_1}x_2^{i_2}, 1 \leq i_1 \leq a_1 - 3; a_2 \leq i_2 \leq 2a_2 - 1; x_1^{i_1}x_3^{i_3}, 1 \leq i_1 \leq 2; 0 \leq i_3 \leq a_3 - 1; x_1x_3^{a_3}; \\ & x_1x_2x_3^{a_3}, x_1^{a_1-1}x_2x_3; x_1^{i_1}x_3^{i_3}, 3 \leq i_1 \leq a_1 - 1; 0 \leq i_3 \leq a_3 - 2; x_1^{a_1}; x_1^{a_1}x_3; x_1^{a_1+1}; x_1^{a_1-2}x_2^{a_2}; \\ & x_1^{a_1-1}x_2^{i_2}, 1 \leq i_2 \leq 2\}. \end{aligned}$$

Using the same steps as, we used in above propositions, we obtain the following description of Lie algebras in question. The derivations represented by the following vector fields form a basis in $\text{Der}A^2(V)$:

$$\begin{aligned}
& x_2^{i_2} x_3^{i_3} \partial_1, \quad 0 \leq i_2 \leq a_2 - 2; a_3 \leq i_3 \leq 2a_3 - 1; x_2^{a_2-1} x_3^{i_3} \partial_1, \quad 1 \leq i_3 \leq a_3 - 1; x_2^{a_2-2} x_3^{a_3-1} \partial_1; \\
& x_2^{a_2} x_3 \partial_1; x_1^{i_1} \partial_1 + \frac{a_1(a_3 - 1) + 1}{a_3(a_2 - 1) + 1} x_1^{i_1-1} x_2 \partial_2 + \frac{a_1(a_2 - 1) + 1}{a_3(a_2 - 1) + 1} x_1^{i_1-1} x_3 \partial_3, \quad 1 \leq i_1 \leq a_1 - 1; \\
& x_2^{i_2} \partial_1, \quad a_2 \leq i_2 \leq 2a_2 - 1; x_1 x_3^{i_3} \partial_1 + \frac{a_1(a_3 - 1) + 1}{a_3(a_2 - 1) + 1} x_2 x_3^{i_3} \partial_2 + \frac{a_1(a_2 - 1) + 1}{a_3(a_2 - 1) + 1} x_3^{i_3+1} \partial_3, \quad 1 \leq i_3 \leq a_3 - 2; \\
& x_1 x_3^{i_3} \partial_1, \quad a_3 - 1 \leq i_3 \leq a_3; x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_1, \quad 1 \leq i_1 \leq a_1 - 2; 1 \leq i_2 \leq a_2 - 1; 0 \leq i_3 \leq a_3 - 2; \\
& x_1^{i_1} x_2^{i_2} \partial_1, \quad 1 \leq i_1 \leq a_1 - 3; a_2 \leq i_2 \leq 2a_2 - 1; x_1 x_2 x_3^{a_3-1} \partial_1; x_1^{i_1} x_3^{i_3} \partial_1, \quad 2 \leq i_1 \leq a_1 - 1; 1 \leq i_3 \leq a_3 - 2; \\
& x_1^{a_1-1} x_2^{i_2} \partial_1, \quad 1 \leq i_2 \leq 2; x_1^2 x_3^{a_3-1} \partial_1; x_1^{a_1-1} x_2 x_3 \partial_1; x_1^{i_1} \partial_1, \quad a_1 \leq i_1 \leq a_1 + 1; x_1^{a_1} x_3 \partial_1; x_1^{a_1-2} x_2^{a_2} \partial_1; \\
& x_2^{i_2} x_3^{i_3} \partial_2, \quad 0 \leq i_2 \leq a_2 - 2; a_3 \leq i_3 \leq 2a_3 - 1; x_2^{i_2} x_3^{i_3} \partial_2, \quad 2 \leq i_2 \leq a_2 - 1; 1 \leq i_3 \leq a_3 - 1; \\
& x_2 x_3^{a_3-1} \partial_2; x_2^{i_2} \partial_2, \quad a_2 \leq i_2 \leq 2a_2 - 1; x_2^{a_2} x_3 \partial_2; x_1^{i_1} x_2^{i_2} \partial_2, \quad 1 \leq i_1 \leq a_1 - 2; 2 \leq i_2 \leq a_2; \\
& x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_2, \quad 1 \leq i_1 \leq a_1 - 2; 1 \leq i_2 \leq a_2 - 1; 1 \leq i_3 \leq a_3 - 2; x_1 x_2 x_3^{a_3-1} \partial_2; \\
& x_1^{i_1} x_2^{i_2} \partial_2, \quad 1 \leq i_1 \leq a_1 - 3; a_2 + 1 \leq i_2 \leq 2a_2 - 1; x_1 x_3^{i_3} \partial_2, \quad a_3 - 1 \leq i_3 \leq a_3; x_1^{a_1-1} x_2^{i_2} \partial_2, \quad 1 \leq i_2 \leq 2; \\
& x_1^{a_1-1} x_2 x_3 \partial_2; x_1^{a_1} \partial_2; x_1^{a_1} x_3 \partial_2; x_1^{a_1+1} \partial_2; x_1^{a_1-1} x_3^{a_3-2} \partial_2; x_1^2 x_3^{a_3-1} \partial_2; x_2^{a_2-2} x_3^{a_3-1} \partial_3; \\
& x_2^{i_2} x_3^{i_3} \partial_3, \quad 0 \leq i_2 \leq a_2 - 2; a_3 \leq i_3 \leq 2a_3 - 1; x_2^{a_2-1} x_3^{i_3} \partial_3, \quad 1 \leq i_3 \leq a_3 - 2; x_2^{i_2} \partial_3, \quad a_2 \leq i_2 \leq 2a_2 - 1; \\
& x_2^{a_2} x_3 \partial_3; x_1^{i_1} x_2^{i_2} x_3^{i_3} \partial_3, \quad 2 \leq i_1 \leq a_1 - 2; 2 \leq i_2 \leq a_2 - 1; 1 \leq i_3 \leq a_3 - 2; \\
& x_1^{i_1} x_2^{i_2} \partial_3, \quad 1 \leq i_1 \leq a_1 - 3; a_2 \leq i_2 \leq 2a_2 - 1; x_1^{a_1} x_3 \partial_3; x_1^{a_1+1} \partial_3; x_1^{a_1-1} x_2 x_3 \partial_3; x_1 x_3^{a_3} \partial_3; \\
& x_1 x_2 x_3^{a_3-1} \partial_3; x_1 x_2^{a_2-1} x_3 \partial_3; x_1^{a_1-2} x_3^{a_3-1} \partial_3; x_1^{a_1-1} x_2^{a_2} \partial_3; x_1^{a_1-1} x_3^{a_3-2} \partial_3; x_1^{a_1-1} x_2^{a_2-2} \partial_3; x_1^{a_1-2} x_2^{a_2-1} \partial_3; \\
& x_1^{a_1-2} x_2 x_3^{a_3-2} \partial_3; x_1 x_2^{i_2} x_3^{a_3-2} \partial_3, \quad 1 \leq i_2 \leq a_2 - 1.
\end{aligned}$$

Therefore we have

$$\lambda^2(V) = 3a_1 a_2 a_3 + 2(a_1 + a_2 + a_3) - 2(a_1 a_2 + a_1 a_3 + a_2 a_3) + 43.$$

In case of $a_1 = 2, a_2 \geq 3, a_3 \geq 3$, we obtain the basis of derivation represented by following derivations form a basis of $\text{Der}A^2(V)$:

$$\begin{aligned}
& x_2^{i_2} x_3^{i_3} \partial_1, \quad 0 \leq i_2 \leq a_2 - 2; a_3 \leq i_3 \leq 2a_3 - 1; x_2^{a_2-2} x_3^{a_3-1} \partial_1; x_2^{a_2-1} x_3^{i_3} \partial_1, \quad 1 \leq i_3 \leq a_3; \\
& x_1^2 x_3 \partial_1; x_1^2 \partial_1; x_1 x_3^{i_3} \partial_1 + \frac{2(a_3 - 1) + 1}{a_3(a_2 - 1) + 1} x_2 x_3^{i_3} \partial_2 + \frac{2(a_2 - 1) + 1}{a_3(a_2 - 1) + 1} x_3^{i_3+1} \partial_3, \quad 0 \leq i_3 \leq a_3 - 3; \\
& x_2^{i_2} \partial_1, \quad a_2 \leq i_2 \leq a_2 + 1; x_2^{a_2} x_3 \partial_1; x_1^3 \partial_1; x_1 x_3^{a_3-2} \partial_1 + \frac{1}{a_3} x_3^{a_3-1} \partial_3; x_1 x_3^{a_3-1} \partial_1 - \frac{2}{a_3} x_1 x_2 \partial_3; \\
& x_1 x_3^{a_3} \partial_1; x_1 x_2^{i_2} \partial_1 + \frac{2(a_3 - 1) + 1}{a_3(a_2 - 1) + 1} x_2^{i_2+1} \partial_2 + \frac{2(a_2 - 1) + 1}{a_3(a_2 - 1) + 1} x_2^{i_2} x_3 \partial_3, \quad 1 \leq i_2 \leq 2; \\
& x_1 x_2 x_3 \partial_1 + \frac{1}{a_3} x_2 x_3^{a_3-3} \partial_3; x_2^{i_2} x_3^{i_3} \partial_2, \quad 0 \leq i_2 \leq a_2 - 2; a_3 \leq i_3 \leq 2a_3 - 1; x_2 x_3^{a_3-2} \partial_2 + \frac{1}{a_3} x_3^{a_3-1} \partial_3;
\end{aligned}$$

$$\begin{aligned}
& x_2 x_3^{a_3-1} \partial_2 - \frac{2}{a_3} x_1 x_2 \partial_3; x_2^{i_2} x_3^{i_3} \partial_2, \quad 2 \leq i_2 \leq a_2 - 1; 1 \leq i_3 \leq a_3 - 1; x_2^{i_2} \partial_2, \quad a_2 - 1 \leq i_2 \leq a_2 + 1; \\
& x_2^{a_2-1} x_3^{a_3} \partial_2; x_2^{a_2} x_3 \partial_2; x_1 x_3^{i_3} \partial_2, \quad a_3 - 2 \leq i_3 \leq a_3; x_1 x_2 \partial_2 - (a_2 - 1) x_1 x_3 \partial_3; x_1 x_2 x_3 \partial_1 - (a_2 - 1) x_1 x_3^2 \partial_3; \\
& x_1 x_2^2 \partial_2; x_1^2 \partial_2 - \frac{a_2}{a_3} x_2^{a_2-2} x_3^{a_3-3} \partial_3; x_1^2 x_3 \partial_2 - \frac{a_2}{a_3} x_2^{a_2-2} x_3^{a_3-2} \partial_3; x_1^3 \partial_2; (2x_1 x_2 + x_3^{a_3}) \partial_3; (2x_1 x_2^2 + x_2 x_3^{a_3}) \partial_3; \\
& x_2^{i_2} x_3^{i_3} \partial_3, \quad 0 \leq i_2 \leq a_2 - 2; a_3 + 1 \leq i_3 \leq 2a_3 - 1; x_2^{i_2} x_3^{a_3} \partial_1, \quad a_2 - 3 \leq i_2 \leq a_2 - 2; x_2^{a_2-2} x_3^{a_3-1} \partial_3; \\
& x_2^{i_2} \partial_3, \quad a_2 - 1 \leq i_2 \leq a_2 + 1; x_2^{a_2-1} x_3^{i_3} \partial_3, \quad 1 \leq i_3 \leq a_3; x_2^{a_2} x_3 \partial_3; x_1 x_3^{i_3} \partial_3, \quad a_3 - 2 \leq i_3 \leq a_3; \\
& x_1 x_2 x_3 \partial_3; x_1^2 \partial_3; x_1^2 x_3 \partial_3; x_1^3 \partial_3.
\end{aligned}$$

Therefore we have

$$\lambda^2(V) = 4a_2 a_3 - 2(a_2 + a_3) + 46.$$

It is easy to see that when $\frac{1+a_1 a_2 a_3}{1-a_3+a_2 a_3} \geq 3$, $\frac{1+a_1 a_2 a_3}{1-a_1+a_1 a_3} \geq 3$, $\frac{1+a_1 a_2 a_3}{1-a_2+a_1 a_2} \geq 3$, then

$$\begin{aligned}
h_2\left(\frac{1}{w_1}, \frac{1}{w_2}, \frac{1}{w_3}\right) &= \frac{3(1+a_1 a_2 a_3)^3}{(1-a_3+a_2 a_3)(1-a_1+a_1 a_3)(1-a_2+a_1 a_2)} + 5\left(\frac{1+a_1 a_2 a_3}{1-a_3+a_2 a_3} + \frac{1+a_1 a_2 a_3}{1-a_1+a_1 a_3} + \frac{1+a_1 a_2 a_3}{1-a_2+a_1 a_2}\right) \\
&- 4\left(\frac{(1+a_1 a_2 a_3)^2}{(1-a_3+a_2 a_3)(1-a_1+a_1 a_3)} + \frac{(1+a_1 a_2 a_3)^2}{(1-a_1+a_1 a_3)(1-a_2+a_1 a_2)} + \frac{(1+a_1 a_2 a_3)^2}{(1-a_3+a_2 a_3)(1-a_2+a_1 a_2)}\right) + 34.
\end{aligned}$$

When $a_1 \geq 3, a_2 \geq 3, a_3 \geq 3$, then need to show that

$$\begin{aligned}
& 3a_1 a_2 a_3 + 2(a_1 + a_2 + a_3) - 2(a_1 a_2 + a_1 a_3 + a_2 a_3) + 43 \leq \frac{3(1+a_1 a_2 a_3)^3}{(1-a_3+a_2 a_3)(1-a_1+a_1 a_3)(1-a_2+a_1 a_2)} \\
& + 5\left(\frac{1+a_1 a_2 a_3}{1-a_3+a_2 a_3} + \frac{1+a_1 a_2 a_3}{1-a_1+a_1 a_3} + \frac{1+a_1 a_2 a_3}{1-a_2+a_1 a_2}\right) - 4\left(\frac{(1+a_1 a_2 a_3)^2}{(1-a_3+a_2 a_3)(1-a_1+a_1 a_3)} + \frac{(1+a_1 a_2 a_3)^2}{(1-a_1+a_1 a_3)(1-a_2+a_1 a_2)}\right) \\
& + \frac{(1+a_1 a_2 a_3)^2}{(1-a_3+a_2 a_3)(1-a_2+a_1 a_2)} + 34.
\end{aligned}$$

After simplification we get

$$\begin{aligned}
& 6(a_1 a_2 + a_2 a_3 + a_1 a_3) + a_1(a_2 - 3) + a_2(a_3 - 3) + a_3(a_1 - 3) + 2a_1^2[a_2(a_3 - 3) + a_3(a_2 - 3)] \\
& + 2a_2^2[a_1(a_3 - 3) + a_3(a_1 - 3)] + 2a_3^2[a_1(a_2 - 3) + a_2(a_1 - 3)] + 2(a_1^2 + a_2^2 + a_3^2) + 2(a_1^3 a_2 + a_2^3 a_3 \\
& + a_3^3 a_1) + a_1^2 a_2^2 a_3^2 + 10(a_1 a_2^2 a_3 + a_1 a_2 a_3^2) + 4a_1^2 a_2 a_3 + 2a_1 a_2 a_3[3a_1 - 8] + a_1^3 a_2 a_3^2(a_3 - 3)(a_2 - 3) \\
& + a_1^2 a_3^2(a_3 - 3)(a_1 a_2 - 2) + 2a_1^2 a_2 a_3^2(a_3 + a_2 - 6) + 2a_1 a_2 a_3^3(a_1 - 3) + a_1^2 a_2^3 a_3(a_3 - 3)(a_1 - 3) \\
& + a_1^2 a_2^2(a_1 - 3)(a_2 a_3 - 2) + 2a_1^3 a_2 a_3(a_2 - 3) + 2a_1^2 a_2^2 a_3(a_1 - 3 + (a_3 - 3)) + a_1 a_2^2 a_3^3(a_2 - 3)(a_1 \\
& - 3) + a_2^2 a_3^2(a_2 - 3)(a_1 a_3 - 2) + 2a_1 a_2^2 a_3^2(a_1 + a_2 - 6) + 2a_2^3 a_3 a_1(a_3 - 3) - 3 \geq 0. \quad \square
\end{aligned}$$

Proposition 3.7. *Let $(V, 0)$ be a fewnomial surface isolated singularity of type 4 which is defined by $f = x_1^{a_1} + x_2^{a_2} + x_3^{a_3} x_2$ ($a_1 \geq 3, a_2 \geq 3, a_3 \geq 2$) with weight type $(\frac{1}{a_1}, \frac{1}{a_2}, \frac{a_2-1}{a_2 a_3}; 1)$.*

Then

$$\lambda^2(V) = \begin{cases} 7a_1 a_3 - 6a_1 - 10a_3 + 48; & a_1 \geq 3, a_2 = 3, a_3 \geq 3 \\ 5a_2 a_3 - 7a_2 - 4a_3 + 53; & a_1 = 3, a_2 \geq 4, a_3 \geq 3 \\ 3a_1 a_2 a_3 + 6a_1 + 5a_2 + 2a_3 - 4a_2 a_3 - 4a_1 a_2 - 2a_1 a_3 + 35; & a_1 \geq 4, a_2 \geq 4, a_3 \geq 3 \\ 11a_1 a_2 - 3a_1 - 15a_2 + 41; & a_1 \geq 4, a_2 \geq 4, a_3 = 5 \\ 46; & a_1 = 3, a_2 = 3, a_3 = 2 \\ 2a_1 a_2 + 2a_1 - 3a_2 + 36; & a_1 \geq 4, a_2 \geq 4, a_3 = 2 \\ 3a_2 + 40; & a_1 = 3, a_2 \geq 4, a_3 = 2 \\ 8a_1 + 26; & a_1 \geq 4, a_2 = 3, a_3 = 2 \end{cases}$$

Furthermore, we need to show that when $a_1 \geq 4, a_2 \geq 4, a_3 \geq 3$, then

$$3a_1a_2a_3 + 6a_1 + 5a_2 + 2a_3 - 4a_2a_3 - 4a_1a_2 - 2a_1a_3 + 35 \leq \frac{3a_1a_2^2a_3}{a_2-1} + 5(a_1 + a_2 + \frac{a_2a_3}{a_2-1}) - 4(a_1a_2 + \frac{a_1a_2a_3}{a_2-1} + \frac{a_2^2a_3}{a_2-1}) + 34.$$

Proof. It is easy to see that the moduli algebra $A^2(V) = \mathbb{C}\{x_1, x_2, x_3\}/(f, m^2J(f))$ has dimension $(a_1a_2a_3 - a_1a_2 - a_2a_3 + a_1 + a_2 + 10)$ and has a monomial basis of the form:

$$\{x_1^{i_1}x_2^{i_2}x_3^{i_3}, 0 \leq i_1 \leq a_1 - 2; 0 \leq i_2 \leq a_2 - 2; 0 \leq i_3 \leq a_3 - 2; x_1^{a_1-1}; x_1^{a_1-1}x_2; x_1^{a_1-1}x_3; x_2x_3^{a_3-1}; x_2x_3^{a_3}; x_2^{a_2-1}x_3; x_2^{a_2-1}; x_2^{a_2}; x_2^{a_2-2}x_3^{a_3-1}; x_1x_3^{a_3-1}; x_1x_2x_3^{a_3-1}; x_1^{i_1}x_3^{i_3}, 0 \leq i_1 \leq a_1 - 2; a_3 - 1 \leq i_3 \leq 2a_3 - 2\}.$$

We obtain the basis of derivation represented by following derivations form a basis of $\text{Der}A^2(V)$:

$$(a_2x_2^2 + x_3^{a_3})\partial_1; (a_2x_2^2x_3 + x_3^{a_3+1})\partial_1; x_2x_3^{i_3}\partial_1, a_3 - 2 \leq i_3 \leq a_3; x_2^3\partial_1; x_1\partial_1 + \frac{a_1}{3}x_2\partial_2 + \frac{2a_1}{3a_3}x_3\partial_3; x_1^{i_1}x_3^{i_3}\partial_1, 1 \leq i_1 \leq a_1 - 2, 1 \leq i_3 \leq 2a_3 - 2; x_1^{i_1}x_2x_3^{i_3}\partial_1, 1 \leq i_1 \leq a_1 - 2, 0 \leq i_3 \leq a_3 - 2; x_1^{a_1-1}\partial_1; x_1^{a_1-1}x_2\partial_1; x_1^{a_1-1}x_3\partial_1; x_1x_2x_3^{a_3-1}\partial_1; x_1x_2^2\partial_1; x_1^{i_1}\partial_1, 2 \leq i_1 \leq a_1 - 2; x_3^{i_3}\partial_2, a_3 \leq i_3 \leq 2a_3 - 1; x_2x_3^{i_3}\partial_2, 1 \leq i_3 \leq a_3; x_1^{i_1}x_3^{i_3}\partial_2, 1 \leq i_1 \leq a_1 - 2, a_3 - 1 \leq i_3 \leq 2a_3 - 2; x_2^2\partial_2; x_2^3\partial_2; x_2^2x_3\partial_2; x_1^{i_1}x_2x_3^{i_3}\partial_2, 1 \leq i_1 \leq a_1 - 2, 0 \leq i_3 \leq a_3 - 2; x_1^{a_1-1}\partial_2; x_1^{a_1-1}x_2\partial_2; x_1^{a_1-1}x_3\partial_2; x_1^{a_1-2}x_3^{a_3-2}\partial_2; x_1x_2x_3^{a_3-1}\partial_2; x_1x_2^2\partial_2; x_3^{i_3}\partial_3, a_3 \leq i_3 \leq 2a_3 - 1; x_2x_3^{i_3}\partial_3, 1 \leq i_3 \leq a_3; x_2^2\partial_3; x_2^2x_3\partial_3; x_2^3\partial_3; x_1^{i_1}x_3^{i_3}\partial_3, 1 \leq i_1 \leq a_1 - 2, a_3 \leq i_3 \leq 2a_3 - 2; x_1^{a_1-3}x_3^{a_3-1}\partial_3; x_1^{a_1-2}x_3^{a_3-2}\partial_3; x_1^{i_1}x_2x_3^{i_3}\partial_3, 1 \leq i_1 \leq a_1 - 4, 1 \leq i_3 \leq a_3 - 2; x_1^{i_1}x_2x_3^{i_3}\partial_3, a_1 - 3 \leq i_1 \leq a_1 - 2, 0 \leq i_3 \leq a_3 - 2; x_1^{a_1-2}x_3^{a_3-1}\partial_3; x_1^{a_1-1}\partial_3; x_1^{a_1-1}x_2\partial_3; x_1^{a_1-1}x_3\partial_3; x_1x_2^2\partial_3; x_1x_2x_3^{a_3-1}\partial_3.$$

Therefore we have

$$\lambda^2(V) = 7a_1a_3 - 6a_1 - 10a_3 + 48.$$

In case of $a_1 = 3, a_2 \geq 4, a_3 \geq 3$, we obtain the basis of derivation represented by following derivations form a basis of $\text{Der}A^2(V)$:

$$(a_2x_2^{a_2-1} + x_3^{a_3})\partial_1; (a_2x_2^{a_2-1}x_3 + x_3^{a_3+1})\partial_1; x_3^{i_3}\partial_1, 2a_3 - 3 \leq i_3 \leq 2a_3 - 2; x_2x_3^{i_3}\partial_1, a_3 - 1 \leq i_3 \leq a_3; x_2^2x_3^{a_3-1}\partial_1; x_2^{a_2-2}x_3^{a_3-2}\partial_1; x_2^{a_2}\partial_1; x_1\partial_1 + \frac{3}{a_2}x_2\partial_2 + \frac{3(a_2-1)}{a_2a_3}x_3\partial_3; x_1x_3^{i_3}\partial_1, 1 \leq i_3 \leq 2a_3 - 2; x_1x_2^{i_2}x_3^{i_3}\partial_1, 1 \leq i_2 \leq a_2 - 2; 0 \leq i_3 \leq a_3 - 2; x_1x_2^{a_2-1}\partial_1; x_1^2\partial_1; x_1^2x_2\partial_1; x_1^2x_3\partial_1; x_1x_2x_3^{a_3-1}\partial_1; x_3^{i_3}\partial_2, a_3 - 1 \leq i_3 \leq 2a_3 - 2; x_2^{i_2}x_3^{i_3}\partial_2, 1 \leq i_2 \leq a_2 - 2; 1 \leq i_3 \leq a_3 - 2; x_2^{i_2}\partial_2, 2 \leq i_2 \leq a_2; x_1x_2^{i_2}x_3^{i_3}\partial_2, 1 \leq i_2 \leq a_2 - 2; 0 \leq i_3 \leq a_3 - 2; x_1^2\partial_2; x_1^2x_2\partial_2; x_1^2x_3\partial_2; x_1x_2x_3^{a_3-1}\partial_2; x_1x_2^{a_2-1}\partial_2; x_1x_3^{i_3}\partial_2, a_3 - 2 \leq i_3 \leq 2a_3 - 2; x_2^2x_3^{a_3-1}\partial_2; x_2x_3^{a_3-1}\partial_2; x_2x_3^{a_3}\partial_2; x_3^{i_3}\partial_3, a_3 - 1 \leq i_3 \leq 2a_3 - 2; x_2^{i_2}x_3^{i_3}\partial_3, 1 \leq i_2 \leq a_2 - 2; 1 \leq i_3 \leq a_3 - 2; x_2^{i_2}\partial_3, a_2 - 2 \leq i_2 \leq a_2; x_2x_3^{i_3}\partial_3, a_3 - 1 \leq i_3 \leq a_3;$$

$x_2^{a_2-1}x_3\partial_2; x_2^{a_2-1}x_3\partial_3; x_1x_3^{i_3}\partial_3, a_3-2 \leq i_3 \leq 2a_3-2; x_1x_2^{i_2}x_3^{i_3}\partial_3, 1 \leq i_2 \leq a_2-4; 1 \leq i_3 \leq a_3-2;$
 $x_1x_2^{i_2}x_3^{i_3}\partial_3, a_2-3 \leq i_2 \leq a_2-2; 0 \leq i_3 \leq a_3-2; x_1x_2^{a_2-1}\partial_3; x_1^2\partial_3; x_1^2x_2\partial_3; x_1^2x_3\partial_3; x_1x_2x_3^{a_3-1}\partial_3.$

Therefore we have

$$\lambda^2(V) = 5a_2a_3 - 7a_2 - 4a_3 + 53.$$

In case of $a_1 \geq 4, a_2 \geq 4, a_3 \geq 3,$

$(a_2x_2^{a_2-1}+x_3^{a_3})\partial_1; (a_2x_2^{a_2-1}x_3+x_3^{a_3+1})\partial_1; x_3^{i_3}\partial_1, 2a_3-3 \leq i_3 \leq 2a_3-2; x_2x_3^{i_3}\partial_1, a_3-2 \leq i_3 \leq a_3;$
 $x_2^{a_2}\partial_1; x_1\partial_1 + \frac{a_1}{a_2}x_2\partial_2 + \frac{a_1(a_2-1)}{a_2a_3}x_3\partial_3; x_1^{i_1}x_3^{i_3}\partial_1, 1 \leq i_1 \leq a_1-2; 1 \leq i_3 \leq 2a_3-2;$
 $x_1^{i_1}x_2^{i_2}x_3^{i_3}\partial_1, 1 \leq i_1 \leq a_1-2; 1 \leq i_2 \leq a_2-2; 0 \leq i_3 \leq a_3-2; x_1^{i_1}\partial_1, 2 \leq i_1 \leq a_1-1; x_1x_2x_3^{a_3-1}\partial_1;$
 $x_1^{a_1-1}x_2\partial_1; x_1^{a_1-1}x_3\partial_1; x_1x_2^{a_2-1}\partial_1; x_3^{a_3-1}\partial_2 + \frac{a_3}{a_2-1}x_2^{a_2-2}\partial_3; x_3^{i_3}\partial_2, a_3 \leq i_3 \leq 2a_3-2;$
 $x_1^{i_1}x_2^{i_2}x_3^{i_3}\partial_2, 0 \leq i_1 \leq a_1-2; 1 \leq i_2 \leq a_2-2; 1 \leq i_3 \leq a_3-2; x_2^{i_2}\partial_2, 2 \leq i_2 \leq a_2;$
 $x_1^{i_1}x_3^{i_3}\partial_2, 1 \leq i_1 \leq a_1-2; a_3-1 \leq i_3 \leq 2a_3-2; x_1^{i_1}x_2^{i_2}\partial_2, 1 \leq i_1 \leq a_1-2; 1 \leq i_2 \leq a_2-2;$
 $x_1^{a_1-1}\partial_2; x_1^{a_1-1}x_2\partial_2; x_1^{a_1-1}x_3\partial_2; x_1x_2x_3^{a_3-1}\partial_2; x_1x_2^{a_2-1}\partial_2; x_1^{a_1-2}x_3^{a_3-2}\partial_2; x_2^{a_2-1}x_3\partial_2; x_2^{a_2-3}x_3^{a_3-1}\partial_2;$
 $x_2x_3^{i_3}\partial_2, a_3-1 \leq i_3 \leq a_3; x_1^{i_1}x_2^{i_2}x_3^{i_3}\partial_3, 0 \leq i_1 \leq a_1-2; 1 \leq i_2 \leq a_2-2; 1 \leq i_3 \leq a_3-2;$
 $x_1^{a_1-1}\partial_3; x_1^{a_1-1}x_2\partial_3; x_1^{a_1-1}x_3\partial_3; x_2^{i_2}\partial_3, a_2-1 \leq i_2 \leq a_2; x_2x_3^{i_3}\partial_3, a_3-1 \leq i_3 \leq a_3; x_2^2x_3^{a_3-1}\partial_3;$
 $x_1^{i_1}x_3^{i_3}\partial_3, 0 \leq i_1 \leq a_1-2; a_3 \leq i_3 \leq 2a_3-2; x_1x_2^{a_2-1}\partial_3; x_1^{a_1-2}x_3^{i_3}\partial_3, a_3-2 \leq i_3 \leq a_3-1;$
 $x_1^{a_1-2}x_2\partial_3; x_1^{a_1-3}x_3^{a_3-1}\partial_3; x_1^{a_1-3}x_2^{a_2-2}\partial_3; x_1^{a_1-2}x_2^{a_2-2}\partial_3; x_1x_2x_3^{a_3-1}\partial_3; x_2^{a_2-1}x_3\partial_3.$

Therefore we have

$$\lambda^2(V) = 3a_1a_2a_3 + 6a_1 + 5a_2 + 2a_3 - 4a_2a_3 - 4a_1a_2 - 2a_1a_3 + 35.$$

In case of $a_1 \geq 4, a_2 \geq 4, a_3 = 5,$

$(a_2x_2^{a_2-1}+x_3^5)\partial_1; (a_2x_2^{a_2-1}x_3+x_3^6)\partial_1; x_3^{i_3}\partial_1, 7 \leq i_3 \leq 8; x_2x_3^{i_3}\partial_1, 3 \leq i_3 \leq 5;$
 $x_2^{a_2}\partial_1; x_1\partial_1 + \frac{a_1}{a_2}x_2\partial_2 + \frac{a_1(a_2-1)}{5a_2}x_3\partial_3; x_1^{i_1}x_3^{i_3}\partial_1, 1 \leq i_1 \leq a_1-2; 1 \leq i_3 \leq 8;$
 $x_1^{i_1}x_2^{i_2}x_3^{i_3}\partial_1, 1 \leq i_1 \leq a_1-2; 1 \leq i_2 \leq a_2-2; 0 \leq i_3 \leq 3; x_1^{i_1}\partial_1, 2 \leq i_1 \leq a_1-1; x_1x_2x_3^4\partial_1;$
 $x_1^{a_1-1}x_2\partial_1; x_1^{a_1-1}x_3\partial_1; x_1x_2^{a_2-1}\partial_1; x_3^4\partial_2 + \frac{5}{a_2-1}x_2^{a_2-2}\partial_3; x_3^{i_3}\partial_2, 5 \leq i_3 \leq 8;$
 $x_1^{i_1}x_2^{i_2}x_3^{i_3}\partial_2, 0 \leq i_1 \leq a_1-2; 1 \leq i_2 \leq a_2-2; 1 \leq i_3 \leq 3; x_2^{i_2}\partial_2, 2 \leq i_2 \leq a_2;$
 $x_1^{i_1}x_3^{i_3}\partial_2, 1 \leq i_1 \leq a_1-2; 4 \leq i_3 \leq 8; x_1^{i_1}x_2^{i_2}\partial_2, 1 \leq i_1 \leq a_1-2; 1 \leq i_2 \leq a_2-2;$
 $x_1^{a_1-1}\partial_2; x_1^{a_1-1}x_2\partial_2; x_1^{a_1-1}x_3\partial_2; x_1x_2x_3^4\partial_2; x_1x_2^{a_2-1}\partial_2; x_1^{a_1-2}x_3^3\partial_2; x_2^{a_2-1}x_3\partial_2; x_2^{a_2-3}x_3^4\partial_2;$
 $x_2x_3^{i_3}\partial_2, 4 \leq i_3 \leq 5; x_1^{i_1}x_2^{i_2}x_3^{i_3}\partial_3, 0 \leq i_1 \leq a_1-2; 1 \leq i_2 \leq a_2-2; 1 \leq i_3 \leq a_3-2;$
 $x_1^{a_1-1}\partial_3; x_1^{a_1-1}x_2\partial_3; x_1^{a_1-1}x_3\partial_3; x_2^{i_2}\partial_3, a_2-1 \leq i_2 \leq a_2; x_2x_3^{i_3}\partial_3, a_3-1 \leq i_3 \leq a_3; x_2^{a_2-3}x_3^4\partial_3;$
 $x_1^{i_1}x_3^{i_3}\partial_3, 1 \leq i_1 \leq a_1-2; 4 \leq i_3 \leq 8; x_2^{a_2-1}x_3\partial_3; x_1^{a_1-2}x_2^{i_2}\partial_3, a_2-3 \leq i_2 \leq a_2-2;$

$$x_1x_2x_3^{a_3-1}\partial_3; x_1^{a_1-2}x_3^{a_3-2}\partial_3; x_1x_2^{i_2}\partial_3; a_2 - 2 \leq i_2 \leq a_2 - 1; x_3^{i_3}\partial_3; a_3 \leq i_3 \leq 2a_3 - 2.$$

Therefore we have

$$\lambda^2(V) = 11a_1a_2 - 3a_1 - 15a_2 + 41.$$

In case of $a_1 = 3, a_2 = 3, a_3 = 2$,

$$\begin{aligned} & x_3^2\partial_1; x_3^3\partial_1; x_2x_3\partial_1; x_2x_3^2\partial_1; x_2^2\partial_1; x_2^2x_3\partial_1; x_2^3\partial_1; x_1\partial_1 + x_2\partial_2 + x_3\partial_3; x_1x_3\partial_1; x_1x_3^2\partial_1; x_1x_2\partial_1; \\ & x_1x_2x_3\partial_1; x_1x_2^2\partial_1; x_1^2\partial_1; x_1^2x_3\partial_1; x_1^2x_2\partial_1; x_3^2\partial_2; x_3^3\partial_2; x_2x_3\partial_2; x_2x_3^2\partial_2; x_2^2\partial_2; x_2^2x_3\partial_2; x_2^3\partial_2; \\ & x_1x_3\partial_2; x_1x_3^2\partial_2; x_1x_2\partial_2; x_1x_2x_3\partial_2; x_1x_2^2\partial_2; x_1^2\partial_2; x_1^2x_3\partial_2; x_1^2x_2\partial_2; x_3^2\partial_3; x_3^3\partial_3; x_2x_3\partial_3; x_2x_3^2\partial_3; \\ & x_2^2\partial_3; x_2^2x_3\partial_3; x_2^3\partial_3; x_1x_3\partial_3; x_1x_3^2\partial_3; x_1x_2\partial_3; x_1x_2x_3\partial_3; x_1x_2^2\partial_3; x_1^2\partial_3; x_1^2x_3\partial_3; x_1^2x_2\partial_3. \end{aligned}$$

Therefore we have

$$\lambda^2(V) = 46.$$

In case of $a_1 \geq 4, a_2 \geq 4, a_3 = 2$,

$$\begin{aligned} & x_3^2\partial_1; x_3^3\partial_1; x_2x_3\partial_1; x_2x_3^2\partial_1; x_2^2x_3\partial_1; x_2^{i_2}\partial_1, a_2 - 2 \leq i_2 \leq a_2; x_1\partial_1 + \frac{a_1}{a_2}x_2\partial_2 + \frac{a_1(a_2 - 1)}{2a_2}x_3\partial_3; \\ & x_1^{i_1}x_3^{i_3}\partial_1, 1 \leq i_1 \leq a_1 - 2; 1 \leq i_3 \leq 2; x_1^{i_1}x_2^{i_2}\partial_1, 1 \leq i_1 \leq a_1 - 2; 1 \leq i_2 \leq a_2 - 2; x_1^{a_1-1}\partial_1; x_1^{a_1-1}x_2\partial_1; \\ & x_1^{a_1-1}x_3\partial_1; x_1x_2x_3\partial_1; x_1x_2^{a_2-1}\partial_1; x_1^{i_1}\partial_1, 2 \leq i_1 \leq a_1 - 2; x_3^2\partial_2; x_3^3\partial_2; x_2x_3^2\partial_2; x_2^{i_2}\partial_2, 2 \leq i_2 \leq a_2; \\ & x_2^2x_3\partial_2; x_1^{i_1}x_3^{i_3}\partial_2, 1 \leq i_1 \leq a_1 - 2; 1 \leq i_3 \leq 2; x_1^{i_1}x_2^{i_2}\partial_2, 1 \leq i_1 \leq a_1 - 2; 1 \leq i_2 \leq a_2 - 2; \\ & x_1x_2x_3\partial_2; x_1^{a_1-1}\partial_2; x_1^{a_1-1}x_2\partial_2; x_1^{a_1-1}x_3\partial_2; x_1x_2^{a_2-1}\partial_2; x_1^{a_1-2}\partial_2; x_3^2\partial_3; x_3^3\partial_3; x_2x_3\partial_3; x_2x_3^2\partial_3; x_2^2x_3\partial_3; \\ & x_2^{i_2}\partial_3, a_2 - 1 \leq i_2 \leq a_2; x_1^{i_1}x_3^2\partial_3, 1 \leq i_1 \leq a_1 - 2; x_1x_2x_3\partial_3; x_1^{a_1-1}\partial_3; x_1^{a_1-1}x_2\partial_3; x_1^{a_1-1}x_3\partial_3; \\ & x_1^{a_1-3}x_2^{a_2-2}\partial_3; x_1x_2^{a_2-1}\partial_3; x_1^{a_1-2}x_2^{i_2}\partial_3, a_2 - 3 \leq i_2 \leq a_2 - 2; x_1^{i_1}x_3\partial_3, a_1 - 3 \leq i_1 \leq a_1 - 2. \end{aligned}$$

Therefore we have

$$\lambda^2(V) = 2a_1a_2 + 2a_1 - 3a_2 + 36.$$

In case of $a_1 = 3, a_2 \geq 4, a_3 = 2$,

$$\begin{aligned} & x_3^2\partial_1; x_3^3\partial_1; x_2x_3\partial_1; x_2x_3^2\partial_1; x_2^2x_3\partial_1; x_2^{i_2}\partial_1, a_2 - 2 \leq i_2 \leq a_2; x_1\partial_1 + \frac{3}{a_2}x_2\partial_2 + \frac{3(a_2 - 1)}{2a_2}x_3\partial_3; \\ & x_1x_3^{i_3}\partial_1, 1 \leq i_3 \leq 2; x_1x_2^{i_2}\partial_1, 1 \leq i_2 \leq a_2 - 2; x_1^2\partial_1; x_1^2x_2\partial_1; x_1^2x_3\partial_1; x_1x_2x_3\partial_1; x_1x_2^{a_2-1}\partial_1; \\ & x_3^2\partial_2; x_3^3\partial_2; x_2x_3\partial_2; x_2x_3^2\partial_2; x_2^{i_2}\partial_2, 2 \leq i_2 \leq a_2; x_2^2x_3\partial_2; x_1x_3\partial_2; x_1x_3^2\partial_2; x_1^2\partial_2; x_1^2x_2\partial_2; x_1^2x_3\partial_2; \\ & x_1x_2x_3\partial_2; x_1x_2^{i_2}\partial_2, 1 \leq i_2 \leq a_2 - 1; x_3^2\partial_3; x_3^3\partial_3; x_2x_3\partial_3; x_2x_3^2\partial_3; x_2^2x_3\partial_3; x_2^{i_2}\partial_3, a_2 - 2 \leq i_2 \leq a_2; \\ & x_1x_3\partial_3; x_1x_3^2\partial_3; x_1x_2x_3\partial_3; x_1^2\partial_3; x_1^2x_2\partial_3; x_1^2x_3\partial_3; x_1x_2^{i_2}\partial_3, a_2 - 3 \leq i_2 \leq a_2 - 1. \end{aligned}$$

Therefore we have

$$\lambda^2(V) = 3a_2 + 40.$$

In case of $a_1 \geq 4, a_2 = 3, a_3 = 2$,

$$x_3^2\partial_1; x_3^3\partial_1; x_2x_3\partial_1; x_2x_3^2\partial_1; x_2^2x_3\partial_1; x_2^2\partial_1; x_2^3\partial_1; x_1\partial_1 + \frac{a_1}{3}x_2\partial_2 + \frac{2a_1}{3}x_3\partial_3; x_1x_2x_3\partial_1;$$

$$\begin{aligned}
& x_1^{i_1} x_3^{i_3} \partial_1, \quad 1 \leq i_1 \leq a_1 - 2; 1 \leq i_3 \leq 2; x_1^{i_1} \partial_1, \quad 2 \leq i_1 \leq a_1 - 2; x_1^{a_1-1} \partial_1; x_1^{a_1-1} x_2 \partial_1; x_1^{a_1-1} x_3 \partial_1; \\
& x_1^{i_1} x_2 \partial_1, \quad 1 \leq i_1 \leq a_1 - 2; x_1 x_2^2 \partial_1; x_3^2 \partial_2; x_3^3 \partial_2; x_2 x_3 \partial_2; x_2^2 x_3 \partial_2; x_2 x_3^2 \partial_2; x_2^2 \partial_2; x_2^3 \partial_2; x_1 x_2 x_3 \partial_2; \\
& x_1^{i_1} x_3^{i_3} \partial_2, \quad 1 \leq i_1 \leq a_1 - 2; 1 \leq i_3 \leq 2; x_1^{a_1-2} \partial_2; x_1^{a_1-1} \partial_2; x_1^{i_1} x_2 \partial_2, \quad 1 \leq i_1 \leq a_1 - 2; x_1 x_2^2 \partial_2; \\
& x_1^{a_1-1} \partial_2; x_1^{a_1-1} x_2 \partial_2; x_3^2 \partial_3; x_3^3 \partial_3; x_2 x_3 \partial_3; x_2 x_3^2 \partial_3; x_2^2 x_3 \partial_3; x_2^2 \partial_3; x_2^3 \partial_3; x_1 x_2 x_3 \partial_3; x_1^{a_1-1} \partial_3; \\
& x_1^{a_1-1} x_2 \partial_3; x_1^{a_1-1} x_3 \partial_3; x_1^{i_1} x_3^2 \partial_3, \quad 1 \leq i_1 \leq a_1 - 2; x_1^{i_1} x_3 \partial_3, \quad a_1 - 3 \leq i_1 \leq a_1 - 2; x_1 x_2^2 \partial_3; \\
& x_1^{i_1} x_2 \partial_3, \quad a_1 - 3 \leq i_1 \leq a_1 - 2.
\end{aligned}$$

Therefore we have

$$\lambda^2(V) = 8a_1 + 26.$$

We have

$$h_2\left(\frac{1}{w_1}, \frac{1}{w_2}, \frac{1}{w_3}\right) = \begin{cases} \frac{3a_1 a_2^2 a_3}{a_2-1} + 5(a_1 + a_2 + \frac{a_2 a_3}{a_2-1}) - 4(a_1 a_2 + \frac{a_1 a_2 a_3}{a_2-1} + \frac{a_2^2 a_3}{a_2-1}) \\ + 34; & a_1 \geq 3, a_2 \geq 3, a_3 \geq \frac{a_2 a_3}{a_2-1}. \end{cases}$$

Case(i): When $a_1 \geq 4, a_2 \geq 4, a_3 \geq 3$, then need to show that

$$3a_1 a_2 a_3 + 6a_1 + 5a_2 + 2a_3 - 4a_2 a_3 - 4a_1 a_2 - 2a_1 a_3 + 35 \leq \frac{3a_1 a_2^2 a_3}{a_2-1} + 5(a_1 + a_2 + \frac{a_2 a_3}{a_2-1}) - 4(a_1 a_2 + \frac{a_1 a_2 a_3}{a_2-1} + \frac{a_2^2 a_3}{a_2-1}) + 34.$$

After simplification we get

$$(a_2 - 1) + (a_1 - 2)(a_2 - 1)(a_3 - 3) + (a_1 - 2)(a_2 - 3)a_3 + 2a_1 + 3a_3 - 8 \geq 0.$$

Case(ii): When $a_1 \geq 3, a_2 = 3, a_3 \geq 3$, then need to show that

$$7a_1 a_3 - 6a_1 - 10a_3 + 48 \leq \frac{27a_1 a_3}{2} + 5(a_1 + 3 + \frac{3a_3}{2}) - 4(3a_1 + \frac{3a_1 a_3}{2} + \frac{9a_3}{2}) + 34.$$

After simplification we get

$$2(a_1 - 2)(a_3 - 3) + 2a_1 + 3a_3 - 8 \geq 0.$$

Case(iii): When $a_1 = 3, a_2 \geq 4, a_3 \geq 3$, then need to show that

$$5a_2 a_3 - 7a_2 - 4a_3 + 53 \leq \frac{9a_2^2 a_3}{a_2-1} + 5(3 + a_2 + \frac{a_2 a_3}{a_2-1}) - 4(3a_2 + \frac{3a_2 a_3}{a_2-1} + \frac{a_2^2 a_3}{a_2-1}) + 34.$$

After simplification we get

$$(a_2 - 1) + (a_2 - 1)(a_3 - 3) + (a_2 - 3)a_3 + 3a_3 - 2 \geq 0.$$

Case(iv): When $a_1 \geq 4, a_2 \geq 4, a_3 = 5$, then need to show that

$$11a_1 a_2 - 3a_1 - 15a_2 + 41 \leq \frac{15a_1 a_2^2}{a_2-1} + 5(a_1 + a_2 + \frac{5a_2}{a_2-1}) - 4(a_1 a_2 + \frac{5a_1 a_2}{a_2-1} + \frac{5a_2^2}{a_2-1}) + 34.$$

After simplification we get

$$(a_2 - 1) + 2(a_1 - 2)(a_2 - 1) + 5(a_1 - 2)(a_2 - 3) + 3a_3 + 3 \geq 0.$$

Case(v): When $a_1 \geq 4, a_2 \geq 4, a_3 = 2$, then need to show that

$$2a_1 a_2 + 2a_1 - 3a_2 + 36 \leq \frac{6a_1 a_2^2}{a_2-1} + 5(a_1 + a_2 + \frac{3a_2}{a_2-1}) - 4(a_1 a_2 + \frac{3a_1 a_2}{a_2-1} + \frac{2a_2^2}{a_2-1}) + 34.$$

After simplification we get

$$(a_1 - 1)(a_2 - 3) + 4 \geq 0.$$

□

Proposition 3.8. *Let $(V, 0)$ be a fewnomial surface isolated singularity of type 5 which is defined by $f = x_1^{a_1}x_2 + x_2^{a_2}x_1 + x_3^{a_3}$ ($a_1 \geq 2, a_2 \geq 2, a_3 \geq 3$) with weight type $(\frac{a_2-1}{a_1a_2-1}, \frac{a_1-1}{a_1a_2-1}, \frac{1}{a_3}; 1)$. Then*

$$\lambda^2(V) = \begin{cases} 3a_1a_2a_3 - 4a_1a_2 - 2a_2a_3 - 2a_1a_3 + 2a_1 + 2a_2 + 6a_3 + 36; & a_1 \geq 4, a_2 \geq 4, a_3 \geq 3 \\ 7a_2a_3 - 10a_2 + 40; & a_1 = 3, a_2 \geq 3, a_3 \geq 3 \\ 4a_2a_3 - 6a_2 + 39; & a_1 = 2, a_2 \geq 4, a_3 \geq 4 \\ 8a_3 + 26; & a_1 = 2, a_2 = 2, a_3 \geq 4 \\ 46; & a_1 = 2, a_2 = 2, a_3 = 3 \\ 12a_3 + 21; & a_1 = 2, a_2 = 3, a_3 \geq 4 \\ 55; & a_1 = 2, a_2 = 3, a_3 = 3 \end{cases}$$

Furthermore, we need to show that when $a_1 \geq 4, a_2 \geq 4, a_3 \geq 3$, then

$$3a_1a_2a_3 - 4a_1a_2 - 2a_2a_3 - 2a_1a_3 + 2a_1 + 2a_2 + 6a_3 + 36 \leq \frac{3(a_1a_2-1)^2}{(a_1-1)(a_2-1)}a_3 + 5\left(\frac{a_1a_2-1}{a_1-1} + \frac{a_1a_2-1}{a_2-1} + a_3\right) - 4\left(\frac{(a_1a_2-1)^2}{(a_1-1)(a_2-1)} + \frac{a_1a_2-1}{a_2-1}a_3 + \frac{a_1a_2-1}{a_1-1}a_3\right) + 34.$$

Proof. It is easy to see that the moduli algebra $A^2(V) = \mathbb{C}\{x_1, x_2, x_3\}/(f, m^2J(f))$ has dimension $(a_1a_2a_3 - a_1a_2 + 11)$ and has a monomial basis of the form:

$$\begin{aligned} & \{x_1^{i_1}x_2^{i_2}x_3^{i_3}, 1 \leq i_1 \leq a_1 - 2; 0 \leq i_2 \leq a_2 - 1; 0 \leq i_3 \leq a_3 - 2; x_2x_3^{a_3-1}; x_1x_2^{a_2}; x_1x_3^{a_3-1}; \\ & x_2^{i_2}x_3^{i_3}, 0 \leq i_2 \leq 2a_2 - 2; 0 \leq i_3 \leq a_3 - 2; x_1^{a_1-1}x_3^{i_3}, 0 \leq i_3 \leq a_3 - 2; x_1^{a_1-1}x_2x_3; \\ & x_1^{a_1-1}x_2^{i_2}, 1 \leq i_2 \leq 2; x_1^{i_1}, a_1 \leq i_1 \leq a_1 + 1; x_1^{a_1}x_3; x_3^{i_3}, a_3 - 1 \leq i_3 \leq a_3\}. \end{aligned}$$

We obtain the following description of Lie algebras in question. The derivations represented by the following vector fields form a basis in $\text{Der}A^2(V)$:

$$\begin{aligned} & x_1^{i_1}x_2^{i_2}x_3^{i_3}\partial_1, 1 \leq i_1 \leq a_1-2, 0 \leq i_2 \leq a_2-1, 0 \leq i_3 \leq a_3-2; x_2x_3^{a_3-1}\partial_1; x_3^{i_3}\partial_1, a_3-1 \leq i_3 \leq a_3; \\ & x_2^{i_2}x_3^{i_3}\partial_1, a_2-2 \leq i_2 \leq 2a_2-2, 0 \leq i_3 \leq a_3-2; x_1x_3^{a_3-1}\partial_1; x_1x_2^{a_2}\partial_1; x_1^{a_1-1}x_3^{i_3}\partial_1, 0 \leq i_3 \leq a_3-2; \\ & x_1^{a_1-1}x_2^{i_2}\partial_1, 1 \leq i_2 \leq 2; x_1^{a_1-1}x_2x_3\partial_1; x_1^{a_1}\partial_1; x_1^{a_1}x_3\partial_1; x_1^{a_1+1}\partial_1; x_3^{i_3}\partial_2, a_3-1 \leq i_3 \leq a_3; \\ & x_2x_3^{a_3-1}\partial_2; x_2^{a_2-2}x_3^{a_3-2}\partial_2; x_2^{i_2}x_3^{i_3}\partial_2, a_2-1 \leq i_2 \leq 2a_2-2, 0 \leq i_3 \leq a_3-2; x_1x_3^{a_3-1}\partial_2; \\ & x_1^{i_1}x_2^{i_2}x_3^{i_3}\partial_2, 1 \leq i_1 \leq a_1-2, 2 \leq i_2 \leq a_2-1, 0 \leq i_3 \leq a_3-2; x_1^{a_1-2}x_3^{a_3-2}\partial_2; \\ & x_1^{a_1-2}x_2x_3^{i_3}\partial_2, a_3-3 \leq i_3 \leq a_3-2; x_1^{a_1-1}x_3^{i_3}\partial_2, a_3-3 \leq i_3 \leq a_3-2; x_1^{a_1-1}x_2\partial_2; x_1^{a_1-1}x_2^2\partial_2; \\ & x_1^{a_1-1}x_2x_3\partial_2; x_1^{a_1+1}\partial_2; x_1^{a_1}\partial_2; x_1^{a_1}x_3\partial_2; x_1x_2^{a_2}\partial_2; x_1^{a_1-3}x_2x_3^{a_3-2}\partial_2; x_3^{i_3}\partial_3, 2 \leq i_3 \leq a_3; \\ & x_2^{i_2}x_3^{i_3}\partial_3, 1 \leq i_2 \leq 2a_2-2, 1 \leq i_3 \leq a_3-2; x_1x_3^{a_3-1}\partial_3; x_1x_2^{a_2}\partial_3; x_1^{a_1-1}x_3^{i_3}\partial_3, 1 \leq i_3 \leq a_3-2; \\ & x_2x_3^{a_3-1}\partial_3; x_1^{i_1}x_2^{i_2}x_3^{i_3}\partial_3, 1 \leq i_1 \leq a_1-2, 0 \leq i_2 \leq a_2-1, 1 \leq i_3 \leq a_3-2; x_1^{a_1-1}x_2x_3\partial_3; \end{aligned}$$

$x_1^{a_1} x_3 \partial_3; x_1^{a_1} \partial_3; x_1^{a_1-2} x_2^{a_2-1} \partial_3; x_1^{a_1-1} x_2^{a_2-2} \partial_3; (a_1 x_1^{a_1-1} x_2 + x_2^{a_2}) \partial_3; x_2^{i_2} \partial_3, a_2 + 1 \leq i_2 \leq a_2 + 2.$

Therefore we have

$$\lambda^2(V) = 3a_1 a_2 a_3 - 4a_1 a_2 - 2a_2 a_3 - 2a_1 a_3 + 2a_1 + 2a_2 + 6a_3 + 36.$$

In case of $a_1 = 3, a_2 \geq 3, a_3 \geq 3$, we have

$$\begin{aligned} & x_3^{i_3} \partial_1, a_3 - 1 \leq i_3 \leq a_3; x_2 x_3^{a_3-1} \partial_1; x_2^{i_2} x_3^{i_3} \partial_1, a_2 - 2 \leq i_2 \leq 2a_2 - 2, 1 \leq i_3 \leq a_3 - 2; \\ & x_2^{i_2} \partial_1, a_2 - 1 \leq i_2 \leq 2a_2 - 2; x_1 x_3^{a_3-1} \partial_1; x_1 x_2^{i_2} x_3^{i_3} \partial_1, 0 \leq i_2 \leq a_2 - 1, 0 \leq i_3 \leq a_3 - 2; \\ & x_1^2 x_3^{i_3} \partial_1, 0 \leq i_3 \leq a_3 - 2; x_1^2 x_2 \partial_1; x_1^2 x_2^2 \partial_1; x_1^2 x_2 x_3 \partial_1; x_1^3 \partial_1; x_1^3 x_3 \partial_1; x_1^4 \partial_1; x_1 x_2^{a_2} \partial_1; \\ & x_3^{i_3} \partial_2, a_3 - 1 \leq i_3 \leq a_3; x_2 x_3^{a_3-1} \partial_2; x_2^{a_2-2} x_3^{a_3-2} \partial_2; x_2^{i_2} x_3^{i_3} \partial_2, a_2 - 1 \leq i_2 \leq 2a_2 - 2, 0 \leq i_3 \leq a_3 - 2; \\ & x_1 x_3^{i_3} \partial_2, a_3 - 2 \leq i_3 \leq a_3 - 1; x_1^2 x_3^{i_3} \partial_2, a_3 - 3 \leq i_3 \leq a_3 - 2; x_1 x_2 x_3^{i_3} \partial_2, a_3 - 3 \leq i_3 \leq a_3 - 2; \\ & x_1 x_2^{a_2} \partial_2; x_1 x_2^{i_2} x_3^{i_3} \partial_2, 2 \leq i_2 \leq a_2 - 1, 0 \leq i_3 \leq a_3 - 2; x_1^2 x_2 \partial_2; x_1^2 x_2 x_3 \partial_2; x_1^2 x_2^2 \partial_2; x_1^3 \partial_2; x_1^3 x_3 \partial_2; \\ & x_1^4 \partial_2; x_3^{i_3} \partial_3, 2 \leq i_3 \leq a_3; x_2^{i_2} x_3^{i_3} \partial_3, 1 \leq i_2 \leq 2a_2 - 2, 1 \leq i_3 \leq a_3 - 2; x_2 x_3^{a_3-1} \partial_3; \\ & x_1^{i_1} x_3^{i_3} \partial_3, 1 \leq i_1 \leq 2, 1 \leq i_3 \leq a_3 - 2; x_1 x_3^{a_3-1} \partial_3; x_1^2 x_2 x_3 \partial_3; x_1^3 \partial_3; x_1^3 x_3 \partial_3; x_1^4 \partial_3; \\ & x_1 x_2^{i_2} x_3^{i_3} \partial_3, 1 \leq i_2 \leq a_2 - 1, 1 \leq i_3 \leq a_3 - 2; x_1 x_2^{a_2-1} \partial_3; x_1 x_2^{a_2} \partial_3; x_1^2 x_2 \partial_3; x_1^2 x_2^2 \partial_3; \\ & x_2^{i_2} \partial_3, 2a_2 - 3 \leq i_2 \leq 2a_2 - 2. \end{aligned}$$

Therefore we have

$$\lambda^2(V) = 7a_2 a_3 - 10a_2 + 40.$$

In case of $a_1 = 2, a_2 \geq 4, a_3 \geq 4$, we have

$$\begin{aligned} & x_3^{i_3} \partial_1, a_3 - 1 \leq i_3 \leq a_3; x_2 x_3^{a_3-1} \partial_1; x_2^{i_2} x_3^{i_3} \partial_1 - \frac{2}{a_2 - 1} x_2^{a_2-3} x_3^{i_3} \partial_1, a_2 \leq i_2 \leq 2a_2 - 2, 0 \leq i_3 \leq a_3 - 2; \\ & x_2^{a_2-2} x_3^{a_3-2} \partial_1 - \frac{2}{a_2 - 1} x_3^{a_3-2} \partial_2; x_2^{a_2-1} x_3^{i_3} \partial_1 - \frac{2}{a_2 - 1} x_2 x_3^{i_3} \partial_2, a_3 - 3 \leq i_3 \leq a_3 - 2; \\ & x_2^{2a_2-1} \partial_1; x_1 x_3^{i_3} \partial_1, 0 \leq i_3 \leq a_3 - 1; x_1 x_2 \partial_1; x_1 x_2^2 \partial_1; x_1 x_2 x_3 \partial_1; x_1^2 \partial_1; x_1^2 x_3 \partial_1; x_1^3 \partial_1; x_2 x_3^{a_3-1} \partial_2; \\ & x_3^{i_3} \partial_2, a_3 - 1 \leq i_3 \leq a_3; x_2^{a_2-2} x_3^{a_3-2} \partial_2; x_2^{i_2} x_3^{i_3} \partial_2, a_2 - 1 \leq i_2 \leq 2a_2 - 2, 1 \leq i_3 \leq a_3 - 2; \\ & x_2^{i_2} \partial_2, a_2 \leq i_2 \leq 2a_2 - 1; x_1 x_3^{i_3} \partial_2, a_3 - 3 \leq i_3 \leq a_3 - 1; x_1 x_2 \partial_2; x_1 x_2^2 \partial_2; x_1 x_2 x_3 \partial_2; x_1^2 \partial_2; x_1^3 \partial_2; x_1^2 x_3 \partial_2; \\ & x_3^{i_3} \partial_3, 2 \leq i_3 \leq a_3; x_2^{i_2} x_3^{i_3} \partial_3, 1 \leq i_2 \leq 2a_2 - 2, 1 \leq i_3 \leq a_3 - 2; x_2 x_3^{a_3-1} \partial_3; x_1 x_3^{i_3} \partial_3, 1 \leq i_3 \leq a_3 - 1; \\ & x_1 x_2 x_3 \partial_3; x_1^2 \partial_3; x_1^2 x_3 \partial_3; x_1^3 \partial_3; x_1 x_2 \partial_3; x_1 x_2^2 \partial_3; x_3^{i_3} \partial_3, a_3 + 1 \leq i_3 \leq a_3 + 3. \end{aligned}$$

Therefore we have

$$\lambda^2(V) = 4a_2 a_3 - 6a_2 + 39.$$

In case of $a_1 = 2, a_2 = 2, a_3 \geq 4$, we have

$$\begin{aligned} & x_3^{i_3} \partial_1, a_3 - 2 \leq i_3 \leq a_3; x_2^{i_2} x_3^{i_3} \partial_1, 1 \leq i_2 \leq 2, 1 \leq i_3 \leq a_3 - 2; x_2 x_3^{a_3-1} \partial_1; x_2^2 \partial_1; x_2^3 \partial_1; \\ & x_1 x_2 \partial_1; x_1 x_2^2 \partial_1; x_1 x_2 x_3 \partial_1; x_1^2 \partial_1; x_1^2 x_3 \partial_1; x_1^3 \partial_1; x_3^{i_3} \partial_2, a_3 - 2 \leq i_3 \leq a_3; x_2 x_3^{i_3} \partial_2, a_3 - 3 \leq i_3 \leq a_3 - 2; \end{aligned}$$

$$x_2x_3^{a_3-1}\partial_2; x_2^2x_3^{i_3}\partial_2, 0 \leq i_3 \leq a_3-2; x_2^3\partial_2; x_1x_3^{i_3}\partial_2, a_3-3 \leq i_3 \leq a_3-1; x_1x_2\partial_2; x_1x_2^2\partial_2; x_1x_2x_3\partial_2; \\ x_1^2\partial_2; x_1^2x_3\partial_2; x_1^3\partial_2; x_3^{i_3}\partial_3, 2 \leq i_3 \leq a_3; x_2x_3^{i_3}\partial_3, 1 \leq i_3 \leq a_3-1; x_2^2x_3^{i_3}\partial_3, 0 \leq i_3 \leq a_3-2; x_2^3\partial_3; \\ x_1x_3^{i_3}\partial_3, 1 \leq i_3 \leq a_3-1; x_1x_2\partial_3; x_1x_2^2\partial_3; x_1x_2x_3\partial_3; x_1^2\partial_3; x_1^2x_3\partial_3; x_1^3\partial_3.$$

Therefore we have

$$\lambda^2(V) = 8a_3 + 36.$$

In case of $a_1 = 2, a_2 = 3, a_3 \geq 4$, we have

$$x_3^{i_3}\partial_1, a_3-1 \leq i_3 \leq a_3; x_2x_3^{i_3}\partial_1, a_3-2 \leq i_3 \leq a_3-1; x_2^2x_3^{i_3}\partial_1, a_3-3 \leq i_3 \leq a_3-2; x_2^5\partial_1; \\ x_2^{i_2}x_3^{i_3}\partial_1, 3 \leq i_2 \leq 4, 0 \leq i_3 \leq a_3-2; x_1x_3^{i_3}\partial_1, 0 \leq i_3 \leq a_3-1; x_1x_2\partial_1; x_1x_2^2\partial_1; x_1x_2x_3\partial_1; x_1^2\partial_1; \\ x_1^3\partial_1; x_3^{i_3}\partial_2, a_3-1 \leq i_3 \leq a_3; x_2x_3^{i_3}\partial_2, a_3-2 \leq i_3 \leq a_3-1; x_2^{i_2}x_3^{i_3}\partial_2, 2 \leq i_2 \leq 4, 1 \leq i_3 \leq a_3-2; \\ x_1^2x_3\partial_1; x_2^{i_2}\partial_2, a_2-2 \leq i_2 \leq a_2; x_1x_3^{i_3}\partial_2, a_3-3 \leq i_3 \leq a_3-1; x_1x_2\partial_2; x_1x_2^2\partial_2; x_1x_2x_3\partial_2; x_1^2\partial_2; \\ x_1^2x_3\partial_2; x_1^3\partial_2; x_3^{i_3}\partial_3, 2 \leq i_3 \leq a_3; x_2^{i_2}x_3^{i_3}\partial_3, 1 \leq i_2 \leq 2, 1 \leq i_3 \leq a_3-2; x_2x_3^{a_3-1}\partial_3; x_1x_2\partial_3; \\ x_2^{i_2}\partial_3, 3 \leq i_2 \leq 5; x_1x_3^{i_3}\partial_3, 1 \leq i_3 \leq a_3-1; x_1x_2x_3\partial_3; x_1x_2^2\partial_3; x_1^2\partial_3; x_1^2x_3\partial_3; x_1^3\partial_3.$$

Therefore we have

$$\lambda^2(V) = 12a_3 + 21.$$

In case of $a_1 = 2, a_2 = 2, a_3 = 3$, we have

$$x_3^2\partial_1; x_3^3\partial_1; x_2x_3\partial_1; x_2x_3^2\partial_1; x_2^2\partial_1; x_2^2x_3\partial_1; x_2^3\partial_1; x_1\partial_1+x_2\partial_2+x_3\partial_3; x_1x_3\partial_1; x_1x_3^2\partial_1; x_1x_2\partial_1; x_1x_2x_3\partial_1; \\ x_1x_2^2\partial_1; x_1^2\partial_1; x_1^2x_3\partial_1; x_1^3\partial_1; x_3^2\partial_2; x_3^3\partial_2; x_2x_3\partial_2; x_2x_3^2\partial_2; x_2^2\partial_2; x_2^2x_3\partial_2; x_3^3\partial_2; x_1x_3\partial_2; x_1x_3^2\partial_2; x_1x_2\partial_2; \\ x_1x_2^2\partial_2; x_1^2\partial_2; x_1^2x_3\partial_2; x_1^3\partial_2; x_3^2\partial_3; x_3^3\partial_3; x_2x_3\partial_3; x_2x_3^2\partial_3; x_2^2\partial_3; x_2^2x_3\partial_3; x_3^3\partial_3; x_1x_3\partial_3; x_1x_3^2\partial_3; x_1x_2\partial_3; \\ x_1x_2x_3\partial_3; x_1x_2^2\partial_3; x_1^2\partial_3; x_1^2x_3\partial_3; x_1^3\partial_3.$$

Therefore we have

$$\lambda^2(V) = 46.$$

In case of $a_1 = 2, a_2 = 3, a_3 = 3$, we have

$$x_3^2\partial_1; x_3^3\partial_1; x_2x_3^2\partial_1; x_2^2x_3\partial_1; x_2^3\partial_1; x_2^3x_3\partial_1; x_2^4\partial_1; x_2^4x_3\partial_1; x_2^5\partial_1; x_1\partial_1+\frac{1}{2}x_2\partial_2+\frac{5}{6}x_3\partial_3; x_1x_3\partial_1; x_1x_3^2\partial_1; \\ x_1x_2\partial_1; x_1x_2x_3\partial_1; x_1x_2^2\partial_1; x_1^2\partial_1; x_1^2x_3\partial_1; x_1^3\partial_1; x_3^2\partial_2; x_3^3\partial_2; x_2x_3\partial_2; x_2x_3^2\partial_2; x_2^2\partial_2; x_2^2x_3\partial_2; x_3^3\partial_2; x_3^3x_3\partial_2; \\ x_2^4\partial_2; x_2^4x_3\partial_2; x_2^5\partial_2; x_1x_3\partial_2; x_1x_3^2\partial_2; x_1x_2\partial_2; x_1x_2x_3\partial_2; x_1x_2^2\partial_2; x_1^2\partial_2; x_1^2x_3\partial_2; x_1^3\partial_2; x_3^2\partial_3; x_3^3\partial_3; x_2x_3\partial_3; \\ x_2x_3^2\partial_3; x_2^2x_3\partial_3; x_2^3\partial_3; x_2^3x_3\partial_3; x_2^4\partial_3; x_2^4x_3\partial_3; x_2^5\partial_3; x_1x_3\partial_3; x_1x_3^2\partial_3; x_1x_2\partial_3; x_1x_2x_3\partial_3; x_1x_2^2\partial_3; x_1^2\partial_3; \\ x_1^2x_3\partial_3; x_1^3\partial_3.$$

Therefore we have

$$\lambda^2(V) = 55.$$

We have

$$h_2\left(\frac{1}{w_1}, \frac{1}{w_2}, \frac{1}{w_3}\right) = \begin{cases} \frac{3(a_1a_2-1)^2}{(a_1-1)(a_2-1)}a_3 + 5\left(\frac{a_1a_2-1}{a_1-1} + \frac{a_1a_2-1}{a_2-1} + a_3\right) - 4\left(\frac{(a_1a_2-1)^2}{(a_1-1)(a_2-1)} + \frac{a_1a_2-1}{a_2-1}a_3 + \frac{a_1a_2-1}{a_1-1}a_3\right) \\ + 34; \quad \frac{a_1a_2-1}{a_1-1} \geq 2, \frac{a_1a_2-1}{a_2-1} \geq 3, a_3 \geq 3. \end{cases}$$

Case(i): When $a_1 \geq 4, a_2 \geq 4, a_3 \geq 3$, then need to show that

$$3a_1a_2a_3 - 4a_1a_2 - 2a_2a_3 - 2a_1a_3 + 2a_1 + 2a_2 + 6a_3 + 36 \leq \frac{3(a_1a_2-1)^2}{(a_1-1)(a_2-1)}a_3 + 5\left(\frac{a_1a_2-1}{a_1-1} + \frac{a_1a_2-1}{a_2-1} + a_3\right) - 4\left(\frac{(a_1a_2-1)^2}{(a_1-1)(a_2-1)} + \frac{a_1a_2-1}{a_2-1}a_3 + \frac{a_1a_2-1}{a_1-1}a_3\right) + 34.$$

After simplification we get

$$[a_1(a_2 - 2)(a_1 - 3) + a_2(a_1 - 2)(a_2 - 3) + (a_1 - 3) + (a_2 - 3)](a_3 - 1) - 2 \geq 0.$$

Case(ii): When $a_1 = 3, a_2 \geq 3, a_3 \geq 3$, then need to show that

$$7a_2a_3 - 10a_2 + 40 \leq \frac{3(3a_2-1)^2}{2(a_2-1)}a_3 + 5\left(\frac{3a_2-1}{2} + \frac{3a_2-1}{a_2-1} + a_3\right) - 4\left(\frac{(3a_2-1)^2}{2(a_2-1)} + \frac{3a_2-1}{a_2-1}a_3 + \frac{3a_2-1}{2}a_3\right) + 34.$$

After simplification we get

$$[a_2(a_2 - 3) + (a_2 - 3)](a_3 - 1) - 4 \geq 0.$$

□

Proposition 3.9. *Let $(V, 0)$ be a weighted homogeneous feynomial isolated singularity which is defined by $f = x_1^{a_1} + x_2^{a_2} + x_3^2$ ($a_1 \geq 3, a_2 \geq 3$) with weight type $(\frac{1}{a_1}, \frac{1}{a_2}, \frac{1}{2}; 1)$.*

Then

$$\lambda^2(V) = \begin{cases} 2a_1a_2 - 3(a_1 + a_2) + 36; & a_1 \geq 5, a_2 \geq 5 \\ 3a_2 + 22; & a_1 = 3, a_2 \geq 4 \\ 30; & a_1 = 3, a_2 = 3 \\ 5a_2 + 23; & a_1 = 4, a_2 \geq 5 \\ 42; & a_1 = 4, a_2 = 4. \end{cases}$$

Furthermore, when $a_1 \geq 2, a_2 = 2$ then

$$\lambda^2(V) = \begin{cases} a_1 + 19; & a_1 \geq 3, \\ 19; & a_1 = 2. \end{cases}$$

Proof. It follows that the generalized moduli algebra

$$A^2(V) = \mathbb{C}\{x_1, x_2, x_3\}/(f, m^2J(f)),$$

has dimension $a_1a_2 - (a_1 + a_2) + 10$ and has a monomial basis of the form:

$$(1) \text{if } a_1 \geq 3, a_2 \geq 3, \{x_1^{i_1}x_2^{i_2}, 1 \leq i_1 \leq a_1 - 2; 0 \leq i_2 \leq a_2 - 2; x_1^{a_1-1}; x_1^{a_1-1}x_2; x_1x_2^{a_2-1}; x_2^{i_2}, 0 \leq i_2 \leq a_2; x_3; x_3^2; x_1x_3; x_2x_3\}, \quad (40)$$

$$(2) \text{if } a_1 \geq 2, a_2 = 2, \{x_1^{i_1}, 0 \leq i_1 \leq a_1 - 1; x_3; x_3^2; x_2; x_2x_3; x_2^2; x_1x_3; x_1x_2\}. \quad (41)$$

The following derivations form a basis of $\text{Der}A^2(V)$:

$$x_1^{i_1}x_2^{i_2}\partial_1, 1 \leq i_1 \leq a_1 - 2, 1 \leq i_2 \leq a_2 - 2; x_1^{a_1-1}x_2\partial_1; x_1x_2^{a_2-1}\partial_1; x_2^{i_2}\partial_1, a_2 - 3 \leq i_2 \leq a_2;$$

$$\begin{aligned}
& x_3\partial_1; x_3^2\partial_1; x_1x_3\partial_1; x_2x_3\partial_1; x_3\partial_2; x_3^2\partial_2; x_1x_3\partial_2; x_2x_3\partial_2; x_1x_2^{a_2-1}\partial_2; x_1\partial_1 + \frac{a_1}{a_2}x_2\partial_2; \\
& x_1^{i_1}\partial_1, 2 \leq i_1 \leq a_1 - 1; x_2^{i_2}\partial_2, 2 \leq i_2 \leq a_2; x_1^{i_1}\partial_2, a_1 - 3 \leq i_1 \leq a_1 - 1; \\
& x_1^{i_1}x_2^{i_2}\partial_2, 1 \leq i_1 \leq a_1 - 2, 1 \leq i_2 \leq a_2 - 2; x_1^{a_1-1}x_2\partial_2; x_3^2\partial_3; x_2x_3\partial_3; x_2^{a_2-1}\partial_3; x_2^{a_2}\partial_3; \\
& x_1x_3\partial_3; x_1^{a_1-1}\partial_3; x_1^{a_1-1}x_2\partial_3; x_1x_2^{a_2-1}\partial_3; x_1^{a_1-3}x_2^{a_2-2}\partial_3; x_1^{a_1-2}x_2^{a_2-3}\partial_3; x_1^{a_1-2}x_2^{a_2-2}\partial_3.
\end{aligned}$$

Therefore we have the following formula

$$\lambda^2(V) = 2a_1a_2 - 3(a_1 + a_2) + 36.$$

In case of $a_1 = 3, a_2 \geq 4$, we have following derivations form a basis of $\text{Der}A^2(V)$:

$$\begin{aligned}
& x_3\partial_1; x_3^2\partial_1; x_1x_3\partial_1; x_1x_2^{i_2}\partial_1, 1 \leq i_2 \leq a_2 - 1; x_1^2x_2\partial_1; x_1^2\partial_1; x_2^{i_2}\partial_1, a_2 - 2 \leq i_2 \leq a_2; \\
& x_2x_3\partial_1; x_3\partial_2; x_1x_2^{i_2}\partial_2, 1 \leq i_2 \leq a_2 - 1; x_1^2x_2\partial_2; x_1^2\partial_2; x_2^{i_2}\partial_2, 2 \leq i_2 \leq a_2; x_1\partial_1 + \frac{3}{a_2}x_2\partial_2; \\
& x_3^2\partial_2; x_1x_3\partial_2; x_2x_3\partial_2; x_3^2\partial_3; x_1^2\partial_3; x_1^2x_2\partial_3; x_2^{a_2-1}\partial_3; x_2^{a_2}\partial_3; x_2x_3\partial_3; x_1x_3\partial_3; x_1x_2^{a_2-2}\partial_3; x_1x_2^{a_2-1}\partial_3.
\end{aligned}$$

Therefore we have the following formula

$$\lambda^2(V) = 3a_2 + 22.$$

In case of $a_1 = 3, a_2 = 3$, we have following derivations form a basis of $\text{Der}A^2(V)$:

$$\begin{aligned}
& x_2^2\partial_1; x_3^2\partial_1; x_1x_2\partial_1; x_1x_2^2\partial_1; x_1^2\partial_1; x_1^2x_2\partial_1; x_2^2\partial_2; x_3^2\partial_2; x_1x_2\partial_2; x_1x_2^2\partial_2; x_1^2\partial_2; x_1^2x_2\partial_2; x_1\partial_1 + x_2\partial_2; \\
& x_3\partial_1; x_3^2\partial_1; x_1x_3\partial_1; x_2x_3\partial_1; x_3\partial_2; x_3^2\partial_2; x_1x_3\partial_2; x_2x_3\partial_2; x_3^2\partial_3; x_2x_3\partial_3; x_2^2\partial_3; x_3^2\partial_3; x_1x_3\partial_3; x_1x_2\partial_3; \\
& x_1x_2^2\partial_3; x_1^2\partial_3; x_1^2x_2\partial_3.
\end{aligned}$$

In case of $a_1 = 4, a_2 \geq 5$, we have following derivations form a basis of $\text{Der}A^2(V)$:

$$\begin{aligned}
& x_2^{i_2}\partial_1, a_2 - 3 \leq i_2 \leq a_2; x_1\partial_1 + \frac{4}{a_2}x_2\partial_2; x_1^{i_1}x_2\partial_1, 1 \leq i_1 \leq 2, 1 \leq i_2 \leq a_2 - 2; x_1x_2^{a_2-1}\partial_1; x_1^2\partial_1; x_1^3\partial_1; \\
& x_1^3x_2\partial_1; x_2^{i_2}\partial_2, 2 \leq i_2 \leq a_2; x_1^{i_1}x_2^{i_2}\partial_2, 1 \leq i_1 \leq 2, 1 \leq i_2 \leq a_2 - 2; x_1x_2^{a_2-1}\partial_2; x_1^2\partial_2; x_1^3\partial_2; x_1^3x_2\partial_2; \\
& x_3\partial_1; x_3^2\partial_1; x_1x_3\partial_1; x_2x_3\partial_1; x_3\partial_2; x_3^2\partial_2; x_1x_3\partial_2; x_2x_3\partial_2; x_3^2\partial_3; x_2x_3\partial_3; x_2^{a_2-1}\partial_3; x_2^{a_2}\partial_3; \\
& x_1x_3\partial_3; x_1^3x_2\partial_3; x_1x_2^{a_2-1}\partial_3; x_1x_2^{a_2-2}\partial_3; x_1^2x_2^{a_2-3}\partial_3; x_1^2x_2^{a_2-2}\partial_3.
\end{aligned}$$

Therefore we have the following formula

$$\lambda^2(V) = 5a_2 + 23.$$

In case of $a_1 = 4, a_2 = 4$, we have following derivations form a basis of $\text{Der}A^2(V)$:

$$\begin{aligned}
& x_2^2\partial_1; x_3^2\partial_1; x_2^4\partial_1; x_1\partial_1 + x_2\partial_2; x_1x_2\partial_1; x_1x_2^2\partial_1; x_1x_2^3\partial_1; x_1^2\partial_1; x_1^2x_2\partial_1; x_1^2x_2^2\partial_1; x_1^3\partial_1; x_1^3x_2\partial_1; x_2^2\partial_2; x_2^3\partial_2; \\
& x_2^4\partial_2; x_1x_2\partial_2; x_1x_2^{i_2}\partial_2; x_1x_2^3\partial_2; x_1^2\partial_2; x_1^2x_2\partial_2; x_1^2x_2^2\partial_2; x_1^3\partial_2; x_1^3x_2\partial_2; x_3\partial_1; x_3^2\partial_1; x_1x_3\partial_1; x_2x_3\partial_1; x_3\partial_2; \\
& x_3^2\partial_2; x_1x_3\partial_2; x_2x_3\partial_2; x_3^2\partial_3; x_2x_3\partial_3; x_2^3\partial_3; x_2^4\partial_3; x_1x_3\partial_3; x_1x_2^2\partial_3; x_1x_2^3\partial_3; x_1^2x_2\partial_3; x_1^2x_2^2\partial_3; x_1^3\partial_3; x_1^3x_2\partial_3.
\end{aligned}$$

Therefore we have the following formula

$$\lambda^2(V) = 42.$$

In case of $a_1 \geq 3, a_2 = 2$, we have following derivations form a basis of $\text{Der}A^2(V)$:

$$x_3^2\partial_1; x_2\partial_1; x_2x_3\partial_1; x_2^2\partial_1; x_1\partial_1 + \frac{7x_2}{2}\partial_2 + \frac{7x_3}{2}\partial_3; x_1x_3\partial_1; x_1x_2\partial_1; x_1^{i_1}\partial_1, 2 \leq i_1 \leq a_1 - 1; x_3\partial_2 - x_2\partial_3; \\ x_3\partial_1; x_3^2\partial_2; x_2x_3\partial_2; x_2^2\partial_2; x_1x_3\partial_2; x_1x_2\partial_2; x_1^{a_1-1}\partial_2; x_3^2\partial_3; x_2x_3\partial_3; x_2^2\partial_3; x_1x_3\partial_3; x_1x_2\partial_3; x_1^{a_1-1}\partial_3.$$

Therefore we have the following formula

$$\lambda^2(V) = a_1 + 19.$$

In case of $a_1 = 2, a_2 = 2$, we have following derivations form a basis of $\text{Der}A^2(V)$:

$$x_3\partial_1; x_3^2\partial_1; x_2\partial_1 - x_1\partial_2; x_2x_3\partial_1; x_2^2\partial_1; x_1\partial_1 + x_2\partial_2 + x_3\partial_3; x_1x_3\partial_1; x_1x_2\partial_1; x_3\partial_2 - x_2\partial_3; x_3^2\partial_3; x_2x_3\partial_3; \\ x_2^2\partial_2; x_1x_3\partial_2; x_1x_2\partial_2; x_3^2\partial_3; x_2x_3\partial_3; x_2^2\partial_3; x_1x_3\partial_3; x_1x_2\partial_3.$$

Therefore we have the following formula

$$\lambda^2(V) = 19.$$

□

Proposition 3.10. *Let $(V, 0)$ be a weighted homogeneous feynomial isolated singularity which is defined by $f = x_1^{a_1}x_2 + x_2^{a_2} + x_3^2$ ($a_1 \geq 2, a_2 \geq 3$) with weight type $(\frac{a_2-1}{a_1a_2}, \frac{1}{a_2}, \frac{1}{2}; 1)$. Then*

$$\lambda^2(V) = \begin{cases} 2a_1a_2 - 2a_1 - 3a_2 + 39; & a_1 \geq 5, a_2 \geq 5 \\ 5a_2 + 31; & a_1 = 4, a_2 \geq 5 \\ 50; & a_1 = 4, a_2 = 4 \\ 4a_1 + 24; & a_1 \geq 3, a_2 = 3 \\ a_2 + 28; & a_1 = 2, a_2 \geq 4 \\ 30; & a_1 = 2, a_2 = 3. \end{cases}$$

Furthermore, we need to show that when $a_1 \geq 5, a_2 \geq 5$, then

$$2a_1a_2 - 2a_1 - 3a_2 + 39 \leq \frac{2a_1a_2^2}{a_2-1} - 3\left(\frac{a_1a_2}{a_2-1} + a_2\right) + 36.$$

Proof. It follows that the generalized moduli algebra

$$A^2(V) = \mathbb{C}\{x_1, x_2, x_3\}/(f, m^2J(f)),$$

has dimension $a_1a_2 - a_2 + 10$ and has a monomial basis of the form:

$$\{x_1^{i_1}x_2^{i_2}, 1 \leq i_1 \leq a_1 - 2; 1 \leq i_2 \leq a_2 - 2; x_1^{i_1}x_2^{a_2-1}, 1 \leq i_1 \leq a_1 - 2; x_1^{a_1-1}x_2^{i_2}, 1 \leq i_2 \leq 2; \\ x_2^{i_2}, 0 \leq i_2 \leq a_2; x_1^{i_1}, 1 \leq i_1 \leq a_1 + 1; x_3; x_3^2; x_1x_3; x_2x_3\}.$$

The following derivations form a basis of $\text{Der}A^2(V)$:

$$\begin{aligned} & x_1^{i_1} x_2^{i_2} \partial_1, 1 \leq i_1 \leq a_1 - 2, 1 \leq i_2 \leq a_2 - 1; x_1^{i_1} \partial_1 + \frac{a_1}{a_2 - 1} x_1^{i_1 - 1} x_2 \partial_2, 1 \leq i_1 \leq a_1 - 3; \\ & x_1^{i_1} \partial_1, a_1 - 2 \leq i_1 \leq a_1 + 1; x_1^{a_1 - 1} x_2^{i_2} \partial_1, 1 \leq i_2 \leq 2; x_1^{i_1} x_2^{i_2} \partial_2, 1 \leq i_1 \leq a_1 - 2, 2 \leq i_2 \leq a_2 - 1; \\ & x_2^{i_2} \partial_1, a_2 - 3 \leq i_2 \leq a_2; x_2^{i_2} \partial_2, 2 \leq i_2 \leq a_2; x_1^{i_1} x_2 \partial_2, a_1 - 3 \leq i_1 \leq a_1 - 1; x_1^{i_1} \partial_2, a_1 - 2 \leq i_1 \leq a_1 + 1; \\ & x_3 \partial_1; x_3^2 \partial_1; x_1 x_3 \partial_1; x_2 x_3 \partial_1; x_3 \partial_2; x_3^2 \partial_2; x_1 x_3 \partial_2; x_2 x_3 \partial_2; x_3^2 \partial_3; x_2 x_3 \partial_3; x_1^{a_1 - 1} x_2 \partial_3; x_1^{a_1 - 1} x_2^2 \partial_3; x_1 x_3 \partial_3; \\ & x_1^{a_1 - 3} x_2^{a_2 - 1} \partial_3; x_1^{a_1 - 2} x_2^{a_2 - 1} \partial_3; x_1^{a_1 - 2} x_2^{a_2 - 2} \partial_3; x_2^{a_2} \partial_3; \left(\frac{x_1^{a_1}}{a_2} + x_2^{a_2 - 1}\right) \partial_3; \left(\frac{x_1^{a_1 + 1}}{a_2} + x_1 x_2^{a_2 - 1}\right) \partial_3. \end{aligned}$$

Therefore we have the following formula

$$\lambda^2(V) = 2a_1 a_2 - 2a_1 - 3a_2 + 39.$$

In case of $a_1 = 4, a_2 \geq 5$, we have following derivations which form a basis of $L^2(V)$:

$$\begin{aligned} & x_2^{i_2} \partial_1, a_2 - 3 \leq i_2 \leq a_2; x_1 \partial_1 + \frac{4}{a_2 - 1} x_2 \partial_1; x_1^{i_1} x_2^{i_2} \partial_1, 1 \leq i_1 \leq 2, 1 \leq i_2 \leq a_2 - 1; x_1^{i_1} \partial_1, 2 \leq i_1 \leq 5; \\ & x_1^3 x_2^{i_2} \partial_1, 1 \leq i_2 \leq 2; x_2^{i_2} \partial_2, 2 \leq i_2 \leq a_2; x_1^{i_1} x_2^{i_2} \partial_2, 1 \leq i_1 \leq 2, 1 \leq i_2 \leq a_2 - 1; x_1^{i_1} \partial_2, 2 \leq i_1 \leq 5; \\ & x_1^3 x_2^{i_2} \partial_2, 1 \leq i_2 \leq 2; x_3 \partial_1; x_3^2 \partial_1; x_1 x_3 \partial_1; x_2 x_3 \partial_1; x_3 \partial_2; x_3^2 \partial_2; x_1 x_3 \partial_2; x_2 x_3 \partial_2; x_3^2 \partial_3; x_2 x_3 \partial_3; \\ & x_1 x_3 \partial_3; x_1^5 \partial_3; x_1^3 x_2 \partial_3; x_1^3 x_2^2 \partial_3; x_1^2 x_2^{a_2 - 2} \partial_2; x_1^2 x_2^{a_2 - 1} \partial_3; x_1 x_2^{a_2 - 1} \partial_3; \left(\frac{x_1^4}{a_2} + x_2^{a_2 - 1}\right) \partial_3; x_2^{a_2} \partial_3. \end{aligned}$$

Therefore we get following formula

$$\lambda^2(V) = 5a_2 + 31.$$

In case of $a_1 = 4, a_2 = 4$, we have following derivations which form a basis of $L^2(V)$:

$$\begin{aligned} & x_2^2 \partial_1; x_2^3 \partial_1; x_2^4 \partial_1; x_1 \partial_1 + \frac{4}{3} x_2 \partial_1; x_1 x_2 \partial_1; x_1 x_2^2 \partial_1; x_1 x_2^3 \partial_1; x_1^2 \partial_1; x_1^2 x_2 \partial_1; x_1^2 x_2^2 \partial_1; x_1^2 x_2^3 \partial_1; x_1^3 \partial_1; x_1^3 x_2 \partial_1; \\ & x_1^3 x_2^2 \partial_1; x_1^4 \partial_1; x_1^5 \partial_1; x_2^2 \partial_2; x_2^3 \partial_2; x_2^4 \partial_2; x_1 \partial_1 + \frac{4}{3} x_2 \partial_2; x_1 x_2 \partial_2; x_1 x_2^2 \partial_2; x_1 x_2^3 \partial_2; x_1^2 \partial_2; x_1^2 x_2 \partial_2; x_1^2 x_2^2 \partial_2; \\ & x_1^2 x_2^3 \partial_2; x_1^3 \partial_2; x_1^3 x_2 \partial_2; x_1^3 x_2^2 \partial_2; x_1^4 \partial_2; x_1^5 \partial_2; x_3 \partial_1; x_3^2 \partial_1; x_1 x_3 \partial_1; x_2 x_3 \partial_1; x_3 \partial_2; x_3^2 \partial_2; x_1 x_3 \partial_2; x_2 x_3 \partial_2; \\ & x_3^2 \partial_3; x_2 x_3 \partial_3; \left(\frac{x_1^4}{4} + x_2^3\right) \partial_3; x_2^4 \partial_3; x_1 x_3 \partial_3; x_1 x_2^3 \partial_3; x_1^2 x_2^2 \partial_3; x_1^2 x_2^3 \partial_3; x_1^3 x_2 \partial_3; x_1^3 x_2^2 \partial_3; x_1^5 \partial_3. \end{aligned}$$

Therefore we get following formula

$$\lambda^2(V) = 50.$$

In case of $a_1 \geq 3, a_2 = 3$, we have following derivations which form a basis of $L^2(V)$:

$$\begin{aligned} & x_2^2 \partial_1; x_2^3 \partial_1; x_1^{i_1} \partial_1 + \frac{a_1}{2} x_1^{i_1 - 1} x_2 \partial_2, 1 \leq i_1 \leq a_1 - 2; x_1^{i_1} x_2^{i_2} \partial_1, 1 \leq i_1 \leq a_1 - 1, 1 \leq i_2 \leq 2; \\ & x_1^{i_1} \partial_1, a_1 - 1 \leq i_1 \leq a_1 + 1; x_2^2 \partial_2; x_2^3 \partial_2; x_1^{i_1} \partial_2, a_1 - 1 \leq i_1 \leq a_1 + 1; x_1^{i_1} x_2^2 \partial_2, 1 \leq i_1 \leq a_1 - 1; \\ & x_1^{i_1} x_2 \partial_2, a_1 - 2 \leq i_1 \leq a_1 - 1; x_3 \partial_1; x_3^2 \partial_1; x_1 x_3 \partial_1; x_2 x_3 \partial_1; x_3 \partial_2; x_3^2 \partial_2; x_1 x_3 \partial_2; x_2 x_3 \partial_2; \\ & x_3^2 \partial_3; x_2 x_3 \partial_3; x_1 x_3 \partial_3; x_1^{a_1 - 1} x_2 \partial_3; x_1^{a_1 - 1} x_2^2 \partial_3; x_1^{a_1 - 2} x_2^2 \partial_3; \left(\frac{x_1^{a_1}}{3} + x_2^2\right) \partial_3; x_2^3 \partial_3; \left(\frac{x_1^{a_1 + 1}}{3} + x_1 x_2^2\right) \partial_3. \end{aligned}$$

Therefore we get following formula

$$\lambda^2(V) = 4a_1 + 24.$$

In case of $a_1 = 2, a_2 \geq 3$, it is easy to check that we get following basis:

$$\begin{aligned} & x_2^{i_2} \partial_1, 2 \leq i_2 \leq a_2; x_1 \partial_1 + \frac{2}{a_2 - 1} x_2 \partial_2; x_2^{i_2} \partial_1, a_2 - 2 \leq i_2 \leq a_2; x_1 x_2 \partial_1; x_1 x_2^2 \partial_1; x_1^2 \partial_1; \\ & x_1^3 \partial_1; x_1 x_2 \partial_2; x_1 x_2^2 \partial_2; x_1^2 \partial_2; x_1^3 \partial_2; x_3 \partial_1; x_3^2 \partial_1; x_1 x_3 \partial_1; x_2 x_3 \partial_1; x_3 \partial_2; x_3^2 \partial_2; x_1 x_3 \partial_2; x_2 x_3 \partial_2; \\ & x_3^2 \partial_3; x_2 x_3 \partial_3; x_1 x_3 \partial_3; x_1^2 \partial_3; x_1^3 \partial_3; x_1 x_2 \partial_3; x_1 x_2^2 \partial_3; x_2^{a_2 - 1} \partial_3; x_2^{a_2} \partial_3. \end{aligned}$$

Therefore

$$\lambda^2(V) = a_2 + 28.$$

In case of $a_1 = 2, a_2 = 3$, we have following derivations which form a basis of $L^2(V)$:

$$\begin{aligned} & x_3 \partial_1; x_3^2 \partial_1; x_2 x_3 \partial_1; x_2^2 \partial_1; x_2^3 \partial_1; x_1 \partial_1 + x_2 \partial_2 + \frac{3x_3}{2} \partial_3; x_1 x_3 \partial_1; x_1 x_2 \partial_1; x_1 x_2^2 \partial_1; x_1^2 \partial_1; x_1^3 \partial_1; x_3 \partial_2; \\ & x_3^2 \partial_2; x_2 x_3 \partial_2; x_2^2 \partial_2; x_2^3 \partial_2; x_1 x_3 \partial_2; x_1 x_2 \partial_2; x_1 x_2^2 \partial_2; x_1^2 \partial_2; x_1^3 \partial_2; x_3^2 \partial_3; x_2 x_3 \partial_3; x_2^2 \partial_3; x_2^3 \partial_3; x_1 x_3 \partial_3; \\ & x_1 x_2 \partial_3; x_1 x_2^2 \partial_3; x_1^2 \partial_3; x_1^3 \partial_3. \end{aligned}$$

Therefore we get following formula

$$\lambda^2(V) = 30.$$

We have

$$\begin{aligned} h_2(a_1, a_2, 2) &= \begin{cases} 2a_1 a_2 - 3(a_1 + a_2) + 36; & a_1 \geq 5, a_2 \geq 5 \\ 3a_2 + 22; & a_1 = 3, a_2 \geq 4 \\ 30; & a_1 = 3, a_2 = 3 \\ 5a_2 + 23; & a_1 = 4, a_2 \geq 5 \\ 42; & a_1 = 4, a_2 = 4. \end{cases} \\ h_2\left(\frac{1}{w_1}, \frac{1}{w_2}, 2\right) &= \begin{cases} \frac{2a_1 a_2^2}{a_2 - 1} - 3\left(\frac{a_1 a_2}{a_2 - 1} + a_2\right) + 36; & \frac{a_1 a_2}{a_2 - 1} \geq 5, a_2 \geq 5 \\ 30; & a_1 = 2, a_2 = 3 \\ 42; & a_1 = 3, a_2 = 4. \end{cases} \end{aligned}$$

It is easy to see that when $a_1 \geq 5, a_2 \geq 5$, then

$$2a_1 a_2 - 2a_1 - 3a_2 + 39 \leq \frac{2a_1 a_2^2}{a_2 - 1} - 3\left(\frac{a_1 a_2}{a_2 - 1} + a_2\right) + 36.$$

After solving above inequality we get $(a_2 - 2)(a_1 - 1) - 3 \geq 0$. It is noted that $(a_2 - 2) \geq 3$ and $(a_1 - 1) \geq 4$. \square

Proposition 3.11. *Let $(V, 0)$ be a weighted homogeneous fewnomial isolated singularity which is defined by $f = x_1^{a_1} x_2 + x_2^{a_2} x_1 + x_3^2$ ($a_1 \geq 2, a_2 \geq 2$) with weight type $(\frac{a_2 - 1}{a_1 a_2 - 1}, \frac{a_1 - 1}{a_1 a_2 - 1}, \frac{1}{2}; 1)$. Then*

$$\lambda^2(V) = \begin{cases} 2a_1a_2 - 2(a_1 + a_2) + 40; & a_1 \geq 4, a_2 \geq 4 \\ 4a_2 + 32; & a_1 = 3, a_2 \geq 4 \\ 2a_2 + 27; & a_1 = 2, a_2 \geq 3 \\ 42; & a_1 = 3, a_2 = 3 \\ 30; & a_1 = 2, a_2 = 2. \end{cases}$$

Furthermore, we need to show that when $a_1 \geq 5, a_2 \geq 5$, then

$$2a_1a_2 - 2(a_1 + a_2) + 40 \leq \frac{2(a_1a_2-1)^2}{(a_1-1)(a_2-1)} - 3\left(\frac{a_1a_2-1}{a_2-1} + \frac{a_1a_2-1}{a_1-1}\right) + 36.$$

Proof. It follows that the generalized moduli algebra

$$A^2(V) = \mathbb{C}\{x_1, x_2, x_3\}/(f, m^2J(f)),$$

has dimension $a_1a_2 + 9$ and has a monomial basis of the form:

$$\{x_1^{i_1}x_2^{i_2}, 0 \leq i_1 \leq a_1 - 2; 0 \leq i_2 \leq a_2 - 1; x_2^{i_2}, a_2 \leq i_2 \leq 2a_2 - 2; x_1^{i_1}, a_1 - 1 \leq i_1 \leq a_1 + 1; x_1x_2^{a_2}; x_1^{a_1-1}x_2; x_1^{a_1-1}x_2^2; x_3; x_3^2; x_1x_3; x_2x_3\}.$$

The following derivations form a basis of $\text{Der}A^2(V)$:

$$\begin{aligned} & x_1^{i_1}x_2^{i_2}\partial_1, 2 \leq i_1 \leq a_1 - 2, 1 \leq i_2 \leq a_2 - 1; x_2^{i_2}\partial_1, a_2 - 2 \leq i_2 \leq 2a_2 - 2; \\ & x_1^{i_1}\partial_1 + \frac{a_2 - 1}{a_1}x_1^{i_1-1}x_2\partial_2, 1 \leq i_1 \leq a_1 - 3; x_1^{a_1-1}x_2\partial_1; x_1^{a_1-1}x_2^2\partial_1; \\ & x_1x_2^{i_2}\partial_1 + \frac{a_2 - 1}{a_1}x_2^{i_2+1}\partial_2, 1 \leq i_2 \leq a_2 - 4; x_1x_2^{i_2}\partial_1, a_2 - 3 \leq i_2 \leq a_2; x_1^{i_1}\partial_1, a_1 - 2 \leq i_1 \leq a_1 + 1; \\ & x_2^{i_2}\partial_2, a_2 - 2 \leq i_2 \leq 2a_2 - 2; x_1^{i_1}x_2^{i_2}\partial_2, 1 \leq i_1 \leq a_1 - 2, 2 \leq i_2 \leq a_2 - 1; \\ & x_1^{i_1}\partial_2, a_1 - 2 \leq i_1 \leq a_1 + 1; x_1^{i_1}x_2\partial_2, a_1 - 3 \leq i_1 \leq a_1 - 1; x_1x_2^{a_2}\partial_2; x_1^{a_1-1}x_2^2\partial_2; x_3\partial_1; \\ & x_3^2\partial_1; x_1x_3\partial_1; x_2x_3\partial_1; x_3\partial_2; x_3^2\partial_2; x_1x_3\partial_2; x_2x_3\partial_2; x_3^2\partial_3; x_2x_3\partial_3; x_1x_3\partial_3; (a_1x_1^{a_1-1}x_2 + x_2^{a_2})\partial_3; \\ & (a_1x_1^{a_1-1}x_2 + x_2^{a_2+1})\partial_3; x_2^{2a_2-3}\partial_3; x_2^{2a_2-2}\partial_3; \frac{x_1^{a_1}}{a_2} + x_1x_2^{a_2-1}\partial_3; x_1x_2^{a_2}\partial_3; \left(\frac{x_1^{a_1+1}}{a_2} + x_1^2x_2^{a_2-1}\right)\partial_3; \\ & x_1^{a_1-2}x_2^{a_2-1}\partial_3. \end{aligned}$$

Therefore we have the following formula

$$\lambda^2(V) = 2a_1a_2 - 2(a_1 + a_2) + 40.$$

In case of $a_1 = 3, a_2 \geq 4$, we have following derivations which form a basis of $L^2(V)$:

$$\begin{aligned} & x_2^{i_2}\partial_1, a_2 - 2 \leq i_2 \leq 2a_2 - 2; x_1\partial_1 + \frac{2}{a_2 - 1}x_2\partial_2; x_1x_2^{i_2}\partial_1 + \frac{2}{a_2 - 1}x_2^{i_2+1}\partial_2, 1 \leq i_2 \leq a_2 - 4; \\ & x_1x_2^{i_2}\partial_1, a_2 - 3 \leq i_2 \leq a_2; x_1^2x_2\partial_1; x_1^2x_2^2\partial_1; x_2^{i_2}\partial_2, a_2 - 2 \leq i_2 \leq 2a_2 - 2; x_1^{i_1}\partial_2, 2 \leq i_1 \leq 4; \\ & x_1^{i_1}\partial_1, 2 \leq i_1 \leq 4; x_1x_2^{i_2}\partial_2, 1 \leq i_2 \leq a_2; x_1^2x_2\partial_2; x_1^2x_2^2\partial_2; x_3\partial_1; x_3^2\partial_1; x_1x_3\partial_1; x_2x_3\partial_1; x_3\partial_2; \\ & x_3^2\partial_2; x_1x_3\partial_2; x_2x_3\partial_2; x_3^2\partial_3; x_1^3\partial_3; x_1^4\partial_3; x_1x_2^{a_2-1}\partial_3; x_1x_2^{a_2}\partial_3; x_1x_3\partial_3; x_2^{2a_2-3}\partial_3; \\ & x_2^{2a_2-2}\partial_3; x_2x_3\partial_3; (3x_1^2x_3 + x_2^{a_2})\partial_3; (3x_1^2x_2^2 + x_2^{a_2+1})\partial_3. \end{aligned}$$

Therefore we get following formula

$$\lambda^2(V) = 4a_2 + 32.$$

In case of $a_1 = 2, a_2 \geq 3$, we have following derivations which form a basis of $L^2(V)$:

$$\begin{aligned} & x_2^{(a_2-2)-i_2} \partial_1 - \frac{2}{a_2-1} x_2^{i_2} \partial_2, 2 \leq i_2 \leq a_2 - 2; x_2^{i_2} \partial_1, 2a_2 - 3 \leq i_2 \leq 2a_2 - 1; \\ & x_1^2 \partial_1; x_1^3 \partial_1; x_2^{i_2} \partial_2, a_2 - 1 \leq i_2 \leq 2a_2 - 1; x_1^2 \partial_2; x_1^3 \partial_2; x_1 x_2 \partial_2; x_1 x_2^2 \partial_2; \\ & x_1 x_2^{i_2} \partial_1 + \frac{x_2^{i_2+1}}{a_2-1} \partial_2, 0 \leq i_2 \leq 2; x_3 \partial_1; x_3^2 \partial_1; x_1 x_3 \partial_1; x_2 x_3 \partial_1; x_3 \partial_2; x_3^2 \partial_2; x_1 x_3 \partial_2; x_2 x_3 \partial_2; \\ & x_3^2 \partial_3; x_2 x_3 \partial_3; x_1 x_3 \partial_3; (2x_1 x_2 + x_2^{a_2}) \partial_3; x_2^{2a_2-2} \partial_3; x_2^{2a_2-1} \partial_3; x_1^2 \partial_3; x_1^3 \partial_3. \end{aligned}$$

Therefore we get following formula

$$\lambda^2(V) = 2a_2 + 27.$$

In case of $a_1 = 3, a_2 = 3$, we have following derivations which form a basis of $L^2(V)$:

$$\begin{aligned} & x_2^2 \partial_1; x_2^3 \partial_1; x_2^4 \partial_1; x_1 \partial_1 + x_2 \partial_2; x_1 x_2 \partial_1; x_1 x_2^2 \partial_1; x_1 x_2^3 \partial_1; x_1^2 \partial_1; x_1^2 x_2 \partial_1; x_1^2 x_2^2 \partial_1; x_1^3 \partial_1; x_1^4 \partial_1; x_2^2 \partial_2; x_2^3 \partial_2; \\ & x_2^4 \partial_2; x_1 x_2 \partial_2; x_1 x_2^2 \partial_2; x_1 x_2^3 \partial_2; x_1^2 \partial_2; x_1^2 x_2 \partial_2; x_1^2 x_2^2 \partial_2; x_1^3 \partial_2; x_1^4 \partial_2; x_3 \partial_1; x_3^2 \partial_1; x_1 x_3 \partial_1; x_2 x_3 \partial_1; x_3 \partial_2; \\ & x_3^2 \partial_2; x_1 x_3 \partial_2; x_2 x_3 \partial_2; x_3^2 \partial_3; x_2 x_3 \partial_3; x_2^2 \partial_3; x_2^3 \partial_3; x_1 x_3 \partial_3; x_1 x_2^2 \partial_3; x_1 x_2^3 \partial_3; x_1^2 x_2 \partial_3; x_1^2 x_2^2 \partial_3; x_1^3 \partial_3; x_1^4 \partial_3. \end{aligned}$$

Therefore we get following formula

$$\lambda^2(V) = 42.$$

In case of $a_1 = 2, a_2 = 2$, we have following derivations which form a basis of $L^2(V)$:

$$\begin{aligned} & x_2^2 \partial_1; x_2^3 \partial_1; x_1 \partial_1 + x_2 \partial_2; x_1 x_2 \partial_1; x_1 x_2^2 \partial_1; x_1^2 \partial_1; x_1^3 \partial_1; x_2^2 \partial_2; x_2^3 \partial_2; x_1 x_2 \partial_2; x_1 x_2^2 \partial_2; x_1^2 \partial_2; x_1^3 \partial_2; \\ & x_3 \partial_1; x_3^2 \partial_1; x_1 x_3 \partial_1; x_2 x_3 \partial_1; x_3 \partial_2; x_3^2 \partial_2; x_1 x_3 \partial_2; x_2 x_3 \partial_2; x_2^2 \partial_3; x_2 x_3 \partial_3; x_2^2 \partial_3; x_2^3 \partial_3; x_1 x_3 \partial_3; x_1 x_2 \partial_3; \\ & x_1 x_2^2 \partial_3; x_1^2 \partial_3; x_1^3 \partial_3. \end{aligned}$$

Therefore we get following formula

$$\lambda^2(V) = 30.$$

We have

$$h_2\left(\frac{1}{w_1}, \frac{1}{w_2}, 2\right) = \begin{cases} \frac{2(a_1 a_2 - 1)^2}{(a_1 - 1)(a_2 - 1)} - 3\left(\frac{a_1 a_2 - 1}{a_2 - 1} + \frac{a_1 a_2 - 1}{a_1 - 1}\right) + 36; & \frac{a_1 a_2 - 1}{a_2 - 1} \geq 5, \frac{a_1 a_2 - 1}{a_1 - 1} \geq 5 \\ 30; & a_1 = 2, a_2 = 2 \\ 42; & a_1 = 3, a_2 = 3. \end{cases}$$

It is easy to see that when $a_1 \geq 5, a_2 \geq 5$, then

$$2a_1 a_2 - 2(a_1 + a_2) + 40 \leq \frac{2(a_1 a_2 - 1)^2}{(a_1 - 1)(a_2 - 1)} - 3\left(\frac{a_1 a_2 - 1}{a_2 - 1} + \frac{a_1 a_2 - 1}{a_1 - 1}\right) + 36.$$

After solving above inequality it is easy to see that

$$a_1(a_2 - 1)(a_1 - 3) + a_2(a_1 - 1)(a_2 - 3) + 2(a_1 - 3) + 2a_1 + 4a_2 - 1 \geq 0.$$

□

4. PROOF OF THEOREM A

Proof. Let $f \in \mathbb{C}\{x_1, x_2, x_3\}$ be a weighted homogeneous fewnomial isolated singularity. Then f can be classified into the following five types:

Type 1. $x_1^{a_1} + x_2^{a_2} + x_3^{a_3}$,

Type 2. $x_1^{a_1}x_2 + x_2^{a_2}x_3 + x_3^{a_3}$,

Type 3. $x_1^{a_1}x_2 + x_2^{a_2}x_3 + x_3^{a_3}x_1$,

Type 4. $x_1^{a_1} + x_2^{a_2} + x_3^{a_3}x_2$,

Type 5. $x_1^{a_1}x_2 + x_2^{a_2}x_1 + x_3^{a_3}$.

It is easy to see from propositions 2.3, 2.4, 2.5, 2.6, 2.7, 3.4, 3.5, 3.6, 3.7 and 3.8, the conjecture $\lambda^{(k+1)}(V) > \lambda^k(V)$, $k = 1$ holds. Hence Theorem A is proved. □

5. PROOF OF THEOREM B

Proof. Let $f \in \mathbb{C}\{x_1, x_2\}$ be a weighted homogeneous fewnomial isolated singularity. Then f can be classified into the following three types:

Type A. $x_1^{a_1} + x_2^{a_2}$,

Type B. $x_1^{a_1}x_2 + x_2^{a_2}$,

Type C. $x_1^{a_1}x_2 + x_2^{a_2}x_1$.

It is easy to see from propositions 3.1, 3.2 and 3.3, the conjecture $\lambda^2(V) \leq h_2(\frac{1}{w_1}, \frac{1}{w_2})$, holds. Hence Theorem B is proved. □

6. PROOF OF THEOREM C

Proof. Let $f \in \mathbb{C}\{x_1, x_2, x_3\}$ be a weighted homogeneous fewnomial isolated singularity. Then f can be classified into the following five types:

Type 1. $x_1^{a_1} + x_2^{a_2} + x_3^{a_3}$,

Type 2. $x_1^{a_1}x_2 + x_2^{a_2}x_3 + x_3^{a_3}$,

Type 3. $x_1^{a_1}x_2 + x_2^{a_2}x_3 + x_3^{a_3}x_1$,

Type 4. $x_1^{a_1} + x_2^{a_2} + x_3^{a_3}x_2$,

Type 5. $x_1^{a_1}x_2 + x_2^{a_2}x_1 + x_3^{a_3}$.

It is easy to see from propositions 3.4, 3.5, 3.6, 3.7 and 3.8, the conjecture $\lambda^2(V) \leq h_2(\frac{1}{w_1}, \frac{1}{w_2}, \frac{1}{w_3})$, holds. Hence Theorem C is proved. □

7. PROOF OF THEOREM D

Proof. It is easy to see from propositions 3.9, 3.10 and 3.11, the following three cases:

- i). $x_1^{a_1} + x_2^{a_2} + x_3^2; a_1, a_2 \geq 3$,
 ii). $x_1^{a_1}x_2 + x_2^{a_2} + x_3^2; a_1 \geq 2, a_2 \geq 3$,
 iii). $x_1^{a_1}x_2 + x_2^{a_2}x_1 + x_3^2; a_1, a_2 \geq 2$,
 satisfy the conjecture $\lambda^2(V) \leq h_2(\frac{1}{w_1}, \frac{1}{w_2}, 2)$. Hence Theorem D is proved. \square

REFERENCES

- [1] M. Benson and S. S.-T. Yau, *Lie algebra and their representations arising from isolated singularities: Computer method in calculating the Lie algebras and their cohomology*, Advance studies in Pure Mathematics 8, Complex Analytic Singularities, (1986), 3-58.
- [2] M. Benson and S. S.-T. Yau, *Equivalence between isolated hypersurface singularities*, Math. Ann. 287 (1990), 107-134.
- [3] R. Block, *Determination of the differentially simple rings with a minimal ideal*, Ann. Math. 90 (1969), 433-459.
- [4] B. Chen, H. Chen, S. S.-T. Yau, and H. Zuo, *The non-existence of negative weight derivations on positive dimensional isolated singularities: generalized Wahl conjecture*, J. Differential Geom. 115(2) (2020), 195-224.
- [5] B. Chen, N. Hussain, S. S.-T. Yau, and H. Zuo, *Variation of complex structures and variation of Lie algebras II: New Lie algebras arising from singularities*, J. Differential Geom. 115(3) (2020), 437-473.
- [6] H. Chen, Y.-J. Xu, and S. S.-T. Yau, *Nonexistence of negative weight derivation of moduli algebras of weighted homogeneous singularities*, J. Algebra. 172 (1995), 243-254.
- [7] H. Chen, S. S.-T. Yau, and H. Zuo, *Non-existence of negative weight derivations on positively graded Artinian algebras*, Trans. Amer. Math. Soc. 372(4) (2019), 2493-2535.
- [8] A. Elashvili and G. Khimshiashvili, *Lie algebras of simple hypersurface singularities*, J. Lie Theory 16 (2006), 621-649.
- [9] W. Ebeling and A. Takahashi, *Strange duality of weighted homogeneous polynomial*, Compositio Math. 147 (2011), 1413-1433.
- [10] G.-M. Greuel, C. Lossen, and E. Shustin, *Introduction to singularities and deformations*, Springer Monographs in Mathematics, Springer, Berlin, 2007.
- [11] N. Hussain, *Survey on derivation Lie algebras of isolated singularities*, Methods Appl. Anal. 25(4) (2018), 307-322.
- [12] C. Hu, S. S.-T. Yau, and H. Zuo, *Torelli Theorem for k -th Yau Algebras over Simple Elliptic Singularities \tilde{E}_6* , 49pp., preprint, submitted.
- [13] N. Hussain, S. S.-T. Yau, and H. Zuo, *On the derivation Lie algebras of fewnomial singularities*, Bull. Aust. Math. Soc. 98(1) (2018), 77-88.
- [14] N. Hussain, S. S.-T. Yau, and H. Zuo, *On the new k -th Yau algebras of isolated hypersurface singularities*, Math. Z. 294(1-2) (2020), 331-358.
- [15] N. Hussain, S. S.-T. Yau, and H. Zuo, *k -th Yau number of isolated hypersurface singularities and an inequality conjecture*, 25pp. in ms, to appear, J. Aust. Math. Soc. (online publication April 2019), doi:10.1017/S1446788719000132.
- [16] N. Hussain, S. S.-T. Yau, and H. Zuo, *New k -th Yau algebras of isolated hypersurface singularities and weak Torelli-type theorem*, 32pp., preprint, submitted.
- [17] J. Mather and S. S.-T. Yau, *Classification of isolated hypersurface singularities by their moduli algebras*, Invent. Math. 69 (1982), 243-251.
- [18] L. Pursell and M. Shanks, *The Lie algebra of a smooth manifold*, Trans. Amer. Math. Soc. 5 (1954), 468-472.
- [19] K. Saito, *Quasihomogene isolierte Singularitäten von Hyperflächen*, Invent. Math. 14 (1971), 123-142.
- [20] C. Seeley, *7-dimensional Nilpotent Lie algebras*, Trans. Amer. Math. Soc. 335, (1993), 479-496.

- [21] C. Seeley and S. S.-T. Yau, *Variation of complex structure and variation of Lie algebras*, Invent. Math. 99 (1990), 545-565.
- [22] Y.-J. Xu and S. S.-T. Yau, *Micro-local characterization quasi-homogeneous singularities*, Amer. J. Math. 118(2) (1996), 389-399.
- [23] S. S.-T. Yau, *Continuous family of finite-dimensional representations of a solvable Lie algebra arising from singularities*, Proc. Natl. Acad. Sci. USA 80 (1983), 7694-7696.
- [24] S. S.-T. Yau, *Solvable Lie algebras and generalized Cartan matrices arising from isolated singularities*, Math. Z. 191 (1986), 489-506.
- [25] S. S.-T. Yau, *Solvability of Lie algebras arising from isolated singularities and nonisolatedness of singularities defined by $sl(2, \mathbb{C})$ invariant polynomials*, Amer. J. Math. 113 (1991), 773-778.
- [26] Y. Yu, *On Jacobian ideals invariant by reducible $sl(2; \mathbb{C})$ action*, Trans. Amer. Math. Soc. 348 (1996), 2759-2791.
- [27] S. S.-T. Yau and H. Zuo, *Derivations of the moduli algebras of weighted homogeneous hypersurface singularities*, J. Algebra 457 (2016), 18-25.
- [28] S. S.-T. Yau and H. Zuo, *A Sharp upper estimate conjecture for the Yau number of weighted homogeneous isolated hypersurface singularity*, Pure Appl. Math. Q. 12(1) (2016), 165-181.

SCHOOL OF DATA SCIENCES, HUASHANG COLLEGE GUANGDONG UNIVERSITY OF FINANCE AND ECONOMICS, GUANGZHOU GUANGDONG, 511300 P. R. CHINA., DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING, 100084, P. R. CHINA.

E-mail address: naveed1573@gmail.com

DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING, 100084, P. R. CHINA.

E-mail address: yau@uic.edu

DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING, 100084, P. R. CHINA.

E-mail address: hqzuo@mail.tsinghua.edu.cn